


Tuning a Three-Component Reaction For Trapping Kinase Substrate Complexes

Alexander V. Statsuk, Dustin J. Maly, Markus A. Seeliger, Miles A. Fabian, William H. Biggs III., David J. Lockhart, Patrick P. Zarrinkar, John Kuriyan, and Kevan M. Shokat

J. Am. Chem. Soc., **2008**, 130 (51), 17568-17574 • DOI: 10.1021/ja807066f • Publication Date (Web): 20 November 2008

Downloaded from <http://pubs.acs.org> on February 8, 2009



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)



ACS Publications
 High quality. High impact.

Tuning a Three-Component Reaction For Trapping Kinase Substrate Complexes

Alexander V. Statsuk,[†] Dustin J. Maly,^{†,||} Markus A. Seeliger,[‡] Miles A. Fabian,[§] William H. Biggs III,[§] David J. Lockhart,[§] Patrick P. Zarrinkar,[§] John Kuriyan,[‡] and Kevan M. Shokat^{*,†}

Howard Hughes Medical Institute, Department of Cellular and Molecular Pharmacology, University of California, San Francisco, California 94107, Department of Molecular and Cell Biology, Department of Chemistry, and Howard Hughes Medical Institute, University of California, Berkeley, California 94720, and Ambit Biosciences, 4215 Sorrento Valley Boulevard, San Diego, California 92121

Received September 5, 2008; E-mail: shokat@cmp.ucsf.edu

Abstract: The upstream protein kinases responsible for thousands of phosphorylation events in the phosphoproteome remain to be discovered. We developed a three-component chemical reaction which converts the transient noncovalent substrate–kinase complex into a covalently cross-linked product by utilizing a dialdehyde-based cross-linker, **1**. Unfortunately, the reaction of **1** with a lysine in the kinase active site and an engineered cysteine on the substrate to form an isoindole cross-linked product could not be performed in the presence of competing cellular proteins due to nonspecific side reactions. In order to more selectively target the cross-linker to protein kinases in cell lysates, we replaced the weak, kinase-binding adenosine moiety of **1** with a potent protein kinase inhibitor scaffold. In addition, we replaced the *o*-phthalaldehyde moiety in **1** with a less-reactive thiophene-2,3-dicarboxaldehyde moiety. The combination of these two structural modifications provides for cross-linking of a cysteine-containing substrate to its corresponding kinase in the presence of competing cellular proteins.

1. Introduction

Protein phosphorylation represents one of the most dominant and evolutionarily conserved post-translational modifications for information transfer in cells and organisms.¹ The human genome encodes 518 eukaryotic protein kinases, and thousands of phosphorylation sites are present in the mammalian proteome.² In order to develop a global view of eukaryotic phosphorylation networks, it will be necessary to map the connectivity between kinases and phosphoproteins. To this end, we and others have developed methods that enable identification of kinase sub-

strates.³ The inverse problem, that of identifying upstream kinases responsible for specific phosphorylation events, has proven much more challenging.⁴ Examples of phosphorylation sites with unknown upstream kinases include Ser-170 of the protein BAD which enhances its antiapoptotic activity,⁵ Ser-497 of the natriuretic peptide receptor, which is responsible for desensitization of the receptor's antihypertensive activity,⁶ and Ser-325 of the tumor suppressor LKB1, which is phosphorylated in response to the biguanide antidiabetic agent metformin.⁷ These examples, as well as the thousands of new "orphan" phosphorylation sites that have been identified in the mammalian phosphoproteome,⁸ highlight the need to develop methods for reverse mapping of phosphorylation networks as a complement to methods for kinase substrate identification.

Several years ago we reported a three-component chemical reaction designed to facilitate identification of the kinase responsible for a given phosphorylation event by covalently cross-linking a protein substrate of interest to its upstream kinase

[†] University of California, San Francisco and Howard Hughes Medical Institute, University of California, Berkeley.

[‡] Department of Molecular and Cell Biology, Department of Chemistry, and Howard Hughes Medical Institute, University of California, Berkeley.

[§] Ambit Biosciences.

^{||} Current address: Department of Chemistry, University of Washington.

(1) (a) Manning, G.; Whyte, D. B.; Martinez, R.; Hunter, T.; Sudarsanam, S. *Science* **2002**, *298*, 1912–1934. (b) Johnson, L. N.; Lewis, R. J. *Chem. Rev.* **2001**, *101*, 2209–2242.

(2) Villén, J.; Beausoleil, S. A.; Gerber, S. A.; Gygi, S. P. *PNAS* **2007**, *104*, 1488–1493.

(3) (a) Manning, B. D.; Cantley, L. C. *Sci. STKE* **2002**, *2002*, pe49. (b) Ptacek, J.; Snyder, M. *Trends Genet.* **2006**, *22*, 545–554. (c) Ptacek, J. *Nature* **2005**, *438*, 679–684. (d) Songyang, Z.; Blechner, S.; Hoagland, N.; Hoekstra, M. F.; Piwnicka-Worms, H.; Cantley, L. C. *Curr. Biol.* **1994**, *4*, 973–982. (e) Obenaus, J. C.; Cantley, L. C.; Yaffe, M. B. *Nucleic Acids Res.* **2003**, *31*, 3635–3641. (f) Berwick, D. C.; Tavaré, J. M. *Trends Biochem. Sci.*, *29*, 227–232. (g) Allen, J. J.; Li, M.; Brinkworth, C. S.; Paulson, J. L.; Wang, D.; Hübner, A.; Chou, W. H.; Davis, R. J.; Burlingame, A. L.; Messing, R. O.; Katayama, C. D.; Hedrick, S. M.; Shokat, K. M. *Nat. Methods* **2007**, *4*, 511–516. (h) Blethrow, J. D.; Glavy, J. S.; Morgan, D. O.; Shokat, K. M. *PNAS* **2008**, *105*, 1442–1447.

(4) (a) Parang, K.; Kohn, J. A.; Saldanha, S. A.; Cole, P. A. *FEBS Lett.* **2002**, *520*, 156–160. (b) Shen, K.; Cole, P. A. *J. Am. Chem. Soc.* **2003**, *125*, 16172–16173.

(5) Dramsi, S.; Scheid, M. P.; Maiti, A.; Hojabrpour, P.; Chen, X.; Schubert, K.; Goodlett, D. R.; Aebersold, R.; Duronio, V. *J. Biol. Chem.* **2002**, *277*, 6399–6405.

(6) Potter, L. R.; Hunter, T. *Mol. Cell. Biol.* **1998**, *18*, 2164–2172.

(7) (a) Forcet, C.; Billaud, M. *Sci. STKE* **2007**, *404*, pe51. (b) Sapkota, G. P.; Boudeau, J.; Deak, M.; Kieloch, A.; Morrice, N.; Alessi, D. R. *Biochem. J.* **2002**, *362*, 481–490.

(8) Beausoleil, S. A.; Jedrychowski, M.; Schwartz, D.; Elias, J. E.; Villén, J.; Li, J.; Cohn, M. A.; Cantley, L. C.; Gygi, S. P. *Proc. Natl. Acad. Sci.* **2004**, *101*, 12130–12135.




Figure 1. Kinase–substrate cross-linking strategy. (A) Phosphorylation of the protein substrate by an upstream kinase. (B) The serine of the kinase is replaced with a reactive cysteine moiety that facilitates kinase–substrate cross-link by cross-linker **1**.

(Figure 1).⁹ This is accomplished by replacing a phosphorylated serine or threonine residue of interest with a cysteine. This substitution provides a nucleophilic residue that is in close proximity to the site of phosphotransfer. To covalently link this cysteine to a lysine in the upstream kinase, a bifunctional, ATP-binding site directed cross-linker (**1**, Figure 1) was developed. This bifunctional reagent contains an *o*-phthaldialdehyde, which can chemoselectively cross-link a cysteine of the substrate to a lysine in the kinase. The irreversible nature of the cross-linking reaction provides the means to trap weak and transient kinase–substrate binding interactions.

Identification of kinase–substrate pairs is ideally carried out in a cellular environment which contains a full complement of cellular proteins and kinases, including scaffolding proteins which enhance kinase–substrate specificity.¹⁰ Successful execution of cross-linking reactions in complex mixtures requires low background labeling of highly abundant cellular proteins that are not relevant to the phosphorylation event being studied. Unfortunately, we have found that cross-linker **1** reacts with many proteins in mammalian cell lysates and is not suitable for use in complex proteomes.

In theory, selectivity can be improved by specifically targeting the cross-linking agent to proteins of interest and by modulating the reactivity of the dialdehyde cross-linker. This strategy is similar to the design of highly selective irreversible enzyme inhibitors,¹¹ with the distinction that the isoindole cross-linking reaction is a three-component reaction rather than a bimolecular one. Yao and co-workers recently reported a modified version of cross-linker **1** that is based on 2,3-naphthalenedicarboxaldehyde rather than on *o*-phthaldialdehyde. This substitution improves the efficiency of the cross-linking reaction in cell

lysates.¹² In this report, we systematically explore two of the structural elements that diminish the utility of cross-linker **1** in cell lysates and describe a new series of cross-linkers with greatly enhanced selectivity for the desired kinase substrate cross-linking reaction. In order to more selectively target protein kinases in cell lysates, we have replaced the weak, kinase-binding adenosine moiety of **1** with a general protein kinase inhibitor. In addition, we have replaced the *o*-phthaldialdehyde cross-linking moiety in **1** with a less-reactive five-membered heterocyclic dialdehyde. The combination of these two structural modifications has led to a compound capable of cross-linking a cysteine containing substrate to its corresponding kinase in the presence of a vast excess of competing cellular proteins without producing significant background cross-linking.

Results and Discussion

In this report, we utilized the serine/threonine kinase Akt as a model kinase for our studies.¹³ Akt is a central node in growth factor signaling, and is hypothesized to be responsible for over 100 substrate phosphorylation events.¹⁴ Although a substrate consensus motif has been identified for Akt (R-X-R-X-X-S/T-B, where X represents any amino acid and B represents bulky hydrophobic residue),¹⁵ 25% of reported Akt substrates do not contain this sequence¹³ making unambiguous determination of Akt–substrate pairs a major challenge. Additionally, phosphorylation of the kinase substrates in vitro may not properly recapitulate phosphorylation events in cells. For example, although Akt was first shown to phosphorylate mTOR on S2448 in vitro,¹⁶ further studies revealed that S6K1, itself a substrate of mTOR, reciprocally phosphorylates mTOR at S2448 in HEK293 cells.¹⁷

To optimize Akt-substrate cross-linking we chose a model peptide substrate sequence (RPRTSSF) derived from GSK3 β .¹⁸ Our previous studies demonstrated that a fluorescein-labeled Akt consensus motif peptide (Fluorescein-ZZRPRTSCF, **2**, Z = aminohexanoic acid), in which the phosphorylated serine residue is replaced with a cysteine, can be cross-linked to Akt1 in the presence of **1** (Figure 2, lane 2). However, the fluorescently labeled Akt1–peptide complex cannot be detected when the cross-linking reaction is conducted in the presence of 7 μ g of HeLa cell lysate, and the observed nonspecific fluorescent products are not discernible from those produced with HeLa cell lysate alone (Figure 2, lanes 3–4). We attribute the absence of Akt1 cross-linked product and abundance of background protein labeling, with proteins with molecular weights much smaller than known kinases, to lysine containing proteins which cross-link to fluorescently labeled peptide, **2**.

In order to reduce the high amount of background labeling, we initially chose to focus on increasing the affinity of the

(9) Maly, D. J.; Allen, J. A.; Shokat, K. M. *J. Am. Chem. Soc.* **2004**, *126*, 9160–1.

(10) Pawson, T.; Nash, P. *Science* **2003**, *300*, 445–452.

(11) Walsh, C. T. *Annu. Rev. Biochem.* **1984**, *53*, 493–535.

(12) Liu, K.; Kalesh, K. A.; Bing Ong, L.; Yao, S. Q. *ChemBiochem* **2008**, In press.

(13) Manning, B. D.; Cantley, L. C. *Cell* **2007**, *129*, 1261–1274.

(14) Bellacosa, A.; Kumar, C. C.; Di Cristofano, A.; Testa, J. R. *Adv. Cancer Res.* **2005**, *94*, 29–86.

(15) Alessi, D. R.; Caudwell, F. B.; Andjelkovic, M. b.; Hemmings, B. A.; Cohen, P. *FEBS Lett.* **1996**, *399*, 333.

(16) (a) Scott, P. H.; Brunn, G. J.; Kohn, A. D.; Roth, R. A.; Lawrence, J. C. *PNAS* **1998**, *95*, 7772–7777. (b) Sekulic, A.; Hudson, C. C.; Homme, J. L.; Yin, P.; Otterness, D. M.; Karnitz, L. M.; Abraham, R. T. *Cancer Res.* **2000**, *60*, 3504–3513.

(17) (a) Chiang, G. G.; Abraham, R. T. *J. Biol. Chem.* **2005**, *280*, 25485–25490. (b) Holz, M. K.; Blenis, J. *J. Biol. Chem.* **2005**, *280*, 26089–26093.

(18) Cross, D. A.; Alessi, D. R.; Cohen, P.; Andjelkovich, M.; Hemmings, B. A. *Nature* **1995**, *378*, 785–789.




Figure 2. Cross-linking of Akt1 with cross-linker **1** and Fluorescein-ZZRPRTSCF-OH **2**, in the presence of HeLa cell lysate proteins. Fluorescent peptide **2** (1 μ M), cross-linker **1** (20 μ M), Akt1 (22 nM), and HeLa cell lysate (7 μ g) were incubated for 20 min at room temperature (rt), followed by SDS PAGE and in gel fluorescent scanning.

adenosine portion of **1** for protein kinases. Adenosine is a weak, micromolar ligand for most protein kinases (Figure 3A) and is capable of binding to many other ATP utilizing proteins outside of the protein kinase family. We reasoned that replacing the adenosine fragment in cross-linker **1** with an alternative molecular scaffold, that has higher specificity and binding affinity for kinase ATP-binding sites would reduce the amount of background labeling. A key requirement for such a molecular scaffold is to maintain promiscuous kinase binding to ensure the resulting cross-linkers would be general and could be used to covalently cross-link any kinase with the substrate in question.

An analysis of nonselective kinase inhibitors revealed a common structural motif that is responsible for promiscuous kinase-binding behavior.¹⁹ Aminopyrazole **3** (Figure 3C) contains all the necessary structural properties for promiscuous inhibition, and therefore, appeared to be a promising molecular replacement for the adenosine fragment in cross-linker **1** (Figure 3B). In order to evaluate ability of aminopyrazole **3** to serve as a general scaffold for an *o*-phthaldialdehyde based cross-linker, we first determined if it is capable of binding to a wide assortment of protein kinases.²⁰ The affinity of aminopyrazole **3** for the ATP-binding site of 402 diverse kinases was assessed with a phage-encoded kinase library based method as described previously.²¹ Compound **3** was screened against a panel of 402 kinases (including 43 clinical mutant kinases) at a single concentration (10 μ M) to identify the number of kinases addressable by **3**. The screen revealed that compound **3** binds 337 out of 359 wild type kinases tested (Supporting Information, Figure 4 and Table 1). A quantitative dissociation constant K_d was then determined for 40 representative kinases that displayed binding interactions with the kinase inhibitor **3**.²² CDK2, CDK5, and JNK3 kinases were among the most potently inhibited kinases with K_d 's of 4, 27, and 26 nM, respectively. Inhibitor **3** has low affinity for several kinases, including Akt1, making cross-linking to this kinase a particularly stringent test of the broad utility of the aminopyrazole based cross-linker.

(19) Aronov, A. M.; Murcko, M. A. *J. Med. Chem.* **2004**, *47*, 5616–5619.

(20) See the Supporting Information for the synthesis of aminopyrazole **3**.

(21) (a) Fabian, M. A.; et al. *Nat. Biotechnol.* **2005**, *23*, 329–336. (b) Karaman, M. W.; et al. *Nat. Biotechnol.* **2008**, *26*, 127–132.

(22) See Supporting Information.

Having established that compound **3** is a promiscuous kinase binder, we turned to identifying the appropriate linkage to the *o*-phthaldialdehyde moiety. We attempted to co-crystallize **3** bound to the tyrosine kinase c-Src²³ to facilitate proper positioning of the *o*-phthaldialdehyde moiety with respect to the kinase active site. Although we solved the crystal structure of the c-Src-**3** cocomplex to 2.0 Å resolution, it was not possible to unambiguously determine the binding orientation of inhibitor **3** in the active site. Thus, we prepared a close structural analogue of **3**, with a quinazoline rather than a pyrimidine core, **4** (Figure 3C), in hopes that an extra phenyl ring would aid in modeling the bound inhibitor **4** into the electron density map, without disrupting binding to the kinase active site.

Kinase inhibitor **4** was prepared starting from 2,4-dichloroquinazoline utilizing the synthetic route used for **3**.²⁴ Under the previously optimized crystallization conditions, we were able to obtain a 2.35 Å resolution crystal structure of compound **4** bound to c-Src and unambiguously visualize its binding mode. The complex of Src with compound **4** crystallized under very similar conditions as the Src·imatinib complex,²⁵ and the unit cell dimensions of the two crystals are also closely related. Because the four Src complexes (which we designate chain A, B, C, and D) with compound **4** in the crystallographic asymmetric unit are highly similar, we will only discuss one of them (chain A).

Compound **4** forms a total of three hydrogen bonds with c-Src. The aminopyrazole forms three hydrogen bonds with the backbone of Glu 339 and Met 341, which mimic the hydrogen bonding pattern of adenine in ATP binding to the kinase. The quinazoline core in compound **4** forms hydrophobic contacts with the side chain of Leu 273 (Figure 3D). Two additional recently deposited structures of compound **4**, co-crystallized with human Ste20-like kinase and human calcium calmodulin dependent protein kinase II δ , show similar binding modes²⁶ which suggests that this class of inhibitors bind to divergent kinases in a consistent manner. In addition, quinazoline based kinase inhibitor **4**, similar to **3**, displayed promiscuous kinase binding properties and bound 317 out of 359 wild type kinases tested (Supporting Information, Table 1). Similar kinase binding profiles for compounds **3** and **4** indicated that the replacement of pyrimidine scaffold in **3** with quinazoline for crystallographic purposes was appropriate and did not significantly impair promiscuous kinase binding properties of the resulting inhibitor **4**. Taken together, these results demonstrate that compound **3** is a promiscuous kinase binder and may serve as alternative molecular scaffold to replace the adenosine fragment in the initially discovered cross-linker **1**.

Structural analysis of **4** bound to the three kinases (c-Src, Ste20-like kinase, and human calcium calmodulin dependent protein kinase II δ) reveals the nitrile group in **4** is in the position corresponding to the 5'-hydroxyl group of ATP. This position is the site of attachment for the *o*-phthaldialdehyde in **1**. We thus replaced the nitrile group of **3** with the *o*-phthaldialdehyde

(23) Benati, D.; Baldari, C. T. *Curr. Med. Chem.* **2008**, *15*, 1154–65.

(24) See Supporting Information.

(25) Seeliger, M. A.; Nagar, B.; Frank, F.; Cao, X.; Henderson, M. N.; Kuriyan, J. *Structure* **2007**, *15*, 299–311.

(26) (a) Pike, A. C.; Rellos, P.; Niesen, F. H.; Turnbull, A.; Oliver, A.; Parker, S. A.; Turk, B. E.; Pearl, L. H.; Knapp, S. *EMBO J.* **2008**, *27*, 704–714. (b) Roos, A. K.; Rellos, P.; Salah, E.; Pike, A. C. W.; Fedorov, O.; Pilka, E. S.; Von Delft, F.; Arrowsmith, C. H.; Weigelt, J.; Edwards, A.; Bountra, C.; Knapp, S.; PDB deposited (PDB 2VN9).

(27) See Supporting Information.




Figure 3. (A) Schematic representation of cross-linker **1** interacting with the hinge region of PKA. (B) Schematic representation of the aminopyrazole scaffold based cross-linker interacting with the hinge region of PKA. (C) Chemical structures of promiscuous kinase inhibitors **3** and **4**. (D) Crystal structure of kinase inhibitor **4** bound to c-Src at 2.35 Å resolution. Key interactions and hydrogen bond distances are highlighted.




Figure 4. (A) Synthesis of cross-linker **10**. (B) Cross-linking Akt1 and fluorescent substrate peptide **2** with or without HeLa cell lysates. Akt1, fluorescent peptide **2** with or without HeLa cell lysates, were treated with either cross-linker **1** or **10** for 20 min, followed by SDS-PAGE and in gel fluorescence scanning. (C) Cross-linking of endogenous HeLa cell lysate proteins with substrate peptide **2**, in the presence of cross-linker **1** or *o*-phthalaldehyde. HeLa cell lysates and substrate peptide **2** were treated with cross-linker **1** or *o*-phthalaldehyde for 20 min at rt, followed by SDS-PAGE and in gel fluorescence scanning.

to produce candidate cross-linker **10**. The synthetic route for **10** is illustrated in Figure 4A. Treatment of the previously synthesized intermediate **5**²⁷ with the commercially available *p*-phenylenediamine in the presence of catalytic amount of HCl in BuOH at 100 °C furnished amine **7**. The final installation of the dialdehyde moiety was conducted by the treatment of amine

7 with aldehyde **8** under reductive amination conditions followed by acidic hydrolysis to yield cross-linker **10**.

The aminopyrazole-based cross-linker **10** efficiently cross-linked Akt1 kinase to cysteine containing peptide **2** with a 2-fold increase in intensity of the cross-linked product band when compared with the adenosine based cross-linker **1** (Figure 4B,

lanes 1–2). In addition the cross-linking reaction was accompanied by the formation of higher molecular weight (~180 kDa) fluorescent species (Figure 4B, lane 2). We attribute these products to further oligomerization of the Akt1-peptide cross-linked product induced by the dialdehyde **10** reacting with lysines on the Akt1 kinase surface. The observation that the cross-linking reaction tolerates significant changes in chemical structure of the initially developed dialdehyde **1** points to the robustness of the core isoindole forming reaction.

We asked if cross-linking of Akt1 with its substrate peptide **2** could be observed in the presence of HeLa cell extract. To determine if the newly designed cross-linker was more selective in the presence of cellular proteins, a mixture of recombinant Akt1 (560 nM), substrate peptide **2** (1 μ M) and HeLa cell lysate (7 μ g) were treated with either cross-linker **1** or **10**. Unfortunately, even the newly designed aminopyrazole based cross-linker suffered a significant decrease in the efficiency of cross-linking of Akt1 with its substrate peptide **2** (Figure 4B, lanes 3–4) when compared with experiments without cell lysate present (Figure 4B, lanes 1–2). The fluorescently labeled reaction products formed a complex pattern of fluorescently labeled bands which spanned the entire molecular weight range of the cell lysate proteome which was similar to that produced with **1** (Figure 4B, lanes 3–4 and Figure 2, lanes 3–4, respectively). The intensity of these background bands from the HeLa cell lysate was somewhat mitigated when cross-linker **10** was employed rather than **1** (Figure 4B, lanes 3–4). Our interpretation is that the kinase-binding portion of the cross-linker reagent is partially responsible for the nonspecific background reactions yet other elements of cross-linker **10** are responsible for the majority of the undesired reactivity of the cross-linker. Therefore, we set out to identify and modify the other remaining elements of **10** in order to obtain a kinase-substrate cross-linker suitable for whole cell lysate studies.

We focused on the *o*-phthalaldehyde moiety since its reactivity is a key determinant of cross-linking reaction. We asked if *o*-phthalaldehyde alone, without any kinase-directed adenosine or aminopyrazole element, is capable of cross-linking cysteine containing peptide **2** to cellular proteins. *o*-Phthalaldehyde has been used to intramolecularly label enzymes that possess proximal lysines and cysteines, such as creatine kinase²⁸ and hexokinase,²⁹ and thus is itself capable of mediating isoindole formation in a cellular context. We asked if *o*-phthalaldehyde would efficiently carry out intermolecular cross-linking of peptide **2** to cellular proteins. HeLa cell lysates were treated with fluorescently labeled peptide substrate **2**, followed by the treatment with cross-linker **1** or with *o*-phthalaldehyde. Surprisingly, *o*-phthalaldehyde and **1** produced similar patterns of cross-linked fluorescent bands, with only modest differences in the fluorescent intensity (Figure 4C). Since *o*-phthalaldehyde contains no particular protein targeting moieties (e.g., adenosine), yet exhibits similar cross-linking to a diverse set of proteins we conclude the broad spectrum of cross-linked products produced by **1** and **10** are a result of the high reactivity of *o*-phthalaldehyde itself. Therefore, we aimed to reduce the reactivity of *o*-phthalaldehyde toward nonkinase proteins in the cell lysates.

The observation that *o*-phthalaldehyde efficiently cross-links background proteins in cell lysates led us to use this assay as a

screen for a second generation of dialdehyde species with reduced reactivity. Consideration of the isoindole forming reaction mechanism and previous structure activity studies of the reaction,³⁰ led us to investigate a ketone functionality as a replacement for one aldehyde group in *o*-phthalaldehyde, to produce *o*-acetylbenzaldehyde **11** (Figure 5A). This analogue is predicted to serve as a less reactive substitute to *o*-phthalaldehyde, since it contains a sterically hindered and less electron deficient acetyl group, while still reacting with thiols and amines to produce isoindoles.³¹ Naphthalene-2,3-dialdehyde **12** was also included in the screen since Yao and co-workers reported an improved version of cross-linker **1**, containing naphthalene-2,3-dialdehyde.¹² Replacement of the acetyl group in *o*-acetylbenzaldehyde **11** with a cyanogroup led to *o*-cyanobenzaldehyde **13** as another possible candidate with reduced reactivity, this cyanoaldehyde forms a thiazolidine based product rather than isoindole.³² We also asked if replacing the six-membered aromatic ring in *o*-phthalaldehyde with a five-membered aromatic heterocycle, such as thiophene, could eliminate nonspecific cross-linking in cell lysates, due to additional steric strain of the resulting thieno[2,3-*c*]pyrrole-based cross-linked product when compared with isoindole. Indeed, commercially available thiophene-2,3-dialdehyde **14** (Figure 5A) has been shown to react with primary thiols and amines to produce thieno[2,3-*c*]pyrroles,³³ which are isoelectronic to isoindoles.

We compared the cross-linking of fluorescently labeled Akt1 substrate peptide **2** to HeLa cell lysate proteins in the presence of each of the unsubstituted cross-linkers, *o*-phthalaldehyde or **11–14** (Figure 5A). *o*-Acetylbenzaldehyde **11** produced a less-intense, but similar pattern of fluorescently labeled bands compared to *o*-phthalaldehyde (Figure 5A, lane 2). Naphthalene-2,3-dialdehyde **12** produced fewer but did not completely eliminate nonspecifically labeled fluorescent cross-linked products when compared with *o*-phthalaldehyde or *o*-acetylbenzaldehyde **11**. However, both *o*-cyanobenzaldehyde **13** and thiophene-2,3-dialdehyde **14** showed dramatic reductions in the nonspecific labeling of cell lysate proteins (Figure 5A, lane 3–4).

On the basis of the dramatic reduction of the background reactions of cellular proteins with the thiophene dialdehyde **14** and its ability to form thieno[2,3-*c*]pyrrole based products upon reaction with amines and thiols,³³ we asked if we could direct thiophene-2,3-dialdehyde to the kinase ATP-binding site and produce a kinase-substrate cross-link. Since aminopyrazole-based cross-linker **10** appeared to produce less background labeling when compared to adenosine-based cross-linker **1**, we chose to append the thiophene dialdehyde moiety directly to aminopyrazole **7**. The required dialdehyde **15** was prepared in two steps from commercially available 2,3-thiophenedialdehyde **14** (Figure 5B).³⁴ Reaction of the aldehyde **15** with the previously synthesized amine **7**, under reductive amination conditions, furnished compound **16** which was subsequently hydrolyzed under mild acidic conditions to afford cross-linker **17**.

(30) Zuman, P. *Chem. Rev.* **2004**, *104*, 3217–3238.

(31) Sternson, L. A.; Stobaugh, J. F.; Repta, A. J. *Anal. Biochem.* **1985**, *233*, 246.

(32) Oliver, G. L.; Dann, J. R.; Gates, J. W. *J. Am. Chem. Soc.* **1958**, *80*, 702–706.

(33) El-borai, M. A.; Aly, E. A.; Gerges, S. S. *Phosphorus, Sulfur Silicon Relat. Elem.* **1995**, *101*, 9–15.

(34) See Supporting Information.

(28) Sheikh, S.; Mukunda, K.; Katiyar, S. S. *Biochim. Biophys. Acta* **1993**, *1203*, 276–81.

(29) Puri, R. N.; Bhatnagar, D.; Roskoski, R., Jr. *Biochim. Biophys. Acta* **1988**, *957*, 34–46.




Figure 5. (A) Cross-linking of endogenous HeLa cell lysate proteins with fluorescent Akt1 substrate peptide **2** in the presence of different cross-linkers OPA or aldehydes **11–14**. HeLa cell lysates and fluorescent Akt1 substrate peptide **2** were treated with cross-linkers OPA or **11–14** for 20 min at rt, followed by SDS-PAGE and in gel scanning fluorescence. (B) Synthesis of the thiophene-2,3-dialdehyde based cross-linker **17**. (C) Cross-linking of Akt1 and fluorescent substrate peptide **2** with cross-linker **17**. Akt1 and fluorescent peptide **2** were treated with cross-linker **17**, for 20 min at rt, followed by SDS-PAGE and in gel scanning fluorescence. (D) Cross-linking of Akt1 and fluorescent peptide substrate **2** with cross-linker **17** in the presence of increasing amounts of HeLa cell lysate proteins. Akt1 and substrate peptide **2** were treated with cross-linker **17** for 20 min at rt, followed by SDS-PAGE and in gel scanning fluorescence.

First, we tested if cross-linker **17** was capable of cross-linking recombinant Akt1 kinase to the fluorescent peptide substrate **2**. Treatment of Akt1 and substrate peptide **2** with cross-linker **17**, produced an intensely fluorescent band of labeled Akt1 kinase, confirming that the newly prepared cross-linker **17** is capable of mediating the desired kinase–substrate covalent cross-link (Figure 5C). However, as in the case of cross-linker **10** (Figure 4B, lane 2), the cross-linking reaction was accompanied by the formation of higher-molecular-weight (\sim 180 kDa) species (Figure 5C, lane 1). We next optimized conditions for kinase substrate cross-linking employing the newly synthesized cross-linker **17**. Studies on the pH dependence of the cross-linking reaction indicate that pH 5 is optimal for the cross-linking reaction (Supporting Information, Figure 1A). The cross-linking reaction is also resistant to the presence of competing thiol nucleophiles, tolerating up to 1000-fold excess of β -mercaptoethanol relative to cysteine containing peptide substrate **2** (Supporting Information, Figure 1B), thus illustrating remarkable chemoselectivity toward the thiol containing peptide. Covalently cross-linked kinase–substrate complex was readily cleaved upon heating as no detectable cross-linked band was observed when samples were boiled prior to SDS-PAGE analysis (Supporting Information, Figure 1C). The reaction showed linear dose dependence, with respect to both cross-linker **17** and fluorescent peptide **2** concentrations (Supporting Information, Figures 2 and 3). Saturation of the cross-linked signal was observed when the concentration of cross-linker **17** and peptide **2** reached \sim 40 and 60 μ M, respectively (Supporting Information, Figures 2 and 3).

Next, we tested if cross-linker **17** could specifically cross-link Akt1 kinase to its substrate peptide **2** in a complex cell lysate. We carried out the cross-linking reaction with Akt1 and its substrate peptide **2** (1 μ M or 49 ng of peptide **2** per reaction) alone (Figure 5D, lane 1), and in the presence of increasing amounts of cell lysate from 1 μ g (lane 2) up to 32 μ g (lane 7) (Figure 5D). Remarkably, fluorescein-containing cross-linked product is readily apparent in the presence of up to 8 μ g of HeLa cell lysate (Figure 5D, lane 6). Thus, thiophene based cross-linker **17** represents a significant improvement over cross-linkers **1** or **10** which produced abundant nonkinase cross-linked products in the presence of cell lysates (Figure 4B, lanes 3–4).

In conclusion, a three-component chemical reaction leading to the formation of the covalently cross-linked kinase–substrate complex in vitro, has been developed to cross-link recombinant Akt kinase with its cysteine-containing peptide substrate in the presence of cell lysates. This was achieved through extensive structural modifications to the adenosine and dialdehyde fragments in cross-linker **1**. The adenosine fragment in cross-linker **1** was replaced with a promiscuous kinase inhibitor that has a nanomolar binding affinity toward a wide range of kinases to enhance targeting to protein kinase active sites. We determined that *o*-phthalaldehyde itself is largely responsible for the nonspecific labeling of cell lysate proteins. We have developed an alternative dialdehyde, thiophene-2,3-dicarboxaldehyde **14**, that shows a remarkable decrease in the nonspecific background labeling of cell lysate proteins. Replacing the *o*-phthalaldehyde derived fragment in our newly designed cross-linker **10** with the thiophene-2,3-dicarboxaldehyde moiety led to cross-linker

17 that allowed selective cross-linking of recombinant Akt1 kinase with its cysteine containing peptide substrate in the presence of competing cellular proteins.

Our findings represent a significant step toward a solution to the challenge of identifying upstream kinases for the given substrate of interest, although further challenges remain. The next step is to perform the cross-linking reaction between the kinase and its full length substrate protein. The replacement of the substrate peptide with the full-length protein may further increase the specificity of the cross-linking reaction in the presence of cell lysates, due to additional recognition elements presented on the protein substrate surface, that are distal from the local phosphorylation site. For example, p38 kinase phosphorylates the protein substrate MAPKAP2 150-fold more efficiently than the corresponding 14-residue substrate peptide resembling local phosphorylation site.³⁵ Once conditions for the cross-linking reaction between full-length kinase and substrate protein are optimized, the next challenge is to perform the cross-linking of the protein substrate with its endogenous kinases in cell lysates. Since many endogenous kinases are present at very low abundance in cells, the detection limits of

the cross-linking reaction must still be addressed. Additional structural modifications to **15** such as optimization of the linker which connects the kinase binding scaffold with the dialdehyde moiety may further increase the efficiency of the reaction in the presence of cellular proteins. The ultimate challenge is to perform the cross-linking reaction in intact cells, as the intact cellular environment provides all the elements necessary encoding kinase substrate specificity via scaffolding proteins, specific protein–protein interaction domains, and subcellular colocalization of kinases and their substrates.

Acknowledgment. M.A.S. was supported by a Johnson & Johnson fellowship of the Life Science Research Foundation, Baltimore, and NIH Grant No. K99 GM080097. K.M.S. thanks NIH for Grant No. RO1-EB001987 for funding, as well as a grant from Eli Lilly.

Supporting Information Available: Complete experimental procedures, kinase binding data for compounds **3** and **4**, X-ray crystallographic data, compound characterization, and complete refs 3c, 21a, and 21b. This material is available free of charge via the Internet at <http://pubs.acs.org/>.

JA807066F

(35) Hawkins, J.; Zheng, S.; Frantz, B.; LoGrasso, P. *Arch. Biochem. Biophys.* **2000**, *382*, 310.