

Reaction Mechanism of the ϵ Subunit of *E. coli* DNA Polymerase III: Insights into Active Site Metal Coordination and Catalytically Significant Residues

G. Andrés Cisneros,^{*,†} Lalith Perera,^{*,†} Roel M. Schaaper,[‡] Lars C. Pedersen,[†] Robert E. London,[†] Lee G. Pedersen,^{†,§} and Thomas A. Darden[†]

Laboratories of Structural Biology and Molecular Genetics, National Institute of Environmental Health Sciences, Research Triangle Park (RTP), North Carolina 27709

Received October 21, 2008; E-mail: cisnero1@niehs.nih.gov; pereral2@niehs.nih.gov

Abstract: The 28 kDa ϵ subunit of *Escherichia coli* DNA polymerase III is the exonucleotidic proofreader responsible for editing polymerase insertion errors. Here, we study the mechanism by which ϵ carries out the exonuclease activity. We performed quantum mechanics/molecular mechanics calculations on the N-terminal domain containing the exonuclease activity. Both the free- ϵ and a complex ϵ bound to a θ homologue (HOT) were studied. For the ϵ -HOT complex Mg²⁺ or Mn²⁺ were investigated as the essential divalent metal cofactors, while only Mg²⁺ was used for free- ϵ . In all calculations a water molecule bound to the catalytic metal acts as the nucleophile for hydrolysis of the phosphate bond. Initially, a direct proton transfer to H162 is observed. Subsequently, the nucleophilic attack takes place followed by a second proton transfer to E14. Our results show that the reaction catalyzed with Mn²⁺ is faster than that with Mg²⁺, in agreement with experiment. In addition, the ϵ -HOT complex shows a slightly lower energy barrier compared to free- ϵ . In all cases the catalytic metal is observed to be pentacoordinated. Charge and frontier orbital analyses suggest that charge transfer may stabilize the pentacoordination. Energy decomposition analysis to study the contribution of each residue to catalysis suggests that there are several important residues. Among these, H98, D103, D129, and D146 have been implicated in catalysis by mutagenesis studies. Some of these residues were found to be structurally conserved on human TREX1, the exonuclease domains from *E. coli* DNA-Pol I, and the DNA polymerase of bacteriophage RB69.

Introduction

The fidelity of DNA replication remains a subject of considerable interest because of its relevance to a large part of biology.¹ Studies of DNA replication of *Escherichia coli* (*E. coli*) suggest that the in vivo substitution error rate for the replication machinery is around 10⁻¹⁰.² The high rate of fidelity is obtained by at least three activities that operate serially: base selection, exonucleolytic proofreading, and postreplicative mismatch repair. Of these three the exonucleolytic proofreading activity has been reported to have an error discrimination in the range of 40–400 fold.²

The replication of DNA in *E. coli* is performed by DNA polymerase III holoenzyme (Pol III). This is a large protein complex that contains 10 distinct subunits.³ In this complex there are two catalytic cores composed of three subunits named α , ϵ , and θ bound in this linear order. The ϵ subunit has been shown to be responsible for the proofreading exonuclease activity in Pol III.⁴

The exonuclease activity removes newly incorporated nucleotides, preferentially excising incorrect nucleotides.⁵ The 186-residue N-terminal domain of ϵ (ϵ 186) has been shown to contain the exonuclease active site.⁶ The reaction catalyzed by ϵ is proposed to involve either one⁷ or two divalent metal ions.⁸ For the two-metal mechanism the catalytic metal (Me1) is proposed to facilitate formation of an attacking hydroxide ion. This hydroxide performs a nucleophilic attack on the α phosphate of the nucleotide base to be excised. The second metal is termed the nucleotide binding metal (Me2). Recent experimental studies have shown that reaction of ϵ in vitro is faster with Mn²⁺ than with Mg²⁺.⁸

Association of ϵ with θ has been proposed to promote exonuclease activity.^{9,10} In addition, it has recently been shown that a homologue of theta (HOT) from bacteriophage P1 can

[†] Laboratory of Structural Biology, National Institute of Environmental Health Sciences.

[‡] Laboratory of Molecular Genetics, National Institute of Environmental Health Sciences.

[§] Department of Chemistry, University of North Carolina, Chapel Hill, NC 27599.

(1) Kunkel, T. A. *J. Biol. Chem.* **2004**, *279*, 16895–16898.

(2) Schaaper, R. M. *J. Biol. Chem.* **1993**, *268*, 23762–23765.

(3) McHenry, C. S. *J. Biol. Chem.* **1991**, *266*, 19127–19130.

(4) Brenowitz, S.; Kwack, S.; Goodman, M. F.; O'Donnel, M.; Echols, H. *J. Biol. Chem.* **1991**, *266*, 7888–7892.

(5) Derbyshire, V.; Pinsonneault, J. K.; Joyce, C. M. *Methods Enzymol.* **2006**, *262*, 363–385.

(6) Perrino, F. W.; Harvey, S.; McNeill, S. M. *Biochemistry* **1999**, *38*, 16001–16009.

(7) Cowan, J. *BioMetals* **2002**, *15*, 225–235.

(8) Hamdan, S.; Bulloch, E. M.; Thompson, P. R.; Beck, J. L.; Yang, J. Y.; Crowther, J. A.; Lilley, P. E.; Carr, P. D.; Ollis, D. L.; Brown, S. E.; Dixon, N. E. *Biochemistry* **2002**, *41*, 5266–5275.

(9) Taft-Benz, S. A.; Schaaper, R. M. *J. Bacteriol.* **2006**, *186*, 2774–2780.

(10) Lehtinen, D. A.; Perrino, F. W. *Biochem. J.* **2004**, *384*, 337–348.

and forming) the 6-31+G* basis was used. All other (nonreactive) QM atoms were represented with 3-21G. The Mn^{2+} atoms were treated with the LANL2DZ pseudopotential. The remaining atoms were included in the MM subsystem and treated with the parm99 force field.²⁶ We previously used these methods to study DNA polymerase λ and observed that the combined basis sets give similar results to those calculated with unmixed (larger) basis sets.¹⁵ After the QM/MM optimization of $\epsilon 186\text{-HOT}(Mg^{2+})$ the free- $\epsilon 186$ system was prepared by replacing HOT with solvent and re-equilibrating. This resulted in a total of 18 043 atoms for free- $\epsilon 186$.

The product structures for all systems ($\epsilon 186\text{-HOT } Mg^{2+}$, Mn^{2+} , and free- $\epsilon 186 \text{ } Mg^{2+}$) were produced in silico from the reactant structure and subjected to QM/MM optimization. Path optimizations were performed following a procedure used previously for the study of the reaction mechanism of DNA polymerase λ .¹⁵ The optimized end points were employed for the reaction path calculation with the quadratic string method (QSM).³² This is a “chain-of-replica” method where the path is represented by a discrete number of structures^{22,33–36} In this method the intermediate structures are obtained by a linear interpolation between the end points, and all structures are optimized to the minimum energy path simultaneously. This results in a highly efficient and unbiased optimization of the path. The initial QSM paths consist of 10 points including end points and were iteratively optimized with constrained MM optimization.^{36–38} After the initial paths were converged the highest energy points were optimized to the closest TS with the quadratic synchronous transit (QST3) method.^{39,40} Following TS optimization the paths between the critical points were calculated with either QSM or reaction coordinate driving (RCD) (see Supporting Information).

All MD simulations were carried out with the AMBER9 suite of programs.²⁶ QM/MM calculations were carried out with modified versions of Gaussian03 and TINKER.^{41,42} The QM/MM optimizations were performed with an iterative method,^{43,44} using the pseudobond model for all boundary atoms.^{45,46} All QM calculations were performed using an effective Hamiltonian (electrostatic embedding).^{43,44}

Results and Discussion

Initial Structure Determination. MD simulations were performed for both $\epsilon 186\text{-HOT}$ and free- $\epsilon 186$ with Mg^{2+} . As mentioned above, the X-ray crystal structures show that the catalytic metal is pentacoordinated with E14 forming a single bond to the metal and a hydrogen bond to the nucleophilic water. Initially the systems were unconstrained during the MD simula-




Figure 2. Atoms included in the QM region. Note that most of the dT nucleotide is not present. Nucleophilic water is labeled “wat”, and pseudobonds are shown in gray.

tions. However, this results in a distortion of the active site because E14 forms a second bond to the catalytic Mg (chelating orientation) and loses the hydrogen bond to the water molecule. This results in an increased intermetal distance. The binding of the substrate is disrupted. The resulting structure contains a differently pentacoordinated catalytic metal that is not conducive to chemistry. This change in coordination may be due to limitations in the classical force field.

A further attempt to regain the hexacoordination involved considering the carbonyl backbone oxygen from T8 as the sixth ligand. This atom is located approximately 3.5 Å from the catalytic metal in the X-ray structure. However, after MD this atom increases the intermetal distance to around 4 Å. Furthermore, its inclusion in the coordination shell would involve displacement of other axial ligands. This would possibly disrupt the orientation of the substrate or the position of E14, resulting in a structure not conducive to catalysis. One last possibility would be for a water molecule to occupy the place of the carbonyl. However, although the carbonyl backbone from T8 moves slightly away, there is still not enough room for a water molecule to fill a sixth coordination. Therefore, in order to retain a catalytically competent structure the MD equilibrations were repeated including a constraint to enforce the single bond of E14 to the catalytic metal.

Subsequently, QM/MM optimizations were carried out with all constraints removed based on the equilibrated systems. The final rmsd of the backbone atoms between the X-ray and optimized structures are 0.4, 0.6, and 0.8 Å for the $\epsilon 186\text{-HOT}(Mg^{2+})$, $\epsilon 186\text{-HOT}(Mn^{2+})$, and free- $\epsilon 186$, respectively. The rmsd of all heavy atoms in the active site between the X-ray and the optimized systems is 0.2 Å for the $\epsilon 186\text{-HOT}$ structures and 0.1 Å for free- $\epsilon 186$.

The QM/MM optimizations do not contain any constraints. In all cases, the five ligands around the catalytic metals were observed to be maintained after the optimizations (see Figure 2 and Table S1 in the Supporting Information for coordination distances). The final calculated structures are consistent with the pentacoordinated structures observed experimentally.^{13,14} The observed geometry for the catalytic metals is distorted trigonal bipyramidal with angles around 110–120° for the axial and around 170° for the equatorial ligands. This is not unexpected for the Mn^{2+} system since Mn^{2+} is known to support this type of coordination; however, this is not common for Mg^{2+} . To investigate whether the Mg^{2+} structures are indeed pentacoordinated, the unconstrained MD structure of $\epsilon 186\text{-HOT}$ (with E14 chelating the catalytic metal) was used for QM/MM

- (32) Burger, S. K.; Yang, W. *J. Chem. Phys.* **2006**, *124*, 054109.
 (33) Elber, R.; Karplus, M. *Chem. Phys. Lett.* **1987**, *139*, 375–380.
 (34) Maragakis, P.; Stefan, A.; Brumer, Y.; Reichman, D.; Kaxiras, E. *J. Chem. Phys.* **2002**, *117*, 4651–4658.
 (35) Liu, H.; Lu, Z.; Cisneros, G. A.; Yang, W. *J. Chem. Phys.* **2004**, *121*, 697–706.
 (36) Cisneros, G. A.; Liu, H.; Lu, Z.; Yang, W. *J. Chem. Phys.* **2005**, *122*, 114502.
 (37) Jhih-Wei, C.; Trout, B.; Brooks, B. *J. Chem. Phys.* **2003**, *119*, 12708–12717.
 (38) Xie, L.; Liu, H.; Yang, W. *J. Chem. Phys.* **2004**, *120*, 8039–8052.
 (39) Gonzalez, C.; Schlegel, H. *J. Phys. Chem.* **1990**, *94*, 5523–5527.
 (40) Schlegel, H. *J. Comput. Chem.* **2003**, *24*, 1514–1527.
 (41) Frisch, M. J., et al. *Gaussian 03*, Revision D.02; Gaussian, Inc.: Wallingford, CT, 2004.
 (42) Ponder, J. *TINKER, Software Tools for Molecular Design*, Version 3.6; Washington University: St. Louis, 1998; the most updated version for the TINKER program can be obtained from J. W. Ponder’s site at <http://dasher.wustl.edu/tinker>.
 (43) Zhang, Y.; Liu, H.; Yang, W. *J. Chem. Phys.* **2000**, *112*, 3483–3491.
 (44) Zhang, Y.; Liu, H.; Yang, W. *Ab Initio QM/MM and Free Energy Calculations of Enzyme Reactions*. In *Computational Methods for Macromolecules-Challenges and Applications*; Springer Verlag: Heidelberg, Germany, 2002.
 (45) Zhang, Y.; Lee, T.; Yang, W. *J. Chem. Phys.* **1999**, *110*, 46–54.
 (46) Zhang, Y. *J. Chem. Phys.* **2005**, *122*, 024114.




Figure 3. HOMO for the optimized ϵ 186-HOT (Mg^{2+}) reactant structure (QM subsystem only), cutoff = 10^{-3} au. Orbital colors denote sign of the respective lobe (red = positive). Note that the orientation is the same as in Figure 2.

optimizations. Three optimizations were performed with different MM environments (taken from different snapshots of the MD simulation) for the reactant and product structures, and in all cases the optimized structures reverted to the single E14– Mg^{2+} (catalytic) bond.

Analysis of the charges on the metals and the location of the highest occupied molecular orbital (HOMO) helps understand the observed pentacoordination for the Mg-containing structures. A significant charge transfer to the metals is observed with charges of 1.3 and 1.5 for the catalytic and nucleotide binding metals, respectively, calculated with Merz–Kollman fitting.⁴⁷ These charges are approximately 0.2–0.4 units less positive than charges calculated by the same procedure for the equivalent metals (both hexacoordinated) in our previous study of DNA polymerase λ .¹⁵ In addition, as seen in Figure 3, the HOMO is centered almost exclusively on the catalytic metal, D12, D167, H162, and the nucleophilic water. This is also observed for free- ϵ 186 (results not shown). Furthermore, cluster calculations on penta- and hexa-hydrated Mg^{2+} give charges close to 2+ on the ion. Taken together, this suggests that the HOMO could provide a channel that enables the catalytic metal to attract charge density from the ligands. This charge transfer helps stabilize the unusual coordination sphere.

Furthermore, pentacoordinated Mg is not exclusive to ϵ . This coordination has been reported in inorganic compounds^{48,49} and other enzymes such as the Klenow fragment of DNA polymerase I,⁵⁰ ribonuclease H (RNaseH),^{51,52} GTPase Ffh,⁵³ and tryptophanyl-tRNA synthetase.^{54–56} Additionally, in a recent simulation of the related RNaseH reaction mechanism this coordination has also been reported.⁵⁷ Interestingly, in the case of RNaseH the pentacoordinated Mg^{2+} is the nucleotide binding and not the catalytic ion.^{51,52,57} These observations together with our results point to a pentacoordinated metal regardless of whether there is a Mn or Mg in the active site. Note that this

only corresponds to the catalytic step, and the octahedral geometry could be restored after catalysis or nucleotide release.

Catalytic Mechanism. On the basis of the optimized structures the corresponding product structures were generated for all three systems. These structures were employed for the reaction path optimizations as explained in the Methods section. In the case of the initial guess for the product only the transfer of the proton to H162 and nucleophilic attack were initially considered. Interestingly, however, the second proton transferred spontaneously to E14 during the optimization procedure to give a lower energy structure.

Figure 4 shows the superposition of the critical points along the reaction path for the ϵ 186-HOT with Mg^{2+} and Mn^{2+} (similar superpositions for free- ϵ 186 are given in the Supporting Information). As can be observed, the reactant structures for both metals are very similar. In all cases, the reaction begins by an initial proton transfer from the nucleophilic water to H162 to form the nucleophilic hydroxyl. Subsequently, the phosphodiester is attacked by the hydroxyl, forming a pentavalent phosphorane transition-state (TS) structure, resulting in an inversion of configuration at the P atom (see Figure 4). Finally, after the phosphate bond has been broken a second proton transfer takes place to E14.

This reaction mechanism is consistent with the proposal by Hamdan et al.⁸ Note that the second proton transfer proposed by Hamdan et al. and observed in the present work only occurs in the second leg of the reaction after the phosphate bond has been broken. Additionally, configuration inversion has been observed experimentally for the analogous reaction catalyzed by RNaseH.⁵⁸

The reaction profiles in Figure 5 show that the reactions for all systems (including both metals) proceed in an approximately similar manner. In the case of ϵ 186-HOT(Mg^{2+}) two shoulders are observed flanking the TS, which correspond to the proton transfers. However, the putative intermediates are not stable structures along the potential-energy surface (PES). This is due to the fact that when these structures were optimized they returned either to the reactant or product minima. In the case of ϵ 186-HOT(Mn^{2+}) and free- ϵ 186 only one TS is observed. In all cases the TS structures were optimized with QST3 and a single imaginary frequency was obtained. The calculated reaction barriers are 16.4, 15.4, and 18.2 kcal/mol for the ϵ 186-HOT(Mg^{2+})-, ϵ 186-HOT(Mn^{2+})-, and free- ϵ 186(Mg^{2+})-catalyzed reactions, respectively.

Experimental results for free- ϵ 186 have been reported for the hydrolysis of the 5'-*p*-nitrophenyl ester of TMP (pNP-TMP) with both Mg^{2+} and Mn^{2+} .⁸ The experimental k_{cat} of 19 and 293 min^{-1} for Mg^{2+} and Mn^{2+} correspond to barriers of 13.5 and 11.9 kcal/mol, respectively. The rate constant for ϵ 186- θ (Mn^{2+}) has been reported in the same work to be 215 min^{-1} , corresponding to a barrier of 12.1 kcal/mol. Furthermore, pre-steady-state kinetic results for degradation of a 10-mer single-stranded DNA (ss-DNA) with Mg^{2+} by ϵ give a k_{cat} of 280 s^{-1} , which corresponds to a barrier of 14.4 kcal/mol.²⁵ These barriers are obtained via the transition-state formula (TST) $k_{\text{cat}} = k_b T/h \exp$

(47) Besler, B. H.; Merz, K. M., Jr.; Kollman, P. J. *Comput. Chem.* **1990**, *11*, 431–439.

(48) Ejfler, J.; Kobylka, M.; Jerzykiewicz, L. B.; Sobota, P. *Dalton Trans.* **2005**, *11*, 2047–2050.

(49) Guzei, I. A.; McGaff, R. W.; Kieler, H. M. *Acta Crystallogr., C* **2005**, *61*, 472–475.

(50) Beese, L. S.; Steitz, T. A. *EMBO J.* **1991**, *10*, 25–33.

(51) Nowotny, M.; Yang, W. *EMBO J.* **2006**, *25*, 1924–1933.

(52) Yang, W.; Lee, J. Y.; Nowotny, M. *Mol. Cell* **2006**, *22*, 5–13.

(53) Focia, P. J.; Alam, H.; Lu, T.; Ramirez, U. D.; Freymann, D. M. *Prot. Struct. Funct. Bioinf.* **2004**, *54*, 222–230.

(54) Kapustina, M.; Carter, C. W. *J. Mol. Biol.* **2006**, *362*, 1159–1180.

(55) Kapustina, M.; Weinreb, V.; Li, L.; Kuhlman, B.; Carter, C. W. *Structure* **2007**, *15*, 1272–1284.

(56) Pham, Y.; Li, L.; Kim, A.; Erdogan, O.; Weinreb, V.; Butterfoss, G. L.; Kuhlman, B.; Carter, C. W. *Cell* **2007**, *25*, 851–862.

(57) De Vivo, M.; Dal Peraro, M.; Klein, M. L. *J. Am. Chem. Soc.* **2008**, *130*, 10955–10969.

(58) Krakowiak, A.; Owczarek, A.; Koziolkiewicz, M.; Stec, W. J. *ChemBioChem* **2002**, *3*, 1242–1250.




Figure 4. Superposition of active sites for the calculated reactants (top), transition states (middle), and products (bottom) from the ϵ 186-HOT structures. Mg^{2+} structure is shown in aquamarine with nucleophilic water in yellow, and Mn^{2+} structure is shown in purple with nucleophilic water in green. The nucleophilic waters are shown in ball and sticks, while divalent metals are rendered as large spheres. Electrophilic phosphate, $\text{O}3'$ on dC, and waters coordinating the nucleotide binding metal are shown as spheres. Hydrogen atoms except for the ones on the nucleophilic waters have been omitted for clarity.

$(-\Delta G^\ddagger/RT)$.⁵⁹ In addition, these barriers assume that the free energy of activation ΔG^\ddagger corresponds to the energy barrier.

Experimental estimates for the energy barriers point to a lower value for Mn^{2+} compared to Mg^{2+} in the free- ϵ case.⁸ This is

similar to the results obtained for the ϵ 186-HOT complex in our calculations, which show a slight preference of Mn^{2+} over Mg^{2+} . Note that the experimental substrate is different from our simulations, which may account for some of the differences. Additionally, as pointed out by Hamdan et al.,⁸ *p*-nitrophenol is a stronger acid than the 3'-OH group of a nucleotide chain and the breaking of the phosphate bond should be easier. Indeed, experimental results for pNP-TMP and ss-DNA (dT_{10}) show that the barrier for the former is around 1 kcal/mol lower than the latter.^{8,25} Moreover, our calculated barriers correspond to potential energies and provide an upper bound to the corresponding free-energy barriers.^{44,60} Therefore, it is possible that a free-energy calculation would lower the value of the barriers closer to the experimental estimate.

Our results suggest that the presence of HOT lowers the energy barrier by ~ 2 kcal/mol. On the other hand, experimental results have reported that addition of θ to ϵ has little measurable effect on the exonuclease activity.¹⁰ Also, free- ϵ 186 has been shown to have a slightly lower energy barrier (0.2 kcal/mol) than ϵ 186- θ with pNP-TMP and Mn^{2+} .⁸ Again, this discrepancy could be a result of the difference in substrates as pNP-TMP is a better leaving group than the $\text{O}3'$ of a nucleotide chain.

Another possible mechanism involves a hydroxyl ion bound to the catalytic metal instead of a water molecule. For this case, when the reactant structure with the hydroxide ion was optimized the structure immediately turned to the product. That is, the reaction immediately takes place in ϵ without a stable reactant structure (results not shown). Therefore, based on these results this reaction scheme is not viable for ϵ . This is not surprising considering what is observed in Figures 4 and 5, where the transition state corresponds to the hydroxyl ion already performing the nucleophilic attack. In contrast, De Vivo et al. have shown that the hydroxyl mechanism is much lower in energy for RNaseH.⁵⁷ In addition, as mentioned above, their structures show that the nucleotide binding metal is the pentacoordinated one not the catalytic one as in the case of ϵ .

Finally, as observed in Figure 5, although the main chemical step has taken place, the calculated products in the reaction mechanism are 5–7 kcal/mol higher than the reactants. One possible reason for this difference is the fact that the $\text{O}3'$ on the dC is not protonated immediately after the breaking of the phosphate bond. To investigate whether this is one of the reasons for the higher energy, a second product with a deprotonated H162 and protonated dC- $\text{O}3'$ was optimized for ϵ 186-HOT(Mg^{2+}) and free- ϵ 186(Mg^{2+}). Indeed, the relative energy of these products is found to be around 2 kcal/mol for ϵ 186-HOT(Mg^{2+}) and -1 kcal/mol for free- ϵ 186.

There are several possible mechanisms to transfer a proton to the $\text{O}3'$ including transfer from solvent,⁵⁷ from H162 or from E14. Two such mechanisms were tested to investigate possible paths. Both mechanisms involve transfer of the proton located on H162 that was originally on the nucleophilic water. The only direct way to transfer this proton to the $\text{O}3'$ is through an oxygen on the phosphate of the excised nucleotide (dT). In both cases the calculated barriers are over 30 kcal/mol, which rule out these paths (see Supporting Information).

Residue Analysis. Further insights into the catalytic mechanism of ϵ can be obtained by understanding the role of individual residues in catalysis. This can be achieved by decomposing the nonbonded intermolecular interaction energy between the MM

(59) Hu, H.; Yang, W. *Annu. Rev. Phys. Chem.* **2008**, *59*, 573–601.

(60) Wang, L.; Yu, X.; Hu, P.; Broyde, S.; Zhang, Y. *J. Am. Chem. Soc.* **2007**, *129*, 4731–4737.




Figure 5. Calculated reaction paths for the ϵ 186-HOT Mg^{2+} - (left), ϵ 186-HOT Mn^{2+} - (middle), and free- ϵ 186 Mg^{2+} - (right) catalyzed reactions.




Figure 6. Positions of residues (red) that have a significant (de)stabilization to the TS during catalysis for any of the three systems. Metal binding residues and H162 are shown in blue. Hydrogens have been omitted for clarity.

environment and the QM region into contributions per residue.^{61,62} This “breakdown” is only obtained in an approximate manner and, therefore, is only qualitative. In this analysis the differences in interaction energies are calculated between individual residues and the QM subsystem when the system goes from reactant to TS. A negative contribution refers to a stabilizing contribution to the TS, and the opposite applies to a positive one. We consider a residue to provide a significant contribution to (de)stabilization if the interaction is greater than 1 kcal/mol; see Supporting Information for details.

Figure 6 shows the residues that have a significant effect on catalysis for all three systems investigated. The Coulomb and van der Waals decomposition analyses show that most of the stabilization is provided by the enzyme with some contribution from the solvent (see Supporting Information). In all cases the nonbonded interactions are electrostatic with no significant contributions from the van der Waals component.

In particular, the residues that contribute to catalysis in all systems include V65, H98, D103, D129, D146, and R159. In addition, for the ϵ 166-HOT(Mg^{2+}) system two residues from HOT are observed to have significant effects: K27 and E42.

Overall the effect of the enzyme environment is stabilizing. In the case of HOT, the overall effect is slightly stabilizing when all contributions from HOT are taken into account.

Of the residues observed from decomposition, H98, D103, D129, and D146 have been studied experimentally. H98 has been shown to be a moderate dominant mutator and is well conserved among Gram-negative and Gram-positive organisms.⁶³ This residue shows a stabilizing contribution in all systems of over -1.5 kcal/mol. D146 is reported to be a weak mutator;⁶⁴ this residue is observed to have a destabilizing contribution in our results. The last two aspartates, D103 and D129, show strong mutator effects.⁶⁴ In our analysis these two residues provide large stabilizations to the TS, around -5 and -2 kcal/mol for D103 and D129, respectively. To our knowledge, none of the remaining residues from the decomposition analysis have been studied experimentally and could be targets for mutagenesis studies.

Finally, three structural alignments have been carried out to further understand the catalytic role of these residues. The alignments compare the X-ray structure of ϵ with a monomer of human TREX1 (see Figure 7),⁶⁵ the exonuclease domain (Klenow fragment) of DNA polymerase I from *E. coli*,⁶⁶ and the exonuclease domain of the DNA polymerase from bacteriophage RB69⁶⁷ (see Figures S9 and S10 in the Supporting Information). In all cases, the sequence homology is very low (below 30%); therefore, the structural alignments were performed by superposing the “DED” catalytic triad in the active site cores. Previous studies have shown that both DNA and RNA exonucleases have a conserved “DEDD” domain.^{68–71} In particular, the comparison with TREX1 is noteworthy since this enzyme has been linked to several autoimmune diseases including systemic lupus erythematosus, familial Chilblain lupus, and dominant Aicardi-Goutières syndrome.^{73–75}

As can be seen from Figures 7, S9 (Supporting Information), and S10 (Supporting Information), there are several residues in

(61) Liu, H.; Zhang, Y.; Yang, W. *J. Am. Chem. Soc.* **2000**, *122*, 6560–6570.

(62) Cisneros, G. A.; Liu, H.; Zhang, Y.; Yang, W. *J. Am. Chem. Soc.* **2003**, *125*, 10348–10393.

(63) Taft-Benz, S. A.; Schaaper, R. M. *Nucleic Acids Res.* **1998**, *26*, 4005–4011.

(64) Slupska, M. M.; King, A. G.; Liu, L. I.; Lin, R. H.; Mao, E. F.; Lackey, C. A.; Chiang, J.-J.; Baikalov, C.; Miller, J. H. *J. Bacteriol.* **1998**, *180*, 5712–5717.

(65) deSilva, U.; Choudhury, S.; Bailey, S. L.; Harvey, S.; Perrino, F. W.; Hollis, T. *J. Biol. Chem.* **2007**, *282*, 10537–10543.

(66) Brautigam, C. A.; Sun, S.; Piccirilli, J. A.; Steitz, T. A. *Biochemistry* **1999**, *38*, 696–704.

(67) Shamoo, Y.; Steitz, T. A. *Cell* **1999**, *99*, 155–166.

(68) Bernad, A.; Blanco, L.; Lázaro, J. M.; Martín, G.; Salas, M. *Cell* **1989**, *59*, 219–228.

(69) Moser, M. J.; Holley, W. R.; Chatterjee, A.; Mian, I. S. *Nucleic Acids Res.* **1997**, *25*, 5110–5118.

(70) Zuo, Y.; Deutscher, M. P. *Nucleic Acids Res.* **2001**, *29*, 1017–1026.

(71) DeRose, E. F.; Li, D.; Darden, T.; Harvey, S.; Perrino, F. W.; Schaaper, R. M.; London, R. E. *Biochemistry* **2002**, *41*, 94–110.

(72) Yang, X. Y.; Wang, M.; Fitzgerald, M. C. *J. Mol. Biol.* **2006**, *363*, 506–519.

(73) Lee-Kirsch, M. A. *Nat. Genet.* **2007**, *39*, 1065–1067.

(74) Rice, G. *Am. J. Hum. Genet.* **2007**, *80*, 811–815.




Figure 7. Structural alignment comparing residues obtained from the decomposition analysis in ϵ 186-HOT (light blue) superposed with the exonuclease domain (red) of TREX1 (red). Residues from energy decomposition for ϵ 186-HOT are shown in green; structurally homologous residues for TREX1 are shown in light blue; metal binding residues (and H162) for ϵ 186-HOT are shown in yellow; structurally homologous residues for TREX1 are shown in red.

the alignments that appear structurally conserved among the exonucleases. In particular, two homologous aspartates are conserved both in structure and position between all three exonucleases. These residues are D103 and D129 in ϵ which correspond to D130 and D154 in TREX1, D424 and D441 in the Klenow fragment, and D222 and D272 in RB69. This is a striking similarity taking into account that these residues are located in the “second” shell around the active site and also the low sequence homology among the exonuclease domains. D103 appears to be conserved among exoribonuclease superfamilies based on sequence alignments and is part of the “DEDD” domain.⁷⁰ As mentioned above, alteration of D103 and D129 have been reported to show strong mutator effects experimentally in ϵ .⁶⁴ Furthermore, D222 in RB69, equivalent to D103 in ϵ , has been shown to be catalytically important.⁷⁶ In this case, mutation of D222 to Ala in RB69 produces a reduction in catalytic activity.

Conclusions

The reaction mechanism for the *E. coli* exonuclease domain of Pol-III, the ϵ subunit, has been determined by means of QM/MM calculations. Three systems were investigated: ϵ 186-HOT

(75) Lee-Kirsch, M. A.; Choudhury, D.; Harvey, S.; Gong, M.; Seneko, L.; Engel, K.; Pfeiffer, C.; Hollis, T.; Perrino, F. W.; Lieberman, J.; Hübner, H. *J. Mol. Med.* **2007**, *85*, 531–537.

(76) Hogg, M.; Cooper, W.; Reha-Krantz, L.; Wallace, S. S. *Nucleic Acids Res.* **2006**, *34*, 2528–2535.

with Mg^{2+} or Mn^{2+} and free- ϵ 186 with Mg^{2+} . In all cases a water molecule bound to the catalytic metal acts as the nucleophile for hydrolysis of the nucleotide. This water is activated by the catalytic metal, which is found to be pentacoordinated. This coordination results in a polarization of the water molecule that promotes the required deprotonation. Initially a proton from this water is transferred to H162 followed by phosphate bond breaking. A second proton transfer to E14 is observed on the downward (second) leg of the reaction. A single transition state is obtained for all systems which corresponds to the phosphate bond-breaking step. This mechanism is consistent with a previously proposed scheme. Our results suggest that ϵ is slightly more active with Mn^{2+} than with Mg^{2+} , in qualitative agreement with experiment. A slightly lower activation barrier for ϵ 186-HOT compared with free- ϵ 186 is observed in our calculations. However, experiments with pNP-TMP point to free- ϵ having a slightly lower barrier.

To explain individual residue contributions to catalysis energy decomposition analysis was carried out using the minimized reactant and transition-state coordinates. This analysis suggests that there are several residues that can have a significant influence on the TS barrier. Among the residues highlighted from the energy decomposition analysis, H98, D103, D129, and D146 have been studied experimentally in ϵ and shown to influence catalysis. The remaining residues could be potentially interesting targets for mutagenesis experiments. Structural superposition with the related TREX1 and exonuclease domains of Klenow fragment and RB69 shows that several residues implicated in catalysis by energy decomposition analysis surrounding the active site are structurally conserved among these exonucleases. In particular, D103 and D129, which are strong mutators in ϵ when altered, are structurally conserved among ϵ , TREX1, and the exonuclease domains of Klenow and RB69.

Acknowledgment. This research was supported by the intramural research program of the NIH and NIEHS Z01 ES090601-11. L.G.P. acknowledges support from NIH HL-06350 and NSF FRG/DMR-0804549. Computing time from the NIEHS and UNC computing facilities are gratefully acknowledged. G.A.C. thanks Drs. J. M. Parks and H. Hu and Prof. W. Yang from Duke University and Prof. Y. Zhang from NYU.

Supporting Information Available: Detailed methods for initial structure, QM/MM structure optimization, and path optimization; metal coordination distances for critical points, free- ϵ structures, second-step QSM paths residue analysis for all residues and waters, structural alignments for Klenow and RB69 and complete citation for refs 41, 73, and 74 This material is available free of charge via the Internet at <http://pubs.acs.org>.

JA8082818