











Oxiranyl Anion Methodology Using Microflow Systems [*J. Am. Chem. Soc.* **2009**, *131*, 1654].
Aiiichiro Nagaki, Eiji Takizawa, and Jun-ichi Yoshida*

Page 1655. Two structures (**t-8** and **t-9**) in Table 2 should be corrected. The correct table is given below.

Table 2. Deprotonation of **8** Followed By Reactions With Electrophiles Using a Microflow System

epoxide	oxyranylithium	electrophile	yield (%) (c/t ratio)	productivity g/h
 c-8^a	 c-9	MeI	 c-10	 t-10
			96 (> 99 : 1) ^c	7.8
 t-8^b	 t-9	MeI	82 (2 : 98) ^c	3.3 ^e
c-8^a	c-9	Me ₃ SiCl	 c-11	 t-11
			97 (> 99 : 1) ^c	9.9
t-8^b	t-9	Me ₃ SiCl	79 (4 : 96) ^c	4.0 ^e
c-8^a	c-9	PhCOPh	 c-12	 t-12
			92 (> 99 : 1) ^d	13
t-8^b	t-9	PhCOPh	72 (4 : 96) ^d	5.1 ^e

JA9008979

10.1021/ja9008979

Published on Web 02/24/2009