

Asymmetric 1,4-Dihydroxylation of 1,3-Dienes by Catalytic Enantioselective Diboration


Heather E. Burks, Laura T. Kliman, and James P. Morken*

Department of Chemistry, Merkert Chemistry Center, Boston College, Chestnut Hill, Massachusetts 02467

Received December 10, 2008; E-mail: morken@bc.edu

Catalytic enantioselective oxidation reactions provide an important strategy for the conversion of simple unsaturated hydrocarbon substrates to synthetically useful organic compounds. While highly selective asymmetric oxidation of hydrocarbons may be accomplished in many ways, the 1,4-dihydroxylation of 1,3-dienes (Scheme 1), a synthetically valuable process, remains a largely unsolved problem. To accomplish this transformation, [4+2] cycloaddition of dienes with $^1\text{O}_2$ is often employed;¹ however, this reaction has not been accomplished under the influence of a chiral catalyst.² Likewise, the Pd(II)-catalyzed 1,4-diacetoxylation and related reactions,³ very promising transformations for asymmetric synthesis, have not yet been accomplished with useful levels of enantioselectivity.⁴ Our research program has focused on the enantioselective diboration of unsaturated hydrocarbons,⁵ and in this communication we describe an enantioselective 1,4-diboration of 1,3-dienes,^{6,7} a transformation that furnishes the above-described 1,4-dihydroxylation products in an economical fashion, with good yields, and with high levels of enantiomeric purity.⁸

Scheme 1



We recently developed a Pd-catalyzed enantioselective diboration of allenes, a process that benefits from accelerated catalysis when done in the presence of monodentate phosphine ligands. Detailed mechanistic studies suggest this reaction proceeds by oxidative addition of the diboron to palladium, a step which is followed by migratory insertion of the terminal alkene into a Pd–B bond (Scheme 2, eq 1).⁹ DFT studies revealed that the insertion step proceeds by an unusual elementary reaction that directly provides a stable, coordinatively saturated η^3 -allyl palladium complex **2** (see structure **1**; rotation of the terminal alkene occurs concomitantly with insertion, allowing π -bonding with the adjacent alkene to develop in the transition state). Considering that olefin insertion involving a 1,3-diene likely benefits from a similar incipient π bond (**3**), it was of interest to determine whether catalysts developed for the asymmetric allene diboration would be effective in catalytic diene diboration.

Scheme 2





Table 1. Catalytic Enantioselective Diboration/Oxidation of 1,3-Dienes^a




entry	diene	product	yield (%) ^b	ee (%) ^c
1			83	84
2			81	84
3			83	91
4			48	91
5			92	86 ^d
6			77	84(91) ^e
7			89	96 ^f
8			70	87 ^g
9			83	88
10			83	86 ^h
11		No Reaction	0	N/A

^a Unless otherwise indicated, (*R,R*)-**L1** was employed with reaction at 60 °C for 12 h, followed by oxidation with 30% H_2O_2 and 3 M NaOH for 3 h. ^b Percent yield of purified material. Value is an average of two experiments. ^c Determined by GC or SFC analysis employing a chiral stationary phase. ^d The product depicted corresponds to that obtained with (*S,S*)-**L2**. ^e Reaction at room temperature; value in parentheses is after a single recrystallization. ^f Reaction at room temperature. ^g 3 equiv of $\text{B}_2(\text{pin})_2$ employed. ^h (*S,S*) enantiomer of ligand **L1** employed for this experiment.

Initial experiments probed the reaction between bis(pinacolato)-diboron ($B_2(\text{pin})_2$) and 1,3-dienes. While no reaction was observed with a catalyst composed of $Pd_2(\text{dba})_3$ and a chiral TADDOL-derived phosphoramidite, the most effective catalyst for enantioselective allene diboration, when the palladium complex was replaced with $Pt_2(\text{dba})_3$, a complex that generally exhibits higher reactivity in diboration reactions,¹⁰ efficient catalysis and modest levels of enantioselectivity were observed. Several permutations of reaction conditions and ligand structure revealed that a catalyst composed of $Pt_2(\text{dba})_3$ and chiral TADDOL-derived phosphonite **L1**¹¹ provides good reactivity for a range of substrates and modest to high levels of enantioselection for many (Table 1). As can be observed in Table 1, acyclic dienes bearing aryl or alkyl substitution at the terminus react efficiently and with high selectivity. With ligand **L1**, butadienes bearing substitution at both C1 and C3 react with lower stereocontrol, however, with **L2** high selectivity is obtained (entry 5). Notably, cyclic dienes can be suitable substrates for the asymmetric diboration/oxidation sequence and provide otherwise difficult-to-access products with high levels of asymmetric induction. As noted in entry 10, the 1,2-diol is the predominant product when one terminus of the substrate is disubstituted, an outcome which likely results from enhanced steric congestion. Lastly, the lack of reaction with *cis*-piperylene (entry 11) suggests that only dienes able to adopt an *S-cis* conformation will participate in the Pt-catalyzed diene diboration. This observation may reflect the importance of structure **3** (Scheme 2) in the reaction mechanism.

In addition to 2-buten-1,4-diols, other important scaffolds can be easily prepared from simple dienes through the asymmetric diboration reaction. Enantiomerically enriched butenolides and derived butyrolactones are prominent structural elements in natural products, and straightforward methods for their preparation are scarce.¹² Diene diboration provides a new approach: subsequent to diboration and oxidation, the unpurified material was subjected to perruthenate-catalyzed oxidation¹³ and furnished the derived butenolide in good yield and without compromising the integrity of the carbinol stereo-center.

Scheme 3




The α -chiral allylboronate functionality embedded in diene diboration products also finds use in stereoselective carbonyl allylation.¹⁴ Critical concerns are whether this transformation proceeds with high chirality transfer and whether it is selective for one constitutional isomer. To address this, benzaldehyde was added to an unquenched diboration reaction. After 12 h of reaction and oxidative workup, this sequence provided a single diastereomer of a single constitutional isomer, with near-perfect chirality transfer. The product structure suggests that this transformation proceeds through chair-like transition state **A** that minimizes A(1,3) interactions and with C=C bond formation occurring at the least hindered carbon of the intermediate diboron.

In conclusion, we have described a catalytic enantioselective diboration of 1,3-dienes, a process which generally provides synthetically useful chiral 2-buten-1,4-diols as the reaction product. Further studies of the substrate scope and reaction utility are in progress and will be reported in the near future.

Acknowledgment. Support by the NIGMS (GM-59417) and Merck is gratefully acknowledged, as is the NSF (DBI-0619576)

Scheme 4



for support of the BC Mass Spectrometry Center. We also thank AllyChem, Co., Ltd. for a generous donation of $B_2(\text{pin})_2$.

Note Added after ASAP Publication. The uncorrected proof version was published June 8, 2009. Minor text corrections have been made in the version published on June 12, 2009.

Supporting Information Available: Characterization and procedures. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) For recent applications of the singlet oxygen [4+2] cycloaddition in synthesis, see: (a) Lee, J. S.; Fuchs, P. L. *J. Am. Chem. Soc.* **2005**, *127*, 13122. (b) Barbarow, J. E.; Miller, A. K.; Trauner, D. *Org. Lett.* **2005**, *7*, 2901. (c) Ono, K.; Nakagawa, M.; Nishida, A. *Angew. Chem., Int. Ed.* **2004**, *43*, 2020. (d) Kusama, H.; Hara, R.; Kawahara, S.; Nishimori, T.; Kashima, H.; Nakamura, N.; Morihira, K.; Kuwajima, I. *J. Am. Chem. Soc.* **2000**, *122*, 3811. (e) Zhou, G.; Gao, X.; Li, W. Z.; Li, Y. *Tetrahedron Lett.* **2001**, *42*, 3101. (f) Izzo, I.; Meduri, G.; Avallone, E.; De Riccardis, F.; Sodano, G. *Eur. J. Org. Chem.* **2000**, 439. (g) For recent reviews, see: (h) Clennan, E. L.; Pace, A. *Tetrahedron* **2005**, *61*, 6665.
- (2) For the enantioselective desymmetrization of $^{18}O_2$ cycloadducts, see: Staben, S. T.; Xin, L.; Toste, F. D. *J. Am. Chem. Soc.* **2006**, *128*, 12658.
- (3) For a recent review of this process: Bäckvall, J.-E. *Palladium-Catalyzed 1,4-Additions to Conjugated Dienes*. In *Metal Catalyzed Cross-Coupling Reactions*, 2nd ed.; De Meijere, A., Diederich, F., Eds.; Wiley-VCH: Weinheim, 2004; pp 479–529.
- (4) (a) Thorarensen, A.; Palmgren, A.; Itami, K.; Bäckvall, J.-E. *Tetrahedron Lett.* **1997**, *38*, 8541. (b) Itami, K.; Palmgren, A.; Thorarensen, A.; Bäckvall, J.-E. *J. Org. Chem.* **1998**, *63*, 6466. (c) Verboom, R. C.; Plietker, B. J.; Bäckvall, J.-E. *J. Organomet. Chem.* **2003**, *687*, 508. (d) El-Qisairi, A. K.; Qaseer, H. A.; Fennelly, J. P.; Chiarelli, M. P.; Henry, P. M. *Catal. Commun.* **2008**, *9*, 1661.
- (5) Recent reviews: (a) Burks, H. E.; Morken, J. P. *Chem. Commun.* **2007**, 4717. (b) Ramirez, J.; Lillo, V.; Segarra, A. M.; Fernandez, E. *Comp. Rend. Chim.* **2007**, *10*, 138. (c) Beletskaya, I.; Moberg, C. *Chem. Rev.* **2006**, *106*, 2320. (d) Ishiyama, T.; Miyaura, N. *Chem. Record* **2004**, *3*, 271.
- (6) Non-enantioselective diene diboration: (a) Ishiyama, T.; Yamamoto, M.; Miyaura, N. *Chem. Commun.* **1996**, 2073. (b) Ishiyama, T.; Yamamoto, M.; Miyaura, N. *Chem. Commun.* **1997**, 689. (c) Clegg, W.; Thorsten, J.; Marder, T. B.; Norman, N. C.; Orpen, A. G.; Peakman, T. M.; Quayle, M. J.; Rice, C. R.; Scott, A. J. *J. Chem. Soc., Dalton Trans.* **1998**, 1431. (d) Morgan, J. B.; Morken, J. P. *Org. Lett.* **2003**, *5*, 2573.
- (7) For asymmetric silylboration of dienes: (a) Gerdin, M.; Moberg, C. *Adv. Synth. Catal.* **2005**, *347*, 749. For diastereoselective diene diboration using chiral diboron reagents, see: ref 6c.
- (8) $B_2(\text{pin})_2$ is a readily available inexpensive reagent (\$800/kg, AllyChem, Co., Ltd. (www.allychem.com)).
- (9) (a) Burks, H. E.; Liu, S.; Morken, J. P. *J. Am. Chem. Soc.* **2007**, *129*, 8766. (b) Pelz, N. F.; Woodward, A. R.; Burks, H. E.; Sieber, J. D.; Morken, J. P. *J. Am. Chem. Soc.* **2004**, *126*, 16328.
- (10) (a) Ishiyama, T.; Matusuda, N.; Miyaura, N.; Suzuki, A. *J. Am. Chem. Soc.* **1993**, *115*, 11018. (b) Ishiyama, T.; Matsuda, N.; Murata, M.; Ozawa, F.; Suzuki, A.; Miyaura, N. *Organometallics* **1996**, *15*, 713. For computational studies comparing Pd and Pt in diboration, see: (c) Cui, Q.; Musaeiev, D. G.; Morokuma, K. *Organometallics* **1998**, *17*, 742. (d) Sakaki, S.; Kikuno, T. *Inorg. Chem.* **1997**, *36*, 226.
- (11) (a) Sakaki, S.; Schweizer, W. B.; Seebach, D. *Helv. Chem. Acta* **1993**, *76*, 2654. (b) Seebach, D.; Hayakawa, M.; Sakaki, S.; Schweizer, W. B. *Tetrahedron* **1993**, *49*, 1711. (c) Alexakis, A.; Burton, J.; Vastra, J.; Benhaim, C.; Fournioux, X.; van den Heuvel, A.; Leveque, J.-M.; Maze, F.; Rosset, S. *Eur. J. Org. Chem.* **2000**, 4011. (d) Bee, C.; Han, S. B.; Hassan, A.; Iida, H.; Krische, M. J. *J. Am. Chem. Soc.* **2008**, *130*, 2746.
- (12) Seitz, M.; Reiser, O. *Curr. Opin. Chem. Biol.* **2005**, *9*, 285.
- (13) Ley, S. V.; Norman, J.; Griffith, W. P.; Marsden, S. P. *Synthesis* **1994**, 639.
- (14) (a) Hoffmann, R. W. *Pure Appl. Chem.* **1988**, *60*, 123. (b) Hoffmann, R. W.; Neil, G.; Schlapbach, A. *Pure Appl. Chem.* **1990**, *62*, 1993.

JA809610H