

Direct Detection of the Oxygen Rebound Intermediates, Ferryl Mb and NO₂, in the Reaction of metMyoglobin with Peroxynitrite

Jia Su and John T. Groves*

Department of Chemistry, Princeton University, Princeton, New Jersey 08544

Received March 27, 2009; E-mail: jtgroves@princeton.edu

Abstract: Oxygenated hemoproteins are known to react rapidly with nitric oxide (NO) to produce peroxynitrite (PN) at the heme site. This process could lead either to attenuation of the effects of NO or to nitrosative protein damage. Peroxynitrite is a powerful nitrating and oxidizing agent that has been implicated in a variety of cell injuries. Accordingly, it is important to delineate the nature and variety of reaction mechanisms of PN reactions with heme proteins. Here, we present direct evidence that ferrylMb and NO₂ are both produced during the reaction of PN and metmyoglobin (metMb). Kinetic evidence indicates that these products evolve from initial formation of a caged radical intermediate [Fe^{IV}=O ·NO₂]. This caged pair reacts mainly via internal return with a rate constant *k_r* to form metMb and nitrate in an oxygen rebound scenario. Detectable amounts of ferrylMb are observed by stopped-flow spectrophotometry, appearing at a rate consistent with the rate, *k_{obs}*, of heme-mediated PN decomposition. Freely diffusing NO₂, which is liberated concomitantly from the radical pair (*k_e*), preferentially nitrates Tyr103 in horse heart myoglobin. The ratio of the rates of in-cage rebound and cage escape, *k_r/k_e*, was found to be ~10 by examining the nitration yields of fluorescein, an external NO₂ trap. This rebound/escape model for the metMb/PN interaction is analogous to the behavior of alkyl hyponitrites and the well-studied geminate recombination processes of deoxymyoglobin with O₂, CO, and NO. The scenario is also similar to the stepwise events of substrate hydroxylation by cytochrome P450 and other oxygenases. It is likely, therefore, that the reaction of metMb with ONOO⁻ and that of oxyMb with NO proceed through the same [Fe^{IV}=O ·NO₂] caged radical intermediate and lead to similar outcomes. The results indicate that while oxyMb may reduce the concentration of intracellular NO, it would not eliminate the formation of NO₂ as a decomposition product of peroxynitrite.

Introduction

Although myoglobin (Mb) is a textbook paradigm for mediating oxygen storage and delivery in vertebrate skeletal and cardiac muscle tissue,¹ important alternative roles as an intracellular nitric oxide (NO) dioxygenase^{2,3} and a nitrite reductase^{4,5} have been recognized only recently. These discoveries have offered many new insights regarding this “old” protein. Nitric oxide (NO) is a central signaling device for both intra- and intercellular communication.⁶ Considering the large amounts of oxyMb in normoxic muscle cells and the compartmentalization of NO production, the scavenging and degradation of NO by oxyMb is probably an important pathway to attenuate the spatial range of these signals. Conversely, the Mb reaction

with nitrite to produce NO may be important under hypoxic conditions.^{4,7}

Peroxynitrite (PN) evolves in vivo from the very fast combination of superoxide ion (O₂^{•-}) and NO, as a consequence of aerobic metabolism and overproduction of NO.^{8,9} Because both peroxynitrite anion (ONOO⁻) and peroxynitrous acid (HOONO) can diffuse readily across biomembranes,^{10,11} PN exhibits a wide range of reactivity in the modifications of many cellular targets, through one- and two-electron oxidations, nitrosation, and nitration of various protein residues.^{12,13} Heme proteins, such as myoglobin,^{14,15} hemoglobin,^{16,17} peroxidases,^{18–20} cytochrome *c*,^{21,22} nitric oxide synthase,²³ cytochrome P450,²⁴


- (1) Wittenberg, B. A.; Wittenberg, J. B. *Annu. Rev. Physiol.* **1989**, *51*, 857–78.
- (2) Brunori, M. *Trends Biochem. Sci.* **2001**, *26*, 209–210.
- (3) Frauenfelder, H.; McMahon, B. H.; Austin, R. H.; Chu, K.; Groves, J. T. *Proc. Natl. Acad. Sci. U.S.A.* **2001**, *98*, 2370–4.
- (4) Hendgen-Cotta, U. B.; Merx, M. W.; Shiva, S.; Schmitz, J.; Becher, S.; Klare, J. P.; Steinhoff, H. J.; Goedecke, A.; Schrader, J.; Gladwin, M. T.; Kelm, M.; Rassaf, T. *Proc. Natl. Acad. Sci. U.S.A.* **2008**, *105*, 10256–10261.
- (5) Cossins, A.; Berenbrink, M. *Nature* **2008**, *454*, 416–417.
- (6) Pacher, P. A.; Beckman, J. S.; Liaudet, L. *Physiol. Rev.* **2007**, *87*, 315–424.

- (7) Fraser, J.; de Mello, L. V.; Ward, D.; Rees, H. H.; Williams, D. R.; Fang, Y. C.; Brass, A.; Gracey, A. Y.; Cossins, A. R. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 2977–2981.
- (8) Beckman, J. S.; Beckman, T. W.; Chen, J.; Marshall, P. A.; Freeman, B. A. *Proc. Natl. Acad. Sci. U.S.A.* **1990**, *87*, 1620–1624.
- (9) Goldstein, S.; Merenyi, G. *Methods Enzymol.* **2008**, *436*, 49–61.
- (10) Marla, S. S.; Lee, J.; Groves, J. T. *Proc. Natl. Acad. Sci. U.S.A.* **1997**, *94*, 14243–8.
- (11) Denicola, A.; Souza, J. M.; Radi, R. *Proc. Natl. Acad. Sci. U.S.A.* **1998**, *95*, 3566–71.
- (12) Szabo, C.; Ischiropoulos, H.; Radi, R. *Nat. Rev. Drug Discovery* **2007**, *6*, 662–680.
- (13) Groves, J. T. *Curr. Opin. Chem. Biol.* **1999**, *3*, 226–35.
- (14) Bourassa, J. L.; Ives, E. P.; Marqueling, A. L.; Shimanovich, R.; Groves, J. T. *J. Am. Chem. Soc.* **2001**, *123*, 5142–3.

and the newly discovered neuroglobins,²⁵ have all been observed to interact with PN and mediate its decomposition.

The reaction of PN with ferric myoglobin (metMb) and the concomitant nitration of protein tyrosine have been reported.^{14,26,27} However, there is no consensus regarding the mechanism of the metMb/PN interaction and, especially, the identification of intermediates formed in the course of the reaction. Herold et al. have provided evidence that the reaction between oxyMb with NO leads to the formation of an intermediate MbFe^{III}-OONO adduct, which ultimately affords metMb and NO₃⁻,^{28,29} although the MbFe^{III}-nitrate complex has been shown to exhibit similar kinetics.³⁰ We have suggested a mechanism for the PN-metMb reaction that involves a protein-caged ferryl intermediate [Fe^{IV}=O •NO₂] deriving from the facile O-O bond homolysis of MbFe^{III}-O-ONO (Scheme 1).¹⁴ The ferryl intermediate has not been unequivocally identified during the oxyMb-NO reaction, although there is suggestive EPR evidence for this process.^{31,32} Reactions of PN with water-soluble Fe^{III} and Mn^{III} porphyrins have been reported by us³³⁻³⁷ and by the Monsanto group.³⁸ Both Fe^{III}TMPS and Fe^{III}TMPyP react rapidly with ONOO⁻ to produce ferryl intermediates.^{34,35} Likewise, the reactions of PN with CYP 119³⁹ and myeloperoxidase²⁰ also lead to buildup of the ferryl species.

Scheme 1



- (15) Herold, S.; Matsui, T.; Watanabe, Y. *J. Am. Chem. Soc.* **2001**, *123*, 4085–6.
- (16) Herold, S.; Shivashankar, K. *Biochemistry* **2003**, *42*, 14036–46.
- (17) Pietraforte, D.; Salzano, A. M.; Scorza, G.; Marino, G.; Minetti, M. *Biochemistry* **2001**, *40*, 15300–9.
- (18) Gebicka, L.; Didik, J. *J. Inorg. Biochem.* **2007**, *101*, 159–164.
- (19) Floris, R.; Piersma, S. R.; Yang, G.; Jones, P.; Wever, R. *Eur. J. Biochem.* **1993**, *215*, 767–775.
- (20) Furtmuller, P. G.; Jantschko, W.; Zederbauer, M.; Schwanninger, M.; Jakopitsch, C.; Herold, S.; Koppenol, W. H.; Obinger, C. *Biochem. Biophys. Res. Commun.* **2005**, *337*, 944–954.
- (21) Bathyany, C.; Souza, J. M.; Duran, R.; Cassina, A.; Cervenansky, C.; Radi, R. *Biochemistry* **2005**, *44*, 8038–8046.
- (22) Cassina, A. M.; Hodara, R.; Souza, J. M.; Thomson, L.; Castro, L.; Ischiropoulos, H.; Freeman, B. A.; Radi, R. *J. Biol. Chem.* **2000**, *275*, 21409–21415.
- (23) Marechal, A.; Mattioli, T. A.; Stuehr, D. J.; Santolini, J. *J. Biol. Chem.* **2007**, *282*, 14101–14112.
- (24) Daiber, A.; Herold, S.; Schoneich, C.; Namgaladze, D.; Peterson, J. A.; Ullrich, V. *Eur. J. Biochem.* **2000**, *267*, 6729–6739.
- (25) Herold, S.; Fago, A.; Weber, R. E.; Dewilde, S.; Moens, L. *J. Biol. Chem.* **2004**, *279*, 22841–22847.
- (26) Herold, S.; Kalinga, S.; Matsui, T.; Watanabe, Y. *J. Am. Chem. Soc.* **2004**, *126*, 6945–55.
- (27) Herold, S.; Shivashankar, K.; Mehl, M. *Biochemistry* **2002**, *41*, 13460–72.
- (28) Herold, S.; Exner, M.; Nauser, T. *Biochemistry* **2001**, *40*, 3385–95.
- (29) Doyle, M. P.; Hoekstra, J. W. *J. Inorg. Biochem.* **1981**, *14*, 351–358.
- (30) Goldstein, S.; Merenyi, G.; Samuni, A. *J. Am. Chem. Soc.* **2004**, *126*, 15694–701.
- (31) Olson, J. S.; Foley, E. W.; Rogge, C.; Tsai, A. L.; Doyle, M. P.; Lemon, D. D. *Free Radical Biol. Med.* **2004**, *36*, 685–697.
- (32) Gardner, P. R.; Gardner, A. M.; Brashear, W. T.; Suzuki, T.; Hvitved, A. N.; Setchell, K. D. R.; Olson, J. S. *J. Inorg. Biochem.* **2006**, *100*, 542–550.
- (33) Groves, J. T.; Marla, S. S. *J. Am. Chem. Soc.* **1995**, *117*, 9578–9579.
- (34) Shimanovich, R.; Groves, J. T. *Arch. Biochem. Biophys.* **2001**, *387*, 307–17.
- (35) Lee, J. B.; Hunt, J. A.; Groves, J. T. *J. Am. Chem. Soc.* **1998**, *120*, 7493–7501.
- (36) Lee, J. B.; Hunt, J. A.; Groves, J. T. *J. Am. Chem. Soc.* **1998**, *120*, 6053–6061.
- (37) Shimanovich, R.; Hannah, S.; Lynch, V.; Gerasimchuk, N.; Mody, T. D.; Magda, D.; Sessler, J.; Groves, J. T. *J. Am. Chem. Soc.* **2001**, *123*, 3613–4.
- (38) Stern, M. K.; Jensen, M. P.; Kramer, K. *J. Am. Chem. Soc.* **1996**, *118*, 8735–8736.
- (39) Sheng, X.; Horner, J. H.; Newcomb, M. *J. Am. Chem. Soc.* **2008**, *130*, 13310–13320.

In the current work, we have used kinetic and spectroscopic techniques to interrogate the interaction between metMb and PN, building on our previously reported studies.¹⁴ The results show that metMb reacts rapidly with PN to form the ferryl species, MbFe^{IV}=O, which has been spectroscopically identified and kinetically characterized. Further, we employed fluorescein as a trap of freely diffusing NO₂ and nano-LC-ESI-MS/MS to confirm that the specific nitration site is Tyr103. Combined with the measured rate constant of ferrylMb formation in the course of PN decay, the results support a mechanism involving a [MbFe^{IV}=O •NO₂] radical pair that dissociates to free MbFe^{IV}=O and •NO₂ or collapses to MbFe^{III} and nitrate ion, in a scenario that is analogous to myoglobin-ligand (NO, CO, and O₂) geminate recombination and cage escape.^{40,41}

Results

Direct Detection of FerrylMb(I) Formed during the Reaction of metMb with PN. PN decomposition was examined by studying the absorbance decay of PN anion at 302 nm by stopped-flow spectrophotometry as we have previously described.¹⁴ The inventory of heme species formed during the reaction of metMb with PN was monitored in the stopped-flow device over the course of PN decomposition. Figure 1A shows the time evolution of optical difference spectra for the Soret region upon reacting 5 μM Fe^{III}Mb with 25 μM PN at 12 °C. A spectral transient of an intermediate (I) was observed to arise within 3 s with increased absorbance at 422 nm, followed by its slower decay to Fe^{III}Mb (λ_{max} = 409 nm) over the following 25 s (Figure 1A, inset). The absorbance at 409 nm showed a concomitant decrease in the absorbance in the first 3 s followed by a modest increase (Figure 1A, inset). An isosbestic point at 417 nm was maintained over the course of the reaction. Difference spectra were also obtained for the characteristic Q-band region by examining the reaction between 50 μM

- (40) Austin, R. H.; Beeson, K. W.; Eisenstein, L.; Frauenfelder, H.; Gunsalus, I. C. *Biochemistry* **1975**, *14*, 5355–5373.
- (41) Jongeward, K. A.; Magde, D.; Taube, D. J.; Marsters, J. C.; Traylor, T. G.; Sharma, V. S. *J. Am. Chem. Soc.* **1988**, *110*, 380–387.




Figure 1. Formation of intermediate **I** from metMb and PN reaction. (A) Soret absorption difference spectra obtained from the reaction of 5 μM metMb and 25 μM PN at pH 7.4 and 12 $^{\circ}\text{C}$. The transients were obtained by subtracting the mixture spectrum at 0.02 s from those at 0.2, 0.5, 1, 2, and 3 s. Inset: Time courses of the reaction at 422 and 409 nm under the above conditions. (B) Q-band absorption difference spectra obtained from the reaction of 50 μM metMb and 1.5 mM PN at pH 7.4 and 25 $^{\circ}\text{C}$. The difference spectra were obtained by subtracting the mixture spectrum at 0.01 s from those at 0.03, 0.06, 0.12, 0.2, and 0.3 s.

metMb and 1.5 mM PN at 25 $^{\circ}\text{C}$. As shown in Figure 1B, distinct absorbance changes were observed at 503, \sim 570, and \sim 590 nm. All of these peaks and valleys in the difference spectra corresponded well with the wavelengths for which metMb and ferrylMb have the largest difference in extinction coefficients. Furthermore, the observed isosbestic points in the difference spectra, including 417 and 524 nm, were also seen when comparing the absorbance spectra of authentic ferrylMb with metMb.⁴² Control experiments were performed accordingly with metMb and decomposed PN (that still contained small amounts of H₂O₂) under the above conditions. No rapidly formed intermediate was observed.

To confirm the identity of the intermediate **I**, double-mixing stopped-flow experiments were carried out by first mixing 20 μM Mb and 100 μM PN for 3 s followed by addition of ascorbate under pseudo-first-order conditions. The decay rate of **I** was found to accelerate from 0.093 to 0.132 s⁻¹ with the addition of 9 mM ascorbate as shown in Figure 2. In separate experiments at 12 $^{\circ}\text{C}$ and pH 7.3, ferrylMb was prepared independently by the reaction of metMb (5 μM) and H₂O₂ (100 μM) followed by adding catalase to quench the excess H₂O₂. The reaction rate of authentic ferrylMb with ascorbate was measured to be 4.0 M⁻¹ s⁻¹ (Figure S1B), consistent both with the reductive trapping of the metMb–PN intermediate described above (4.2 M⁻¹ s⁻¹, Figure S1A) and with the literature value, 2.7 \pm 0.8 M⁻¹ s⁻¹.⁴³ Hence, intermediate **I** formed during




Figure 2. Decay of intermediate **I** in the absence and presence of 9 mM ascorbate monitored at 422 nm. The experiments were carried out in a double-mixing stopped-flow spectrometer by first mixing 20 μM Mb and 100 μM PN for 3 s followed by addition of buffer (black trace) or ascorbate solution (red trace) under pseudo-first-order conditions at pH 7.4 and 12 $^{\circ}\text{C}$. First exponential decay equations were used to fit the experimental curves. The values of k_{obs} from those two fittings (green) are 0.093 s⁻¹ (black) and 0.132 s⁻¹ (red), respectively.

metMb/PN interaction exhibits the same kinetic behavior in reacting with electron donor ascorbate as that of an authentic ferrylMb.

The decay of Mb intermediate **I** back to metMb was studied by following the absorbance changes at 422 nm. The lifetime of **I** at 12 $^{\circ}\text{C}$ (30 s, Figure 2) is distinctly longer than that of PN (20 s). This behavior indicates that the decay of **I** is mainly (75% ferryl was reduced during 20 s) dependent on its reduction by PN, as well as several other pathways, such as reduction by tyrosine,⁴² NO₂,³⁰ nitrite, and NO.⁴⁴ To clarify the kinetics of this main reduction pathway, ferrylMb was prepared fresh in one syringe by treatment of metMb with H₂O₂ followed by catalase quenching and then rapidly mixed with PN in the other syringe. A first-order treatment in terms of PN could not be applied for the entire process because the concentration of PN was not sufficiently constant at the end of the reaction. Fitting the first 0.5 s decay of ferrylMb at 422 nm to a linear function, the initial rates (v_0) of this bimolecular reaction with different concentrations of PN could be determined. The slope of v_0 versus PN concentration gave a value of $(1.53 \pm 0.06) \times 10^4$ M⁻¹ s⁻¹ at 25 $^{\circ}\text{C}$ (Figure S2A), in good agreement with an earlier report by Herold et al.⁴⁵ The rate constant of ferrylMb reduction by PN was measured to be $(3.8 \pm 0.3) \times 10^3$ M⁻¹ s⁻¹ at 12 $^{\circ}\text{C}$ by the same method (Figure S2B). Likewise, the rate constant of nitrite reduction of authentic ferrylMb was measured to be 30 M⁻¹ s⁻¹ at 25 $^{\circ}\text{C}$.

MetMb Tyrosine Is Nitrated by Peroxynitrite. To achieve an efficient and reproducible mixing of PN with Mb, we used the stopped-flow spectrometer as a mixing device to determine the nitration extents of metMb and cyanide-bound metMb (metMbcN) by PN. We quantified the yields of nitrated metMb by electrospray mass spectrometry (ESI-MS). MetMb, metMbcN, and nitrated myoglobin (nitroMb) were found to have equal ionization efficiency by examining standard mixtures. Direct MS detection of the entire reaction mixture avoids errors associated with enzymatic digestion and sample handling when analyzing for the extent of nitration. Reaction of 100 μM metMb with 10 and 30 equiv (1 and 3 mM) of PN resulted in 8 \pm 1 and 10 \pm 1 μM mononitrated Mb at 25 $^{\circ}\text{C}$ and pH 7.4,

(42) Lardinois, O. M.; Ortiz de Montellano, P. R. *Biochemistry* **2004**, *43*, 4601–4610.

(43) KrogerOhlsen, M.; Skibsted, L. H. *J. Agric. Food Chem.* **1997**, *45*, 668–676.

(44) Herold, S.; Rehmman, F. J. *J. Biol. Inorg. Chem.* **2001**, *6*, 543–55.

(45) Exner, M.; Herold, S. *Chem. Res. Toxicol.* **2000**, *13*, 287–93.




Figure 3. Nitration of fluorescein by PN. (A) Interaction of FI and PN produces nitroFI. (B) Stopped-flow rapid-scan UV-vis spectra of the reaction between 15 μM FI and 800 μM PN in 0.1 M phosphate buffer at pH 7.4 and 25 $^{\circ}\text{C}$. Traces shown were taken at 0, 2, 5, 10 and 15 s. The directions of spectral changes are indicated by arrows. (C) The traces exhibit PN decomposition (302 nm, solid line) and, concurrently, nitroFI formation (520 nm, dashed line) observed by stopped-flow spectrophotometry upon mixing equal volumes of 50 μM fluorescein with 920 μM peroxyinitrite at pH 7.4 and 25 $^{\circ}\text{C}$.

respectively, as determined by inspection of the mass spectral data (Figure S3A). The small amounts of Mb nitration observed with decomposed PN solutions were subtracted from the yields observed with active PN (H_2O_2 was <5% relative to PN concentration). The yields of nitroMb in heme-blocked metMbCN with 1 and 3 mM PN were 21 ± 1 and 27 ± 1 μM , respectively. NitroMb was purified by anion exchange HPLC followed by trypsin or Glu-C endoproteinase digestion. The specific nitration site of the nitroMb was identified by LC-MS/MS. In the MS/MS spectra, both Y*LEFISDAIHVLHSK from trypsin digestion (Figure S3B) and LKPLAQSHATKHKIPIKY*LE from Glu-C digestion displayed a 45 Da molecular weight increase due to the modification of tyrosine 103, whereas tyrosine 146 was unchanged. These results unambiguously identify Tyr103 as the preferential site of nitration in the PN-mediated horse heart metMb nitration.

Fluorescein Traps NO_2 Evolved from metMb/PN Interactions. Fluorescein (FI) reacts with PN to form two nitrated isomers, 2'- and 4'-nitrofluorescein (FINO₂), the structures of which were confirmed by mass spectrometry and ¹H NMR.^{46,47} The reaction can be followed conveniently by monitoring the shift of λ_{max} from 490 to 494 nm and the significant absorbance increase at 520 nm ($\epsilon = 1.13 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$) (Figure 3). As FI has negligible absorbance at 520 nm, the total nitration yields of FI could be determined from the absorbance increase at 520 nm. When mixing 50 μM metMb with 0–50 μM FI solutions followed by an addition of 30 equiv (1.5 mM) PN at 25 $^{\circ}\text{C}$, we found that 25 μM FI entirely suppressed the nitration of 50 μM metMb by 1.5 mM PN.

The mechanism of the metMb/PN interaction was further examined by stopped-flow rapid mixing of metMb, FI, and PN solutions. The experiments were performed by varying the concentration of metMb while keeping the concentrations of




Figure 4. (A) Overlay of the experimental and simulated time courses of nitroFI formation in the presence and absence of metMb. The parameters were set as shown in eqs 1–4 and Scheme S1. Experiments (E) and simulations (S) were done under the same reaction conditions: 30 μM FI, 520 μM PN, 0, 20 and 50 μM metMb. From top to bottom: 0 μM metMb (a, E in black and S in blue), 20 μM metMb (b, E in black and S in red), 50 μM metMb (c, E in black and S in green), 20 μM metMb with $k_c = 0 \text{ s}^{-1}$ (d, S in dashed line and red), and 50 μM metMb with $k_c = 0 \text{ s}^{-1}$ (e, S in dashed line and blue). (B) The formation and decay of intermediate I during the reaction of 10 μM (final concentration) metMb and 100 μM PN in the absence (black) and presence (red) of 10 μM FI at pH 7.4 and 12 $^{\circ}\text{C}$.





FI and PN constant. Figure 4A shows a representative set of time profiles of PN-mediated FINO₂ formation in the presence of metMb (0, 20, and 50 μM). Control experiments with metMbCN carried out under the same conditions at pH 7.6 showed very little variation in nitroFI yields (data not shown). As seen in Figure 4A (traces a–c in black), addition of 20 and 50 μM metMb to FI reduced the yields of FINO₂ from 26.4 μM to 13.6 and 10.2 μM , respectively. Importantly, the initial FINO₂ formation rates (0–2.5 s) were found to increase with added metMb (Figure 4A). Furthermore, the time courses for the formation and decay of the intermediate I during the metMb/PN reaction in the presence and absence of FI were measured at 426 nm at pH 7.4 and 12 $^{\circ}\text{C}$ (Figure 4B). With 10 μM FI in the medium, the maximum concentration of I was noticeably increased.

Numerical Simulation Validates the Radical Rebound Mechanism in the metMb/PN Interaction. Kinetic simulations of intermediate I formation and PN-mediated FI nitration in the presence and absence of metMb were performed with Berkeley Madonna Modeling and Analysis software. Kinetic modeling was initiated by inputting the reaction equations (eqs 1–4 and measured or literature rate constants as shown in Scheme S1). The simulations yielded individual time courses for the concentrations of ferrylMb, NO₂, PN, and nitroFI under different conditions. From the simplest model of PN-mediated FI nitration in the absence of metMb, which is the same as the mechanism of phenol nitration⁹ (Scheme S1), we conclude that PN

(46) Shimanovich, R. Ph.D. Thesis, Princeton University, 2003.

(47) Jiang, Q.; Hurst, J. K. *J. Biol. Chem.* **1997**, *272*, 32767–72.

spontaneous decay produced about 10% NO₂. For the simulations of FI nitration in the presence of various metMb concentrations (Figure 4A), the competition between tyrosine and FI was explicitly considered (eqs 5 and 7 in Scheme S1). As FI proved to be a better trap of NO₂ than the tyrosines in the metMb protein by at least 10-fold, we set the values of tyrosine reactions at 1 order of magnitude smaller than those for FI radical reactions ($k_5 > 10k_7$). The four core equations as shown below were used to model the effect of metMb on NO₂/FINO₂ formation.



The value of k_e^{-1} ($1.2 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$), representing second-order NO₂ rebinding from solvent, was adopted from the measurements of Goldstein et al.,³⁰ k_3 ($(1.53 \pm 0.06) \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$) was measured by experiment as described above. Because k_e and k_r must be much larger than k_{cat} (the metMb-PN turnover rate: $1.03 \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$ at 25 °C),¹⁴ and k_3 represents only the reduction of cage-escaped ferrylMb (~10% of total Mb), the rate-determining step of this catalytic cycle is the formation of $[\text{Fe}^{\text{IV}}=\text{O} \cdot \text{NO}_2]$. Hence, the rate constant of the first step ($[\text{Fe}^{\text{IV}}=\text{O} \cdot \text{NO}_2]$ formation via eq 1) is equal to k_{cat} . We found by kinetic simulation that it is the ratio of k_e and k_r that determines the yields of ferrylMb as well as the rates of its generation. The time courses of intermediate ferrylMb formation at 25 °C (Figure 5A) were fit well with $k_e/k_r = \sim 0.1$ using the above equations. The observed yields of ferrylMb computed from the known extinction coefficients were ~3% of the total Mb with 240 μM PN and ~4% of the total Mb with 450 μM PN. As seen in Figure 5A, these yields can also be estimated well by the simulation model described as above.

The rate of intermediate I (i.e., ferrylMb) formation as a function of initial PN concentration at constant metMb concentration at pH 7.4 and 25 °C was fit to eq 5 (Figure 5B). With $k_r/k_e = 15$, the bimolecular rate constant of metMb/PN reaction can be deduced to be $1.0 \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$. This value matched our measured k_{cat} under these conditions.

$$\frac{d[\text{Fe}^{\text{IV}}\text{O}]_0}{dt} = \frac{k_e k_{\text{cat}} [\text{Fe}^{\text{III}}][\text{PN}]_0}{k_e + k_r} \quad (5)$$

The data for reactions between 520 μM PN and 0, 20, 50 μM metMb in the presence of 30 μM FI as described above were nicely simulated on the basis of eqs 1–4 with ~5% trappable NO₂ (Figure 4A, traces a–c in blue, red, and green). When simulating the metMbCN-mediated reactions under the same conditions, good fits were achieved by setting k_{cat} to zero, while all other reactions were retained. As shown in Figure 4A, if the cage escape of NO₂ from $[\text{ferrylMb} \cdot \text{NO}_2]$ was not considered ($k_e = 0$) in the simulation, the simulated time courses of FINO₂ formation in the presence of 20 and 50 μM metMb




Figure 5. (A) Experimental and simulated time courses of intermediate ferrylMb formation at pH 7.4 and 25 °C. Reaction conditions: 8.4 μM metMb with 450 μM PN (experimental trace in blue, simulated in black) or with 240 μM PN (experimental trace in red, simulated in black). The parameters used are: $k_{\text{cat}} = 1.0 \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$, $k_3 = 1.53 \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$, and $k_e/k_r = 1/10$. Tyrosine nitration was not considered in this set of simulations. (B) Plot of intermediate I initial formation rates observed at 422 nm vs initial PN concentrations. The reactions were conducted by mixing 8.4 μM metMb with various concentrations of PN in the stopped-flow spectrophotometer at pH 7.4 and 25 °C.

deviated widely from the experimental results (Figure 4A, traces d and e, respectively).

Discussion

The results presented here provide the first direct evidence that the metmyoglobin–peroxynitrite interaction yields ferrylMb ($\text{MbFe}^{\text{IV}}=\text{O}$) concurrently with the formation of freely diffusing $\cdot\text{NO}_2$. The data indicate a mechanism that involves rate-limiting O–O bond homolysis in a peroxynitrito– Fe^{III} –heme species ($\text{MbFe}^{\text{III}}-\text{O}-\text{ONO}$) to afford a caged $[\text{Fe}^{\text{IV}}=\text{O} \cdot \text{NO}_2]$ intermediate (Scheme 1), as we had suggested¹⁴ and in analogy to the process we had earlier delineated for synthetic iron and manganese porphyrins.^{34–37} The major pathway to close the catalytic cycle is cage collapse of $[\text{Fe}^{\text{IV}}=\text{O} \cdot \text{NO}_2]$ to form an Fe^{III} –heme and nitrate, a process analogous to ferryl oxygen rebound to carbon radical intermediates.⁴⁸ The results also show that $\cdot\text{NO}_2$ can diffuse away from the heme site to induce nitration elsewhere. To confirm this latter pathway, we employed fluorescein as a chemical trap of $\cdot\text{NO}_2$. The yield and kinetics of both ferrylMb formation and FINO₂ formation are used to quantitate the degree of dynamic cage escape of $\cdot\text{NO}_2$ into the protein interior and the medium during the catalytic decomposition of PN. Overall, the behavior of NO₂ within the internal Mb cavities is reminiscent of the recombination and escape scenarios observed for photodissociated O₂, NO, and CO and is further evidence of directed, small molecule processing within the myoglobin structure.³

FerrylMb Detection and Identification. The direct detection of ferrylMb and the kinetics of its formation provide strong evidence for the mechanism of metMb/PN interaction proposed

(48) Groves, J. T. *Proc. Natl. Acad. Sci. U.S.A.* **2003**, *100*, 3569–74.


in Scheme 1. Reasonable candidates for the transient intermediate (**I**) include $\text{MbFe}^{\text{III}}\text{OONO}$, $\text{MbFe}^{\text{IV}}=\text{O}$, $\text{MbFe}^{\text{III}}\text{ONO}_2$, $\text{MbFe}^{\text{III}}\text{NO}$, and the nitrito- Fe^{III} adduct,⁴⁹ $\text{MbFe}^{\text{III}}\text{NO}_2$. As shown in Figure 1A, the Soret band of metMb shifted from 409 to 425 nm for **I** with an isosbestic point at 417 nm. Further, the Q-band absorbance at 503 nm decreased with concurrent increases at 570 and 590 nm (Figure 1B). Isosbestic points were apparent at 525 and ~ 620 nm. This pattern explicitly excludes the possibility of **I** being $\text{MbFe}^{\text{III}}\text{OONO}$, $\text{MbFe}^{\text{III}}\text{ONO}_2$, or $\text{MbFe}^{\text{III}}\text{NO}_2$, because all of these iron(III) species have Soret maxima below 417 nm,²⁸ which is the isosbestic point between metMb and **I**.

$\text{MbFe}^{\text{III}}\text{NO}$, which is also a transient species,⁵⁰ has a UV-vis absorption spectrum similar to ferrylMb and intermediate **I** observed here.⁵¹ However, both the absorbance changes in the Q-band region and the kinetic behavior of **I** exclude $\text{MbFe}^{\text{III}}\text{NO}$. Authentic ferrylMb has a broad Q-band centered around 590 nm and an isosbestic point with metMb near 620 nm.⁴² In the difference spectrum of authentic 5 μM ferrylMb and metMb, maxima are observed at 560 and 588 nm and a zero-crossing (isosbestic) at ~ 625 nm, similar to what is seen in Figure 1B. By contrast, $\text{MbFe}^{\text{III}}\text{NO}$ has an isosbestic point with metMb at 590 nm.⁵⁰ Further, the rate constant for NO dissociation from $\text{MbFe}^{\text{III}}\text{NO}$ has been reported to be very fast, 43 s^{-1} under conditions similar to ours.⁵⁰ Given the known rate constant for the PN reaction with metMb, and supposing even 100% efficiency for $\text{MbFe}^{\text{III}}\text{NO}$ formation, that species would not build up significantly, and the half-time for decay would be 1 s. Our results show the ferrylMb decay half-time for intermediate **I** to be 10 s, and it persists after the completion of PN decay. This is much too slow to be $\text{MbFe}^{\text{III}}\text{NO}$ and is consistent with the known behavior of ferrylMb. Also, the on-rate constant for $[\text{MbFe}^{\text{III}} + \text{NO} \rightarrow \text{MbFe}^{\text{III}}\text{NO}]$ is only $5 \times 10^{-4} \text{ M}^{-1} \text{ s}^{-1}$, a very different situation from the diffusion-controlled binding rate of NO to Fe^{II} .⁵⁰

To further differentiate these two possibilities and to identify **I** as ferrylMb and not $\text{MbFe}^{\text{III}}\text{NO}$, we examined the reaction of **I** with ascorbate. Accelerated decay of **I** to MbFe^{III} was observed with increasing concentrations of ascorbate. The reduction of ferrylMb to metMb by ascorbate is reported to have a rate constant of $2.7 \text{ M}^{-1} \text{ s}^{-1}$,⁴³ while metMb/ $\text{MbFe}^{\text{III}}\text{NO}$ is too short-lived to react under such conditions. Significantly, the bimolecular rate constant observed for the reaction of **I** with ascorbate was found to be $4.2 \text{ M}^{-1} \text{ s}^{-1}$ at 12 °C and pH 7.4, which matched the measured rate constant between authentic ferrylMb and ascorbate ($4.0 \text{ M}^{-1} \text{ s}^{-1}$) under the same conditions. On this basis, we identified intermediate **I** as ferrylMb ($\text{MbFe}^{\text{IV}}=\text{O}$).

We also found that the yield of **I** was enhanced in the presence of fluorescein (Figure 4B) by a small but distinct amount. One can anticipate from Scheme 1 that the efficient capture of NO_2 in the medium by a trapping agent would increase the steady-state concentration of $\text{MbFe}^{\text{IV}}=\text{O}$ by decreasing the amount of NO_2 return to the active site ferryl. Taken together, the direct detection of ferrylMb UV-vis absorbances during the metMb/PN reaction, the kinetics of its reduction by ascorbate, and the effect of NO_2 trapping described here provide

Scheme 2. Core Catalytic Cycle of metMb-Catalyzed Peroxynitrite Decay



strong evidence for the formation of ferrylMb in the metMb/PN reaction.

Analysis of the kinetics of ferrylMb formation provides a revealing perspective of the metMb/PN reaction. The rate of $\text{MbFe}^{\text{IV}}=\text{O}$ formation was found to be highly dependent on the temperature and PN concentration. To observe ferrylMb, these variables needed to be carefully chosen. At 25 °C and with 20 equiv of PN, the peak ferrylMb concentration was achieved within 0.3–1.0 s, while at 12 °C, 4% ferrylMb was formed within 2–3 s with the same amount of PN. Significantly, the overall metMb–PN turnover rate could be well deduced to be $1.0 \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$ from the second-order rate constant for the appearance of ferrylMb. This observation again shows that the detected ferrylMb comes from the PN reaction. The alternate possibility that ferrylMb derives instead from the reaction of the small amounts of hydrogen peroxide in the PN can be excluded because the flux through this reaction would be 400-fold too slow (5% H_2O_2 ; $k_{\text{H}_2\text{O}_2} = 5 \times 10^2 \text{ M}^{-1} \text{ s}^{-1}$),^{42,52} even in the higher concentration regime used to analyze the changes in the Q-band region. Consistent with this expectation, solutions of decomposed peroxynitrite containing the same amount ($\sim 5\%$) of hydrogen peroxide did not show any rapid buildup of **I**.

The spectral transient giving rise to ferrylMb can be treated as an approach to a quasi-steady-state as $\cdot\text{NO}_2$ is released from the Mb active site, according to the simplified core catalytic cycle in Scheme 2. The caged radical pair $[\text{Fe}^{\text{IV}}=\text{O} \cdot \text{NO}_2]$ is not expected to build up to detectable concentrations due to rapid in-cage rebound and cage escape of NO_2 ($k_r \approx k_e \gg k_1$). Similarly, freely diffusing $\cdot\text{NO}_2$ also reaches a low, steady-state concentration. The degree of ferrylMb formation depends upon the recombination efficiency with respect to diffusional cage escape of $\cdot\text{NO}_2$ (k_e/k_r) and the rate of ferrylMb reduction by PN (k_3).

In its simplest form, the extent and time dependence of ferrylMb buildup in this situation can be understood from Scheme 2, which reflects the balance between the oxidation rate k_1 , the fraction of NO_2 cage escape $k_e/(k_e + k_r)$, and the reduction rate, k_3 . Kinetic simulations of the measured rates and yields according to Scheme 2 gave $k_e/k_r = 0.10$ (vide infra). We have determined k_3 to be $(1.53 \pm 0.06) \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$ at 25 °C and pH 7.4, which is quite close to the measured k_{obs} and consistent with the original report.⁴⁵ FerrylMb can also be reduced by electron transfer from Tyr103⁴² and reaction with freely diffusing NO_2 from the medium by a reencounter mechanism (k_2).³⁰ The NO_2 -mediated ferrylMb reduction produces a

(49) Copeland, D. M.; Soare, S. A. S.; West, A. H.; Richter-Addo, G. B. *J. Inorg. Biochem.* **2006**, *100*, 1413–1425.


(50) Laverman, L. E.; Wanat, A.; Oszejka, J.; Stochel, G.; Ford, P. C.; van Eldik, R. *J. Am. Chem. Soc.* **2001**, *123*, 285–93.

(51) Behan, R. K.; Hoffart, L. M.; Stone, K. L.; Krebs, C.; Green, M. T. *J. Am. Chem. Soc.* **2007**, *129*, 5855–5859.

(52) Matsui, T.; Ozaki, S.; Liang, E.; Phillips, G. N., Jr.; Watanabe, Y. *J. Biol. Chem.* **1999**, *274*, 2838–44.

Fe^{III}–nitrate complex, which has been found to decompose rapidly to MbFe^{III}–OH + NO₃[−] with a rate of 190 ± 20 s^{−1}.³⁰ The reduction of ferrylMb by nitrite is negligible under these conditions ($k = 30 \text{ M}^{-1} \text{ s}^{-1}$).

At what point does free •NO₂ emerge from the metMb/ peroxynitrite interaction? This process can be considered in analogy to geminate recombination of diatomic molecules (NO, O₂, and CO) with deoxy myoglobin. The details of such diffusive cage phenomena within heme proteins have been extensively studied by photophysical techniques that allow the rapid formation of the dissociated heme–ligand intermediate [Mb L] and spectrophotometric monitoring of ligand rebinding (eq 6).^{40,41}



Here, Mb–L is the ligand-bound state, [Mb L] is a caged state with the dissociated ligand (L) located close to the heme iron in the distal pocket, while in the Mb + L state the ligand molecule has diffused from the distal pocket into the hydrophobic (xenon) cavities within the protein.^{53,54} Small molecules can then diffuse from the Xe cavities into the surrounding medium. This latter process may be conformationally gated because there is no obvious ligand entry/exit pathway in the myoglobin fold.³ The results have shown that ligand recombination (k_{BA}) in myoglobin competes with cage escape (k_{BC}), whereby the ligand L diffuses first into the Xe pockets and then into the medium (k_{out}). Typically, the relative efficiencies of in-cage ligand capture to cage escape ($k_{\text{BA}}/k_{\text{BC}}$) are within an order of magnitude of each other.^{40,41} Similar results and conclusions have been reported to describe the behavior of small molecules in solution.^{55,56} Ligand recombination in deoxyMb (Fe^{II}) is relatively slow for CO, resulting in diffusive release to the medium after photolysis.⁵⁷ By contrast, the NO binding process, which is nearly barrierless, leads to efficient ligand capture by the ferro-heme, while for O₂, rebinding and cage escape occur with similar efficiency in deoxyMb.⁴¹

The kinetics of NO rebinding to deoxyMb have been shown to be biphasic. This feature of the data has been interpreted to be the result of very rapid NO recombination from the distal pocket within 10 ps and a slower (200 ps) binding process of NO molecules from the Xe cavities.⁵⁸ Statistical mechanics simulations have shown that 5% of the NO molecules were still found in the distal pocket and Xe cavities even after 1 ns.⁵⁹ The kinetic barrier to this slower recombination phase has been associated with entropic considerations, solvent effects, and the dynamics of conformational gating for diffusive return to the distal pocket.^{57,58,60}




Figure 6. Proposed distal pocket structure of the caged intermediate [Fe^{IV}=O •NO₂] in horse heart myoglobin (PDB entry: 2V1F, a myoglobin compound II crystal structure). The caged radical pair [Fe^{IV}=O •NO₂] is not expected to build up to detectable concentrations due to rapid in-cage rebound (k_r) and cage escape of NO₂ (k_e), which may include the adjacent xenon cavity Xe4. NO₂ that diffuses away from the heme center can induce nitration of Tyr103.

The results for the reaction of myoglobin with peroxynitrite indicate that this ligand capture and escape scenario applies as well for the behavior of [MbFe^{IV}=O •NO₂] (Figure 6). Here, the NO₂ formed at the active site by O–O bond homolysis in the rate-limiting step can react with the ferryl oxygen to form nitrate (k_r) or retreat into the protein interior (k_e). Moreover, Figure 6 illustrates that Tyr103 is preferentially nitrated, as was unambiguously determined from mass spectroscopic peptide sequence analysis on the purified mono-nitroMb. This nitration selectivity is likely caused by better solvent accessibility of this tyrosine as well as its coplanar donor-to-acceptor orientation with the heme.⁶¹ Interestingly, the degree of cage escape of NO₂ observed for the myoglobin reaction is similar to the behavior of alkyl peroxynitrites upon rearrangement to alkyl nitrates.⁶² In further analogy to this stochastic behavior of incipient radical pairs, we have recently shown that cage escape competes with cage recombination during the C–H bond hydroxylation reaction of the diiron hydroxylase AlkB,⁶³ suggesting that this phenomenon may be more general in metalloenzyme catalysis.

Fluorescein as a Probe of NO₂ Cage Escape. The observed nitration of fluorescein by peroxynitrite in the presence of metMb confirmed the release of freely diffusing NO₂ from the protein during PN turnover. The signature spectral changes accompanying F•NO₂ formation allowed both a convenient assay of NO₂ produced and a sensitive measure of the rate of its evolution (Figure 3). The data showing the overall yield of

(53) Olson, J. S.; Phillips, G. N. *J. Biol. Chem.* **1996**, *271*, 17593–17596.

(54) Schoenborn, B. P.; Watson, H. C.; Kendrew, J. C. *Nature* **1965**, *207*, 28–&.

(55) Oelkers, A. B.; Scatena, L. F.; Tyler, D. R. *J. Phys. Chem. A* **2007**, *111*, 5353–5360.

(56) Oelkers, A. B.; Tyler, D. R. *Photochem. Photobiol. Sci.* **2008**, *7*, 1386–1390.

(57) Goldbeck, R. A.; Bhaskaran, S.; Ortega, C.; Mendoza, J. L.; Olson, J. S.; Soman, J.; Kliger, D. S.; Esquerra, R. M. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 1254–1259.

(58) Ionascu, D.; Gruia, F.; Ye, X.; Yu, A.; Rosca, F.; Beck, C.; Demidov, A.; Olson, J. S.; Champion, P. M. *J. Am. Chem. Soc.* **2005**, *127*, 16921–16934.

(59) Nutt, D. R.; Meuwly, M. *Biophys. J.* **2006**, *90*, 1191–1201.

(60) Frauenfelder, H.; McMahon, B. H.; Fenimore, P. W. *Proc. Natl. Acad. Sci. U.S.A.* **2003**, *100*, 8615–8617.

(61) Makinen, M. W.; Schichman, S. A.; Hill, S. C.; Gray, H. B. *Science* **1983**, *222*, 929–931.

(62) Goldstein, S.; Lind, J.; Merenyi, G. *J. Phys. Chem. A* **2004**, *108*, 1719–1725.

(63) Austin, R. N.; Luddy, K.; Erickson, K.; Pender-Cudlip, M.; Bertrand, E.; Deng, D.; Buzdygon, R. S.; van Beilen, J. B.; Groves, J. T. *Angew. Chem., Int. Ed.* **2008**, *47*, 5232–5234.

FINO₂ and the time course of its production in the presence of metMb (Figure 4A) show distinctly faster FI nitration with 50 μ M metMb than was observed with 20 μ M metMb. Further, the decrease in FINO₂ yield upon doubling the metMb concentration was only 20%. The presence of equal amounts of FI in the medium completely suppressed protein tyrosine nitration. This observation shows that FI is at least a 10-fold more efficient trap of NO₂ than tyrosine within Mb. Further, protection of tyrosine by FI shows that both probes are sampling the same pool of NO₂. The increase in the amount of ferrylMb detected during PN decay in the presence of FI, noted above, is a further confirmation that freely diffusing NO₂ is present in the system, some of which re-encounters ferrylMb at the active site. PN spontaneous decay produced about 10% FINO₂, which is consistent with the measurements of available NO₂ from PN decay reported by Ingold.⁶⁴ By comparison, the yields of FINO₂ indicated about 5% trappable NO₂ from the Mb-mediated route. This value is in qualitative agreement with the 10% efficiency of NO₂ cage escape derived from the ferrylMb kinetics but less than the 20% estimate described in our initial study.¹⁴ The fact that fluorescein nitration detects only one-half of the total NO₂ produced is suggestive of the amount of NO₂ in the xenon cavities^{58,59} that returns to react with ferrylMb without diffusing into the medium. It is also clear from these results that myoglobin reduces the net amount of trappable NO₂ by only a factor of 2.

The unambiguous observation of the ferrylMb intermediate provides strong confirmation of the proposed mechanism, in which the transient intermediate [Fe^{IV}=O •NO₂] can either proceed to cage escape or rebound to form nitrate and ferric Mb. The decomposition of PN catalyzed by water-soluble iron(III) porphyrins (e.g., FeTMPS and FeTMPyP) affords significant amounts of oxoFe^{IV} porphyrins.^{34,35} In particular, the fast rebound of NO₂ to Fe^{IV}TMPS accounts for an important pathway for oxoFe^{IV} reduction to Fe^{III}. The rate constant of this rebound can be as fast as $1.7 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$,³⁴ which is the same as the experimental result obtained by Goldstein et al. with ferrylMb and NO₂ ($1.2 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$).³⁰

Kinetic Simulation of the Peroxynitrite–Myoglobin Reaction.

Because all of the pertinent rate constants for spontaneous and Mb-catalyzed PN decay are known, it was possible to explicitly simulate the reaction mechanism discussed above (Scheme 2) and the effect of added fluorescein. The enhanced initial rate of FINO₂ formation, the crossing of the yield curves at 2.5 s, and the degree of product yield decrease with increasing [metMb] were all reproduced very closely by the kinetic simulation (Figure 4A, traces a–c). The observed amount of ferrylMb under turnover conditions required 10% escape of NO₂ from the distal pocket ($k_e/k_r = 0.1$), while the kinetics and yield of FINO₂ led to a best fit with 5% NO₂ escape into the medium. By contrast, the experimental data for FI nitration (Figure 4A traces a–c in black) were completely inconsistent with the simulation (Figure 4A, traces d and e) if k_e was not included. The data showed 60–100% more FINO₂ observed experimentally than predicted if there was no cage escape of NO₂. The simulation also allowed an estimation of the formation and decay of NO₂ through the various reaction channels. The generation of NO₂ results mainly from PN spontaneous decomposition and cage escape. In a metMb/PN (1:30) reaction, NO₂ is consumed predominantly by dimerization and hydrolysis (50%), reen-

counter with the ferrylMb (k_2) to form nitrate (k_r) (30%), and tyrosine nitration (20%).

The results illuminated here for the reaction of PN with metMb are pertinent to the discussion regarding the roles of oxyMb and oxyHb in trapping NO in vivo,^{31,65} because the two processes intersect at the Fe(III)–OONO intermediate. Recently, this NO dioxygenase activity has been extended to other myoglobin-like proteins, such as neuroglobin,⁶⁶ cytoglobin,⁶⁵ and bacterial hemoglobins.⁶⁷ One would expect, a priori, that ferrylMb and NO₂ would be formed to the same extent from the oxyMb/NO reaction as from metMb and PN. Olson and Gardner have recently described a careful analysis of the isotopic content of nitrate resulting from the reaction of ¹⁸O₂-oxyMb with NO.³² The results showed 99% dioxygenation of NO to afford ¹⁸O₂-¹⁶O-nitrate via rebound to ferrylMb. Significantly, when we simulated the fate of ferrylMb and NO₂ formed from the single-turnover reaction of ¹⁸O₂-oxyMb with NO using the same kinetic scheme described above (Scheme 2) and the known rate constant for oxyMb + NO,²⁸ we also find only ~1% singly labeled nitrate, quantitatively consistent with the experimental result. The cage recombination mechanism affords little singly labeled nitrate in the oxyMb/NO reaction because the concentration of NO₂ is low under these conditions, making the bimolecular dimerization-hydrolysis pathway inefficient as compared to NO₂ rebound and tyrosine nitration.

Interestingly, computational studies of the NO reaction with oxyMb by Siegbahn⁶⁸ and the truncated hemoglobin (HbN) by Crespo⁶⁹ also find stepwise processes involving a caged [Fe^{IV}=O •NO₂] intermediate analogous to Scheme 2. The free energy barrier to O–O bond cleavage in the Fe^{III}–O–ONO intermediate for myoglobin was found to be only 5.2 kcal/mol, and formation of the [Fe^{IV}=O •NO₂] intermediate was exergonic by 6.6–11.2 kcal/mol, with the range depending on the entropy considerations. Oxygen rebound to form nitrate was found to be highly exergonic (–24.7 kcal/mol) with a negligible enthalpic barrier to capture of NO₂ by Fe^{IV}=O. The computed free energy barrier to this NO₂ rebound was found to lie between 2.7 and 7.2 kcal/mol depending on how the entropy of NO₂ was treated. Overall, the experimental results for the metMb PN reaction described here are in remarkable agreement with the calculated reaction energy landscape. The detection of ferrylMb and freely diffusing NO₂ suggests that the larger of the calculated exothermicities for NO₂ formation and the larger of the entropy barriers, both resulting from the more loosely bound NO₂, are closer to the real situation.

Conclusions

The reaction of metMb with PN has been shown by direct detection to afford a ferrylMb intermediate and freely diffusing NO₂. This mechanism is similar overall to the reaction of PN with water-soluble iron porphyrins. Rapid O–O bond homolysis of an intermediate peroxynitrito–Fe^{III} complex affords a

(64) Richeson, C. E.; Mulder, P.; Bowry, V. W.; Ingold, K. U. *J. Am. Chem. Soc.* **1998**, *120*, 7211–7219.

(65) Mammen, P. P.; White, J.; McGrath, A. J.; Kanatous, S. B.; Horton, J. W.; Garry, D. J. *Circulation* **2003**, *108*, 291–291.

(66) Brunori, M.; Giuffrè, A.; Nienhaus, K.; Nienhaus, G. U.; Scandurra, F. M.; Vallone, B. *Proc. Natl. Acad. Sci. U.S.A.* **2005**, *102*, 8483–8488.

(67) Ascenzi, P.; Bocedi, A.; Bolognesi, M.; Fabozzi, G.; Milani, M.; Visca, P. *Biochem. Biophys. Res. Commun.* **2006**, *339*, 450–456.

(68) Blomberg, L. M.; Blomberg, M. R. A.; Siegbahn, P. E. M. *J. Biol. Inorg. Chem.* **2004**, *9*, 923–935.

(69) Crespo, A.; Marti, M. A.; Kalko, S. G.; Morreale, A.; Orozco, M.; Gelpi, J. L.; Luque, F. J.; Estrin, D. A. *J. Am. Chem. Soc.* **2005**, *127*, 4433–4444.

caged [Fe^{IV}=O • NO₂] ensemble from which •NO₂ may either rebound to form nitrate (90%) or escape the cage (5–10%) to nitrate tyrosine or other targets such as added FI. The amounts of NO₂ and free ferrylMb liberated from this radical cage are dependent on the internal structure and gating dynamics of the protein in a manner similar to cage recombination of NO to the Fe(II) heme. The less obvious ferryl intermediate in the horse heart metMb/PN interaction as compared to the ones observed from the synthetic porphyrins FeTMPS, FeTMPyP, or myeloperoxidase reactions can be attributed to its compact heme cavity and peptide structure. Overall, the results show that the protein fold of myoglobin modifies the fates of the ferryl and NO₂ intermediates that would derive from nitric oxide scavenging in vivo; nonetheless, freely diffusing NO₂ does emerge from this encounter in amounts similar to that of spontaneous PN decay.

Experimental Procedures

Reagents. Horse heart myoglobin, catalase, potassium ferricyanide, sodium nitrite, sodium nitrate, sodium hydroxide, fluorescein, and L-ascorbic acid were purchased from Sigma. Hydrogen peroxide (30%) and perchloric acid (70%) were obtained from J.T. Baker. Diethylenetriamine pentaacetic acid (DTPA) was obtained from Alfa Aesar. Peroxynitrite was synthesized from hydrogen peroxide and nitrous acid as described^{70,71} using a sp250i syringe pump (KD scientific). To avoid any contamination with bicarbonate from ambient air, all reagents for peroxynitrite synthesis were degassed with argon thoroughly before using. Synthesized peroxynitrite was collected and kept under argon. Contaminating hydrogen peroxide was reduced to less than 5% (molar ratio) of peroxynitrite by manganese dioxide treatment. Peroxynitrite concentrations were determined at 302 nm ($\epsilon_{302} = 1670 \text{ M}^{-1} \text{ cm}^{-1}$).⁷² Except where noted otherwise, PN solutions were 1 mM or less to minimized second-order decay.⁷³ Peroxynitrite solutions were prepared by diluting the stock solution immediately before use with 0.01 M NaOH to achieve the required concentrations. All of the solutions involving the reactions with peroxynitrite were purged with argon vigorously before mixing.

After myoglobin (horse heart, Sigma) was dissolved in 0.1 M (pH 7.2) phosphate buffer, a small amount of potassium ferricyanide was added to oxidize the possible remaining oxyMb in solution. The protein was then purified chromatographically over an Econo-Pac 10DG column (Bio-Rad). The concentration of metMb was determined by measuring the absorbance at 409 nm ($\epsilon_{408} = 188 \text{ mM}^{-1} \text{ cm}^{-1}$).⁷⁴ The cyanide-bound form of metMb (metMbCN) was prepared by adding a small excess of KCN to the metMb stock solution. The concentrations of metMbCN solutions were determined by measuring the absorbance at 422 nm ($\epsilon_{422} = 116 \text{ mM}^{-1} \text{ cm}^{-1}$).⁷⁴ UV/vis spectra were obtained on a HP 8452A diode array spectrometer. All UV/vis spectra were recorded at room temperature.

Stopped-Flow Kinetic Analysis and metMb Nitration by PN. Stopped-flow kinetic studies were carried out using a Hi-Tech SF-61DX2 double mixing stopped-flow spectrophotometer (Hi-Tech, Salisbury, UK). The reactions were monitored either at a selected wavelength (single wavelength mode) or by using a diode array detector to record the entire visible range at each time point. For all studies of the reactions under CO₂-free conditions, the

stopped-flow lines were washed with argon-purged doubly deionized water immediately before the experiments. Thoroughly degassed solutions of metMb, PN, and other reactants were transferred in Hamilton gastight syringes right before mixing. Several shots were collected as fast as possible. With all of these efforts, essentially no CO₂ was involved during the mixing of fluorescein/metMb and PN. The traces were averaged from at least three shots. Kinetic data were analyzed and fitted on KinetAsyst software provided by Hi-Tech. The nitrated Mb solutions were collected at the outlet of the stopped-flow. When Mb samples were treated with PN in the presence of fluorescein, proteins were separated from fluorescein by Microcon YM-3 (MWCO 3000) centrifugal filter devices (Millipore) and subsequently recovered prior to MS determination.

To confirm the identity of intermediate I, double mixing stopped-flow experiments were carried out to study its decomposition process in the absence and presence of ascorbate. In the first mixing, 20 μM metMb in syringe A was mixed with an equal volume of 100 μM PN in syringe B. The reaction mixture was aged for 3 s to allow the full development of intermediate I (intermediate I was observed to reach its maximum around 2.5 s under the same conditions). The products of the first mixing were subsequently mixed with an equal volume of ascorbate solution from syringe C. The double mixing experiments were studied at wavelengths 422, 409, and 302 nm.

Calculations and data treatment were performed using Microsoft Excel and a commercial graphics and data analysis software (OriginPro 7.5, OriginLab).

HPLC-MS Analysis. Nitrated protein mixtures were thoroughly dialyzed (Slide-A-Lyzer mini Dialysis units, 7000 MWCO, Pierce) against water prior to further analysis by an Agilent 6510 LC Q-TOF MS. This Q-TOF was coupled with an Agilent HPLC-chip (G4240-62001, Zorbax 300SB-C18) for sample loading and separation. The proteins were eluted over a course of 15 min by using a nonlinear gradient of 3% to 90% acetonitrile containing 0.1% formic acid at a flow rate of 0.5 $\mu\text{L}/\text{min}$. Proteins eluted from HPLC-chip were directly injected into the coupled Q-TOF MS. MS data were acquired and processed using an Agilent MassHunter workstation and included analysis software. LC/MS analysis of fluorescein and its nitrated products was performed with an Agilent MSD apparatus with a UV/vis detector, equipped with a Mercury MS (Phenomenex) column (20 \times 2.0 mm).

NanoLC-MS/MS Analysis. Nitrated Mb was purified using anion exchange semipreparative HPLC (column: PL-SAX, 1000 Å, 8 μM 150 \times 7.5 mm, Polymer Laboratories). Solvent A is 20 mM Tris-HClO₄ (pH 8.0), and solvent B is 20 mM Tris-HClO₄/0.5 M NaCl (pH 8.0). Nitrated Mb and native Mb were eluted using an increasing linear gradient of solvent B from 0–80% in 35 min with a flow rate of 0.8 mL/min. The 409 nm-active fractions were collected and identified by MS. Purified mononitrated myoglobin was digested overnight at 37 °C with modified trypsin (Promega, Madison, WI) or GluC endoproteinase (Roche, Germany) in NH₄HCO₃ buffer (50 mM, pH 8). A fused-silica capillary nanocolumn packed with reverse phase Eclipse XDB-C18 packing materials was employed to separate digested peptides. Peptides were eluted over the course of 150 min by using a nonlinear gradient of 5% to 80% acetonitrile containing 0.1% formic acid, at a flow rate of 120 nL/min. Tandem mass spectra were collected by an LQC DECA XP PLUS mass spectrometer controlled by Xcalibur software (ThermoFisher). The acquired MS/MS spectra were automatically searched against a protein database for horse heart myoglobin using SEQUEST algorithm to identify the nitrotyrosine-containing peptides.

Computer Simulations. Kinetic simulations were performed with Berkeley Madonna Modeling and Analysis of Dynamic Systems software, version 8.3.22 (<http://www.berkeleymadonna.com>).

(70) Saha, A.; Goldstein, S.; Cabelli, D.; Czapski, G. *Free Radical Biol. Med.* **1998**, *24*, 653–9.

(71) Uppu, R. M.; Squadrito, G. L.; Cueto, R.; Pryor, W. A. *Methods Enzymol.* **1996**, *269*, 285–96.

(72) Koppenol, W. H.; Kissner, R.; Beckman, J. S. *Methods Enzymol.* **1996**, *269*, 296–302.

(73) Kissner, R.; Koppenol, W. H. *J. Am. Chem. Soc.* **2002**, *124*, 234–239.

(74) Rossifanelli, A.; Antonini, E.; Caputo, A. *Adv. Protein Chem.* **1964**, *19*, 73–222.

Acknowledgment. We are grateful for support of this research by the National Institutes of Health (2R37 GM036298). We thank Drs. Dorothy Little and Saw Kyin for technical assistance with the MS measurements.

Supporting Information Available: Kinetic plots for ferrylMb reduction catalyzed by ascorbate; kinetic plots for ferrylMb reduction by peroxynitrite; mass spectra of metMb/PN reaction

mixture; additional equations used in the mechanistic simulations; experimental and simulated nitrofluorescein formation and concurrent PN decomposition in the absence of proteins; and simulations of intermediate **I** formation and decay in the absence and presence of fluorescein. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JA902473R