


Programmed Synthesis of Tetraarylthiophenes through Sequential C–H Arylation

Shuichi Yanagisawa, Kirika Ueda, Hiromi Sekizawa, and Kenichiro Itami*

Department of Chemistry, Graduate School of Science, Nagoya University, Nagoya 464-8602, Japan

Received July 24, 2009; E-mail: itami@chem.nagoya-u.ac.jp

Multiply arylated thiophenes are privileged structures with many interesting functions, including optoelectronic¹ and biological² properties. In order to accelerate the discovery and structure–property relationship studies of new functional molecules of this class, a flexible method for accessing all possible isomers in a programmable format is much needed. We herein report a general protocol for the programmed synthesis of tetraarylthiophenes^{3,4} through the sequential C–H/C–O arylation of 3-methoxythiophene (**1**). The development of regioselective C–H bond arylation catalysts has been a key to realizing our concept.



To realize a programmed synthesis of tetraarylthiophenes, a general method allowing regioselective arylations onto a thiophene core had to be developed. As a means for installing aryl groups onto the core, we selected the metal-catalyzed C–H bond arylation of thiophenes in view of synthetic expediency.^{5,6} Though this approach was conceptually attractive, obstacles to overcome included insufficient reactivity and the control of regioselectivity of thienyl C–H bonds toward catalytic C–H arylation reactions.⁵ For example, several groups have demonstrated that certain Pd,^{5,7} Cu,⁸ Rh,⁹ and Ir¹⁰ catalysts can promote the C2/C5 diarylation of thiophenes, but arylation at the C3/C4 position is very difficult. Very recently, the Miura group has demonstrated that 3-thiophenecarboxylic acid undergoes Pd-catalyzed tetraarylation with bromoarenes, accompanied by cleavage of three C–H bonds and decarboxylation.¹¹ However, the installation of four different aryl groups onto thiophene ring has not been realized.³

We selected 3-methoxythiophene (**1**) as our first-generation platform¹² for tetraarylthiophene synthesis because of its high reactivity and selectivity in catalytic C–H bond arylation with iodoarenes.^{9,10} For example, **1** reacts with iodobenzene in the presence of RhCl(CO){P[OCH(CF₃)₂]₃}₂ catalyst (**Cat-A**) and Ag₂CO₃ to afford 3-methoxy-2-phenylthiophene (**2a**) in 80% yield with virtually complete regioselectivity. This C2-selective arylation occurs with various iodoarenes.⁹ Thus, the focal point of this study was to establish a C4- and/or C5-selective catalyst for the arylation of 2-aryl-3-methoxythiophenes **2**. However, we identified that **Cat-A** is problematic for further arylations. For example, the reaction of **2a** (1.5 equiv) and iodobenzene (1 equiv) in the presence of **Cat-A** (3 mol %) and Ag₂CO₃ (1 equiv) in *m*-xylene at 150 °C furnished the C5 phenylation product (**4aa**) in only 38% yield. Even more critically, further arylation (C4 arylation) of **4aa** did not take place with **Cat-A**. Thus, the need for a new catalytic system was obvious at this point. In particular, the

development of a catalyst promoting the hard-to-achieve arylation at the β-position of thiophene ring was crucial. Extensive screening led to the development of two catalysts (**Cat-B** and **Cat-C**) for the second and third arylations of **1**, respectively.

Table 1. Ligand-Controlled Regiodivergent Arylation of **2a**^a

entry	Ar ²	ligand	major product	% yield (3:4) ^b
1	C ₆ H ₅ (a)	P[OCH(CF ₃) ₂] ₃	3aa	80 (97:3)
2	C ₆ H ₅ (a)	2,2'-bipyridyl	4aa	89 (1:99)
3	<i>o</i> -MeC ₆ H ₄ (b)	P[OCH(CF ₃) ₂] ₃	3ab	88 (98:2)
4	<i>o</i> -MeC ₆ H ₄ (b)	2,2'-bipyridyl	4ab	98 (2:98)
5 ^c	<i>p</i> -MeC ₆ H ₄ (c)	P[OCH(CF ₃) ₂] ₃	3ac	83 (96:4)
6	<i>p</i> -MeC ₆ H ₄ (c)	2,2'-bipyridyl	4ac	91 (1:99)
7 ^c	<i>p</i> -MeOC ₆ H ₄ (d)	P[OCH(CF ₃) ₂] ₃	3ad	75 (98:2)
8	<i>p</i> -MeOC ₆ H ₄ (d)	2,2'-bipyridyl	4ad	83 (1:>99)
9	<i>p</i> -AcC ₆ H ₄ (e)	P[OCH(CF ₃) ₂] ₃	3ae	71 (95:5)
10	<i>p</i> -AcC ₆ H ₄ (e)	2,2'-bipyridyl	4ae	80 (1:>99)
11	<i>p</i> -NO ₂ C ₆ H ₄ (f)	P[OCH(CF ₃) ₂] ₃	3af	69 (93:7)
12	<i>p</i> -NO ₂ C ₆ H ₄ (f)	2,2'-bipyridyl	4af	69 (2:98)
13	<i>p</i> -CF ₃ C ₆ H ₄ (g)	P[OCH(CF ₃) ₂] ₃	3ag	77 (95:5)
14	<i>p</i> -CF ₃ C ₆ H ₄ (g)	2,2'-bipyridyl	4ag	77 (1:>99)

^a Conditions: **2a** (1.5 equiv), Ar²I (1 equiv), PdCl₂ (5 mol %), ligand (10 mol %), Ag₂CO₃ (1 equiv), *m*-xylene, 120 °C. ^b Isolated yield. Isomer ratio was determined by ¹H NMR and GC analyses. ^c PdCl₂ (10 mol %) and P[OCH(CF₃)₂]₃ (20 mol %) were employed.

We first found that the otherwise difficult C4-selective arylation of **2a** with iodoarenes can be promoted by the catalyst PdCl₂/P[OCH(CF₃)₂]₃ (**Cat-B**) and Ag₂CO₃ in *m*-xylene at 120 °C, furnishing **3** with high regioselectivity (93–98% C4) (Table 1). Given the C5 selectivity of **Cat-A** for the arylation of **2a** (see above), an interesting metal-controlled regiodivergency¹³ (C5 for Rh^I, C4 for Pd^{II}) has been identified in P[OCH(CF₃)₂]₃-bound metal catalysts. More interestingly, we found that the C4 selectivity of **Cat-B** can be switched to C5 selectivity (98 to >99%) by changing the supporting neutral ligand to 2,2'-bipyridyl¹⁴ (**Cat-C**). This ligand-controlled regiodivergent C–H arylation¹³ (C4 for P[OCH(CF₃)₂]₃, C5 for 2,2'-bipyridyl) turned out to be general for PdCl₂/Ag₂CO₃ catalysis (Table 1).¹⁵

With two regiodivergent C–H arylation catalysts in hand, the third C–H arylation for making 2,4,5-triaryl-3-methoxythiophenes **5** was next examined for two possible routes via either **3** or **4**. Gratifyingly, the **Cat-C** system promoted the C5 arylation of 2,4-diaryl-3-methoxythiophenes **3** with iodoarenes, giving **5** in good yields (Table 2). Under otherwise identical conditions, the **Cat-B** system was unable to promote the arylation for both **3** and **4**.


Finally, the fourth arylation to obtain the targeted tetraarylthiophenes **7** was investigated. Extensive screening of reaction conditions established a high-yield procedure involving Suzuki–Miyaura coupling¹⁶ (Table 3). Thus, the BBr₃-promoted demethylation of **5** followed by treatment of the thus-obtained crude alcohols with Tf₂O/*i*-Pr₃NEt/

Table 2. Arylation of **3** Catalyzed by Pd/bipy/Ag₂CO₃^a

entry	Ar ²	Ar ³	5 (% yield)
1	C ₆ H ₅ (a)	C ₆ H ₅ (a)	5aaa (87)
2	<i>p</i> -MeC ₆ H ₄ (c)	<i>p</i> -NO ₂ C ₆ H ₄ (f)	5acf (61)
3 ^b	<i>p</i> -MeC ₆ H ₄ (c)	<i>p</i> -CF ₃ C ₆ H ₄ (g)	5acg (70)
4 ^b	<i>p</i> -AcC ₆ H ₄ (e)	<i>p</i> -MeC ₆ H ₄ (c)	5aec (65)
5	<i>p</i> -AcC ₆ H ₄ (e)	<i>p</i> -NO ₂ C ₆ H ₄ (f)	5aef (73)
6 ^b	<i>p</i> -CF ₃ C ₆ H ₄ (g)	<i>p</i> -MeC ₆ H ₄ (c)	5agc (70)

^a Conditions: **3** (1 equiv), Ar³I (3 equiv), PdCl₂ (5 mol %), 2,2'-bipyridyl (10 mol %), Ag₂CO₃ (2 equiv), *m*-xylene, 120 °C. ^b Using 1 equiv of Ag₂CO₃.

DMAP produced the corresponding triflates **6** in high yields (>70%).¹⁷ The triflates were cross-coupled with arylboronic acids in the presence of Pd(PPh₃)₄ catalyst and Ba(OH)₂ to finally afford tetraarylthiophenes **7** in good yields with virtually complete isomeric purities (Table 3).¹⁸

Table 3. Synthesis of Tetraarylthiophenes **7**^a

^a Conditions: (a) **5**, BBr₃, CH₂Cl₂, then Tf₂O, *i*-Pr₂NEt, DMAP, CH₂Cl₂ (see the Supporting Information for details); (b) **6** (1 equiv), Ar⁴B(OH)₂ (3 equiv), Pd(PPh₃)₄ (5 mol %), Ba(OH)₂ (2 equiv), 1-butanol/H₂O (5:4), 65 °C.

In summary, we have established a general protocol for the programmed synthesis of tetraarylthiophenes. The utilization of three catalysts, RhCl(CO){P[OCH(CF₃)₂]₂}, PdCl₂/P[OCH(CF₃)₂]₃, and PdCl₂/bipy, enables regioselective sequential arylations at the three C–H bonds of 3-methoxythiophene with iodoarenes. Interesting metal- and ligand-controlled regiodivergent C–H arylations have been uncovered during this study. Noteworthy features of the present method are that (i) all of the aryl groups assembled on the thiophene core stem from readily available aryl iodides or boronic acids, (ii) the installation of aryl groups at the desired positions can be achieved, and (iii) simple alteration of the application order of aryl reagents in the sequence results in the production of all possible isomers of tetraarylthiophenes. Although we focused on the synthesis of tetraarylthiophenes in this work, our strategy is also applicable to the regioselective synthesis of di- or triarylthiophenes by skipping one or two of the C–H arylation steps prior to the final C–O bond arylation. The present strategy should find many uses for combinatorial lead-structure identification and optimization in the development of

functional organic materials where the structure–property relationships are often not easily predictable. These studies as well as the mechanistic investigations of the newly developed catalysts are now the focus of our ongoing efforts.

Acknowledgment. This work was supported in part by a Grant-in-Aid for Scientific Research from MEXT and JSPS. S.Y. is a recipient of a JSPS Predoctoral Fellowship.

Supporting Information Available: Experimental procedures and characterization data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) Selected recent examples: (a) Mori, A.; Sugie, A. *Bull. Chem. Soc. Jpn.* **2008**, *81*, 548. (b) Su, F.-K.; Hong, J.-L.; Lin, L.-L. *Synth. Met.* **2004**, *142*, 63. (c) Geramita, K.; McBee, J.; Shen, Y.; Radu, N.; Tilley, T. D. *Chem. Mater.* **2006**, *18*, 3261. (d) Xing, Y.; Xu, X.; Wang, F.; Lu, P. *Opt. Mater.* **2006**, *29*, 407. (e) Dang, Y.; Chen, Y. *J. Org. Chem.* **2007**, *72*, 6901. (f) Kojima, T.; Nishida, J.; Tokito, S.; Tada, H.; Yamashita, Y. *Chem. Commun.* **2007**, 1430.
- (2) Recent examples: (a) Lauth, M.; Bergström, Å.; Shimokawa, T.; Toftgård, R. *Proc. Natl. Acad. Sci. U.S.A.* **2007**, *104*, 8455. (b) Fournier dit Chabert, J.; Marquez, B.; Neville, L.; Joucla, L.; Broussous, S.; Bouhours, P.; David, E.; Pellet-Rostaing, S.; Marquet, B.; Moreau, N.; Lemaire, M. *Bioorg. Med. Chem.* **2007**, *15*, 4482. (c) Bey, E.; Marchais-Oberwinkler, S.; Werth, R.; Negri, M.; Al-Soud, Y. A.; Kruchten, P.; Oster, A.; Frotscher, M.; Birk, B.; Hartmann, R. W. *J. Med. Chem.* **2008**, *51*, 6725.
- (3) Synthesis of tetraarylthiophenes: (a) Reference 1b. (b) Reference 1d. (c) Reference 1e. (d) Dang, T. T.; Rasool, N.; Dang, T. T.; Reinke, H.; Langer, P. *Tetrahedron Lett.* **2007**, *48*, 845. (e) Nakano, M.; Tsurugi, H.; Satoh, T.; Miura, M. *Org. Lett.* **2008**, *10*, 1851.
- (4) Programmed synthesis of multiply arylated heterocycles: (a) Itami, K.; Yamazaki, D.; Yoshida, J. *J. Am. Chem. Soc.* **2004**, *126*, 15396. (b) Campeau, L.-C.; Bertrand-Laperle, M.; Leclerc, J.-P.; Villemure, E.; Gorelsky, S.; Fagnou, K. *J. Am. Chem. Soc.* **2008**, *130*, 3276. (c) Goikman, R.; Jacques, T. L.; Sames, D. *J. Am. Chem. Soc.* **2009**, *131*, 3042. (d) Reference 1a.
- (5) Reviews of C–H bond arylation of arenes: (a) Alberico, D.; Scott, M. E.; Lautens, M. *Chem. Rev.* **2007**, *107*, 174. (b) Seregin, I. V.; Gevorgyan, V. *Chem. Rev.* **2007**, *36*, 1173. Selected recent breakthroughs: (c) Phipps, R. J.; Gaunt, M. J. *Science* **2009**, *323*, 1593. (d) Stuart, D. R.; Fagnou, K. *Science* **2007**, *316*, 1172. (e) Giri, R.; Maugel, N.; Li, J.-J.; Wang, D.-H.; Breazzano, S. P.; Saunders, L. B.; Yu, J.-Q. *J. Am. Chem. Soc.* **2007**, *129*, 3510. (f) Gorelsky, S. I.; Lapointe, D.; Fagnou, K. *J. Am. Chem. Soc.* **2008**, *130*, 10848. (g) Lewis, J. C.; Bergman, R. G.; Ellman, J. A. *Acc. Chem. Res.* **2008**, *41*, 1013. (h) Berman, A. M.; Lewis, J. C.; Bergman, R. G.; Ellman, J. A. *J. Am. Chem. Soc.* **2008**, *130*, 14926. (i) Ackermann, L.; Althammer, A.; Fenner, S. *Angew. Chem., Int. Ed.* **2009**, *48*, 201. (j) Liegault, B.; Lapointe, D.; Caron, L.; Vlassova, A.; Fagnou, K. *J. Org. Chem.* **2009**, *74*, 1826. (k) Campeau, L.-C.; Stuart, D. R.; Leclerc, J.-P.; Bertrand-Laperle, M.; Villemure, E.; Sun, H.-Y.; Lasserre, S.; Guimond, N.; Lecavallier, M.; Fagnou, K. *J. Am. Chem. Soc.* **2009**, *131*, 3291. (l) Wang, C.; Piel, I.; Glorius, F. *J. Am. Chem. Soc.* **2009**, *131*, 4194. (m) Hull, K. L.; Sanford, M. S. *J. Am. Chem. Soc.* **2009**, *131*, 9651.
- (6) Reviews of transition-metal-catalyzed C–C bond formation via C–H bond cleavage: (a) Kakiuchi, F.; Kochi, T. *Synthesis* **2008**, 3013. (b) Kakiuchi, F.; Chatani, N. *Adv. Synth. Catal.* **2003**, *345*, 1077.
- (7) (a) Yokooji, A.; Satoh, T.; Miura, M.; Nomura, M. *Tetrahedron* **2004**, *60*, 6757. (b) Borghese, A.; Geldhof, G.; Antoine, L. *Tetrahedron Lett.* **2006**, *47*, 9249. (c) Mohanakrishnan, A. K.; Amaladass, P.; Clement, J. A. *Tetrahedron Lett.* **2007**, *48*, 539.
- (8) Do, H.-Q.; Khan, R. M. K.; Daugulis, O. *J. Am. Chem. Soc.* **2008**, *130*, 15185.
- (9) Yanagisawa, S.; Sudo, T.; Noyori, R.; Itami, K. *J. Am. Chem. Soc.* **2006**, *128*, 11748.
- (10) Join, B.; Yamamoto, T.; Itami, K. *Angew. Chem., Int. Ed.* **2009**, *48*, 3644.
- (11) (a) Reference 3e. Also see: (b) Okazawa, T.; Satoh, T.; Miura, M.; Nomura, M. *J. Am. Chem. Soc.* **2002**, *124*, 5286.
- (12) Platform synthesis: Itami, K.; Yoshida, J. *Chem.—Eur. J.* **2006**, *12*, 3966.
- (13) Examples of regiodivergent C–H arylation of heterocycles: (a) Pivsa-Art, S.; Satoh, T.; Kawamura, Y.; Miura, M.; Nomura, M. *Bull. Chem. Soc. Jpn.* **1998**, *71*, 467. (b) Kondo, Y.; Komine, T.; Sakamoto, T. *Org. Lett.* **2000**, *2*, 3111. (c) Glover, B.; Harvey, K. A.; Liu, B.; Sharp, M. J.; Tymoschenko, M. *Org. Lett.* **2003**, *5*, 301.
- (14) Pd/bipy catalyst in C–H arylation: (a) Sugie, A.; Kobayashi, K.; Suzuki, Y.; Osakada, K.; Mori, A. *Chem. Lett.* **2006**, *35*, 1100. (b) Reference 1a.
- (15) The C4 selectivity exerted by the PdCl₂/P[OCH(CF₃)₂]₂/Ag₂CO₃ catalytic system appears to be unprecedented and is the subject of further investigation. A full study of this newly developed catalyst will be reported in due course.
- (16) Miyaura, N.; Suzuki, A. *Chem. Rev.* **1995**, *95*, 2457.
- (17) See the Supporting Information for details.
- (18) For the use of Ba(OH)₂ in Suzuki–Miyaura coupling, see: Watanabe, T.; Miyaura, N.; Suzuki, A. *Synlett* **1992**, 207.

JA906215B