

Intermediate Rate Atomic Trajectories of RNA by Solid-State NMR Spectroscopy

Greg L. Olsen,[†] Michael F. Bardaro, Jr.,[†] Dorothy C. Echodu,[†] Gary P. Drobny,^{*,†} and Gabriele Varani^{*,‡}

Department of Chemistry, Box 351700, and Department of Biochemistry, Box 357350, University of Washington, Seattle, Washington 98195

Received September 4, 2009; E-mail: drobnay@chem.washington.edu; varani@chem.washington.edu

Abstract: Many RNAs undergo large conformational changes in response to the binding of proteins and small molecules. However, when RNA functional dynamics occur in the nanosecond–microsecond time scale, they become invisible to traditional solution NMR relaxation methods. Residual dipolar coupling methods have revealed the presence of extensive nanosecond–microsecond domain motions in HIV-1 TAR RNA, but this technique lacks information on the rates of motions. We have used solid-state deuterium NMR to quantitatively describe trajectories of key residues in TAR by exploiting the sensitivity of this technique to motions that occur in the nanosecond–microsecond regime. Deuterium line shape and relaxation data were used to model motions of residues within the TAR binding interface. The resulting motional models indicate two functionally essential bases within the single-stranded bulge sample both the free and Tat-bound conformations on the microsecond time scale in the complete absence of the protein. Thus, our results strongly support a conformational capture mechanism for recognition: the protein does not induce a new RNA structure, but instead captures an already-populated conformation.

Introduction

Functional dynamics in HIV-1 TAR occur on a time scale slower than ~ 10 ns^{1,2} but faster than ~ 100 μ s.^{3,4} These motions allow the RNA to adapt its structure to generate the conformation recognized by Tat and cyclin T1 proteins^{5–7} (Figure 1a), a key regulatory checkpoint in the viral lifecycle. Dynamics occurring on the nanosecond–microsecond time scale are invisible to conventional solution-state NMR relaxation measurements because they are masked by global rotational diffusion.¹ Additionally, such motions are too fast to be detected by relaxation dispersion methods that are powerful in studies of slower protein dynamics.^{8–13} Residual dipolar couplings (RDCs)

provide rich information on conformational dynamics in this time scale,¹⁴ and their application to RNA has recently demonstrated that the helical domains of TAR experience extensive motions that connect the conformations of TAR observed in the absence and presence of protein and small molecule ligands.^{1,2} However, RDCs do not provide information on rates; therefore, the time scale of motion can only be broadly and indirectly inferred by a process of elimination when faster (subnanosecond) and slower (millisecond) motions are not observed.^{1,2,14}

Unlike these solution-state methods, solid-state deuterium NMR is very sensitive to motions within the intermediate (nanosecond–microsecond) time scale.¹⁵ By studying samples in a highly hydrated solid-state gel that preserves the conformational dynamics, but avoids line shape sharpening by molecular tumbling, information on anisotropic motions is retained. In addition, solid-state relaxation measurements provide information on the rates or time scales of motions faster than those that affect the line shapes. Thus, the combined measurement of quadrupole echo line shape, Zeeman spin–lattice relaxation (T_{1Z}), and quadrupolar order relaxation (T_{1Q}) allows the quantitative characterization of atomic motions occurring at rates between 10^4 and 10^{10} s⁻¹.^{16,17}

[†] Department of Chemistry.

[‡] Department of Biochemistry.

- (1) Zhang, Q.; Sun, X.; Watt, E.; Al-Hashimi, H. M. *Science* **2006**, *311*, 653–656.
- (2) Zhang, Q.; Stelzer, A. C.; Fisher, C. K.; Al-Hashimi, H. M. *Nature* **2007**, *450*, 1263–1268.
- (3) Hansen, A. L.; Al-Hashimi, H. M. *J. Am. Chem. Soc.* **2007**, *129*, 1–11.
- (4) Bardaro, M. F., Jr.; Shajani, Z.; Patora-Komisarska, K.; Robinson, J. A.; G., V. *Nucleic Acids Res.* **2009**, *37*, 1529–1540.
- (5) Puglisi, J. D.; Chen, L.; Frankel, A. D.; Williamson, J. R. *Proc. Natl. Acad. Sci. U.S.A.* **1993**, *90*, 3680–3684.
- (6) Aboul-ela, F.; Karn, J.; Varani, G. *Nucleic Acids Res.* **1996**, *24*, 3974–81.
- (7) Aboul-ela, F.; Karn, J.; Varani, G. *J. Mol. Biol.* **1995**, *253*, 313–32.
- (8) Karn, J. *J. Mol. Biol.* **1999**, *293*, 235–254.
- (9) Korzhnev, D. M.; Salvatella, X.; Vendruscolo, M.; Nardo, A. A. D.; Davidson, A. R.; Dobson, C. M.; Kay, L. E. *Nature* **2004**, *430*, 586–590.
- (10) Palmer, A. G. I.; Kroenke, C. D.; Loria, J. P. *Methods Enzymol.* **2001**, *339*, 204–238.
- (11) Henzler-Wildman, K. A.; Lei, M.; Thai, V.; Kerns, S. J.; Karplus, M.; Kern, D. *Nature* **2007**, *450*, 913–916.
- (12) Boehr, D. D.; McElheny, D.; Dyson, H. J.; Wright, P. E. *Science* **2006**, *313*, 1638–1642.

(13) Sprangers, R.; Kay, L. E. *Nature* **2007**, *445*, 618–622.

(14) Lange, O. F.; Lakomek, N. A.; Fares, C.; Schroder, G. F.; Walter, K. F. A.; Becker, S.; Meiler, J.; Grubmuller, H.; Griesinger, C.; de Groot, B. L. *Science* **2008**, *320*, 1471–1475.

(15) Olsen, G.; Echodu, D.; Shajani, Z.; Bardaro, M.; Varani, G.; Drobny, G. P. *J. Am. Chem. Soc.* **2008**, *130*, 2896–2897.

(16) Spiess, H. W. In *NMR Basic Principles and Progress*, D. N. S., *Rotation of Molecules and Nuclear Spin Relaxation*; Diehl, P., Fluck, E., Kosfeld, R., Eds.; Springer, Verlag: Berlin, 1978; Vol. 15, pp 59–214.




Figure 1. (a) HIV-1 TAR changes conformation when Tat protein or small molecules bind to it, forming a structure conducive to binding of the protein and activation of transcription. (b) Location of the three nucleotides whose motions were probed using solid-state NMR. U23 (teal) is critical for binding of Tat; U25 (blue) is a single-stranded residue whose conformation changes considerably when tat binds; U38 (green) belongs to the upper helix and is base paired with A27. Sites of deuterium labeling (the 5 and 6 base positions) are shown in red.




Figure 2. (a) Experimental deuterium solid-state NMR spectra of selectively 5,6 deuterated U23 (left), U25 (center), and U38 (right) in TAR RNA. The curves through the experimental data represent best fits using the motional models and parameters described in the text; the line in the center of each spectrum corresponds to the residual deuterium signal from the deuterium-depleted solvent water. (b) Experimental relaxation times (left side of each column) are compared to those calculated using the motional models described in the text (right side of each column) for both T_{1z} and T_{1Q} .

In this work, we apply solid-state NMR measurements to obtain motional rates and amplitudes in the intermediate nanosecond–microsecond regime typically inaccessible to solution NMR. Using this approach, we find that the extrahelical residues, U23 and U25, experience significant conformational exchange over the nanosecond–microsecond time scale. Additionally, by probing the helical site U38, we determine that TAR experiences significant interhelical motions even in the solid state on an intermediate time scale, coincident with the results of RDC measurements.²

Results

We probed motions of 5,6-²H-labeled uridine bases at three positions in TAR RNA (U23, U25, and U38, Figure 1b) at high levels of hydration to reproduce conditions observed in solution.¹⁸ The line shapes for all three sites clearly reveal modulations by intermediate time scale (nanosecond–microsecond) motions¹⁵ (Figure 2a). To identify the underlying atomic motions and define their rates and amplitudes, we modeled dynamics at each site by generating model trajectories; the procedure by which these were obtained is described in detail in the Materials and Methods. Briefly, for each site, modeling was performed by hypothesizing that the motions that modulate the line shape correspond to transitions between the two conformations observed for these single-stranded nucleotides in free TAR, and

in the complex of TAR with peptide mimics of Tat protein.^{19,20} For U38, we assumed that this residue might experience helical domain motions similar to those observed using RDCs.^{1,2} Once initial trajectories were generated under these assumptions, motional rates and amplitudes were defined by systematically fitting the experimental line shapes and relaxation rates.

For all three nucleotides, we were able to obtain high-quality fits to the line shapes (Figure 2a) and to the two independent relaxation rates (Figure 2b) simultaneously. As described in the Materials and Methods and presented graphically in Supporting Information Figures S1–S3, the experimental data constrain the parameters of the fit within very narrow boundaries because multiple observables are fit simultaneously. Furthermore, because two deuterated sites per residue were monitored simultaneously, a model capable of producing a satisfactory fit to the data must define a single trajectory for the base plane, which simultaneously reproduces superposition line shape data, superposition T_{1z} data, and superposition T_{1Q} data arising from two labeled sites. Therefore, while we cannot formally exclude the possibility that other models may also describe the dynamic processes presented here, the model trajectories we report robustly satisfy multiple independent experimental observations.

Amplitudes and Rates of Motions for the Nonhelical Bases. The spectroscopic data for U23 are well-modeled by a large-amplitude (24°) intermediate-time scale ($6.7 \times 10^7 \text{ s}^{-1}$) hop of the base, in addition to a rapid (10^{10} s^{-1}) small-amplitude ($\pm 11^\circ$) twisting of the base plane about the glycosidic bond occurring at both ends of the hop (Figure 3a). This trajectory

(17) Vold, R. R.; Vold, R. L. Deuterium Relaxation in Molecular Solids. *Advances in Magnetic and Optical Resonance*; Academic Press: San Diego, CA, 1991; Vol. 16, pp 85–171.

(18) Olsen, G. L., Jr.; Echodu, B. M. C. D.; Drobny, G. P.; Varani, G. *J. Biomol. NMR* **2009**, *45*, 133–142.

(19) Aboul-ela, F.; Karn, J.; Varani, G. *J. Mol. Biol.* **1995**, *253*, 313–332.

(20) Long, K. S.; Crothers, D. M. *Biochemistry* **1999**, *38*, 10059–10069.




Figure 3. (a) The motions of U23 (in teal with sites monitored by selective deuteration in red) can be accurately modeled by a 24° jump linking the conformation observed for this residue in the free structure of TAR⁶ (at the bottom) to its position observed in the complex of TAR with peptides derived from the tat protein (at the top);¹⁹ in addition to these large scale motions, we observe much faster $\pm 11^\circ$ rotations of the base with respect to the sugar. (b) The motions of U25 (green with sites monitored by selective deuteration in red) can be accurately modeled as a slow 30° outward jump to generate the extrahelical conformation observed in many TAR complexes, together with a slower $\pm 40^\circ$ rotation about the glycosidic bond.

would correspond to transient unstacking of U23 from A22, as observed in ligand-free TAR;⁶ it is also consistent with sampling of positions near A27 and U38, the base-triple partners which U23 must find to form the tat- and argininimide-bound structures.^{19,21} The NMR data for U25 are well-modeled by a 30° jump at a rate of $6 \times 10^7 \text{ s}^{-1}$; such a motion is consistent with an outward and downward rotation of the backbone away from the intrahelical conformation observed in free TAR⁶ as it is flipped out of the helix, as observed in bound TAR¹⁹ (Figure 3b). In addition, U25 experiences much slower, large-amplitude twisting of the base ($6 \times 10^5 \text{ s}^{-1}$; $\pm 40^\circ$). Extensive picosecond–nanosecond motions were previously observed for this same residue³ and for C24 immediately adjacent to it.⁴ However, the large amplitude motions described here for both U23 and U25 occur at rates that make them undetectable to traditional solution-state NMR relaxation experiments, as well as $T_{1\rho}$ power-dependence⁴ experiments.

Amplitudes and Rates of Motions for the Helical U38 Base. The spectroscopic data for U38 are well described by a combination of 13° twisting and 13° bending, corresponding to a movement of the entire upper helix that carries the U38:A27 base pair, occurring at a rate of $1.4 \times 10^6 \text{ s}^{-1}$. In addition, we observe small-amplitude local motions ($\pm 4^\circ$) at a much faster rate of $2.2 \times 10^8 \text{ s}^{-1}$ (Figure 4). There was no evidence of substantial fast (nanosecond–picosecond) motions in this residue when solution relaxation data were analyzed,⁴ but these motions are similar to the global motions of the upper helix observed using RDCs.^{1,2} A model for the overall interhelical bending and twisting motions of the TAR construct in the solid state cannot be constructed on the basis of the motions of the upper helix alone. However, due to the absence of an apical loop, the more compact lower helix likely undergoes motions of amplitude similar to or even larger than those we observe for the upper helix. These combined helical motions in the solid samples will correspond to significantly larger interhelical excursions, perhaps comparable in amplitude to those obtained using RDCs under solution conditions. Thus, while the amplitude we find for the helical motion carrying the A27:U38 base pair is relatively small, the motions we see are likely more representative of those in solution than this would suggest.

Consistent with the global nature of the motions responsible for the U38 line shape, the spectral signature of these slow collective motions only becomes visible as hydration increases to the point where sample conditions approach those observed in solution;¹⁸ the more localized motions of U23 and U25 are instead observed even at low hydration levels.¹⁸

Discussion

The essential functional dynamics of TAR RNA occur on a time scale (nanosecond–microsecond) that is invisible to current solution relaxation NMR techniques^{3,4} and that is also too fast to be studied by relaxation dispersion.¹⁰ Thus, the motions we have reported for U23 and U25 were never described before. The helical domain motions we observe for U38 are analogous to those previously described using RDCs,^{1,2} although possibly of reduced amplitude. However, the RDC technique cannot provide information on the rates of these motions; instead they are only broadly and indirectly inferred by a process of elimination when motions on other time scales are not observed. In contrast, our approach yields both amplitudes and rates of motion.

The sensitivity of the deuterium solid-state line shape to anisotropic motions allows us to extract time-dependent trajectories for atomic motions in TAR RNA and to quantitatively characterize conformational exchange processes occurring on a microsecond time scale. Conformational exchange has been extensively characterized in proteins on the much slower time window (millisecond) that is accessible via relaxation dispersion experiments,^{9–11,13} but not at microsecond rates. The widespread observation of extensive conformational rearrangement in RNA recognition and function makes it highly probable that nanosecond–microsecond conformational exchange will be found in other RNAs. Similarly, there is emerging evidence¹⁴ that functional motion in proteins is also rich in the nanosecond–microsecond time window. Solid-state NMR approaches provide a bridge in covering RNA and protein dynamics²² by providing atomic-level information on motions occurring in this intermediate time window.

Our data strongly suggest the recruitment of Tat occurs by a conformational capture mechanism (Figure 5), as was previously suggested on the basis of data on the motions of the TAR helices.^{2,23} In this picture, the protein does not need to alter the conformation of TAR, but instead selects a conformation in the RNA that is already substantially populated in its absence. The motions we have identified in U23 and U25 are consistent with the very motions that would connect the structure of TAR in its ligand-free form⁶ to that observed in the presence of peptide mimics of Tat proteins.^{7,20,21} This second structure is essential to provide a binding site to recruit the Tat⁵ and cyclin T1 regulatory proteins to TAR; thus these motions likely underlie the functional rearrangements critical for transcription and thus for HIV replication.⁸ The motions of U38 are thus quite likely to be essential as well, because they correspond to the helical domain motions allowing TAR to sample the manifold of conformational states observed in its complexes with small molecules and, in particular, with Tat.² Global reorientation of the upper helix, as monitored at U38, occurs on a very similar time scale to the hopping of U23, suggesting

(21) Puglisi, J. D.; Tan, R.; Calnan, B. J.; Frankel, A. D.; Williamson, J. R. *Science* **1992**, *257*, 76–80.

(22) Lorieau, J. L.; McDermott, A. E. *J. Am. Chem. Soc.* **2006**, *128*, 11505–11512.

(23) Leulliot, N.; Varani, G. *Biochemistry* **2001**, *40*, 7947–7956.




Figure 4. (a) Residue U38 (green) is represented inside a pseudocylindrical plane (blue) representative of the upper helical domain of TAR, while the lower pseudocylindrical plane (red) represents the lower helix. (b) The motions of U38 correspond to a 13° twist and 13° bend of the entire upper helix, in addition to much faster $\pm 4^\circ$ librational motions of the base.




Figure 5. (a) In the absence of ligands, the three residues in TAR experience a wide range of conformations (indicated by the clouds colored according to Figure 1), which include the conformation observed in the complex with peptides derived from Tat protein. The two conformations of TAR are represented as the two alternative structures of U23 (in teal) and U25 (green); the complex with a 37-mer peptide derived from Tat protein is shown in (b), emphasizing the conformational capture mechanism of the Tat protein.

that these motions may be correlated and, together, may create the unique RNA structure required to recruit Tat protein to TAR.

Materials and Methods

Preparation of RNA Samples for Solid-State Investigation.

[5,6- ^2H]-labeled uridine nucleotides for deuterium dynamics experiments were prepared in house using well-established methods.²⁴ Conversion of the deuterated uridine nucleotide to phosphoramidites for RNA synthesis was done by Dharmacon (Lafayette, CO). RNA oligonucleotides were deprotected and desalted by Dharmacon. Each sample was checked for homogeneity using analytical denaturing polyacrylamide gel electrophoresis and was used without further purification. The following three TAR RNA

29-mer constructs were used: 5'-GGCCAGA- $^{2\text{F}}\text{U}$ -CUG(pS)AGCGAAAGC*UCUCUGGCC-3' ("U38"), 5'-GGCCAGA*U- $^{2\text{F}}\text{C}$ -UG(pS)AGCGAAAGCUCUCUGGCC-3' ("U23"), and 5'-GGC-CAGA- $^{2\text{F}}\text{U}$ -C*UG(pS)AGCGAAAGCUCUCUGGCC-3' ("U25").

Here, pS indicates a phosphorothioate label, $^{2\text{F}}\text{U}$ - or $^{2\text{F}}\text{C}$ -indicates a 2' fluorine substitution, and *U indicates a [5,6- ^2H] base-labeled uridine. Previous work has demonstrated that phosphorothioate and 2'-fluorine substitutions at those positions do not perturb the structure of TAR.^{25–27} These labels were added to allow us to verify the conformation of TAR RNA using REDOR

(25) Cruse, W. B. T.; Salisbury, S. A.; Brown, T.; Cosstick, R.; Eckstein, F.; Kennard, O. *J. Mol. Biol.* **1986**, *192*, 891–905.

(26) Merritt, M. E.; Sigurdsson, S. T.; Drobny, G. P. *J. Am. Chem. Soc.* **1999**, *121*, 6070–6071.

(27) Olsen, G. L.; Edwards, T. E.; Deka, P.; Varani, G.; Sigurdsson, S. T.; Drobny, G. P. *Nucleic Acids Res.* **2005**, *33*, 3447–54.

(24) Huang, X.; Yu, P.; LeProust, E.; Gao, X. *Nucleic Acids Res.* **1997**, *25*, 4758–4763.

methods.²⁷ Although perturbations to the dynamics arising from these modifications cannot be ruled out completely, the combination of the high degree of structural similarity between wild-type and substituted TAR, as well as our focus on base motions, suggest that perturbations to the base dynamics by these substitutions are not substantial.

NMR Sample Preparation. 5.70 μmol of the U38 oligonucleotide was dissolved in buffer (50 mM NaCl, 10 mM sodium cacodylate, pH 5.5), then frozen using liquid nitrogen and lyophilized (final composition 10% NaCl, and 4.7% cacodylate respectively, by weight upon lyophilization). To remove residual H₂O and D₂O, samples were redissolved in deuterium-depleted water, and then lyophilized 3–4 times. Each sample was then redissolved a final time in deuterium-depleted water, annealed in a 90 °C water bath for 5 min, allowed to cool to room temperature for 30 min, and then frozen using liquid nitrogen. All samples were again lyophilized extensively to remove residual water (~80 mTorr, 7–15 days) before transfer to the NMR sample rotor. The dry sample was packed into a 5 mm Kel-F sample chamber, and re-lyophilized to remove water absorbed during sample packing.

Samples were hydrated to 16 waters per nucleotide in sealed chambers over saturated salt solutions containing deuterium-depleted water.²⁸ Water addition to the samples was monitored gravimetrically. Samples were then sealed and allowed to equilibrate an additional week before use in NMR experiments. The hydration levels were chosen after carefully conducting a thorough investigation of the line shape as a function of water content.¹⁸ Water introduced to salted nucleic acid samples progressively populates the grooves, bases, and backbone in a well-characterized fashion, beginning with the phosphates and eventually reaching limiting local hydration so that subsequent addition of water contributes primarily to the surrounding water shells.²⁹ At $W = 16$, local hydration of the nucleic acid backbone and bases is substantially complete.^{18,29–31} Following that hydration level, the solid-state NMR spectra display much smaller changes indicative of relatively small changes in dynamics between $W = 16$ and the highest hydration we have investigated, $W = 30$.¹⁸

Samples for analysis of U23 (4.3 μmol) and U25 (6.6 μmol) were prepared using the procedure described above, with different sample sizes as indicated.

Solid-State NMR Experiments. All experiments were performed at 11.7 T field (76.76 MHz deuterium resonance frequency), using a home-built single-channel static deuterium probe. Pulses were generated using an AMT (American Microwave Technologies) M3446 (1 kW and 10–130 MHz) amplifier. Typical 90° pulse times were 2.0–2.4 μs . All measurements were made at room temperature (26 °C), with continuous application of 20 °C cooling air. 5 mm kel-F sample rotors were made in-house. Dwell time for all experiments was 0.2 μs . Recycle delay for all experiments was 1 s.

Deuterium line shapes were recorded using a quadrupole echo pulse sequence³² ($90^\circ_x - \tau - 90^\circ_y - \tau - \text{echo}$), where the second half of the echo signal was acquired and Fourier-transformed following left-shift of all time-domain data sets to their respective echo maxima. Quadrupole echo experiments used eight-step phase cycling, with an echo delay of 40 μs . Typical spectra resulted from acquisition of 300 000–500 000 scans, collected as 80 000-scan data sets, then coadded.

Spin–lattice relaxation times were determined using saturation-recovery experiments. An initial train of five 90_x pulses at 2 ms intervals was used, followed by a variable relaxation delay and

quadrupole echo detection ($(90^\circ_x)n - \text{delay} - 90^\circ_x - \tau - 90^\circ_y - \tau - \text{echo}$; $n = 5$). Typical delay times for hydrated samples were 1, 100, 500, 1000, and 2000 ms.

T_{1Q} relaxation times were determined using the broadband Jeener–Broekaert pulse sequence,^{33,34} with eight-step phase cycling. The pulse sequence used is as follows: $90 - \tau_1 - 67 - \tau_1 - 45 - \tau_2 - 45 - [\text{variable relaxation delay}] - 45 - \text{echo delay} - 90 - \text{echo delay} - \text{acquisition}$. Typical delays were 5 μs (τ_1), 2.5 μs (τ_2), and 1, 100, 500, 1000, and 2000 ms (variable relaxation delay), with 40 μs echo delays.

Data Processing. Data processing was performed using in-house written NMR processing software. Data acquisition was begun prior to the echo maximum for all experiments, and all time-domain data sets were left-shifted prior to Fourier Transformation. Line shape spectra were multiplied with an exponential corresponding to 500 kHz line broadening. Relaxation data were analyzed using a nonlinear least-squares fit to a single exponential decay to obtain powder-averaged spin–lattice relaxation times.

Modeling RNA Dynamics. To construct the models, we make the assumption that the motions are composed of independent components, which can be represented using multi-axis models. This permits a representation of the complex motions in which local motions are represented by jumps of the principal axis system with respect to an intermediate frame, while conformational exchange jumps of the bulge residues or collective motions of the helix (which would carry the U38:A27 pair) are treated by jumps of this intermediate frame with respect to the crystal frame. Motions of the C–D bond vector are then described by two successive independent transformations, where one set of angles describes the local motion and a second set describes the motions of the intermediate frame, representing a conformational exchange motion in each model.

Motional models for each site were constructed on the basis of known structural constraints. Each base is confined to move within a steric “box” defined by surrounding residues, which limit its available motional space: U38 local motion, for example, is restricted to positions between base pairs above and below the A27:U38 pair; motions of this local jumping frame itself thus represent motions of the upper helix. For bulge residues U23 and U25, the picture is more complicated. These two residues reside in the single-stranded bulge linking two helices, and thus both should be directly dependent, both structurally and dynamically, on the motions of the adjacent helices. This dependence should be transmitted by fluctuations of the sugar–phosphate backbone, involving numerous torsion angles. Motions for U23, however, must be sterically bounded from below by A22 and from above by the upper helix; thus we can treat motions of this site as delimited by its stacking interactions atop A22 and base-pairing interactions with A27:U38 of the upper helix, while restricting lateral motion of the residue to rotations of the bulge backbone. Therefore, to construct a model for the conformational exchange motions of U23, we first extracted coordinates for the C1′–N1 bond vector in bound and unbound TAR (1ANR, 1ARJ), to define likely end points for conformational exchange. Vectors representing the C1′–N1 bond axis were defined with respect to a frame attached to the A27:U38 base pair for each TAR conformation; these were then used to construct initial-guess jump sites representing the conformational exchange motion. Remarkably, refinement of this initial model to fit the observed solid-state NMR data required a reduction in jump amplitude of only 3°.

The motions of residue U25 are not nearly so well-defined, and this is reflected in the models we report here. It has been shown that U25 can be substituted by a short linker without significant impairment of TAR function, suggesting that it functions largely as a spacer residue. Solution NMR structures suggest that, while

(28) Weast, R. C. *CRC Handbook of Chemistry and Physics*, 60th ed.; CRC Press: Boca Raton, FL, 1979; p E-46.

(29) Falk, M.; Hartman, K. A.; Lord, R. C. *J. Am. Chem. Soc.* **1962**, *84*, 3843–3846.

(30) Wang, A. C.; Kennedy, M. A.; Reid, D. R.; P., D. G. *J. Magn. Reson., Ser. B* **1994**, *105*, 1–10.

(31) Schurr, J. M.; Fujimoto, B. S.; Diaz, R.; Robinson, B. H. *J. Magn. Reson.* **1999**, *140*, 404–431.

(32) Davis, J. H. *Biophys. J.* **1979**, *27*, 339–358.

(33) Wimperis, S.; Bodenhausen, G. *Chem. Phys. Lett.* **1986**, *132*, 194–199.

(34) Hoatson, G. L. *J. Magn. Reson.* **1990**, *94*, 152–159.

motions of this residue must be bounded from above by adjacent residue G26 and laterally by the conformation of the adjacent bulge backbone, it is not restrained by either stacking or base-pairing interactions. Because the bulge is known to be extruded from between the two helices during tat recognition, it would be expected that motions of U25 would be relatively unrestricted and determined largely by those of the local backbone alone.³⁵ This is consistent with the large-amplitude unrestrained motions we have used to model the U25 data.

Angular excursions corresponding to structural transitions bounded by those constraints were represented as site geometries and used to construct multi-axis jump models. Line shape and T_{1Z} simulations were then performed using the programs MXET1,³⁶ EXPRESS,³⁷ and Turbopowder.³⁸ Fitting of the model trajectories against the experimental results was performed iteratively, by systematically varying jump rates and trajectories (amplitudes of motions) in the simulations until a satisfactory fit was obtained (examples are shown in the Supporting Information).

To generate the corresponding T_{1Q} values, model parameters were input to correlation functions derived from the formalism of Torchia and Szabo,³⁹ specific to the motional model derived for each site. Relaxation times corresponding to the resulting spectral densities for each model were then obtained using the following relationships:^{40,41}

$$R_{1Z} = \frac{1}{T_{1Z}} = \frac{3\pi^2}{2} \left(\frac{e^2qQ}{h} \right)^2 [J(\omega_o) + 4J(2\omega_o)] \quad (1)$$

$$R_{1Q} = \frac{1}{T_{1Q}} = \frac{9\pi^2}{2} \left(\frac{e^2qQ}{h} \right)^2 [J(\omega_o)] \quad (2)$$

where $((e^2qQ)/h)$ is the static quadrupolar coupling constant.

The requirement that candidate motional models fit the combined relaxation and line shape data restricts possible motional models drastically. Separately, relaxation or line shape data for each site could typically be fit by more than one model; however, systematic grid searches of the parameter space revealed that, for U38,

concatenated two-site jump models simultaneously consistent with both the observed line shape and the relaxation data could be restricted to a single motional model. For U23, we are able to identify two models consistent with the data. In addition to the model described above, a second model in which the motional rates are approximately reversed can also reproduce the data. In such a scenario, however, the rate of the conformational exchange motion must be increased to $2.9 \times 10^{10} \text{ s}^{-1}$ (and the local twisting about the glycosidic bond is slowed to $5.4 \times 10^7 \text{ s}^{-1}$). Given the large amplitude of the deformations of the backbone required for conformational exchange in TAR, such a rapid rate for this component of the motion is physically implausible. Moreover, it is also unlikely that the local motions within the respective potential minima at each end of the exchange jump should be so slow.

While the fit for U25 is less satisfactory, as seen in Figure S3, this model too is tightly constrained by the data. Given the lack of restraints on the dynamics of this residue, and the resulting absence of a well-defined potential landscape, definition of more finely detailed motional trajectories than that which we provide here would be difficult to support. Obtaining a full representation of the more complicated motions undergone by U25 would likely require incorporation of additional motional axes. Thus far, our attempts to add a third dynamic axis have not produced improvement in the fits for this residue.

The goodness of fit is shown by χ^2 plots for fits to the combined T_{1Z} and T_{1Q} relaxation data for each of the three sites, as shown in Supporting Information Figures S1–S3 for each of the three residues. Figure S4 shows mean quadrupole echo amplitudes for the three sites at 16 waters per nucleotide.

Acknowledgment. This work was supported by grants from the NSF (MCB-0642253) and NIH (RO1-EB03152) to G.P.D. and G.V. We are grateful to Dr. Zahra Shajani for many stimulating discussions and to members of the Varani research group for reading the manuscript.

Supporting Information Available: Illustration of the fitting procedure used to obtain the models presented in the main text. χ^2 plots for fits to the combined T_{1Z} and T_{1Q} relaxation data for each of the three sites, as a function of model parameters (rates or amplitudes), and quadrupole echo amplitudes for the three sites at 16 waters per nucleotide. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JA907515S

- (35) Aboul-ela, F.; Varani, G. *J. Mol. Struct.* **1998**, *423*, 29–39.
 (36) Greenfield, M. S.; Ronemus, A. D.; Vold, R. L.; Vold, R. R.; Ellis, P. D.; Raidy, T. E. *J. Magn. Reson.* **1987**, *72*, 89–107.
 (37) Vold, R. L.; Hoatson, G. L. *J. Magn. Reson.* **2009**, *198*, 57–72.
 (38) Wittebort, R. J.; Olejniczak, E. T.; Griffin, R. G. *J. Chem. Phys.* **1987**, *86*, 5411–5420.
 (39) Torchia, D. A.; Szabo, A. *J. Magn. Reson.* **1982**, *49*, 107–121.
 (40) Palmer, A. G., III; Williams, J.; McDermott, A. *J. Phys. Chem.* **1996**, *100*, 13293–13310.
 (41) Alam, T. M.; Drobny, G. P. *Chem. Rev.* **1991**, *91*, 1545–1590.