

The Cold Denatured State of the C-terminal Domain of Protein L9 Is Compact and Contains Both Native and Non-native Structure

Bing Shan,[†] Sebastian McClendon,[‡] Carla Rospigliosi,[‡] David Eliezer,^{*,‡} and Daniel P. Raleigh^{*,†,§}

Department of Chemistry, State University of New York at Stony Brook, Stony Brook, New York 11794-3400, Department of Biochemistry and Program in Structural Biology, Weill Cornell Medical College, New York, New York 10065, and Graduate Program in Biochemistry and Structural Biology, and Graduate Program in Biophysics, State University of New York at Stony Brook, Stony Brook, New York 11794

Received September 23, 2009; E-mail: draleigh@notes.cc.sunysb.edu; dae2005@med.cornell.edu

Abstract: Cold denaturation is a general property of globular proteins, and the process provides insight into the origins of the cooperativity of protein folding and the nature of partially folded states. Unfortunately, studies of protein cold denaturation have been hindered by the fact that the cold denatured state is normally difficult to access experimentally. Special conditions such as addition of high concentrations of denaturant, encapsulation into reverse micelles, the formation of emulsified solutions, high pressure, or extremes of pH have been applied, but these can perturb the unfolded state of proteins. The cold denatured state of the C-terminal domain of the ribosomal protein L9 can be populated under native-like conditions by taking advantage of a destabilizing point mutation which leads to cold denaturation at temperatures above 0 °C. This state is in slow exchange with the native state on the NMR time scale. Virtually complete backbone ¹⁵N, ¹³C, and ¹H as well as side-chain ¹³C_β and ¹H_β chemical shift assignments were obtained for the cold denatured state at pH 5.7, 12 °C. Chemical shift analysis, backbone N–H residual dipolar couplings, amide proton NOEs, and R₂ relaxation rates all indicate that the cold denatured state of CTL9 (the C-terminal domain of the ribosomal protein L9) not only contains significant native-like secondary structure but also non-native structure. The regions corresponding to the two native α-helices show a strong tendency to populate helical Φ and Ψ angles. The segment which connects α-helix 2 and β-strand 2 (residues 107–124) in the native state exhibits a significant preference to form non-native helical structure in the cold denatured state. The structure observed in the cold denatured state of the I98A mutant is similar to that observed in the pH 3.8 unfolded state of wild type CTL9 at 25 °C, suggesting that it is a robust feature of the denatured state ensemble of this protein. The implications for protein folding and for studies of cold denatured states are discussed.

Introduction

Cold denaturation is a general property of globular proteins, as predicted by the Gibbs–Helmholtz equation, and is believed to be driven by the increased hydration of nonpolar groups of proteins at low temperatures, as well as by the decrease in hydrophobic interactions within the protein.^{1–4} Thus the cold induced unfolding of a protein is sometimes thought to proceed via a different mechanism from thermal unfolding. The process of cold denaturation has been studied for a wide range of proteins, but structural characterization of the cold unfolded state is less extensive than studies of

heat or denaturant induced unfolded states. Investigating the structure of the cold denatured state and the cooperativity of cold denaturation is crucial for a complete understanding of protein folding and cooperativity.^{3,5,6} An increased knowledge of the cold denatured state is also important for understanding the molecular basis for cold adaptation in psychrophilic organisms.^{7,8} Most studies of cold unfolding have focused on thermodynamic aspects of the process or have used relatively low resolution spectroscopic methods.

Unfortunately, the study of cold denaturation is difficult because the midpoint of the cold unfolding transition is typically well below 0 °C.^{3,4} As a consequence studies have been limited to a relatively small number of proteins compared to studies of thermal unfolding. Furthermore the majority of cases require

[†] Department of Chemistry, State University of New York at Stony Brook.

[‡] Weill Cornell Medical College.

[§] Graduate Program in Biochemistry and Structural Biology, and Graduate Program in Biophysics, State University of New York at Stony Brook.

(1) Clark, J. H. *J. Gen. Physiol.* **1945**, *28*, 539–545.
 (2) Brandts, J. F. *J. Am. Chem. Soc.* **1964**, *86*, 4291–4301.
 (3) Privalov, P. L. *Crit. Rev. Biochem. Mol. Biol.* **1990**, *25*, 281–305.
 (4) Privalov, P. L.; Gill, S. J. *Adv. Protein Chem.* **1988**, *39*, 191–234.

(5) Sanfelice, D.; Tancredi, T.; Politou, A.; Pastore, A.; Temussi, P. A. *J. Am. Chem. Soc.* **2009**, *131*, 11662–11663.

(6) Martin, S. R.; Esposito, V.; Rios, P. D. L.; Pastore, A.; Temussi, P. A. *J. Am. Chem. Soc.* **2008**, *130*, 9963–9970.

(7) Lonhienne, T.; Gerday, C.; Feller, G. *Biochim. Biophys. Acta* **2000**, *1543*, 1–10.

(8) Feller, G.; Gerday, C. *Nat. Rev. Microbiol.* **2003**, *1*, 200–208.




Figure 1. The I98A mutant undergoes cold denaturation. (A) Ribbon diagram of CTL9 (residues 58–149 of protein L9), Protein data bank entry 1DIV. The hydrophobic core residue I98 as well as the N and C-termini are labeled. The ribbon diagram was made using PyMol. (B) Thermal denaturation curve of I98A CTL9 monitored by CD at 222 nm in H₂O at pH 5.7. (C) 1D ¹H NMR spectra of I98A CTL9 at different temperatures in 100% D₂O at pD 6.0 (uncorrected pH meter reading). Only the aromatic region is shown, resonance assignments are labeled.

special conditions such as encapsulation,^{9–11} high pressure,^{12–14} emulsified samples,¹⁵ extremes of pH,^{2–4,16} or the addition of high concentrations of denaturant.^{2,3,17} All of these methods have the potential to offer insight into the structural and thermodynamic aspects of cold denaturation, although some approaches may perturb the system in complicating ways. For example, encapsulation in reverse micelles has been used to study cold unfolding, but our previous NMR diffusion measurements have shown that cold denatured proteins expand as the temperature is lowered¹⁸ and other groups have shown that under certain circumstances the reverse micelle systems used can shed water and contract as the temperature is decreased.^{9,15,19,20} The combination could lead to significant protein micelle interactions at low temperatures in certain cases. There have been very few reported high resolution studies of the cold denatured state in the absence of encapsulation, denaturant, or extremes of pH or

pressure.²¹ Here, we report the detailed characterization of the structural and dynamic properties of the cold denatured state using NMR spectroscopy. We use a destabilizing point mutant of the C-terminal domain of the ribosomal protein L9 (CTL9) to populate the cold unfolded state in homogeneous solution in the absence of denaturants at near neutral pH. The implications for the folding of CTL9 are discussed as are the implications for the nature of the cold denatured state.

CTL9 is a 92 residue globular protein which adopts a relatively rare fold made up of two α -helices and a three stranded mixed parallel, antiparallel β -sheet (Figure 1A).²² An I98A (the isoleucine to alanine point mutant of CTL9 at residue 98) point mutation destabilizes the protein and reduces ΔH° of unfolding at the midpoint of the thermal unfolding while increasing ΔC_p° for unfolding. As expected, this increases the temperature of cold unfolding and, in the present case, leads to observable cold denaturation at temperatures above 0 °C in the absence of denaturant or extreme pH values¹⁸ (Figure 1B and 1C). Data presented here together with earlier hydrodynamic measurements show that the mutation does not alter the overall structure of the protein.¹⁸ We have characterized the hydrodynamic properties of the cold induced unfolded state of the I98A mutant of CTL9 under native and near native conditions using pulsed field gradient NMR diffusion experiments, and we have shown that the cold denatured state is compact relative to the acid or urea unfolded states but expands as the temperature is lowered.¹⁸

(9) Van Horn, W. D.; Simorellis, A. K.; Flynn, P. F. *J. Am. Chem. Soc.* **2005**, *127*, 13553–13560.

(10) Babu, C. R.; Hilsner, V. J.; Wand, A. J. *Nat. Struct. Mol. Biol.* **2004**, *11*, 352–357.

(11) Pometun, M. S.; Peterson, R. W.; Babu, C. R.; Wand, A. J. *J. Am. Chem. Soc.* **2006**, *128*, 10652–10653.

(12) Jonas, J. *Biochim. Biophys. Acta* **2002**, *1595*, 145–159.

(13) Kunugi, S.; Tanaka, N. *Biochim. Biophys. Acta* **2002**, *1595*, 329–344.

(14) Kitahara, R.; Okuno, A.; Kato, M.; Taniguchi, Y.; Yokoyama, S.; Akasaka, K. *Magn. Reson. Chem.* **2006**, *44*, S108–S113.

(15) Davidovic, M.; Mattea, C.; Qvist, J.; Halle, B. *J. Am. Chem. Soc.* **2009**, *131*, 1025–1036.

(16) Privalov, P. L.; Griko, Y. V.; Venyaminov, S. Y.; Kutysenko, V. P. *J. Mol. Biol.* **1986**, *190*, 487–498.

(17) Wong, K. B.; Freund, S. M. V.; Fersht, A. R. *J. Mol. Biol.* **1996**, *259*, 805–818.

(18) Li, Y.; Shan, B.; Raleigh, D. P. *J. Mol. Biol.* **2007**, *368*, 256–262.

(19) Van Horn, W. D.; Ogilvie, M. E.; Flynn, P. F. *J. Am. Chem. Soc.* **2009**, *131*, 8030–8039.

(20) Simorellis, A. K.; Van Horn, W. D.; Flynn, P. F. *J. Am. Chem. Soc.* **2006**, *128*, 5082–5090.

(21) Pastore, A.; Martin, S. R.; Politou, A.; Kondapalli, K. C.; Stemmler, T.; Temussi, P. A. *J. Am. Chem. Soc.* **2007**, *129*, 5374–5375.

(22) Hoffman, D. W.; Cameron, C. S.; Davies, C.; White, S. W.; Ramakrishnan, V. *J. Mol. Biol.* **1996**, *264*, 1058–1071.

Materials and Methods

Protein Expression and Purification. ^{15}N -labeled CTL9 and ^{13}C , ^{15}N -labeled CTL9 and the labeled I98A mutant were overexpressed and purified as previously described.^{23,24} The identity of the proteins was confirmed using MALDI-TOF mass spectroscopy, and the purity was checked using analytical HPLC. The yield of pure protein was 30 mg per liter of M9 minimal media.

Circular Dichroism Spectroscopy. CD experiments were carried out using a Chirascan CD spectrometer. The protein was dissolved in 20 mM acetate, 100 mM NaCl buffer, pH 5.7 at a concentration of $\sim 18 \mu\text{M}$. Thermal denaturation experiments were performed by monitoring the ellipticity at 222 nm with a heating rate of $1 \text{ }^\circ\text{C}/\text{min}$. Data were collected from 2 to $94 \text{ }^\circ\text{C}$, with a $2 \text{ }^\circ\text{C}$ increment. The experiments were repeated 3 times with independently prepared samples to access the reproducibility of the results and the experimental uncertainties. Additional CD melting experiments were conducted as a function of pH between pH 4.7 and pH 8.0.

Nuclear Magnetic Resonance Experiments. Protein samples for NMR experiments were prepared in 90% H_2O and 10% D_2O at pH 5.7, with 20 mM sodium acetate and 100 mM NaCl, unless otherwise specified. The protein concentration was $\sim 1 \text{ mM}$. All NMR experiments for the cold denatured state of I98A CTL9 were recorded on 800 or 900 MHz Bruker spectrometers equipped with cryogenic probes at the New York Structural Biology Center. Triple resonance experiments for establishment of the assignments for wild-type CTL9 were performed on a 600 MHz Varian spectrometer at the University of Connecticut Health Center. The ^1H dimension was centered at the water resonance, and the ^{15}N offset frequency was set to 118.0 ppm for all heteronuclear NMR experiments.

^{15}N - ^1H correlated heteronuclear single coherence (HSQC) spectra for the I98A mutant were recorded using 1024×512 complex points with 8 scans per increment and spectral widths of 9920.6 and 2189.6 Hz for the ^1H and ^{15}N dimensions, respectively. The HSQC spectra of wild-type CTL9 sample had spectral widths of 6600.7 Hz (^1H) and 1920.0 Hz (^{15}N), respectively.

The HNC0/HNCAC0^{25–27} and the HNCACB^{28,29} experiments were performed to enable sequence-specific backbone and side chain $^{13}\text{C}_\beta$ assignments. The HBHACONH^{29,30} experiment was carried out to obtain side chain $^1\text{H}_\alpha$ and $^1\text{H}_\beta$ assignments. The watergate sequence was employed to suppress water signals in all triple resonance experiments. All spectra were processed using NMRPipe software,³¹ and chemical shift assignments were made using NMRViewJ.³² All the chemical shifts were referenced to the absolute frequency of DSS at 0.00 ppm. The random coil chemical shift values of Wishart were used to calculate the secondary chemical shifts.³³ SSP (secondary structure propensity) scores were calculated based on $^{13}\text{C}_\alpha$, $^{13}\text{C}_\beta$, and $^1\text{H}_\alpha$ shifts, using the software provided by Professor Forman-Kay at her Website (<http://pound.med.utoronto.ca/software.html>).³⁴

A 3D ^{15}N HSQC-NOESY-HSQC experiment was performed on a ^{15}N -labeled sample of I98A CTL9. The spectrum was acquired

with spectral widths of 8012.8 (NH), 2108.8 (^{15}N), and 2108.8 (^{15}N) Hz and in a data matrix of $1024 \times 128 \times 138$ complex points, with a mixing time of 300 ms. Coherence selection with gradients and sensitivity enhancement were used, and water suppression was achieved using a water flip back pulse sequence. NOESY peak volumes were measured using NMRViewJ. The NOE peak volumes were normalized as the ratio of the $NN(i, i+1)$ cross peak to the $NN(i, i)$ diagonal peaks to compensate for any potential relaxation effects on the NOE intensities.

^{15}N - ^1H one bond residual dipolar couplings (RDCs) were measured in weakly aligned bicelle media. The bicelles were prepared by mixing $450 \mu\text{L}$ of NMR buffer with $50 \mu\text{L}$ of pentaethylene glycol octylether (C8E5). The mixture was gently vortexed until the solution became clear. Then 1-octanol was added in $2 \mu\text{L}$ increments for a total additional volume of 10 to $11 \mu\text{L}$. The bicelles were left at $10 \text{ }^\circ\text{C}$ overnight. Then the protein solution was added at a ratio of 1:1 to the bicelle solution and gently vortexed. The alignment of the protein with the bicelles was checked by measuring the ^2H quadrupolar splitting of the water line. A ^{15}N - ^1H HSQC experiment was performed on the protein sample in the bicelles. There was no significant change in either the chemical shift or peak intensity observed in any of the resonances, indicating the bicelles do not perturb the protein conformation significantly. To measure the ^{15}N - ^1H RDCs, 2D inphase- and antiphase-HSQC (IPAP-HSQC) spectra³⁵ were recorded for the sample in the bicelles and in isotropic buffer solutions. The IPAP-HSQC spectra were acquired with 2048×512 complex points and with spectral widths of 8012.8 and 2270.7 Hz for the ^1H and ^{15}N dimension, respectively. The spectra were processed using NMRPipe software, and the ^{15}N - ^1H splittings were measured using NMRViewJ. The RDCs were calculated as the difference between the scalar coupling splitting in isotropic solution and the observed splitting in the anisotropic solution.

Transverse relaxation (R_2) experiments were performed on a ^{15}N -labeled sample as described using a Carr–Purcell–Meiboom–Gill (CPMG) sequence.³⁶ The pulse spacing in the CPMG sequence was 0.9 ms. The spectra were recorded at 10 delay times: 16.96, 33.92, 50.88, 67.84, 84.8, 101.76, 118.72, 135.68, 152.64, and 169.6 ms, with a recycle delay of 3 s. To estimate the uncertainty, duplicate spectra were acquired at 50.88, 101.76, and 152.64 ms delay times. Each spectrum was collected with 4 scans and 1024×256 complex points. The spectral widths were 8012.8 Hz (^1H) and 2027.7 (^{15}N). R_2 rates were determined by fitting the NMR resonance intensities as a function of delay times to a single exponential decay using NMRViewJ. The expected intrinsic R_2 rates for a completely unstructured protein were obtained by fitting the experimental R_2 rates to the phenomenological model proposed by Schwalbe and co-workers.³⁷

Results

The I98A mutant of CTL9 adopts the same structure as wild-type CTL9 but undergoes cold denaturation above $0 \text{ }^\circ\text{C}$. Our previous thermodynamic studies indicated that replacing the hydrophobic core residue I98 in CTL9 by an alanine causes a decrease in $\Delta H^\circ(T_m)$ and an increase in ΔC_p° , thus the temperature of the midpoint of cold denaturation increases accordingly.¹⁸ At pH values near 6.0 cold denaturation was observed at temperatures above $0 \text{ }^\circ\text{C}$ by both CD monitored temperature melting experiments and by NMR monitored melting studies (Figure 1B and 1C and Supporting Information). As shown in Figure 1C, resonances from both the native and

(23) Sato, S.; Raleigh, D. P. *J. Mol. Biol.* **2002**, *318*, 571–582.

(24) Shan, B.; Bhattacharya, S.; Eliezer, D.; Raleigh, D. P. *Biochemistry* **2008**, *47*, 9565–9573.

(25) Grzesiek, S.; Bax, A. *J. Magn. Reson.* **1992**, *96*, 432–440.

(26) Kay, L. E.; Xu, G. Y.; Yamazaki, T. *J. Magn. Reson. A* **1994**, *109*, 129–133.

(27) Clubb, R. T.; Thanabal, V.; Wagner, T. *J. Magn. Reson.* **1992**, *97*, 213–217.

(28) Wittekind, M.; Mueller, L. *J. Magn. Reson. B* **1993**, *101*, 201–205.

(29) Muhandiram, D. R.; Kay, L. E. *J. Magn. Reson. B* **1994**, *103*, 203–216.

(30) Grzesiek, S.; Bax, A. *J. Biomol. NMR* **1993**, *3*, 185–204.

(31) Delaglio, F.; Grzesiek, S.; Vuister, G. W.; Zhu, G.; Pfeifer, J.; Bax, A. *J. Biomol. NMR* **1995**, *6*, 277–293.

(32) Johnson, B. A. *Methods Mol. Biol.* **2004**, *278*, 313–352.

(33) Wishart, D. S.; Bigam, C. G.; Holm, A.; Hodges, R. S.; Sykes, B. D. *J. Biomol. NMR* **1995**, *5*, 67–81.

(34) Marsh, J. A.; Singh, V. K.; Jia, Z.; Forman-Kay, J. D. *Protein Sci.* **2006**, *15*, 2795–2804.

(35) Ottiger, M.; Delaglio, F.; Bax, A. *J. Magn. Reson.* **1998**, *131*, 373–378.

(36) Farrow, N. A.; Zhang, O.; Forman-Kay, J. D.; Kay, L. E. *Biochemistry* **1995**, *34*, 868–878.

(37) Schwalbe, H.; Fiebig, K. M.; Buck, M.; Jones, J. A.; Grimshaw, S. B.; Spencer, A.; Glaser, S. J.; Smith, L. J.; Dobson, C. M. *Biochemistry* **1997**, *36*, 8977–8991.




Figure 2. ^{15}N -HSQC spectrum of the cold denatured state of I98A CTL9 with the assignments indicated. The spectrum was recorded at pH 5.7, 12 °C.

denatured state can be detected at 25 °C. As the temperature decreases, the resonances of the cold denatured state increase while the peaks from the native state diminish gradually. Similar to the wild-type protein, the stability of I98A CTL9 is strongly dependent on pH due to the protonation of one or more of the three histidine side chains.²³ At pH 5.7, the cold denatured state is 79% populated at 12 °C, making NMR studies of the cold denatured state feasible. Thus, we chose pH 5.7, 12 °C as experimental conditions to explore the structural and dynamic properties of the cold denatured state of I98A CTL9.

The HSQC spectrum of the cold denatured I98A CTL9 is typical for an unfolded protein (Figure 2), and one major set of peaks are observed, indicating that the cold denatured state is the dominant conformer, a result which is in agreement with the 1D NMR experiments. Peaks from the native state are observed at lower contour levels. All of the peaks are sharp as expected for a monomeric protein, which is not experiencing exchange broadening. We used pH jump fluorescence detected stopped flow measurements to directly measure the exchange rate at 12 °C (Supporting Information). The exchange rate, $k_{\text{ex}} = k_{\text{folding}} + k_{\text{unfolding}}$ is 0.49 s^{-1} , confirming that the system is in slow exchange on the chemical shift time scale. The folding rate is $8.8 \times 10^{-2} \text{ sec}^{-1}$ and the unfolding rate is 0.40 s^{-1} , corresponding to life times of 11 s and 2.5 s for the cold denatured state and the native state, respectively.

Using triple resonance HNCACB, HNCO, and HNCACO experiments, virtually complete backbone ^{13}C , ^1H , and ^{15}N assignments were obtained. Side chain $^1\text{H}_\alpha$ and $^1\text{H}_\beta$ assignments were made using the HBHACONH experiment. The assignments have been deposited in the Biological Magnetic Resonance data Bank (BMRB).

Chemical shift analysis provides evidence of native and non-native secondary structure in the cold denatured state. Chemical shifts are valuable probes of protein structure because of the strong correlation between secondary chemical shifts, i.e., the deviations of observed chemical shifts from random coil

Table 1. Average Secondary Shifts and SSP Values in Secondary Structural Elements of Wild-Type CTL9 and I98A CTL9

	wild-type CTL9 at pH 5.7, 12 °C					
	$^1\text{H}_\alpha$	$^{13}\text{C}_\alpha$	$^{13}\text{C}_\beta$	^{13}CO	$^{13}\text{C}_\alpha - ^{13}\text{C}_\beta$	SSP
Helix 1	-0.35	2.80	-0.67	1.95	3.47	0.91
Helix 2	-0.28	2.91	0.70	1.90	2.21	0.78
Loop	-0.09	0.88	0.67	0.08	0.21	0.09
Strand 1	0.16	-0.51	1.28	-0.89	-1.79	-0.51
Strand 2	0.50	0.04	1.57	-0.98	-1.53	-0.42
Strand 3	0.48	-1.05	1.30	-1.51	-2.35	-0.28
	I98A CTL9 at pH 5.7, 12 °C					
	$^1\text{H}_\alpha$	$^{13}\text{C}_\alpha$	$^{13}\text{C}_\beta$	^{13}CO	$^{13}\text{C}_\alpha - ^{13}\text{C}_\beta$	SSP
Helix 1	-0.10	1.23	-0.03	0.76	1.26	0.33
Helix 2	-0.08	0.96	-0.05	0.83	1.01	0.25
Loop	-0.08	0.87	-0.04	0.43	0.91	0.20
Strand 1	-0.01	0.06	0.23	-0.65	-0.17	-0.17
Strand 2	-0.03	0.27	0.23	-0.83	0.04	-0.18
Strand 3	-0.01	0.23	0.26	-0.33	-0.03	0.00

chemical shifts, with protein structure. $^{13}\text{C}_\alpha$ and ^{13}CO shifts are shifted downfield in α -helices and upfield in β -sheets, while the opposite trend is usually observed for $^1\text{H}_\alpha$ and $^{13}\text{C}_\beta$ shifts.^{38–40} The $^{13}\text{C}_\alpha$, ^{13}CO , $^1\text{H}_\alpha$, and $^{13}\text{C}_\beta$ secondary shifts of the cold denatured state of I98A CTL9 were calculated using the random coil values reported by Wishart.³³ The data are presented in Table 1A and Figure S1. The various observed trends are broadly self-consistent and indicate that the two α -helical regions (residues 58–74 and residues 95–106) in the native state have a strong tendency to form an α -helical structure in the cold denatured state. In addition, the loop region which connects α -helix 2 and β -strand 2, i.e., residues 107–124 in

(38) Eliezer, D. *Methods Mol. Biol.* **2007**, 350, 49–67.

(39) Spera, S.; Bax, A. *J. Am. Chem. Soc.* **1991**, 112, 5490–5492.

(40) Wishart, D. S.; Sykes, B. D.; Richards, F. M. *J. Mol. Biol.* **1991**, 22, 311–333.




Figure 3. Plots of the difference in the $^{13}\text{C}_\alpha$ secondary shifts and the $^{13}\text{C}_\beta$ secondary shifts for (A) The native state of wild-type CTL9, (B) the cold denatured state of I98A CTL9. Note that the scale is different in panel A and panel B. Assignments were obtained at pH 5.7, 12 °C. A schematic diagram of the elements of secondary structure of the native state of CTL9 is shown at the top of each panel, with solid bars, hashed bars, and open arrows indicating α helix, 3–10 helix, and β -strand structure, respectively. The color coding corresponds to that used in Figure 1.

the native state, also exhibits a clear trend to populate helical Φ and Ψ angles. This is of interest because it implies that the non-native structure is formed.

Individual secondary shifts are sensitive to protein secondary structure, but their predictive value strongly depends on the accuracy of the chemical shift standardization. Combinations of secondary shift data can often provide a more accurate picture of the structural propensities for unfolded and partially unfolded proteins. We calculated the difference between $^{13}\text{C}_\alpha$ secondary shifts and $^{13}\text{C}_\beta$ secondary shifts, as plotted in Figure 3. The figure compares the $\Delta\delta\text{C}_\alpha - \Delta\delta\text{C}_\beta$ values for the native state (Figure 3A) and the cold denatured state (Figure 3B). Positive $\Delta\delta\text{C}_\alpha - \Delta\delta\text{C}_\beta$ values indicate α -helical structure, and negative values indicate a propensity to adopt a β -strand structure.³⁴ Positive $\Delta\delta\text{C}_\alpha - \Delta\delta\text{C}_\beta$ values are detected for the two helical regions as well as for the loop between α -helix 2 and β -strand 2 in the cold denatured state with average values of 1.26, 1.01, and 0.91, respectively. For comparison, the values for folded wild-type CTL9 under the same conditions are 3.47, 2.21, and 0.21 (Figure 3A and Table 1A), respectively. Negative $\Delta\delta\text{C}_\alpha - \Delta\delta\text{C}_\beta$ values, with an average of -0.17 , are observed in residues 78–82, which reside in the first β -strand in the native state, while the average is -1.79 in the wild-type CTL9 under the same conditions.

Forman-Kay and co-workers have developed the secondary structure propensity (SSP) algorithm to detect secondary structure propensities in partially unfolded and natively disordered proteins.³⁴ The SSP score is the weighted average of the chemical shifts from different nuclei in a given residue, with the relative weighting reflecting the sensitivity of different




Figure 4. SSP analysis of wild-type CTL9 (red ●) and I98A CTL9 (blue ●) at pH 5.7, 12 °C. A schematic diagram of the elements of secondary structure of the native state of CTL9 is shown at the top of the figure.

secondary shifts to structure. An SSP score of 1.0 suggests a fully formed α -helix, while a value of -1.0 suggests a fully formed β -strand. Figure 4 displays the SSP scores of the cold denatured state of I98A at pH 5.7, 12 °C and those of the wild-type CTL9 under the same conditions. Although the SSP algorithm was developed for the analysis of secondary structure in non-native protein conformations, the SSP values for wild-type CTL9 clearly reflect the native secondary structure of the protein (Table 1A), consistent with the wild-type protein being well folded under these conditions. The cold denatured state of I98A, in comparison, is less structured and, as expected, exhibits SSP scores with smaller amplitudes. Nevertheless, some obvious structural trends are detected. Positive values are observed for α -helix 1 and α -helix 2. The average is 0.33 ± 0.08 for α -helix 1 and 0.25 ± 0.06 for α -helix 2. Large positive SSP scores are also found in the region which connects α -helix 2 and β -strand 2 (residues 107–124), with an average of 0.20 ± 0.11 . For comparison, the values for the folded state of the wild-type CTL9 are 0.91 ± 0.28 , 0.78 ± 0.31 , and 0.09 ± 0.50 , respectively. Negative SSP scores are observed in the cold denatured state for residues 78–83 and residues 128–133, which are located in the first and second β -strand, respectively, in the native state. The average SSP values for these two segments are -0.17 ± 0.10 and -0.18 ± 0.06 . The corresponding values for folded wild-type CTL9 are -0.51 ± 0.32 and -0.42 ± 0.20 under the same conditions. Fractional SSP values are normally interpreted in terms of fractional populations of secondary structures. However, the estimates of secondary structure do not have to agree with those obtained from CD because of the ways the respective spectroscopic signals depend on structure. NMR secondary shifts are primarily local in origin, and a residue which populates the helical region of the ϕ – Ψ map will have helical secondary shifts. In contrast, there is a length dependence to the CD signal of α -helices with the rotational strength (intensity) of the spectrum falling off for short helices. Thus, the CD spectrum of a system which has a tendency to transiently populate short helical segments may suggest, incorrectly, that there is very low helical content while an NMR secondary shift analysis (such as the SSP analysis) indicates a higher level of structure. The estimated fractional helical content of the cold unfolded state derived from the SSP analysis is 0.13. This can be compared to that estimated from the mean residue ellipticity at 222 nm. The observed CD signal is the sum of the signal from the folded and cold denatured states. An estimate of the signal due to the cold folded state can be obtained from the measured fractional populations and the known ellipticity of the fully folded state (Supporting

Information). Using $-11\,200\text{ deg cm}^{-1}\text{ dmol}^{-1}$ for the ellipticity of the folded state and a fractional population of 0.21 for the folded state yields a value of $-4230\text{ deg cm}^{-1}\text{ dmol}^{-1}$ for the mean residue ellipticity at 222 nm for the cold unfolded state at 12 °C. This corresponds to an estimated apparent fraction of helix ranging from 6 to 12%. The two values reflect the choice for the expected ellipticity of a fully unfolded state with no helical structure. The larger value results if the empirical relationship proposed by Baldwin and co-workers⁴¹ is used for the nonhelical value, and the smaller estimate is obtained if the actual ellipticity of CTL9 under strongly denaturing conditions is used (pH 2.0, 8 M urea). In any case, the estimated helical content obtained from CD is in reasonable agreement with that obtained via the SPP analysis. This is consistent with the notation that the helical structure detected by the chemical shift analysis corresponds to segments of the chain in which consecutive residues adopt helical conformations. NOE measurements, described in subsequent subsections, are consistent with this interpretation.

The SSP analysis is in excellent agreement with the secondary shifts analysis, which argues that the cold denatured state of I98A CTL9 contains both native-like and non-native α -helical as well as a tendency to sample native-like β -strand or extended structures.

Residual dipolar couplings confirm α -helical secondary structure in the cold denatured state. Residual dipolar couplings (RDCs) are playing an increasingly important role in structural determination of macromolecules.^{42,43} For well folded globular proteins, the magnitude and sign of the RDCs depend on the orientation of the bond vectors relative to the alignment tensor of the entire molecule. In contrast, unfolded proteins generally cannot be described by a global alignment tensor. Instead, the polypeptide chain can be viewed as behaving like a collection of short segments of polypeptide in which each short segment transiently aligns relative to the external magnetic field. The length of the segment, often termed the persistence length, reflects the number of amino acid residues within which the polypeptide chain has a correlated orientation. Previous work has suggested that the NH bond vectors adopt alignments that are approximately perpendicular for α -helical and β -strand or extended structures. Thus the NH RDCs in independently aligning α -helices and a β -strand have opposite signs since the NH vector is aligned parallel to the long axis of an α -helix but perpendicular to the long axis of a β -strand.^{44–47} As a result, NH RDCs can report on the residual secondary structure in unfolded proteins. In contrast, the RDCs of highly unfolded proteins usually have the same sign. Recent work has shown that RDCs for denatured states with little or no residual structure can be reasonably well predicted using ensembles constructed from dihedral angle choices derived from certain regions of protein structure contained in the protein data bank.^{48,49} The predicted RDCs are dominated by those that come from highly




Figure 5. Residual dipolar couplings for CTL9 and the I98A mutant. Measurements were made at pH 5.7, 12 °C. (A) Plot of D_{NH} for the wild-type CTL9 versus residue number. (B) Plot of D_{NH} (black bars) and SSP scores (●) for I98A CTL9 versus residue number. A schematic diagram of the elements of secondary structure of the native state of CTL9 is shown at the top of each panel.

extended conformations, since these align most strongly with the field and thus explain the observation that highly unfolded proteins exhibit strong RDCs with mostly the same sign. If there is local residual structure such as helices in an otherwise extended polypeptide, then the alignment of the NH bond vectors in the helical segments will tend to be perpendicular to that of the NH bond vectors in the extended regions. Irrespective of whether one wishes to describe this as local alignment of persistence-length segments or alignment of highly extended conformations, the end result is that RDCs are well suited for probing the residual structure, and especially the residual helical structure, in unfolded proteins. In the current study, NH RDCs of the cold denatured state of I98A CTL9 were measured in weakly aligned bicelles. The HSQC spectra obtained in the bicelle media (data not shown) are identical to those collected in isotropic solution, indicating that the alignment media does not significantly perturb the conformation. Figure 5A displays the NH RDCs measured for the wild-type CTL9 at pH 5.7, 12 °C. Although the globular wild-type protein aligns in the bicelle medium as a rigid structure, one can note that the NH RDCs are positive for α -helix 1 and negative in α -helix 2, consistent with the native structure in which the two helices are oriented perpendicular to each other. All of the NH RDCs are of the same sign (negative) in the β -strand regions, consistent with the β -strands being oriented either parallel or antiparallel to each other. Thus, the RDC data of the native state of CTL9 agree qualitatively with the crystal structure. The NH-RDCs for the cold denatured state of I98A CTL9 are plotted in Figure 5B. The segments which are predicted to populate the α -helical structure based on the chemical shift analysis, including the loop connecting α -helix 2 and β -strand 2, exhibit RDCs which have opposite signs relative to the β -strand regions or extended

(41) Luo, P.; Baldwin, R. L. *Biochemistry* **1997**, *36*, 8413–8421.

(42) Tjandra, N.; Bax, A. *Science* **1997**, *278*, 1111–1114.

(43) Tolman, J. R.; Flanagan, J. M.; Kennedy, M. A.; Prestegard, J. H. *Proc. Natl. Acad. Sci. U. S. A.* **1995**, *92*, 9279–9283.

(44) Mohana-Borges, R.; Goto, N. K.; Kroon, G. J. A.; Dyson, H. J.; Wright, P. E. *J. Mol. Biol.* **2004**, *340*, 1131–1142.

(45) Louhivuori, M.; Paakkonen, K.; Fredriksson, K.; Permi, P.; Lounila, J.; Annala, A. *J. Am. Chem. Soc.* **2003**, *125*, 15647–15650.

(46) Ohnishi, S.; Lee, A. L.; Edgell, M. H.; Shortle, D. *Biochemistry* **2004**, *43*, 4064–4070.

(47) Ohnishi, S.; Shortle, D. *Proteins* **2004**, *50*, 546–551.

(48) Jha, A. K.; Colubri, A.; Freed, K. F.; Sosnick, T. R. *Proc. Natl. Acad. Sci. U.S.A.* **2005**, *102*, 13099–13104.

(49) Bernado, P.; Blanchard, L.; Timmins, P.; Marion, D.; Ruigrok, R. W. H.; Blackledge, M. *Proc. Natl. Acad. Sci. U.S.A.* **2005**, *102*, 17002–17007.




Figure 6. Summary of NOEs observed for I98A CTL9 at pH 5.7, 12 °C. (A) Plot of the volume of the sequential amide proton NOEs versus residue number. (B) Plot of normalized sequential amide proton NOEs versus residue number. Peak volumes are normalized as the ratio of observed $d_{NN}(i,i+1)$ NOE crosspeaks to the diagonal peaks.

regions. The three β -strand segments display negative RDCs. It is worth noting that β -strand 3 (residues 138–147) displays significant negative RDCs in the cold denatured state in spite of the small SSP scores in this region. A plausible explanation is that the alignments of the NH vectors are similar in both β -strands and the extended structure in unfolded proteins.⁴⁴ Other than the β -strand 3 region, the pattern of the RDC data is in excellent agreement with the SSP scores and the RDC data offer additional evidence that there is significant helical structure in the cold denatured state. It is noteworthy that all RDCs have much smaller amplitudes in the cold denatured state than in the native state. Since the experimentally measured RDCs are population weighted averages of an ensemble of conformations, the small magnitudes of the RDCs reveal that the residual secondary structure in the cold denatured state is only partially developed. Note that the measured RDCs for the cold denatured state do not include any contributions from averaging from the native structure since the native and the cold denatured states are in slow exchange.

Amide proton NOEs are consistent with native-like structure in the cold denatured state. Proton NOEs provide valuable structural probes: Short-range and medium range NOEs are widely used to delineate protein secondary structure, and long-range NOEs are particularly useful for defining the global topology of proteins.⁵⁰ However, measurements of NOEs in unfolded proteins are quite difficult due to the flexible nature

of the polypeptide chain in the unfolded state ensemble and the high degeneracy of the resonances. Nevertheless, sufficient spectral resolution can be obtained using the ^{15}N -HSQC-NOESY-HSQC experiment, provided that the population of a conformation with a defined secondary structure is high enough. Sequential amide proton NOEs, which are often used to distinguish α -helices (stronger NOEs) from β -strand and extended structures (weaker or absent NOEs), are detected throughout the protein in the cold denatured state (Figure 6A and 6B). The two α -helical regions, as well as the loop region connecting α -helix 2 and β -strand 2, display sequential amide NOEs which have higher intensities than the NOEs observed for other segments of the polypeptide chain. The average intensities of the normalized NOEs are 1.7 times as large in these three regions as those observed for other residues. Thus, the observed NOEs are consistent with the chemical shift and RDC measurements.

Because a low population of native state I98A CTL9 is present under our conditions, the possibility exists that some of the observed NOEs could in fact be transferred from this native state population. Transferred NOE experiments are typically measured for systems in fast exchange^{51,52} but can also be observed for proteins in slow exchange.^{53–55} In the present case the NOE mixing time was 300 ms and the measured exchange rate between the cold denatured state and the native

(50) Wuthrich, K. *NMR of proteins and nucleic acids*; John Wiley & Sons, Inc.: New York, 1986.

(51) Clore, G. M.; Gronenborn, A. M. *J. Magn. Reson.* **1983**, *53*, 423–442.

(52) Post, C. B. *Curr. Opin. Struct. Biol.* **2003**, *13*, 581–588.




Figure 7. Plots of ^{15}N R_2 relaxation rates for wild-type CTL9 (red ●) and the cold denatured state of I98A CTL9 (blue ●). The black solid line (—) is the best fit to the phenomenological model of Schwalbe and co-workers. Data were collected at 12 °C for both proteins.

state is 0.49 s^{-1} , and the calculated life times are 2.5 s for the folded state and 11 s for the cold denatured state. In principle, some degree of NOE transfer could potentially contribute to the measured data. However, the observation of strong NOEs for residues 120–126, a region with no regular secondary structure in the native state of the protein, suggests that the major contribution to the observed NOEs is from the cold denatured state. In addition, the data are consistent with the chemical shift and RDC data, and those measurements are not affected under conditions of slow exchange.

^{15}N Transverse Relaxation Rate Analysis of the Cold Denatured State. Relaxation rate measurements have been widely used to study the structure and dynamics of unfolded proteins.^{56,57} Transverse ^{15}N relaxation rates (R_2) are particularly useful because they are very sensitive probes of deviations from random coil behavior in unfolded proteins. For a completely unstructured peptide with only local interactions, the ^{15}N R_2 relaxation rates are expected to be uniform in the center of the polypeptide chain and slightly smaller at the ends.^{37,58} The observation of significantly larger R_2 rates is thought to reflect the formation of local residual structure such as hydrophobic clusters.^{37,38,58,59} Figure 7 plots the measured relaxation rates for wild type CTL9 and I98A CTL9 at 12 °C and the relaxation rates calculated based on the Schwalbe phenomenological model.³⁷ The slow exchange between the folded and the cold denatured state will not affect the measured R_2 rates. The calculated relaxation rates follow the expected inverted U-shaped profile. The fit yields an apparent persistence length of 8 residues, which is consistent with previous estimates⁶⁰ and supports the analysis of the RDC data. Not surprisingly, the native state displays large systematic deviations from the values calculated using the Schwalbe random coil model. The ^{15}N relaxation rates for the cold denatured state exhibit smaller but

still significant positive deviations from the random coil model. In particular, the largest deviations are found in the two α -helical regions as well as in the loop connecting α -helix 2 and β -strand 2. No significant deviations from the model random coil relaxation rates are found in the β -strand regions, suggesting that the α -helical structure is better formed in the cold denatured state than the β -strand structure.

Discussion

Studies of the cold denatured state have often relied on special conditions such as addition of denaturants, high pressure, encapsulation, emulsified samples, or extremes of pH, under which the structural and dynamic properties of the protein may be perturbed. Nevertheless these investigations have provided valuable information about unfolded proteins. There have also been a few reported studies of cold denaturation under near native conditions.^{5,21,61,62} The current study provides a detailed residue specific characterization of the cold denatured state of a protein under near physiological conditions. We have obtained essentially complete ^1H , ^{13}C , and ^{15}N backbone assignments as well as $^{13}\text{C}_\beta$ and $^1\text{H}_\beta$ side chain assignments for the cold denatured state of I98A CTL9 at pH 5.7 and 12 °C. The chemical shift analysis, NOE analysis, RDC data, and transverse relaxation rates all yield a consistent picture. Residues in the two α -helical regions in the native state have a strong tendency to sample the α -helical region of the Ramachandran plot. Residues 107–124, which form a short 3_{10} helix and the connecting loop linking α -helix 2 and β -strand 2 in the native state, also preferentially form an α -helical structure in the cold denatured state. The R_2 values for each of these three segments exhibit deviations from the phenomenological model for R_2 values in fully unfolded states.

Previous studies of the structure of cold denatured states, particularly pressure dependent studies, have suggested that cold denatured states either contain significant residual structure, which often mimics early intermediates that are populated during the kinetic refolding pathway, or are largely unstructured.^{12,63–65} The cold denatured state of CTL9 exhibits behavior intermediate between these two extremes. The folding of CTL9 has been well documented to be two-state. The work presented here provides additional evidence that the protein folds from a state which contains significant residual secondary structure.

The observed secondary structure and the deviations of the R_2 values from the random coil model are consistent with previous hydrodynamic measurements, which have demonstrated that the cold denatured state of the I98A mutant is more compact than the highly denatured pH 2.0 state or the urea denatured state of the wild-type.¹⁸ NMR characterization of the acid denatured state and the urea denatured state revealed that significantly less residual structure is present.

The cold denatured state of the I98A mutant is more similar to the pH 3.8 unfolded state of wild type CTL9: both are compact and both contain native and non-native secondary structure in the same regions of the chain. Indeed, there is a strong correlation between the SSP scores of these two unfolded

- (53) Bodner, C. R.; Dobson, C. M.; Bax, A. *J. Mol. Biol.* **2009**, *390*, 775–790.
 (54) Mok, Y. K.; Kay, C. M.; Kay, L. E.; Forman-Kay, J. D. *J. Mol. Biol.* **1999**, *289*, 619–638.
 (55) Mok, Y. K.; Kay, C. M.; Kay, L. E.; Forman-Kay, J. D. *J. Mol. Biol.* **2003**, *329*, 185–187.
 (56) Palmer, A. G. *Curr. Opin. Struct. Biol.* **1997**, *7*, 732–737.
 (57) Kay, L. E. *Nat. Struct. Biol.* **1998**, *5*, 513–517.
 (58) Klein-Seetharaman, J.; Oikawa, M.; Grimshaw, S. B.; Wirmer, J.; Duchardt, E.; Ueda, T.; Imoto, T.; Smith, L. J.; Dobson, C. M.; Schwalbe, H. *Science* **2002**, *295*, 1719–1722.
 (59) Platt, G. W.; McParland, V. J.; Kalverda, A. P.; Homans, S. W.; Radford, S. E. *J. Mol. Biol.* **2005**, *346*, 279–294.
 (60) Shan, B.; Eliezer, D.; Raleigh, D. P. *Biochemistry* **2009**, *48*, 4707–4719.

- (61) Antonino, L. C.; Kautz, R. A.; Nakano, T.; Fox, R. O.; Fink, A. L. *Proc. Natl. Acad. Sci. U.S.A.* **1991**, *88*, 7715–7718.
 (62) Szyperki, T.; Mills, J. L.; Perl, D.; Balbach, J. *Eur. Biophys. J.* **2006**, *35*, 363–366.
 (63) Jonas, J.; Jonas, A. *Annu. Rev. Biophys. Biomol. Struct.* **1994**, *23*, 287–318.
 (64) Jonas, J. *ACS Symp. Ser.* **1997**, *676*, 310–323.
 (65) Jonas, J.; Ballard, L.; Nash, D. *Biophys. J.* **1998**, *75*, 445–452.




Figure 8. Comparison of the SSP scores determined for the pH 3.8 unfolded state of wild type CTL9 with those determined for the cold denatured state of I98A CTL9. The SSP scores for the pH 3.8 unfolded state were previously reported.⁶⁰

states, with an r^2 value of 0.84 (p -value $< 10^{-4}$) (Figure 8). The similarity between these two unfolded states of CTL9 strongly suggests that the residual structure observed is robust and is not an artifact of the conditions used to populate the unfolded states. It is difficult to detect experimentally when a mutation alters the unfolded state structure or energetics, but the observation of similar helical structural propensities in the cold denatured state of the I98A mutant and the pH 3.8 unfolded state of the wild-type⁶⁰ argues that the isoleucine to alanine mutation at residue 98 may have little impact on the unfolded state of CTL9. Thus the native and non-native structure observed in the I98A mutant is highly likely to be formed in the unfolded state of the wild type, which is populated under refolding conditions.

Nascent residual structure in the unfolded state can reduce the conformational search in the early stages of protein folding. As a consequence, structural information about the unfolded state of CTL9 may have important implications for the folding of this protein.¹⁷ A previously reported φ -value analysis of the

transition state for the folding of CTL9 revealed small φ -values for the α -helices and somewhat larger but still modest φ -values in β -strand 2, β -strand 3, and the loop which connects the two strands.⁶⁶ φ -Value analysis reports on the development of interactions in the transition state relative to the unfolded state, and given the significant helical structure in the unfolded state of CTL9, the transition state for the folding of CTL9 likely contains significant helical structure as well. One should also bear in mind that a mutation which destabilizes the unfolded state may decrease the φ -value relative to the value which would be measured for the case where the mutation does not affect unfolded state energetics.⁶⁷

Acknowledgment. This work was supported by NIH Grants GM070941 and NSF MCB0919860 (D.P.R.), AG019391 and AG025440 (D.E.) and NS058137 (S.M.), and by the Irma T. Hirsch Foundation and a gift from Herbert and Ann Siegel (D.E.). D.P.R. and D.E. are members of the New York Structural Biology Center, which is a STAR center supported by the New York State Office of Science, Technology and Academic Research, by NIH Grant P41 GM66354, and which received funds from NIH, USA, the Keck Foundation, New York State, and the NYC Economic Development Corporation for the purchase of 900 MHz spectrometers. We thank Dr. Shibani Bhattacharya and Mr. Mark W. Maciejewski for help with NMR experiments and Mr. Wenli Meng for comments on the manuscript. Prof. Jeff Hoch is greatly appreciated for generously providing NMR time at the NMR facility at the University of Connecticut Health Center.

Supporting Information Available: A plot of CD detected thermal unfolding curves for I98A CTL9 at different pH values; a plot of the fluorescence detected stopped-flow pH jump experiment, and plots of $^{13}\text{C}_\alpha$, $^{13}\text{C}_\beta$, ^{13}CO , and $^1\text{H}_\alpha$ secondary shifts for the cold denatured state of I98A CTL9 at pH 5.7, 12 °C. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JA908104S

(66) Li, Y.; Gupta, R.; Cho, J. H.; Raleigh, D. P. *Biochemistry* **2007**, *46*, 1013–1021.

(67) Cho, J.-H.; Raleigh, D. P. *J. Am. Chem. Soc.* **2006**, *128*, 16492–16493.