


## Iridium-Catalyzed Kinetic Asymmetric Transformations of Racemic Allylic Benzoates

Levi M. Stanley, Chen Bai, Mitsuhiro Ueda, and John F. Hartwig\*

Department of Chemistry, University of Illinois, 600 South Mathews Avenue, Urbana, Illinois 61801

Received May 3, 2010; E-mail: jhartwig@illinois.edu

Allylic substitution catalyzed by metallacyclic iridium complexes derived from phosphoramidites<sup>1</sup> has become a valuable method to prepare enantioenriched small molecules with broad scope by the addition of carbon<sup>2</sup> and heteroatom<sup>3</sup> nucleophiles to allylic esters. Although many linear allylic carbonates react to form products with high enantiomeric excess (eq 1), iridium-catalyzed allylic substitution of racemic, branched allylic esters typically occurs with much lower enantioselectivity (eq 2).<sup>4</sup> Such allylic esters are attractive substrates for metal-catalyzed allylic substitution because they are readily prepared from a wide range of aldehydes and a vinyl Grignard reagent. Moreover, these allylic esters are less substituted at the alkene unit and typically react with rates that are faster than those of the corresponding linear isomers.<sup>4b,c</sup>



To date, enantioselective substitution of racemic, branched allylic electrophiles has been limited primarily to reactions of stabilized carbon nucleophiles catalyzed by complexes of Mo,<sup>5</sup> Pd,<sup>6</sup> or Rh<sup>7</sup> and a few examples of allylic amination catalyzed by complexes of Rh<sup>8</sup> and Pd.<sup>9</sup> Examples of iridium-catalyzed allylic substitution of racemic, branched allylic electrophiles that occur with high enantioselectivity are rare. Helmchen reported allylic alkylation of racemic allylic acetates catalyzed by an iridium–phosphoramidite complex, but only one substrate reacted in greater than 80% ee.<sup>4a,b</sup> In addition, Carreira<sup>4c</sup> and Alexakis<sup>4d</sup> reported kinetic resolutions by etherification and alkylation of racemic, branched allylic carbonates or esters with catalysts generated in situ from [Ir(COE)<sub>2</sub>Cl]<sub>2</sub> and a chiral diene ligand or [Ir(COD)Cl]<sub>2</sub> and a chiral phosphoramidite ligand. However, these allylic etherifications occurred with widely varying levels of enantioselectivity, and the one example of allylic alkylation occurred in 84% ee. Palladium-catalyzed isomerization of branched allylic esters to linear allylic esters, followed by iridium-catalyzed enantioselective, allylic substitution of the linear isomer, has been reported,<sup>10</sup> but this strategy requires removal of the palladium catalyst prior to the allylic substitution step and is limited to reactions of aromatic allylic esters that fully rearrange to the linear isomer.

Here, we report enantioselective allylic substitution of racemic, branched allylic benzoates with a variety of anionic carbon and heteroatom nucleophiles in the presence of a single-component, metallacyclic iridium complex to form substitution products with high enantioselectivity. These kinetic asymmetric processes occur with distinct stereochemical courses for racemic aliphatic and aromatic allylic benzoates, and the higher reactivity of the branched allylic esters enables transformations of sterically demanding electrophiles.

Recently published mechanistic studies on allylic amination showed that enantiomeric, branched allylic amines coordinate to a

metallacyclic iridium catalyst with much different binding constants.<sup>11</sup> These data implied that the metallacyclic iridium catalyst should select for reaction of one enantiomer of a racemic, branched allylic ester and form a substitution product with high enantiomeric excess, as long as the allylic substitution occurs with a high degree of retention of configuration. Prior reactions of enantioenriched allylic esters with iridium catalysts formed the substitution product with a modest enantiomeric excess,<sup>4a,b</sup> but reactions of enantioenriched allylic esters catalyzed by metallacyclic iridium complexes could occur with a higher degree of retention of configuration. Such substitutions of racemic, branched allylic esters would contrast with typical kinetic resolutions that leave behind an enantioenriched reactant but form products possessing modest enantiopurity.





Figure 1. Metallacyclic iridium–phosphoramidite complexes **1a** and **1b**.

Table 1. Ir-Catalyzed Allylic Substitution of Racemic Aliphatic Allylic Benzoates<sup>a</sup>



entry	R (2)	nucleophile	product	yield (%) <sup>b</sup>	ee (%) <sup>c</sup>
1	BnCH <sub>2</sub> ( <b>2a</b> )	NaOPh	<b>3a</b>	83	95
2	<i>n</i> -Pr ( <b>2b</b> )	NaOPh	<b>3b</b>	86	92
3	<i>i</i> -Pr ( <b>2c</b> )	NaOPh	<b>3c</b>	76	98
4	Cy ( <b>2d</b> )	NaOPh	<b>3d</b>	86	96
5 <sup>d</sup>	<i>t</i> -Bu ( <b>2e</b> )	NaOPh	<b>3e</b>	88	96
6	BnCH <sub>2</sub> ( <b>2a</b> )	LiN(Boc) <sub>2</sub>	<b>4a</b>	96	93
7	BnCH <sub>2</sub> ( <b>2a</b> )	KNHC(O)CF <sub>3</sub>	<b>5a</b>	74	98
8	BnCH <sub>2</sub> ( <b>2a</b> )	NaBzIm	<b>6a</b>	84	97
9	BnCH <sub>2</sub> ( <b>2a</b> )	NaTs	<b>7a</b>	80	94
10	BnCH <sub>2</sub> ( <b>2a</b> )	NaCH(CO <sub>2</sub> Me) <sub>2</sub>	<b>8a</b>	82	94
11	BnCH <sub>2</sub> ( <b>2a</b> )	NaCH(CN) <sub>2</sub>	<b>9a</b>	77	88

<sup>a</sup> See Supporting Information for experimental details. <sup>b</sup> Isolated yield of products **3–9**. Branched-to-linear selectivities were >95:5. <sup>c</sup> The enantiomeric excess of **3–9** was determined by chiral HPLC methods. <sup>d</sup> Reaction was run in the presence of 4 mol % **1b** at 50 °C for 24 h.


To test if we could obtain substitution products having high enantiomeric excess from the selective reaction of one enantiomer of a racemic mixture of allylic esters, we conducted reactions of the

racemic, branched allylic esters derived from 5-phenylpent-1-en-3-ol with sodium phenoxide (NaOPh) in the presence of our single-component, metallacyclic Ir–phosphoramidite complexes **1a**<sup>11</sup> and **1b**<sup>3d</sup> (Figure 1). After testing several allylic carbonates and esters, we found that the reaction of NaOPh with 2.2 equiv of allylic benzoate **2a** in the presence of 2 mol % **1b** formed allyl phenyl ether **3a** in 83% yield and 95% ee at 0 °C (Table 1, entry 1).<sup>12</sup>

Reactions of a series of anionic nucleophiles with aliphatic allylic benzoates are summarized in Table 1. As shown in entries 2–4, NaOPh reacted with branched allylic benzoates **2b–2d** (R = *n*-Pr, *i*-Pr, or *c*-hexyl) to form allyl aryl ethers **3b–3d** in 76–86% yield and 92–98% ee. The reactions of the branched allylic esters occur faster than the analogous reactions of the linear isomers. This higher reactivity allowed for an unusual asymmetric allylic substitution process. The reaction of NaOPh with allylic benzoate **2e** in which the allyl unit is substituted with a *t*-Bu group occurred at 50 °C to set a neopentyl stereocenter in the allyl aryl ether product **3e** in 96% ee (entry 5). The reaction of NaOPh with the linear isomer of **2e** did not occur under identical or more forcing conditions. Nucleophilic substitutions at neopentyl centers are challenging, and enantioselective substitutions at such a site are particularly rare.<sup>13</sup>

The enantioselective allylic substitution also occurred with anionic nitrogen, sulfur, and carbon nucleophiles in the presence of catalyst **1b**, as summarized in entries 6–11. The reactions of **2a** with the ammonia equivalents LiN(Boc)<sub>2</sub> and KNHC(O)CF<sub>3</sub>, the heteroaromatic derivative sodium benzimidazolite (NaBzIm), the sulfur nucleophile sodium *p*-toluenesulfinate (NaTs), and the sodium salts of the carbon nucleophiles dimethyl malonate and malononitrile occurred with high enantioselectivity (88–98% ee). The resulting products **4a–9a** were isolated in good yields (74–96%).

**Scheme 1.** Strategy for Kinetic Asymmetric Allylic Substitution of Racemic, Branched Aromatic Allylic Benzoates<sup>a</sup>




<sup>a</sup> [Ir] = [Ir(COD)(κ<sup>2</sup>-L2)(ethylene)] (**1b**),  $K_1 = k_1/k_{-1}$ , and  $K_2 = k_2/k_{-2}$ .

The reactivity of branched aromatic allylic benzoates was distinct from that of branched aliphatic allylic benzoates. Allylic substitution of branched aromatic allylic benzoates in the presence of catalyst **1b** occurred in competition with isomerization to the linear isomer. The isomerization of one enantiomer of the allylic benzoate was faster than isomerization of the other, but the competitive processes resulted in reaction mixtures. These mixtures consisted of (1) the (*R*)-allylic substitution product with modest enantiopurity, (2) the achiral linear aromatic allylic benzoate, and (3) the (*S*)-enantiomer of the branched allylic benzoate possessing high enantiopurity.

These data are consistent with a system that reacts with the relative rates and equilibria in Scheme 1. The rate constant for nucleophilic attack ( $k_5$ ) on allyliridium complex **A** is competitive with the rate constant for collapse ( $k_3$ ) to form the linear allylic benzoate from **A** ( $k_5 \approx k_3$ ); the rate (controlled by  $K_1k_3$ ) for the formation of the linear allylic benzoate from (*R*)-**2** is greater than the rate for the formation (controlled by  $K_2k_4$ ) of the linear allylic

benzoate from (*S*)-**2** ( $K_1k_3 \gg K_2k_4$ );<sup>14</sup> the rate for nucleophilic attack ( $k_6$ ) on allyliridium complex **B** is greater than the rate constant for collapse ( $k_4$ ) to form the linear allylic benzoate from **B**. We sought to exploit these relative rates to convert the more reactive (*R*)-enantiomer of the branched allylic benzoate **2** to the linear isomer, followed by allylic substitution of the remaining (*S*)-enantiomer of **2** with retention of configuration. This process would lead to the product of allylic substitution with high enantiomeric excess.

To establish the foundation for this strategy, we evaluated the selectivity factors (*s*) for kinetic resolution of racemic aromatic allylic benzoates in the absence of a nucleophile and the stereochemical outcome from allylic substitutions of the less reactive enantiomer of the branched allylic benzoates. As anticipated from the preliminary experiment, reaction of the racemic allylic ester from 1-phenylprop-2-en-1-ol formed a mixture of the (*S*)-enantiomer of the starting material and the achiral linear allylic benzoate with a high *s*-value. The kinetic resolution of ( $\pm$ )-**2f** in the presence of 0.5 mol % of **1b** occurred with a selectivity factor of 50. The linear allylic benzoate **10a** formed in 49% yield, and the remaining branched allylic benzoate (*S*)-**2f** was isolated in 45% yield with 98% ee (eq 3).<sup>15</sup> In addition, allylic substitution of the (*S*)-enantiomer of a representative branched allylic benzoate in the presence of **2b** occurred with essentially complete retention of configuration. The reaction of (*S*)-**2f** (98% ee) with NaOPh and 2 mol % of **2b** formed allyl aryl ether (*S*)-**3f** in 91% yield and 98% ee (eq 4).



Having established that kinetic resolution of racemic aromatic allylic benzoates by metallacycle **1b** is highly selective and that the allylic substitution of the less reactive enantiomer occurs with complete retention of configuration, we developed a protocol to convert these racemic electrophiles to enantioenriched substitution products by sequential kinetic resolution and allylic substitution. The kinetic resolution of ( $\pm$ )-**2f** in the presence of 2 mol % of **1b** at 0 °C in THF occurred in approximately 2 h. Addition of NaOPh to the resulting solution containing the less reactive enantiomer of the branched allylic benzoate (*S*)-**2f** and the linear cinnamyl benzoate **10a** formed allyl aryl ether (*S*)-**3f** in 81% yield and 96% ee (Table 2, entry 1). The absolute stereochemistry of the product shows that the allyl aryl ether (*S*)-**3f** was formed by reaction of the phenoxide with the less reactive *S*-enantiomer of the allylic benzoate **2f**, not by reaction with the linear allylic benzoate. The absolute stereochemistry of the product was the opposite of that resulting from etherification of linear aromatic allylic electrophiles.<sup>3g</sup>

The scope of this combined enantioselective isomerization and allylic substitution is shown in Table 2. This process encompasses reactions of branched allylic benzoates containing a variety of aryl groups. The reactions of NaOPh with electron-rich, electron-neutral, and electron-deficient aromatic allylic benzoates containing para, meta, and ortho substitution as well as reactions of heteroaromatic allylic benzoates occurred in 83–90% yield and in 84–98% ee (entries 1–8).

Reactions of nitrogen, sulfur, and carbon nucleophiles with branched aromatic allylic benzoates also occurred in high yields and enantioselectivities. The reactions of LiN(Boc)<sub>2</sub> and KNH-

**Table 2.** Ir-Catalyzed Allylic Substitution of Racemic Aromatic Allylic Benzoates<sup>a</sup>

entry	R (2)	nucleophile	product	yield (%) <sup>b,c</sup>	ee (%) <sup>d</sup>
1	Ph ( <b>2f</b> )	NaOPh	<b>3f</b>	81	96
2	4-MeO-C <sub>6</sub> H <sub>4</sub> ( <b>2g</b> )	NaOPh	<b>3g</b>	90	95
3	4-F <sub>3</sub> C-C <sub>6</sub> H <sub>4</sub> ( <b>2h</b> )	NaOPh	<b>3h</b>	86	95
4	4-Br-C <sub>6</sub> H <sub>4</sub> ( <b>2i</b> )	NaOPh	<b>3i</b>	84	98
5	3-Br-C <sub>6</sub> H <sub>4</sub> ( <b>2j</b> )	NaOPh	<b>3j</b>	95	98
6	2-F-C <sub>6</sub> H <sub>4</sub> ( <b>2k</b> )	NaOPh	<b>3k</b>	87	94
7 <sup>e</sup>	2-Me-C <sub>6</sub> H <sub>4</sub> ( <b>2l</b> )	NaOPh	<b>3l</b>	83	-84
8	3-pyridyl ( <b>2m</b> )	NaOPh	<b>3m</b>	86	92
9	Ph ( <b>2f</b> )	LiN(Boc) <sub>2</sub>	<b>4b</b>	92	93
10 <sup>f</sup>	Ph ( <b>2f</b> )	KNHC(O)CF <sub>3</sub>	<b>5b</b>	76	92
11	Ph ( <b>2f</b> )	NaTs	<b>7b</b>	75	96
12 <sup>e</sup>	2-Me-C <sub>6</sub> H <sub>4</sub> ( <b>2l</b> )	NaTs	<b>7c</b>	93	-94
13	Ph ( <b>2f</b> )	NaCH(CO <sub>2</sub> Me) <sub>2</sub>	<b>8b</b>	87	98
14	Ph ( <b>2f</b> )	NaCH(CN) <sub>2</sub>	<b>9b</b>	83	91

<sup>a</sup> See Supporting Information for experimental details. <sup>b</sup> Isolated yield of products **3–9**. Branched-to-linear selectivities were >95:5. <sup>c</sup> The yields of **10** (based on **2**) ranged from 48–52% (entries 1–6, 8–11, and 13–14). The yield of **10** was 5–8% for reactions of **2l** (entries 7 and 12). See Table S2 in the Supporting Information. <sup>d</sup> Enantiomeric excess of **3–9** determined by chiral HPLC methods. <sup>e</sup> Nucleophile was added prior to isomerization. <sup>f</sup> Allylic substitution was run at 0 °C to rt in the presence of 4 mol % **1b**.

C(O)CF<sub>3</sub> with **2f** formed protected allylamines **4b** and **5b** in 93% and 92% ee (entries 9 and 10). The reactions of NaTs with allylic benzoates **2f** and **2l** generated (*S*)-**7b** (R = Ph) and (*R*)-**7c** (R = 2-Me-C<sub>6</sub>H<sub>4</sub>) in good yields with excellent enantioselectivities (entries 11 and 12). Furthermore, the sodium salts of dimethyl malonate and malononitrile reacted with **2f** to form allylic alkylation products **8b** and **9b** in 98% and 91% ee (entries 13 and 14).

In principle, enantioselective allylic substitution without isomerization is possible if the rate of nucleophilic attack on complex **A** (*k*<sub>5</sub>) greatly exceeds that for collapse to the linear allylic benzoate (*k*<sub>3</sub>) (Scheme 1). The identity of the nucleophile does not substantially alter the ratio of rate constants *k*<sub>5</sub>/*k*<sub>3</sub>. However, ortho substitution on the aromatic group (R) of **2** did substantially change the ratio of rate constants *k*<sub>5</sub>/*k*<sub>3</sub>. The branched aromatic allylic benzoate **2k** containing a small ortho-fluoro group (entry 6) reacted like para- and meta-substituted branched allylic benzoates to form (*S*)-**3k** with high enantioselectivity. In contrast, branched aromatic allylic benzoates containing larger ortho substituents rearranged to the linear isomer much more slowly than did the ortho-fluoro-, meta-, and para-substituted substrates. Thus, ortho-methyl derivative **2l** underwent substitution in a manner that was similar to that of the aliphatic electrophiles; the reactions of NaOPh and NaTs with aromatic allylic benzoate **2l** (R = *o*-Me-C<sub>6</sub>H<sub>4</sub>) formed allyl aryl ether (*R*)-**3l** in 83% yield and 84% ee and allylic sulfone (*R*)-**7c** in 93% yield and 94% ee by addition of the nucleophile at the same time as the catalyst (entry 7). The enantioselectivities of the reactions of the branched cinnamyl carbonates containing ortho substituents were substantially higher than those of the linear isomer, further underscoring the improved transformations that can be realized in some cases by use of the branched isomers.

In summary, we have developed versatile methods for iridium-catalyzed, kinetic asymmetric substitution of racemic allylic electrophiles. These reactions occur between a variety of aliphatic, aromatic, and heteroaromatic allylic benzoates and a range of anionic heteroatom and carbon nucleophiles to form the corre-

sponding allylic substitution products with good to excellent enantioselectivity. The high reactivity of branched allylic benzoates enables enantioselective allylic substitutions that are slow or poorly selective with linear allylic electrophiles. Moreover, the high-yield, one-pot synthesis of the branched allylic esters makes these reactions practical with 2.2 equiv of this reagent, particularly for processes in which the nucleophile is a valuable component. Efforts to extend the current methods to neutral nucleophiles and to reactions of additional allylic electrophiles are ongoing.

**Acknowledgment.** We thank the NIH for financial support of this work (GM55382 and GM58108 to J.F.H. and GM84584 to L.M.S.), Johnson-Matthey for gifts of IrCl<sub>3</sub> and [Ir(COD)Cl]<sub>2</sub>, and Dr. Klaus Dittrich and BASF for gifts of chiral amines. M.U. thanks the JSPS for a fellowship.

**Supporting Information Available:** Experimental procedures and characterization data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References

- (1) (a) Kiener, C. A.; Shu, C. T.; Incarvito, C.; Hartwig, J. F. *J. Am. Chem. Soc.* **2003**, *125*, 14272. (b) Leitner, A.; Shekar, S.; Pouy, M. J.; Hartwig, J. F. *J. Am. Chem. Soc.* **2005**, *127*, 15506.
- (2) For selected examples, see: (a) Alexakis, A.; Polet, D. *Org. Lett.* **2004**, *6*, 3529. (b) Lipowsky, G.; Miller, N.; Helmchen, G. *Angew. Chem., Int. Ed.* **2004**, *43*, 4595. (c) Graening, T.; Hartwig, J. F. *J. Am. Chem. Soc.* **2005**, *127*, 17192. (d) Weix, D. J.; Hartwig, J. F. *J. Am. Chem. Soc.* **2007**, *129*, 7720. (e) Liu, W.-B.; He, H.; Dai, L.-X.; You, S.-L. *Org. Lett.* **2008**, *10*, 1815. (f) Polet, D.; Rathgeb, X.; Falciola, C. A.; Langlois, J.-B.; El Hajjaji, S.; Alexakis, A. *Chem.—Eur. J.* **2009**, *15*, 1205.
- (3) For selected examples, see: (a) Ohmura, T.; Hartwig, J. F. *J. Am. Chem. Soc.* **2002**, *124*, 15164. (b) Shu, C.; Leitner, A.; Hartwig, J. F. *Angew. Chem., Int. Ed.* **2004**, *43*, 4797. (c) Weihofen, R.; Tverskoy, O.; Helmchen, G. *Angew. Chem., Int. Ed.* **2006**, *45*, 5546. (d) Stanley, L. M.; Hartwig, J. F. *J. Am. Chem. Soc.* **2009**, *131*, 8971. (e) Pouy, M. J.; Stanley, L. M.; Hartwig, J. F. *J. Am. Chem. Soc.* **2009**, *131*, 11312. (f) Stanley, L. M.; Hartwig, J. F. *Angew. Chem., Int. Ed.* **2009**, *48*, 7841. (g) Lopez, F.; Ohmura, T.; Hartwig, J. F. *J. Am. Chem. Soc.* **2003**, *125*, 3426. (h) Shu, C.; Hartwig, J. F. *Angew. Chem., Int. Ed.* **2004**, *43*, 4794. (i) Lyothier, I.; Defieber, C.; Carreira, E. M. *Angew. Chem., Int. Ed.* **2006**, *45*, 6204. (j) Ueda, M.; Hartwig, J. F. *Org. Lett.* **2009**, *12*, 92.
- (4) (a) Bartels, B.; Helmchen, G. *Chem. Commun.* **1999**, 741. (b) Bartels, B.; García-Yebra, C.; Rominger, F.; Helmchen, G. *Eur. J. Inorg. Chem.* **2002**, 2569. (c) Fischer, C.; Defieber, C.; Suzuki, T.; Carreira, E. M. *J. Am. Chem. Soc.* **2004**, *126*, 1628. (d) Polet, D.; Alexakis, A.; Tissot-Croset, K.; Corminboeuf, C.; Dittrich, K. *Chem.—Eur. J.* **2006**, *12*, 3596.
- (5) (a) Trost, B. M.; Hachiya, I. *J. Am. Chem. Soc.* **1998**, *120*, 1104. (b) Malkov, A. V.; Spoor, P.; Vinader, V.; Kocovsky, P. *Tetrahedron Lett.* **2001**, *42*, 509. (c) Glorius, F.; Neuburger, M.; Pfaltz, A. *Helv. Chim. Acta* **2001**, *84*, 3178. (d) Belda, O.; Moberg, C. *Acc. Chem. Res.* **2004**, *37*, 159.
- (6) (a) Prétôt, R.; Pfaltz, A. *Angew. Chem., Int. Ed.* **1998**, *37*, 323. (b) Pàmies, D.; eguez, M.; Claver, C. *J. Am. Chem. Soc.* **2005**, *127*, 3646.
- (7) Hayashi, T.; Okada, A.; Suzuka, T.; Kawatsura, M. *Org. Lett.* **2003**, *5*, 1713.
- (8) Vrieze, D. C.; Hoge, G. S.; Hoerter, P. Z.; Van Haitsma, J. T.; Samas, B. M. *Org. Lett.* **2009**, *11*, 3140.
- (9) (a) Hayashi, T.; Kishi, K.; Yamamoto, A.; Ito, Y. *Tetrahedron Lett.* **1990**, *31*, 1743. (b) You, S.-L.; Zhu, X.-Z.; Luo, Y.-M.; Hou, X.-L.; Dai, L.-X. *J. Am. Chem. Soc.* **2001**, *123*, 7471.
- (10) Shekhar, S.; Trantow, B.; Leitner, A.; Hartwig, J. F. *J. Am. Chem. Soc.* **2006**, *128*, 11770.
- (11) Marković, D.; Hartwig, J. F. *J. Am. Chem. Soc.* **2007**, *129*, 11680.
- (12) The remaining branched allylic benzoate **2a** was isolated in 52% yield (based on 2.2 equiv of **2a**) and 78% ee. The ee of the remaining **2a–e** is necessarily modest in reactions conducted with 2.2 equiv of **2a–e** that occur to form products **3–9** with high ee. However, the ee of remaining **2a** was 94–97% with concomitant decreasing ee of **3a** (75–93%) when performing the reactions of NaOPh with less than 2.2 equiv of **2a** (1.8–2.0 equiv). See Table S1 in the Supporting Information for these data.
- (13) For recent examples of enantioselective ligand-accelerated allylic alkylation with Grignard reagents to set a neopentyl stereocenter, see: Jackowski, O.; Alexakis, A. *Angew. Chem., Int. Ed.* **2010**, *49*, 3346.
- (14) The rates of formation of allyliridium complexes **A** and **B** from the linear allylic benzoate are considered to be negligible in this analysis because the linear allylic benzoate does not undergo conversion under these reaction conditions.
- (15) The selectivity factor (*s*) = ln{(1-*c*)(1-*ee*)} / ln{(1-*c*)(1+*ee*)} where *c* is the conversion of (±)-**2f** and *ee* is the enantiomeric excess of the remaining branched allylic benzoate **2f**. Kagan, H. G.; Fiaud, J. C. *Topics in Stereochemistry*; John Wiley & Sons: New York, 1988; Vol. 18, p 249.

JA103779E