

## Synthesis of Quaternary Carbon Stereogenic Centers through Enantioselective Cu-Catalyzed Allylic Substitutions with Vinylaluminum Reagents

Fang Gao, Kevin P. McGrath, Yunmi Lee, and Amir H. Hoveyda\*

*Department of Chemistry, Merkert Chemistry Center, Boston College,  
Chestnut Hill, Massachusetts 02467*

Received July 30, 2010; E-mail: amir.hoveyda@bc.edu

**Abstract:** Catalytic enantioselective allylic substitution (EAS) reactions, which involve the use of alkyl- or aryl-substituted vinylaluminum reagents and afford 1,4-dienes containing a quaternary carbon stereogenic center at their C-3 site, are disclosed. The C–C bond-forming transformations are promoted by 0.5–2.5 mol % of sulfonate bearing chiral bidentate N-heterocyclic carbene (NHC) complexes, furnishing the desired products efficiently (66–97% yield of isolated products) and in high site (>98%  $S_N2'$ )- and enantioselectivity [up to 99:1 enantiomer ratio (er)]. To the best of our knowledge, the present report puts forward the first cases of allylic substitution reactions that result in the generation of all-carbon quaternary stereogenic centers through the addition of a vinyl unit. The aryl- and vinyl-substituted vinylaluminum reagents, which cannot be prepared in high efficiency through direct reaction with diisobutylaluminum hydride, are accessed through a recently introduced Ni-catalyzed reaction of the corresponding terminal alkynes with the same inexpensive metal–hydride agent. Sequential Ni-catalyzed hydrometalations and Cu-catalyzed C–C bond-forming reactions allow for efficient and selective synthesis of a range of enantiomerically enriched EAS products, which cannot be accessed by previously disclosed strategies (due to inefficient vinylmetal synthesis or low reactivity and/or selectivity with Si-substituted derivatives). The utility of the protocols developed is demonstrated through a concise enantioselective synthesis of natural product bakuchiol.

### Introduction


A significant challenge in chemical synthesis concerns the development of efficient catalytic enantioselective reactions that furnish all-carbon quaternary stereogenic centers.<sup>1</sup> One general strategy for promoting such processes relates to additions of C-based nucleophiles to electrophilic carbon sites. The corresponding transformations with vinylmetal reagents represent an attractive option: the products, bearing a quaternary carbon stereogenic center at the allylic position, can be functionalized and/or utilized in the synthesis of natural products. Nonetheless, catalytic enantioselective protocols delivering quaternary carbon stereogenic centers by addition of a vinylmetal are scarce.<sup>2,3</sup> There has been only one method developed in which an alkene-

based electrophile is utilized in conjunction with a vinylboronic acid;<sup>4</sup> related catalytic enantioselective allylic substitution (EAS) reactions that generate quaternary carbons<sup>5–7</sup> and involve addition of a vinyl group<sup>8</sup> have not yet been introduced (eq 1). Herein, we report the first examples of catalytic EAS processes with alkyl- or aryl-substituted vinylaluminums, affording 1,4-dienes that contain a quaternary carbon stereogenic center at their C-3 site. Transformations are promoted by 0.5–2.5 mol % of chiral bidentate N-heterocyclic carbene (NHC) complexes, furnishing the desired products efficiently and in high site (>98%  $S_N2'$ )- and enantioselectivity [up to 99:1 enantiomer ratio (er)]. Whereas alkyl-substituted vinylaluminums are obtained through reaction of the corresponding alkynes with diisobutylaluminum hydride (dibal-H),<sup>8a</sup> the aryl- or vinyl-substituted vinylmetal variants are accessed through a recently introduced Ni-catalyzed


- (1) (a) *Quaternary Stereocenters: Challenges and Solutions for Organic Synthesis*; Christophers, J., Baro, A., Eds.; Wiley-VCH: Weinheim, 2006. (b) Douglas, C. J.; Overman, L. E. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101*, 5363–5367. (c) Cozzi, P. G.; Hilgraf, R.; Zimmermann, N. *Eur. J. Org. Chem.* **2007**, 5969–5994.
- (2) For Ti-catalyzed enantioselective additions of vinylzinc reagents to ketones, see: (a) Li, H.; Walsh, P. J. *J. Am. Chem. Soc.* **2004**, *126*, 6538–6539. (b) Li, H.; Walsh, P. J. *J. Am. Chem. Soc.* **2005**, *127*, 8355–8361. For Ti-catalyzed enantioselective additions of vinylaluminums to aryl ketones, see: (c) Biradar, D. B.; Gau, H.-M. *Org. Lett.* **2009**, *11*, 499–502.
- (3) Molecules that bear a benzylic vinyl unit at an all-carbon quaternary stereogenic center have been synthesized enantioselectively through Ni-catalyzed reactions of styrene precursors with ethylene. See: (a) Shi, W.-J.; Zhang, Q.; Xie, J.-H.; Zhu, S.-F.; Hou, G.-H.; Zhou, Q.-L. *J. Am. Chem. Soc.* **2006**, *128*, 2780–2781. (b) Zhang, A.; RajanBabu, T. V. *J. Am. Chem. Soc.* **2006**, *128*, 5620–5621. (c) Smith, C. R.; Lim, H. J.; Zhang, A.; RajanBabu, T. V. *Synthesis* **2009**, 2089–2900.

- (4) For Rh-catalyzed additions of vinylboronic acids to  $\alpha,\beta$ -unsaturated pyridylsulfones, see: (a) Mauleón, P.; Carretero, J. C. *Chem. Commun.* **2005**, 4961–4963. Two isolated cases of alkyl-containing vinylaluminum conjugate additions to  $\beta$ -substituted cyclic enones, promoted by Feringa-type chiral phosphoramidite ligands and which proceed in 75:25 and 86.5:13.5 er, have been reported. See: (b) Vuagnoux-d'Augustin, M.; Alexakis, A. *Chem.-Eur. J.* **2007**, *13*, 9647–9662. (c) Palais, L.; Alexakis, A. *Chem.-Eur. J.* **2009**, *15*, 10473–10485.
- (5) For reviews on catalytic allylic alkylation reactions with “hard” C-based nucleophilic reagents, see: (a) Hoveyda, A. H.; Hird, A. W.; Kacprzynski, M. A. *Chem. Commun.* **2004**, 1779–1785. (b) Yorimitsu, H.; Oshima, K. *Angew. Chem., Int. Ed.* **2005**, *44*, 4435–4439. (c) Alexakis, A.; Bäckvall, J. E.; Krause, N.; Pàmies, O.; Diéguez, M. *Chem. Rev.* **2008**, *108*, 2796–2823.

**Scheme 1.** Proposed Synthesis of Bakuchiol through Site- and Enantioselective NHC–Cu-Catalyzed Allylic Substitution with a Vinylmetal Reagent



hydroalumination.<sup>9,10</sup> The sequential Ni- and Cu-catalyzed processes allow for efficient and selective synthesis of a range of enantiomerically enriched EAS products, which cannot be accessed by previously disclosed strategies (due to inefficient vinylmetal synthesis or low reactivity and/or selectivity with Si-substituted derivatives).



We have designed methods for enantioselective formation of tertiary C–C bonds through additions of vinylaluminums to allylic phosphates, catalyzed by chiral bidentate NHC–Cu complexes.<sup>6,8</sup> The requisite alkyl-substituted vinylmetals were accessed by hydroalumination of terminal alkynes with dibal-

H.<sup>8a</sup> We later addressed the problem of inefficiency in preparing aryl-substituted vinylaluminums through hydrometalations of the derived silyl-containing aryl alkynes;<sup>8b</sup> desilylation with a protic acid subsequent to Cu-catalyzed alkylation delivers the desired enantiomerically enriched 1,4-diene. However, the more sterically demanding Si-containing aryl-substituted vinylaluminums cannot be utilized effectively in the synthesis of the more congested quaternary carbon centers. An alternative approach for efficient preparation of aryl-substituted vinylaluminums, as will be described below, had to be introduced. From inception, we considered Cu-catalyzed EAS reactions of aryl-substituted vinylmetals to be a key component of our investigations; the present studies were partly driven by the question as to whether a concise synthesis of enantiomerically enriched bakuchiol and related natural products might be devised by the use of the targeted class of reactions (Scheme 1).<sup>11</sup>



## Results and Discussion

### 1. NHC–Cu-Catalyzed Enantioselective Allylic Substitution (EAS) Reactions with Vinylaluminum Reagents Derived from Alkyl-Substituted Alkynes.

We first established the feasibility of the proposed catalytic transformations. We focused our attention on alkyl-substituted vinylaluminums, because such entities can be easily accessed through site-selective hydroalumination with dibal-H. Initial screening pointed to Cu complexes derived from sulfonate-containing bidentate Ag carbenes **1a–c**<sup>12</sup> (Scheme 2) as optimal (see below for further discussion).

- (6) For catalytic enantioselective allylic substitutions promoted by chiral bidentate NHC–Cu complexes developed in these laboratories, see: (a) Larsen, A. O.; Leu, W.; Nieto Oberhuber, C.; Campbell, J. E.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2004**, *126*, 11130–11131. (b) Van Veldhuizen, J. J.; Campbell, J. E.; Giudici, R. E.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2005**, *127*, 6877–6882. (c) Kacprzynski, M. A.; May, T. L.; Kazane, S. A.; Hoveyda, A. H. *Angew. Chem., Int. Ed.* **2007**, *46*, 4554–4558. For application of such transformations to natural product synthesis, see: (d) Gillingham, D. G.; Hoveyda, A. H. *Angew. Chem., Int. Ed.* **2007**, *46*, 3860–3864. For reactions promoted by the corresponding Mg(II)-based complexes, see: (e) Lee, Y.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2006**, *128*, 15604–15605. For transformations promoted by the corresponding Zn(II)- and Al(III)-based complexes, see: (f) Lee, Y.; Li, B.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2009**, *131*, 11625–11633. For EAS reactions involving aryl- and hetero-arylmethyl reagents and promoted by the same class of NHC–Cu complexes, see: (g) Gao, F.; Lee, Y.; Mandai, K.; Hoveyda, A. H. *Angew. Chem., Int. Ed.* **2010**, *49*, in press.
- (7) For Cu-catalyzed EAS reactions involving amino acid-based ligands developed in these laboratories, see: (a) Luchaco-Cullis, C. A.; Mizutani, H.; Murphy, K. E.; Hoveyda, A. H. *Angew. Chem., Int. Ed.* **2001**, *40*, 1456–1460. (b) Kacprzynski, M. A.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2004**, *126*, 10676–10681. (c) Murphy, K. E.; Hoveyda, A. H. *Org. Lett.* **2005**, *7*, 1255–1258.
- (8) For Cu-catalyzed EAS reactions with vinylaluminum reagents that afford tertiary C–C bonds and are promoted by sulfonate-based bidentate chiral NHC–Cu complexes, see: (a) Lee, Y.; Akiyama, K.; Gillingham, D. G.; Brown, M. K.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2008**, *130*, 446–447. (b) Akiyama, K.; Gao, F.; Hoveyda, A. H. *Angew. Chem., Int. Ed.* **2010**, *49*, 419–423.
- (9) Gao, F.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2010**, *132*, 10961–10963.

- (10) For a review on hydroaluminations of alkynes and alkenes, see: Eisch, J. J. In *Comprehensive Organic Synthesis*; Trost, B. M., Fleming, I., Schreiber, S. L., Eds.; Pergamon: Oxford, 1991; Vol. 8, pp 733–761.
- (11) For previous enantioselective syntheses of bakuchiol, see: (a) Takano, S.; Shimazaki, Y.; Ogasawara, K. *Tetrahedron Lett.* **1990**, *31*, 3325–3326. (b) Du, X.-L.; Chen, H.-L.; Feng, H.-J.; Li, Y.-C. *Helv. Chim. Acta* **2008**, *91*, 371–378. (c) Esumi, T.; Shimizu, H.; Kashiya, A.; Sasaki, C.; Toyota, M.; Fukuyama, Y. *Tetrahedron Lett.* **2008**, *49*, 6846–6849. (d) Bequette, J. P.; Jungong, C. S.; Novikov, A. V. *Tetrahedron Lett.* **2009**, *50*, 6963–6964.
- (12) For initial reports on sulfonate-based bidentate NHC–Cu complexes, see: (a) Brown, M. K.; May, T. L.; Baxter, C. A.; Hoveyda, A. H. *Angew. Chem., Int. Ed.* **2007**, *46*, 1097–1100. (b) May, T. L.; Brown, M. K.; Hoveyda, A. H. *Angew. Chem., Int. Ed.* **2008**, *47*, 7358–7362. For applications in complex molecule synthesis, see: (c) ref 6d. (d) Brown, M. K.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2008**, *130*, 12904–12906. (e) Peese, K. M.; Gin, D. Y. *Chem.-Eur. J.* **2008**, *14*, 1654–1665.

**Scheme 2.** Chiral Bidentate NHC–Ag(I) Complexes Used in This Study<sup>a</sup><sup>a</sup> Mes = 2,4,6-Me<sub>3</sub>C<sub>6</sub>H<sub>2</sub>.**Table 1.** NHC–Cu-Catalyzed Additions of an Alkyl-Substituted Vinylaluminum Reagent to Allylic Phosphates Bearing a Trisubstituted Alkene<sup>a</sup>


entry	substrate (G)	mol % 1a	temp (°C); time	S <sub>N</sub> 2' <sup>b</sup> (%) <sup>b</sup>	yield (%) <sup>c</sup>	er <sup>d</sup>
1	Ph	0.5	–15; 3 h	>98	82	93.5:6.5
2	<i>o</i> -BrC <sub>6</sub> H <sub>4</sub>	1.0	22; 10 min	>98	92	96.5:3.5
3	<i>o</i> -CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	2.5	–15; 3 h	>98	96	98:2
4	<i>o</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	2.0	22; 10 min	>98	89	96.5:3.5
5	<i>o</i> -MeOC <sub>6</sub> H <sub>4</sub>	0.5	22; 30 min	>98	83	97.5:2.5
6	<i>o</i> -MeC <sub>6</sub> H <sub>4</sub>	2.0	–15; 3 h	>98	87	96.5:3.5
7	<i>p</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	0.5	22; 10 min	>98	91	94.5:5.5
8	Cy	2.0	–50; 6 h	>98	91	95:5
9	Me <sub>2</sub> CCH(CH <sub>2</sub> ) <sub>2</sub>	1.0	–50; 24 h	>98	77	92.5:7.5
10	CO <sub>2</sub> <i>t</i> -Bu	0.5	–15; 3 h	>98	97	90:10
11	SiMe <sub>2</sub> Ph	0.5	–15; 3 h	>98	85	95.5:4.5

<sup>a</sup> Reactions were performed under N<sub>2</sub> atmosphere; >98% conversion in all cases, except 91% conv. in entry 9; >98% *E* product isomer in all cases. <sup>b</sup> Determined by analysis of 400 MHz <sup>1</sup>H NMR of unpurified mixtures. <sup>c</sup> Yields of isolated and purified products. <sup>d</sup> Determined by HPLC analysis; see the Supporting Information for details.


A considerable range of allylic phosphates undergo facile EAS reactions in the presence of 0.5–2.5 mol % **1a**<sup>13</sup> with *n*-hexyl-substituted vinylaluminum **2** to furnish the desired 3,3-disubstituted 1,4-dienes in 90:10–98:2 er and 77–97% yield (Table 1). Aryl-substituted substrates (entries 1–7) bearing electron-donating (entry 5) and -withdrawing substituents (entries 2–4 and 7), as well as those that carry an alkyl group (entries 8–9), serve as effective starting materials. Reactions with aryl-substituted allylic phosphates are sufficiently selective to be performed at –15 to 22 °C and are complete in 3 h. Transformations that involve alkyl-substituted electrophiles (entries 8–9, Table 1) must be performed at –50 °C for maximum enantioselectivity<sup>14</sup> and might require longer reaction times (6–24 h vs 10 min–3 h), unless higher catalyst loading is used (e.g., 2 mol %, –50 °C, 6 h, entry 8, Table 1). The Cu-catalyzed process can be used to access 1,4-dienes containing a carboxylic ester or silyl-substituted quaternary carbon stereogenic center at the C3 position (entries 10–11).

(13) Similar efficiencies and selectivities are observed with complexes derived from **1b** and **1c**. Use of the NHC–Cu derived from **1a** is due to the slightly shorter route used for the preparation of the latter Ag complex.


(14) For example, reaction in entry 9 of Table 1, when performed at 22 °C, affords the desired product in 88.5:11.5 er.

**Scheme 3****Sterically hindered alkyne:**


**3**  
(0.5 mol % **1b**; –15 °C, 12 h)  
>98% yield (<2% alkynyl addn)  
91:9 er, >98% *E*

**Halogen-containing alkyne:**


**4**  
(0.5 mol % **1b**; –15 °C, 3 h)  
91% yield (<2% alkynyl addn)  
96:4 er, >98% *E*

**Heteroatom-containing alkyne:**


**5**  
(0.5 mol % **1b**; –15 °C, 3 h)  
91% yield (<2% alkynyl addn)  
96:4 er, >98% *E*

**Propargyl ether:**


**6**  
(0.5 mol % **1b**; –50 °C, 24 h)  
82% yield (<2% alkynyl addn)  
94.5:5.5 er, >98% *Z*

**Scheme 4.** NHC–Cu-Catalyzed EAS with Aryl Alkynes or Enynes Leads to Significant Amounts of Alkynyl Addition Products

**7**  
(0.5 mol % **1b**; –15 °C, 12 h)  
>98% conv. (vinyl:alkynyl addn = 71:29)  
66% yield of pure vinyl addn product  
97.5:2.5 er (vinyl addn product)



**8**  
(1.0 mol % **1a**; –15 °C, 12 h)  
72% conv. (vinyl:alkynyl addn = 55:45)  
yield not determined (inseparable products)  
96.5:3.5 er (vinyl addn product)




As illustrated through Cu-catalyzed synthesis of 1,4-dienes **3–6** (Scheme 3), a variety of vinylaluminum reagents, including one that contains a sterically demanding *tert*-butyl substituent (**3**), a halide (**4**), or heteroatom units (**5–6**),<sup>15</sup> can be used. Such transformations proceed with similarly high efficiency and stereoselectivity (single olefin isomer and 91:9–96:4 er) as observed in processes that involve alkyl-substituted alkynes (Table 1).

**2. NHC–Cu-Catalyzed Enantioselective Allylic Substitution (EAS) Reactions with Vinylaluminum Reagents Derived from Vinyl- or Aryl-Substituted Alkynes. i. Inefficient Synthesis of Vinylaluminum Reagents.** In contrast to NHC–Cu-catalyzed reactions that involve alkyl-substituted vinylaluminum reagents, when the vinylmetal is derived from hydrometalation of an enyne or an aryl alkyne (e.g., **7** and **8**, Scheme 4), although high er values are obtained, product mixtures are contaminated with alkynyl addition products (up to 45%).<sup>16</sup> Presumably, during hydroalumination, the vinylmetal serves as an efficient base to deprotonate the starting alkyne, affording alkynylalu-

(15) For directed (*Z*-selective) hydroalumination of terminal propargyl ethers (cf., **6**), see: Alexakis, A.; Duffault, J. M. *Tetrahedron Lett.* **1988**, *29*, 6243–6246.

(16) The ability of NHC–Cu complexes to promote addition of an alkyne unit is noteworthy. Development of the corresponding enantioselective variants is in progress and will be disclosed shortly.

**Scheme 5.** Use of a Si-Substituted Vinylaluminum Reagent Derived from an Aryl Alkyne in NHC–Cu-Catalyzed EAS



with **1a**: >98% conv, **11:12** = 83:17, **11** in 75% yield, 52.5:47.5 er

with **1b**: >98% conv, **11:12** = 67:33, **11** in 60% yield, 61:39 er


with **1c**: >98% conv, **11:12** = 55:45, **11** in 51% yield, 56.5:43.5 er

minum and the corresponding protonated alkene. The above complication is exacerbated by reaction of the alkynylaluminum, which appears to be somewhat more efficient relative to addition of the vinylmetal reagent. Control experiments indicate that, whereas 25% alkyne deprotonation occurs in hydroalumination of phenylacetylene, 45% alkynyl addition product<sup>17</sup> is generated along with 1,4-diene **8** (1.5 equiv of hydroalumination product mixture used; see Scheme 4).


**ii. The First Approach: Hydroalumination of Silyl-Substituted Alkynes.** Our first attempt to bypass the inefficiency of the aforementioned hydroaluminations led us to examine the formerly reported catalytic EAS reactions with silyl-substituted vinylaluminum,<sup>8b</sup> the outcome of these investigations is shown in Scheme 5. In a Cu-catalyzed process with allylic phosphate **10**, applicable to enantioselective synthesis of bakuchiol (Scheme 1), use of trisubstituted vinylaluminum **9**, generated from reaction of the corresponding trimethylsilylethynyl-4-anisole with dibal-H, gives rise to substantial amounts of the *i*-Bu addition product and furnishes 1,4-diene **11** with low enantioselectivity (52.5:47.5–61:39 er). Stabilization of electron density at the vinylic carbon by the silyl substituent likely retards the rate of vinyl transfer, resulting in the formation of **11**–**45%** of **12** in the product mixture.

**iii. The Second Approach: Efficient Ni-Catalyzed Hydroalumination of Vinyl- or Aryl-Substituted Terminal Alkynes.** In the course of searching for an effective solution to the problem of efficient vinylaluminum synthesis with minimal alkynylaluminum contamination, we discovered that in the presence of 3.0 mol % of commercially available Ni(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub>, reaction of dibal-H with enynes or aryl-substituted alkynes proceeds efficiently (within 2–12 h at 4–22 °C) to afford the desired organometallic reagent with high selectivity [85% to >98% terminal:internal ( $\beta$ : $\alpha$ )] and with <5% alkyne deprotonation. Thus, as depicted in Scheme 6, 1,4-diene **7** can be obtained in 89% yield when the Ni-catalyzed hydroalumination is utilized in conjunction with the NHC–Cu-catalyzed process (vs 66% yield with uncatalyzed hydroalumination in Scheme 4); the amount of alkynyl addition side product is reduced to 5% (vs

**Scheme 6.** Ni-Catalyzed Hydroalumination of an Enyne and in Situ Use in an NHC–Cu-Catalyzed EAS Reaction



**Table 2.** NHC–Cu-Catalyzed Additions of Aryl-Substituted Vinylaluminum Reagents to Allylic Phosphates Containing a Trisubstituted Alkene<sup>a</sup>



entry	substrate (R)	alkyne (Ar)	time (h)	$\beta$ : $\alpha$ <sup>b</sup>	$S_N2'$ : $S_N2$ <sup>b</sup>	yield (%) <sup>c</sup>	er <sup>d</sup>
1	Ph	Ph	3	95:5	>98	78	96:4
2	Ph	<i>o</i> -MeC <sub>6</sub> H <sub>4</sub>	6	>98:2	>98	88 <sup>e</sup>	95:5
3	Ph	<i>p</i> -MeOC <sub>6</sub> H <sub>4</sub>	6	>98:2	>98	84	96.5:3.5
4	Ph	<i>p</i> -CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	3	87:13	>98	82	94:6
5	<i>o</i> -BrC <sub>6</sub> H <sub>4</sub>	Ph	3	96:4	>98	91	98:2
6	<i>o</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	Ph	24	>98:2	>98	92	98:2
7	<i>o</i> -MeC <sub>6</sub> H <sub>4</sub>	<i>p</i> -MeOC <sub>6</sub> H <sub>4</sub>	3	>98:2	>98	82	98:2
8	<i>p</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	Ph	3	92:8	>98	84	93:7
9	<i>p</i> -CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	<i>p</i> -MeOC <sub>6</sub> H <sub>4</sub>	3	>98:2	>98	89	94:6
10	Me <sub>2</sub> CCH(CH <sub>2</sub> ) <sub>2</sub>	Ph	3	93:7	>98	81	90:10
11	Me <sub>2</sub> CCH(CH <sub>2</sub> ) <sub>2</sub>	<i>o</i> -MeC <sub>6</sub> H <sub>4</sub>	6	>98:2	>98	85 <sup>e</sup>	91:9
12	Me <sub>2</sub> CCH(CH <sub>2</sub> ) <sub>2</sub>	<i>p</i> -MeOC <sub>6</sub> H <sub>4</sub>	6	>98:2	>98	90	91:9
13	Me <sub>2</sub> CCH(CH <sub>2</sub> ) <sub>2</sub>	<i>p</i> -CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	3	85:15	>98	79	87:13


<sup>a</sup> Reactions were performed under N<sub>2</sub> atmosphere; >98% conversion in all cases. <sup>b</sup> Determined by analysis of 400 MHz <sup>1</sup>H NMR spectra of the unpurified mixtures. <sup>c</sup> Yields of isolated and purified  $\beta$  products. <sup>d</sup> Enantiomer ratio of the  $\beta$  product, determined by HPLC analysis; see the Supporting Information for details. <sup>e</sup> Alkynyl product (~10%) present in the product mixture.

29%). Formation of the desired 1,4-diene (**7**) in 99:1 er indicates that the presence of the Ni salt in the reaction mixture does not adversely effect on the EAS process; this contention is substantiated by reactions of aryl alkynes, described below.

As the data in Table 2 illustrate, an assortment of aryl-substituted alkynes is readily and selectively converted to the derived  $\beta$ -vinylaluminums, which are then used in situ for highly efficient as well as site- and enantioselective EAS reactions (78–92% yield of pure  $\beta$  product, >98%  $S_N2'$ , 87:13–98:2 er). In stark contrast to the process involving uncatalyzed hydroalumination (45% alkynyl addition; Scheme 4), the transformation illustrated in entry 5 of Table 2 does not afford any products derived from alkynylaluminum addition (<2% by 400 MHz <sup>1</sup>H NMR analysis of the unpurified mixture). Only in cases where the aryl unit of the alkyne bears an electron-withdrawing *p*-CF<sub>3</sub> substituent is >10% of the  $\alpha$ -vinylaluminum adduct observed; otherwise, the desired  $\beta$ -substituted 1,4-diene products are obtained in  $\geq 92\%$  selectivity. It is worthy of note that the

(17) The higher preponderance of alkynyl product (45%) versus alkyne deprotonation (25%) is because 1.5 equiv of alkyne substrate (and 1.5 equiv of dibal-H) is used, which indicates that Cu-catalyzed addition of alkynylaluminum is faster than that of the vinylaluminum reagent.

**Scheme 7.** Application of Sequential Ni-Catalyzed Alkyne Hydroalumination/NHC–Cu-Catalyzed EAS to Enantioselective Synthesis of Bakuchiol




presence of the Ni catalyst enhances the rate of hydroalumination such that higher conversion is attained for the overall process (vinylaluminum formation/EAS). As an example, only 72% conversion is achieved in formation of **8** under the conditions shown in Scheme 4, largely due to lower amounts of available vinylaluminum reagent, whereas complete consumption of the allylic phosphate is observed in reaction in entry 5 of Table 2. In cases where the alkyne substrate contains a sterically demanding *o*-substituted aryl group (entries 2 and 11, Table 2), ~10% of the alkynyl addition product is formed, likely as a result of diminution in the rate of hydroalumination.

The utility of the sequential  $\beta$ -selective Ni-catalyzed terminal alkyne hydroalumination/NHC–Cu-catalyzed EAS is highlighted in the synthesis of enantiomerically enriched bakuchiol shown in Scheme 7. The three-vessel process, involving geraniol and a readily available terminal alkyne as starting materials, proceeds in 72% overall yield. The route depicted in Scheme 7 is substantially more concise than the most efficient of the previously reported approaches, the shortest of which requires 10 steps and delivers the target in 49% yield.<sup>11b</sup>

**3. Mechanistic Considerations and the Significance of Various Structural Features of Bidentate Sulfonate-Containing NHC–Cu Complexes.** The efficiency and high site- and enantioselectivities in the catalytic EAS reactions described above arise from unique catalytic abilities of sulfonate-based bidentate NHC–Cu complexes **1a–c**,<sup>18</sup> as shown in Scheme 8, phenoxy-bridged or monodentate NHC–Cu variants are ineffective catalysts. The above findings can be explained through mode of reaction **I** (Scheme 8). The syn relationship between the phenyl backbone of the NHC and the sulfonate, which is in contrast to the anti stereochemistry in the phenoxy-bridged systems, has been

substantiated by X-ray structures and rationalized in a previous report regarding the corresponding Zn and Al complexes.<sup>6f</sup> The above stereochemical attribute allows the vinyl unit to be positioned properly for S<sub>N</sub>2' addition to the coordinated alkene,<sup>19</sup> while the equatorially positioned sulfonate oxygen facilitates addition by Lewis acid activation of the allylic phosphate.




Such features are absent in phenoxy-bridged or monodentate NHC–Cu complexes. High enantioselectivity likely arises, because mode of addition **II** engenders unfavorable steric interactions, as shown in Scheme 8; furthermore, examination of molecular models indicates that the aforementioned Lewis acid activation involving the phosphate group (cf., **I**) cannot be easily assisted by the sulfonate unit in **II**. Consistent with the proposed model, when the corresponding *Z*-trisubstituted allylic phosphates are used, the opposite product enantiomer is generated predominantly but in low enantioselectivity, because reaction via **I** would lead to unfavorable steric interactions shown in **II**; the example in eq 2 is representative.

## Conclusions

Among various types of C-based nucleophiles used in C–C bond-forming reactions, products arising from transformations with vinylmetal reagents are among the most versatile and useful. Nonetheless, related catalytic enantioselective protocols,<sup>20–22</sup> particularly those that furnish quaternary carbon stereogenic centers,<sup>2,4</sup> continue to be relatively uncommon. In addition to the challenges associated with the design of effective chiral catalysts that promote vinylmetal additions to a sterically

- (19) An NHC–Cu-catalyzed EAS process might proceed through a Cu(I) mechanism (direct transfer of the vinyl unit) or a pathway that involves a Cu(III) complex (cuprate addition followed by alkyl–vinyl reductive elimination). Either scenario benefits from the Lewis acid activation proposed, whereas the possibility of situating the vinyl group vis-à-vis the coordinated substrate in the manner shown in **I** would facilitate the Cu(I) pathway (proper alignment of C–Cu and alkene  $\pi^*$ ). It is not clear at present which pathway is energetically preferred. Although previous mechanistic studies point to Cu(III) mechanism being operative, such investigations were in connection to alkyl- or allyl-copper complexes, considered allyl halides as substrates, and did not involve a catalyst. The more polarized nature of a Cu–C bond in a vinylmetal complex, particularly a strongly Lewis base-activated NHC–Cu–vinyl complex (see ref 6f), and the associated steric demands of forming a Cu(III)-substituted quaternary carbon, could favor the Cu(I) pathway. For recent reports regarding the mechanism of non-catalytic allylic substitution reactions with alkyl- and allylcopper reagents, see: (a) Sofia, A.; Karlström, E.; Bäckvall, J.-E. *Chem.-Eur. J.* **2001**, *7*, 1981–1989. (b) Yoshikai, N.; Zhang, S.-L.; Nakamura, E. *J. Am. Chem. Soc.* **2008**, *130*, 12862–12863. (c) Bartholomew, E. R.; Bertz, S. H.; Cope, S.; Murphy, M.; Ogle, C. A. *J. Am. Chem. Soc.* **2008**, *130*, 11244–11245.
- (20) For examples of catalytic enantioselective vinyl conjugate additions to unsaturated carbonyls, affording tertiary C–C bonds, see: (a) Oi, S.; Taira, A.; Honma, Y.; Inoue, Y. *Org. Lett.* **2003**, *5*, 97–99. (b) Oi, S.; Sato, T.; Inoue, Y. *Tetrahedron Lett.* **2004**, *45*, 5051–5055. (c) Otomaru, Y.; Hayashi, T. *Tetrahedron: Asymmetry* **2004**, *15*, 2647–2651. (d) Nicolaou, K. C.; Tang, W.; Dagneau, P.; Faraoni, R. *Angew. Chem., Int. Ed.* **2005**, *44*, 3874–3879. (e) Nakao, Y.; Chen, J.; Imanaka, H.; Hiyama, T.; Ichikawa, Y.; Duan, W.-L.; Shintani, R.; Hayashi, T. *J. Am. Chem. Soc.* **2007**, *129*, 9137–9143. (f) Lee, K.-s.; Hoveyda, A. H. *J. Org. Chem.* **2009**, *74*, 4455–4462.

(18) Cu complexes derived from NHC–Ag complexes **1a–c** afford the desired EAS products in similar yields ( $\pm 5\%$ ) and enantioselectivities ( $\pm 3\%$ ).

**Scheme 8.** Proposed Mechanistic Model and Key Features of the Chiral Bidentate NHC–Cu Complex

congested electrophilic site, identification of a class of vinylmetal reagents, which can be easily and efficiently accessed and are only sufficiently nucleophilic to allow the catalytic process to predominate, has been a longstanding complication. The present investigations put forth protocols that address both of the aforementioned challenges: a Ni-catalyzed reaction involving readily available alkynes and inexpensive dibal-H, generating vinylaluminums that are used in situ for efficient and highly site- and enantioselective EAS reactions promoted by chiral NHC–Cu complexes.

Together with previous disclosures regarding metal-catalyzed EAS as well as conjugate additions involving alkyl- and aryl-substituted zinc and aluminum reagents,<sup>12</sup> and the more recent examples delivering boron-substituted quaternary carbon stereogenic centers,<sup>23</sup> the present findings further underline the unique catalytic activity of sulfonate-based bidentate chiral NHC–Cu complexes. The design of additional classes of NHC-

based catalysts that exhibit exceptional reactivity and deliver high selectivity and the development of corresponding catalytic enantioselective protocols, including those that involve various vinyl units, are in progress and will be disclosed in due course.

**Acknowledgment.** The NIH (GM-47480) provided financial support. Y.L. is grateful for an AstraZeneca graduate fellowship. We thank Tricia L. May for helpful discussions. Mass spectrometry facilities at Boston College are supported by the NSF (DBI-0619576).

**Supporting Information Available:** Experimental procedures and spectral data for substrates and products. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JA106829K

(21) For examples of catalytic enantioselective vinyl additions to carbonyls, affording tertiary C–C bonds, see: (a) Oppolzer, W.; Radinov, R. N. *J. Am. Chem. Soc.* **1993**, *115*, 1593–1594. (b) Miller, K. M.; Huang, W.-S.; Jamison, T. F. *J. Am. Chem. Soc.* **2003**, *125*, 3442–3443. (c) Tomita, D.; Wada, R.; Kanai, M.; Shibasaki, M. *J. Am. Chem. Soc.* **2005**, *127*, 4138–4139. (d) Yang, Y.; Zhu, S.-F.; Zhou, C.-Y.; Zhou, Q.-L. *J. Am. Chem. Soc.* **2008**, *130*, 14052–14053. (e) Kerrigan, M. H.; Jeon, S.-J.; Chen, Y. K.; Carroll, P. J.; Walsh, P. J. *J. Am. Chem. Soc.* **2009**, *131*, 8434–8445.

(22) For examples of catalytic enantioselective vinyl additions to aldimines, affording tertiary C–C bonds, see: (a) Patel, S. J.; Jamison, T. F. *Angew. Chem., Int. Ed.* **2004**, *43*, 3941–3944. (b) Kong, J.-R.; Cho, C.-W.; Krische, M. J. *J. Am. Chem. Soc.* **2005**, *127*, 11269–11276. (c) Ngai, M.-Y.; Barchuk, A.; Krische, M. J. *J. Am. Chem. Soc.* **2007**, *129*, 12644–12645. (d) Lou, S.; Schaus, S. E. *J. Am. Chem. Soc.* **2008**, *130*, 6922–6923. (e) Nakao, Y.; Takeda, M.; Chen, J.; Salvi, L.; Hiyama, T.; Ichikawa, Y.; Shintani, R.; Hayashi, T. *Chem. Lett.* **2008**, *37*, 290–291.

(23) Guzman-Martinez, A.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2010**, *132*, 10634–10637.