

## Holo-Ni(II)*HpNikR* Is an Asymmetric Tetramer Containing Two Different Nickel-Binding Sites

Abby L. West, Franz St. John, Pedro E. M. Lopes, Alexander D. MacKerell, Jr.,\*  
Edwin Pozharski,\* and Sarah L. J. Michel\*

Department of Pharmaceutical Sciences, School of Pharmacy, University of Maryland,  
Baltimore, Maryland 21201-1180, United States

Received May 13, 2010; E-mail: smichel@rx.umaryland.edu

**Abstract:** The metalloregulatory protein NikR from *Helicobacter pylori* (*HpNikR*) is a master regulator of gene expression which both activates and represses specific genes in response to nickel availability. Here, we report the first crystal structure (at 2.37 Å resolution) of Ni(II)*HpNikR* prepared directly from the holo protein. The protein contains four nickel ions located in two distinct coordination environments. Two nickel ions are bound to sites in a four-coordinate square-planar geometry as predicted on the basis of the structures of NikR from *Escherichia coli* and *Pyrococcus horikoshii*. The remaining two nickel ions are bound to sites with unexpected 5- or 6-coordinate geometries which were previously thought to be involved in nickel incorporation into the protein. The nickel with 5-/6-coordinate geometry utilizes three histidines from two separate monomeric *HpNikR* units along with two or three water molecules as ligands. The spatial location of the nickel in the 5-/6-coordinate site is within approximately 5 Å of the expected site if a 4-coordinate square-planar geometry occurred. Two of the histidines that participate as ligands in the 5-/6-coordinate site would also participate as ligands if the 4-coordinate site was occupied, making it impossible for both sites to be occupied simultaneously. DFT calculations show that the 5-/6-coordinate geometries are energetically favorable when the local protein environment is included in the calculations. The presence of two distinct coordination environments in *HpNikR* is suggested to be related to the specificity and binding affinity of this transcription factor for DNA.

### Introduction

*Helicobacter pylori* (*H. pylori* or *Hp*) is a virulent bacterium that colonizes the highly acidic, mucosal layer (or gastric epithelium) of the human stomach, causing gastritis and in some cases gastric cancer.<sup>1–4</sup> *H. pylori* infection constitutes a significant public health threat with more than half of the world's population infected. There is an urgent need for the development of novel anti-*Helicobacter* agents.<sup>5</sup> *H. pylori* are among a select number of bacteria that are able to survive in the highly acidic environment of the gastric epithelium. The pH of the gastric epithelium ranges from 4 to 6.5 with periodic acid shocks of pH < 2.<sup>6</sup>

One feature that enables *H. pylori* to survive under acidic conditions is that it releases large quantities of ammonia to neutralize its immediate environment.<sup>6,7</sup> Much of this ammonia is produced by urease, which is a nickel-dependent enzyme that converts urea to ammonia and bicarbonate.<sup>6,8</sup> *H. pylori* contains high levels of urease (~10% of total protein content), and the

metalloregulatory protein NikR regulates its expression.<sup>6,9,10</sup> *HpNikR* activates the transcription of urease in response to nickel by binding to a specific region of the promoter of *ureA* that encodes for urease.<sup>11–14</sup> In addition to regulating urease production, *HpNikR* regulates the transcription of a large number of genes either directly or indirectly.<sup>15</sup> The genes for which direct regulation have been established contain variable recognition sequences;<sup>11–14,16</sup> and the physical basis for DNA recognition and discrimination by *HpNikR* is currently unresolved.

*HpNikR* has homologues in both Gram-negative and Gram-positive bacteria.<sup>17</sup> The best studied homologue of NikR is from

- (1) Marshall, B. J.; Warren, J. R. *Lancet* **1984**, *1*, 1311–1315.
- (2) Cover, T. L.; Blaser, M. J. *Annu. Rev. Med.* **1992**, *43*, 135–145.
- (3) Sepulveda, A. R.; Coelho, L. G. *Helicobacter* **2002**, *7* (Suppl. 1), 37–42.
- (4) Kusters, J. G.; van Vliet, A. H.; Kuipers, E. J. *Clin. Microbiol. Rev.* **2006**, *19*, 449–490.
- (5) Loughlin, M. F. *Expert. Opin. Ther. Targets* **2003**, *7*, 725–735.
- (6) van Vliet, A. H.; Ernst, F. D.; Kusters, J. G. *Trends Microbiol.* **2004**, *12*, 489–494.
- (7) van Vliet, A. H.; Kuipers, E. J.; Stoof, J.; Poppelaars, S. W.; Kusters, J. G. *Infect. Immun.* **2004**, *72*, 766–773.

- (8) Carter, E. L.; Flugge, N.; Boer, J. L.; Mulrooney, S. B.; Hausinger, R. P. *Metallomics* **2009**, *1*, 207–221.
- (9) Li, Y.; Zamble, D. B. *Chem. Rev.* **2009**, *109*, 4617–4643.
- (10) van Vliet, A. H.; Poppelaars, S. W.; Davies, B. J.; Stoof, J.; Bereswill, S.; Kist, M.; Penn, C. W.; Kuipers, E. J.; Kusters, J. G. *Infect. Immun.* **2002**, *70*, 2846–2852.
- (11) Benanti, E. L.; Chivers, P. T. *J. Biol. Chem.* **2007**, *282*, 20365–20375.
- (12) Delany, I.; Ieva, R.; Soragni, A.; Hilleringmann, M.; Rappuoli, R.; Scarlato, V. *J. Bacteriol.* **2005**, *187*, 7703–7715.
- (13) Dosanjh, N. S.; Hammerbacher, N. A.; Michel, S. L. *Biochemistry* **2007**, *46*, 2520–2529.
- (14) Abraham, L. O.; Li, Y.; Zamble, D. B. *J. Inorg. Biochem.* **2006**, *100*, 1005–1014.
- (15) Contreras, M.; Thiberge, J. M.; Mandrand-Berthelot, M. A.; Labigne, A. *Mol. Microbiol.* **2003**, *49*, 947–963.
- (16) Dosanjh, N. S.; West, A. L.; Michel, S. L. *Biochemistry* **2009**, *48*, 527–536.

*Escherichia coli*, EcNikR,<sup>18,19</sup> which has low sequence similarity with HpNikR (21% identity and 54% homology). Unlike HpNikR, which regulates multiple genes as either an activator or a repressor,<sup>20</sup> EcNikR regulates a single gene as a repressor.<sup>21</sup> Several crystal structures of EcNikR have been reported.<sup>22–24</sup> These structures revealed that EcNikR folds as a symmetrical tetramer with a central tetrameric metal-binding domain (MBD) flanked by two dimeric DNA-binding domains (DBD). Nickel binds to the metal-binding domains via three ligands from one monomer (His<sub>2</sub>Cys) and one ligand from another monomer (His) in a 4-coordinate square-planar geometry.<sup>22–24</sup> The four nickel ions each occupy the analogous site on each monomer. Structures of NikR from *Pyrococcus horikoshii* have also been reported, and again the protein is a symmetrical tetramer with four nickel ions per tetramer in a square-planar geometry.<sup>25</sup> The histidine and cysteine ligands that bind nickel in *E. coli* NikR and *P. horikoshii* NikR are conserved in *H. pylori* NikR; thus, it was presumed that HpNikR would also bind four nickel ions at four identical square-planar sites.

There are no reported structures of holo-Ni(II)HpNikR, although there are two crystal structures of nickel bound to HpNikR that were generated by soaking excess nickel (20–100 mM) into crystals of apo-HpNikR.<sup>26</sup> In one of these structures, named Ni1-HpNikR (PDB code 2CAD), three nickel sites are observed with six nickel atoms per HpNikR tetramer. One of the sites is the predicted square-planar site called final (“F”), the second is a novel octahedral nickel site that is thought to be an intermediate nickel-binding site (“I”), and the third is another novel octahedral site, referred to as an external binding site (“X”), made up of ligands from the protein and citrate from the crystallization conditions. In the second structure, Ni2-HpNikR (pdb code 2CAJ) nickel is bound at a site between the “I” and the “X” sites of Ni1-HpNikR with just two histidine ligands. Because these two structures were generated by soaking apo-HpNikR with excess nickel, the unexpected nickel sites were proposed to represent ‘snapshots’ of nickel incorporation serving as sites for nickel to bind enroute to its final holo-Ni(II)HpNikR state. The true holo-Ni(II)HpNikR structure was predicted to house a total of four nickel ions in a square-planar coordination geometry at identical sites.

In this contribution we report the 2.37 Å resolution crystal structure of holo-Ni(II)HpNikR. While initial crystallization efforts targeted the holo-Ni(II)HpNikR-DNA complex structure, we obtained a crystal structure of holo-Ni(II)HpNikR in the absence of DNA. While unexpected, this structure allowed us to test the hypothesis that holo-Ni(II)HpNikR, like its homologues in *E. coli* and *P. horikoshii*, contains four square-planar nickel sites. Surprisingly, the structure does not contain nickel

bound to four identical square-planar nickel sites as predicted. Instead, the protein has only two of the nickel ions bound to the predicted square-planar coordination sites. The additional two nickel ions, while bound in the vicinity of the two remaining predicted sites, assume a 5 and a 6-coordinate geometry. These nickels are in locations analogous to site “I” in the Ni1HpNikR structure and are coordinated to three histidine ligands and two water molecules forming a square-pyramidal geometry in one case and to three histidine ligands and three water molecules forming an octahedral geometry in the other case, respectively. When nickel is bound to these sites in either the 5- or 6-coordinate geometry, it is in a location that is approximately 5 Å removed from where the nickel is predicted to be located if bound in a square-planar geometry. To verify the geometry and investigate the energetics of the nickel sites, we have performed DFT calculations. Based on this combination of experimental and computational methods it is shown that two of the four nickel ions found in holo-Ni(II)HpNikR bind in either 5- or 6-coordination geometries, in locations adjacent to but distinct from the predicted square-planar site; in the remainder of this manuscript these two sites when discussed collectively will be referred to as the 5-/6-site. The implications of the presence of two of the bound nickels in 5 or 6 coordinate geometries in addition to the two nickels in the 4-coordinate geometry on Ni(II)HpNikR’s propensity to recognize multiple, different DNA sequences are discussed.

## Materials and Methods

**Holo-Ni(II)HpNikR Overexpression and Purification.** A pET15b construct containing HpNikR was transformed into *E. coli* BL21(DE3) cells and grown in LB medium containing 100 µg/mL ampicillin.<sup>13</sup> Purification of Holo-Ni(II)HpNikR was carried out using a nickel-loaded 5 mL Hitrap-HP (Amersham) affinity column followed by further purification using a Hitrap Q-Sepharose (Amersham) column. Holo-Ni(II)HpNikR was exchanged into final buffer (20 mM Hepes, 100 mM NaCl pH 8.0) and concentrated to 10 mg/mL for initial crystal screening. HpNikR as isolated was tested for metal content with inductively coupled plasma mass spectroscopy (ICP-MS) and found to contain 4 nickel ions/tetramer. Oligonucleotides with the sequence, 5′-AATATAACACTAATTCATTTTAAATAATAATTAG-3′ were purchased from Integrated DNA Technologies (IDT) in the desalted form and used in crystallization trials. The oligonucleotides were purified by reverse-phase HPLC (Waters) using a Symmetry 300 C18 column and applying a linear gradient. Purified DNA was lyophilized, resuspended in 20 mM Hepes, 100 mM NaCl pH 7.5, and annealed at 95 °C.

**Fluorescence Anisotropy To Verify DNA Binding.** To confirm that holo-Ni(II)HpNikR bound DNA, a fluorescence anisotropy titration of the protein with PureA (which corresponds to the ureA promoter) was performed. A 10 nM solution of fluorescently labeled DNA (PureA-F-5′-CTTCAAGATATAACACTAATT[F]CATTTTAAATAATAATTAGTTAATGAA-3′) in 20 mM HEPES, 100 mM NaCl, and 20 mM glycine (pH 7.5) was added to a 0.5 cm PL Spectrosil far-UV quartz window fluorescence cuvette (Starna Cells) pretreated with 5 mM BSA to prevent adherence of either the protein or DNA to the cuvette walls. The anisotropy, *r*, of the free DNA oligonucleotide was measured. Holo-Ni(II)HpNikR was titrated and the change in anisotropy measured. Data were fit to a 1:1 binding equilibrium as previously reported.<sup>13</sup>

**Initial Crystal Screening.** Crystal screens were performed using 96-well sitting drop crystallization trays (Art Robbins Instruments, Sunnyvale, California, U.S.A.) with the OryxNano crystallization robot (Douglas Instruments Ltd., Berkshire, UK) at the X-ray Crystallography Core Facility at the University of Maryland, Baltimore. Crystallization drops were set as 50% protein/DNA (protein/DNA 1:1.2 equivalents) to an initial drop volume of 1.4

- (17) Dosanjh, N. S.; Michel, S. L. *Curr. Opin. Chem. Biol.* **2006**, *10*, 123–130.
- (18) Chivers, P. T.; Sauer, R. T. *Protein Sci.* **1999**, *8*, 2494–2500.
- (19) de Pina, K.; Desjardin, V.; Mandrand-Berthelot, M. A.; Giordano, G.; Wu, L. F. *J. Bacteriol.* **1999**, *181*, 670–674.
- (20) Ernst, F. D.; Kuipers, E. J.; Heijens, A.; Sarwari, R.; Stoof, J.; Penn, C. W.; Kusters, J. G.; van Vliet, A. H. *Infect. Immun.* **2005**, *73*, 7252–7258.
- (21) Chivers, P. T.; Sauer, R. T. *Chem. Biol.* **2002**, *9*, 1141–1148.
- (22) Schreiter, E. R.; Sintchak, M. D.; Guo, Y.; Chivers, P. T.; Sauer, R. T.; Drennan, C. L. *Nat. Struct. Biol.* **2003**, *10*, 794–799.
- (23) Schreiter, E. R.; Wang, S. C.; Zamble, D. B.; Drennan, C. L. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 13676–13681.
- (24) Phillips, C. M.; Schreiter, E. R.; Guo, Y.; Wang, S. C.; Zamble, D. B.; Drennan, C. L. *Biochemistry* **2008**, *47*, 1938–1946.
- (25) Chivers, P. T.; Tahirov, T. H. *J. Mol. Biol.* **2005**, *348*, 597–607.
- (26) Dian, C.; Schauer, K.; Kapp, U.; McSweeney, S. M.; Labigne, A.; Terradot, L. *J. Mol. Biol.* **2006**, *361*, 715–730.

$\mu\text{L}$ . A crystal was obtained in condition No. 5 of the NeXtal Nucleix Suite (0.05 M MES, 0.2 M potassium chloride, 0.01 M magnesium sulfate and 5% PEG 8000, pH 5.6) after 4 days of incubation at 21.5 °C. This crystal was cryoprotected in mother liquor supplemented with 12% glycerol and flash cooled in liquid nitrogen prior to data collection.

**Data Collection, Processing, and Model Building.** X-ray diffraction data were collected remotely at the Stanford Synchrotron Radiation Lightsource (SSRL, beamline 12-2). Data processing was done with DENZO/SCALEPACK.<sup>27</sup> The structure was solved by molecular replacement using the nickel-binding and DNA-binding domains as separate models<sup>26</sup> (PDB ID 2CA9). PHASER<sup>28</sup> was used for molecular replacement in CCP4.<sup>29,30</sup> Surprisingly, no DNA was found in the asymmetric unit, which contains four metal-binding domains and only two DNA-binding domains. Iterative model building was performed using Coot for manual model building<sup>31</sup> with REFMAC for restrained refinement.<sup>32,33</sup> Tight NCS restraints were applied between identical monomers in the nickel-binding domain core tetramer. Final rounds of refinement included NCS with TLS/restrained refinement.<sup>34–36</sup> Anomalous scattering was performed on a holo-Ni(II)HpNikR crystal in order to confirm the presence of nickel. Distance restraints were applied for nickel-coordinating side chains and waters. Anisotropic temperature factors were refined for nickel ions with the rest of the model treated isotropically. Figures were generated using PyMOL (www.pymol.org).

During the refinement of the structure, it was observed that DNA was absent in the crystal. In addition, it appeared that the cleavage of the protein may have occurred. This was confirmed by N-terminal sequencing of a protein/DNA solution that had been used to grow crystals, which revealed that the protein had been cleaved between residues Trp54 and Ala55 (Trp54 is at the interface of the DNA-binding domain and the metal-binding domain).

**pH-Dependent Ni(II)-Binding Titrations.** Apo-HpNikR was prepared as previously described.<sup>13</sup> Metal-binding titrations were performed in screw-capped quartz cuvettes using a Perkin Elmer Lambda 25 UV-vis spectrometer. Titrations were carried out in 20 mM potassium phosphate (and 20 mM glycine at pH 5.6) using metal-free reagents and water that had been purified using a Milli-Q purification system and passed over Sigma Chelex resin. Upon their preparation, buffers were purged with helium to degas and transferred into a Coy inert atmospheric chamber. Ni(II)sulfate hexahydrate (99.99% Ni) from Puratrem (Strem Chemicals) was used as the nickel source. The binding interaction between apo-HpNikR and Ni(II) was measured at four pH conditions: pH 5.6, 6.5, 7.5, and 8.5 (typically with 45  $\mu\text{M}$  apo-HpNikR). Titrations were performed in triplicate under anaerobic conditions.

**Density Functional Theory Calculations.** Model complexes representative of the coordination modes of the Ni atoms in HpNikR were used to study the electronic and geometric structures of the experimentally determined 5- and 6-coordinate metal sites as well

as the 4-coordinate sites. Large clusters obtained from the crystallographic coordinates included all coordinating residues, neighboring residues with a structural role in defining the coordination geometries, for example Gln87, and all residues connecting them as well as the two or three water molecules directly coordinated to the metal when present. Residues that do not play a direct structural role at the Ni centers were modeled as glycines. Anchor atoms were selected and kept frozen during the minimization of the geometries of the large clusters to mimic the structural impact of the protein on the residues adjacent to the nickel ions. These atoms were either the carbonyl C or N of the protein backbone or the terminal  $-\text{COO}^-$  and  $-\text{NH}_3^+$  groups, respectively. Arrows in Figure S5 (Supporting Information) indicate the anchor atoms. The use of large clusters allows inclusion of the effects of the protein medium in the calculations in a more rigorous way. This comes with the penalty of additional computational cost since the size of the largest model is 193 atoms. Smaller complexes of the 5- and 6-coordinated Ni(II) centers consisting of Ni and just the coordinating ligands: two or three waters and three imidazole rings (to represent histidine) were also optimized to study the electronic structures.

NWChem version 5.1.1, as developed and distributed by Pacific Northwest National Laboratory, P.O. Box 999, Richland, Washington 99352, U.S.A., and funded by the U.S. Department of Energy, was used to obtain the computational results.<sup>37,38</sup> All calculations were based on density functional theory (DFT)<sup>39–41</sup> and employed the B3LYP (Becke 3-parameter, Lee, Yang and Parr, exchange-correlation) hybrid functional<sup>42–44</sup> with the LANL2TZ basis set which includes effective core potentials for Ni<sup>45</sup> and the 6-31G\* and 3-21G basis sets, respectively, for the lighter atoms (H, C, N, O, S).<sup>46–49</sup> In the smaller model complexes, all lighter atoms were described by the 6-31G\* basis set. In the large clusters only atoms coordinating Ni or participating in hydrogen bonds with coordinating ligands were described by the 6-31G\* basis, the remaining being described by the 3-21G basis. All structures were fully geometry optimized at the B3LYP level of theory until convergence to a root-mean-square value of the energy gradient and nuclear step of  $4.5 \times 10^{-4}$  hartree/bohr and  $1.8 \times 10^{-3}$  bohr, respectively. Contour plots were made with gOpenMol 3.00 (gOpenMol can be downloaded from <http://www.csc.fi/english/pages/gOpenMol>), and optimized geometries were plotted using a version of Ortep-3 for Microsoft Windows.<sup>50</sup> VMD was used to plot structures of the large models superimposed on the protein.<sup>51</sup>

## Results and Discussion

**Overall Structure.** Crystal trays were set up with holo-Ni(II)HpNikR and *PureA* DNA in a 1:1.2 ratio. The crystals

- (27) Otwinowski, Z.; Minor, W. *Macromol. Cryst., Pt. A* **1997**, *276*, 307–326.
- (28) McCoy, A. J.; Grosse-Kunstleve, R. W.; Adams, P. D.; Winn, M. D.; Storoni, L. C.; Read, R. J. *J. Appl. Crystallogr.* **2007**, *40*, 658–674.
- (29) Bailey, S. *Acta Crystallogr., Sect. D* **1994**, *50*, 760–763.
- (30) Potterton, E.; Briggs, P.; Turkenburg, M.; Dodson, E. *Acta Crystallogr., Sect. D* **2003**, *59*, 1131–1137.
- (31) Emsley, P.; Cowtan, K. *Acta Crystallogr., Sect. D* **2004**, *60*, 2126–2132.
- (32) Murshudov, G. N.; Vagin, A. A.; Dodson, E. J. *Acta Crystallogr., Sect. D* **1997**, *53*, 240–255.
- (33) Vagin, A. A.; Steiner, R. A.; Lebedev, A. A.; Potterton, L.; McNicholas, S.; Long, F.; Murshudov, G. N. *Acta Crystallogr., Sect. D* **2004**, *60*, 2184–2195.
- (34) Painter, J.; Merritt, E. A. *Acta Crystallogr., Sect. D* **2006**, *62*, 439–450.
- (35) Winn, M. D.; Isupov, M. N.; Murshudov, G. N. *Acta Crystallogr., Sect. D* **2001**, *57*, 122–133.
- (36) Winn, M. D.; Murshudov, G. N.; Papiz, M. Z. *Macromol. Cryst., Pt. D* **2003**, *374*, 300–321.

- (37) Kendall, R. A.; Aprà, E.; Bernholdt, D. E.; Bylaska, E. J.; Dupuis, M.; Fann, G. I.; Harrison, R. J.; Ju, J.; Nichols, J. A.; Nieplocha, J.; Straatsma, T. P.; Windus, T. L.; Wong, A. T. *Comput. Phys. Commun.* **2000**, *128*, 260–283.
- (38) Bylaska, E. J.; et al. *NW Chem, A Computational Chemistry Package for Parallel Computers*, Version 5.1.1; Pacific Northwest National Laboratory: Richland, WA, 2007.
- (39) Hohenberg, P.; Kohn, W. *Phys. Rev.* **1964**, *136*, B864.
- (40) Kohn, W.; Sham, L. J. *Phys. Rev.* **1965**, *140*, A1133.
- (41) Parr, R. G.; Yang, W. *Density-Functional Theory of Atoms and Molecules*; Oxford University Press: New York, 1989.
- (42) Becke, A. D. *J. Chem. Phys.* **1993**, *98*, 1372–1378.
- (43) Kim, K.; Jordan, K. D. *J. Phys. Chem.* **1994**, *98*, 10089–10094.
- (44) Stephens, P. J.; Devlin, F. J.; Chabalowski, C. F.; Frisch, M. J. *J. Phys. Chem.* **1994**, *98*, 11623–11627.
- (45) Hay, P. J.; Wadt, W. R. *J. Chem. Phys.* **1985**, *82*, 299.
- (46) Binkley, J. S.; Pople, J. A.; Hehre, W. J. *J. Am. Chem. Soc.* **1980**, *102*, 939–947.
- (47) Ditchfield, R.; Hehre, W. J.; Pople, J. A. *J. Chem. Phys.* **1971**, *54*, 724–728.
- (48) Hariharan, P. C.; Pople, J. A. *Theor. Chim. Acta* **1973**, *28*, 213–222.
- (49) Hehre, W. J.; Ditchfield, R.; Pople, J. A. *J. Chem. Phys.* **1972**, *56*, 2257–2261.
- (50) Barnes, C. J. *J. Appl. Crystallogr.* **1997**, *30*, 568.

that formed were found to contain just the protein (Figure 1A). A similar result was reported by O'Halloran and Mondragon when they sought to obtain crystals of the zinc metalloregulators CueR and ZntR in complex with their DNA partners.<sup>52</sup> The structure of holo-Ni(II)*Hp*NikR was determined at 2.37 Å resolution, and data processing and refinement statistics are shown in Table 1. The asymmetric unit (Figure 1B) contained four NikR monomers, two of which were missing the N-terminal fragment that forms the DNA-binding domains (DBD). The metal-binding domain (MBD) is of ferredoxin-like ( $\beta\alpha\beta\beta\alpha\beta$ ) fold, and the DBD may be described as a ribbon–helix–helix fold.<sup>22,25,26,53</sup>

**Domain Swapping.** The unit cell contained density for a tetrameric metal-binding subunit but only two of the four DNA-binding domains, which was unexpected. The two missing DNA-binding domains had been cleaved during crystallization. A complete tetrameric structure is observed due to domain swapping of the DNA-binding domains between adjacent unit cells. The tetrameric structure is seen by taking two MBDs from one asymmetric unit and two MBDs from the adjacent unit cell with two symmetrical faces of the DBDs that make identical contacts with respective MBDs. This strings together unit cells throughout the crystal yielding the pseudo-tetrameric structures (Figure 1C). The domain swapping does not appear to affect the structure of the metal-binding domains as we observe low RMSDs (<1.3 Å for C $\alpha$  atoms) with respect to the MBDs of previously reported apo-*Hp*NikR structures. In addition, we have recently solved the X-ray structure of just the metal-binding domain of holo-Ni(II)*Hp*NikR (data not shown) to 3.1 Å resolution. From this structure, we are able to determine that the metal-binding domains and the nickel sites are identical to those in the crystal structure of holo-Ni(II)*Hp*NikR that we report here.

**Ni(II)-Binding Sites.** In the presence of four ligands, Ni(II) preferentially adopts a square-planar coordination geometry while most other divalent metal ions adopt a tetrahedral geometry.<sup>54</sup> The presence of four sets of ligands available to bind nickel in a square-planar geometry in the apo*Hp*NikR structure and the observation that nickel is bound to all four of these sets of ligands in a square-planar geometry in the structures of holo-*Ec*NikR and holo-*Ph*NikR, led us to predict that holo-Ni(II)*Hp*NikR would also house four nickel ions in four square-planar sites. Surprisingly, we found that holo-Ni(II)*Hp*NikR is a heterotetramer with two unique classes of nickel sites. Nickels Ni1 and Ni2 are found in square-planar sites (Figure 2A), consistent with that seen in the *Ec* and *Ph* structures. Each nickel ion (Ni1 and Ni2) is ligated to His99, His101, and Cys107 from one MBD and His88 from a second MBD, as expected.

The remaining two nickel ions (Ni3 and Ni4) present in the holo-Ni(II)*Hp*NikR structure are coordinated to three histidine residues (His 74 and His 101 from one MBD and His 88 from the second MBD) as well as 2 or 3 water molecules making a 5-coordinate, square-pyramidal geometry in one case (Ni3) and an 6-coordinate, octahedral geometry (Ni4) in the other case (Figure 2B and C, respectively). This second site is analogous

to the site that was named “I” for intermediate in the nickel-soaked structure, Ni1-*Hp*NikR.<sup>26</sup> The second site has two ligands, His 101 and His 88, in common with the anticipated square-planar geometry, making it prohibitive for two nickel ions to simultaneously participate in the two coordination geometries at this second class of sites (Figure S1, Supporting Information). Inspection of the apo-*Hp*NikR structure indicates that the ligands that coordinate nickel at the 5-/6-coordinate sites are preorganized, which may facilitate binding in this geometry.

**Hydrogen-Bonding Interactions.** One of the water molecules at each of the 5-/6-coordinate sites is stabilized by a hydrogen-bonding interaction with Gln87 (Figure 2B and C), (also seen in the “I” site of the nickel-soaked apo-*Hp*NikR protein, Ni1-*Hp*NikR)<sup>26</sup> which DFT calculations (described below) posit as important for stabilizing the nickel–aquo bond. These interactions appear to lead to two of the Ni(II)-binding sites assuming the unexpected 5- or 6-coordinate geometries. Ligation in either of these geometries locates the nickel approximately 5 Å from its position if ligated in the anticipated square-planar geometry. While this result was unanticipated, the presence of different metal sites in families of metallotranscription factors is not without precedent. For example, Glasfeld and co-workers recently reported a novel metal site in ScaR, a member of the DtxR/MntR family of metalloregulators.<sup>55</sup>

Hydrogen bonding between the occupied 5-/6-coordinate sites and the occupied 4-coordinate sites in *H. pylori* NikR is limited. This contrasts *E. coli* NikR where an extensive hydrogen-bonding network linking two individual square-planar sites was mapped out by Drennan and co-workers.<sup>22</sup> The *E. coli* NikR hydrogen-bonding network is composed of a Glu97, Gln75, and Tyr58 (Glu109, Gln87, and Val70 in *Hp*NikR). Replacement of the tyrosine involved in *Ec*NikR hydrogen bonding by valine in *Hp*NikR disrupts the hydrogen bonding between the two sites.

**Relationship between Coordination Chemistry and the Method Used To Generate Crystals.** The two previously reported structures of Ni(II)*Hp*NikR were generated by soaking apo-Ni(II)*Hp*NikR crystals with excess nickel sulfate or nickel chloride (20–100 mM).<sup>26</sup> In these structures, nickel is found bound to multiple different sites including the 5-/6-coordinate site, in addition to the square-planar sites, and neither of the structures has four 4-coordinate sites.<sup>26</sup> Because the structures were generated by soaking nickel with apo-*Hp*NikR that was already crystallized, it was thought that the structures did not represent the ground state associated with the ‘holo’ form of NikR. However, as we report here, holo-Ni(II)*Hp*NikR does not contain four 4-coordinate sites but rather two 4-coordinate sites and two 5-/6-coordinate sites, with the latter two sites lying approximately 5 Å from what would correspond to the 4-coordinate ligation geometry. Because this structure was obtained directly from holo-Ni(II)*Hp*NikR, it is more likely representative of the conformation of the holo form of the protein. Thus, we propose that holo-Ni(II)*Hp*NikR in the absence of DNA contains two distinct and stable classes of high affinity sites—the expected square-planar sites and the unanticipated 5-/6-coordinate sites.

**Comparison to Isothermal Titration Calorimetry Data.** There is solution-based evidence to support the presence of two different high-affinity nickel-binding sites in Ni(II)*Hp*NikR. Ciurli, Scarlato, and co-workers used isothermal titration


(51) Humphrey, W.; Dalke, A.; Schulten, K. *J. Mol. Graph.* **1996**, *14*, 33–38.

(52) Changela, A.; Chen, K.; Xue, Y.; Holschen, J.; Outten, C. E.; O'Halloran, T. V.; Mondragon, A. *Science* **2003**, *301*, 1383–1387.

(53) Schreiber, E. R.; Drennan, C. L. *Nat. Rev. Microbiol.* **2007**, *5*, 710–720.

(54) Cotton, F. A.; Wilkinson, G. *Advanced Inorganic Chemistry*, 5th ed.; John Wiley and Sons: New York, 1988.

(55) Stoll, K. E.; Draper, W. E.; Kliegman, J. I.; Golynskiy, M. V.; Brew-Appiah, R. A.; Phillips, R. K.; Brown, H. K.; Breyer, W. A.; Jakubovics, N. S.; Jenkinson, H. F.; Brennan, R. G.; Cohen, S. M.; Glasfeld, A. *Biochemistry* **2009**, *48*, 10308–20.




**Figure 1.** (A) Ribbon diagram of holo-Ni(II)HpNikR tetramer with the metal-binding domains (MBD) colored red and the DNA-binding domains (DBD) colored blue (dark blue DBD from one asymmetric unit and teal blue from a second asymmetric unit). The 4-coordinate, square-planar nickel atoms are colored dark green and labeled Ni1 and Ni2. The 5-/6-coordinate nickel atoms are colored light green and labeled Ni3 for the 6-coordinate site and Ni4 for the 5-coordinate site. The locations of the sites named “I” and “X” are indicated by arrows. (B) Asymmetric unit of holo-Ni(II)HpNikR. Only the symmetry mates completing the MBDs are shown. (C) Structure of holo-Ni(II)HpNikR the showing domain-swapping phenomenon at the DBD. All figures were prepared using PyMol.

**Table 1.** Data Collection and Model Refinement Statistics

parameter	<i>HpNikR</i> PDB code 3LGH
Data Collection	
wavelength	0.97946
space group	$P2_12_12_1$
unit cells $a$ (Å), $b$ (Å), $c$ (Å)	89.5, 105.1, 48.2
resolution range (Å)	50.0–2.37
redundancy	4.3 (1.8)
no. of unique reflections	16660 (1,316)
$R_{\text{merge}}/R_{\text{pim}}$ (%) <sup>60,61</sup>	18.6/9.4
completeness (%)	90.5 (73.6)
mean $[I/\sigma_I]$	7.0 (1.0)
Matthews coefficient	2.22
Refinement Statistics	
resolution range	39.3–2.37
no. of reflections	16595
$R_{\text{cryst}}/R_{\text{work}}$ (%) <sup>62,63</sup>	27.1/26.8
$R_{\text{free}}$ no. of reflections)	31.8 (857)
no. of protein atoms	3472
no. of water molecules	62
average $B_f$ (Å)	42.8
Ramachandran Statistics	
favoured (%)	95.7
outliers (%)	0.9
RMSD from Ideality	
bond lengths (Å)	0.011
bond angles (deg)	1.326

calorimetry (ITC) to measure nickel binding to apo-*HpNikR* and reported that two of the nickel ions bound with 12 nM affinity and the remaining two nickel ions bound with 125 nM affinity at pH 7.0.<sup>56</sup> We note that Zambale and co-workers used a competitive titration with EGTA and UV visible spectroscopy to measure nickel affinity for *HpNikR* and only observed one high-affinity site (pM binding).<sup>14</sup> Ciurli, Scarlato, and co-workers argue that the EGTA approach yields inaccurate binding constants because EGTA can form polynuclear complexes with Ni(II) and suggest that the binding affinities determined from the direct titration with Ni(II) using ITC are more accurate.<sup>56</sup>

**Role of pH in Nickel-Binding Stoichiometry.** The structures of *EcNikR* have all been reported at pH 7.5–8.5, whereas the structure that we report here was determined at pH 5.6. To determine whether the pH affected the nickel sites in *HpNikR*, titrations of Ni(II) with apo-*HpNikR* at a range of pH's were performed (pH 5.6, 6.5, 7.5, 8.5). The spectra all exhibited an absorbance maximum at 304 nm which saturated upon addition of 1 nickel ion/*HpNikR* (4 nickel ions/tetramer). An overlay of the titrations is shown in Figure S2, Supporting Information. An overlay of the absorbance spectra at all four pH values as a function of extinction coefficient shows no differences in the spectra as a function of pH as shown in Figure S3, Supporting Information. This suggests that the nickel sites are not affected by pH in *HpNikR*. Zambale and co-workers and Michaud-Soret and co-workers have also reported that they do not observe any changes in nickel stoichiometry and extinction coefficient of Ni(II)-*HpNikR* over a similar range of pH.<sup>14,57,58</sup> Interestingly, Michaud-Soret and co-workers do observe a pH dependence




**Figure 2.** (A) Representation of the square-planar nickel site in holo-Ni(II)*HpNikR*. (B) Representation of the 5-coordinate nickel site in holo-Ni(II)*HpNikR*. The electron density around the water molecules is shown. (C) Representation of the 6-coordinate nickel site in holo-Ni(II)*HpNikR*. The electron density around the water molecules is shown.

on nickel binding for *E. coli* *NikR* suggesting that *E. coli* *NikR* may be more sensitive to pH.<sup>57</sup>

**Role of Tryptophan (W54) in Locking the DBD.** The two DNA-binding domains are oriented in a ‘trans’ conformation relative to the MBD in the holo-Ni(II)-*HpNikR* structure (Figure

(56) Zambelli, B.; Bellucci, M.; Danielli, A.; Scarlato, V.; Ciurli, S. *Chem. Commun. (Cambridge)* **2007**, 3649–3651.



**Figure 3.** Representation of the (A) protein backbone and the pocket formed at the interface of the MBD and DBD. (B) Trp54 from the DBD is stabilized by a stacking interaction with Lys64 and Lys140.

3A). Close inspection of the interface reveals that Trp54, which is part of the DBD, lies in a pocket that is stabilized by two lysine residues (Lys64 and Lys140) in the MBD (Figure 3A and B). Due to domain swapping in the holo-Ni(II)HpNikR structure both Trp54 residues from the DBD are located in pockets formed by Lys64/Lys140. We propose that the tryptophan serves to orient the DNA-binding domain perpendicular to the MBD. We also speculate that the tryptophan may play a role in directing the DNA-binding domain to the correct orientation to recognize its partner DNA. This tryptophan is not conserved in *E. coli* NikR and the structures of *E. coli* NikR all have the DNA-binding domains in a linear conformation relative to the MBD.<sup>23</sup> *P. horikoshii* NikR contains this tryptophan, but the protein's DNA targets are not known, limiting any comparison.<sup>25</sup>

**DFT Calculations of the Nickel Sites.** Density functional calculations were performed on the nickel sites of holo-Ni(II)HpNikR using models of the active site to understand the geometries observed in the crystal structures. Ni(II) can form complexes with coordination numbers ranging from 3 to 6.<sup>54,59</sup> Qualitative rationalization of the geometrical preferences of these complexes can be made with the help of the molecular orbital (MO) interaction diagram in Figure S4, Supporting Information. Square-planar coordination is a consequence of the  $d^8$  configuration, since the  $D_{4h}$  ligand field causes one of the d orbitals,  $b_{1g}$  ( $x^2 - y^2$ ), to be high in energy, while another d orbital,  $a_{1g}$  ( $z^2$ ) lowers its energy and approaches the nonbonding manifold,  $e_g + b_{2g}$  ( $xy, xz, yz$ ). Eight electrons can occupy the  $b_{2g}$ ,  $e_g$ , and  $a_{1g}$  orbitals, thus originating diamagnetic complexes. This is

**Table 2.** Selected Bond Distances and Angles for the DFT Optimized Complexes and Experimental X-ray Structure<sup>a</sup>

	6-coordinated (Ni3)		5-coordinated (Ni4)	
	model complex	exper.	model complex	exper.
Ni–O <sub>eq</sub>	2.12	2.1	2.11	2.1
Ni–O <sub>ax</sub>	2.14	2.1	2.07	2.1
Ni–O <sub>ax,2</sub>	2.22	2.1	NA	NA
Ni–N1	2.14	2.1	2.08	2.1
Ni–N2	2.12	2.1	2.15	2.1
Ni–N3	2.13	2.1	2.01	2.1
O <sub>eq</sub> –O1	2.59	2.5	2.64	3.2
O <sub>eq</sub> –O2	NA	NA	2.81	2.6
O1–O3	NA	NA	2.53	3.3
O <sub>ax</sub> –Ni–O <sub>ax,2</sub>	171.69	175.05	NA	NA
O <sub>ax</sub> –Ni–O <sub>eq</sub>	92.29 (79.84)	82 (93)	95.03	71
O <sub>ax</sub> –Ni–N1	86.72 (89.42)	94 (87)	96.05	79
O <sub>ax</sub> –Ni–N2	95.84 (92.02)	89 (96)	92.78	102
O <sub>ax</sub> –Ni–N3	88.40 (94.35)	92 (87)	95.82	91

	4-coordinated (Ni1)		4-coordinated (Ni2)	
	model complex	exper.	model complex	exper.
Ni–S	2.25	2.3	2.26	2.3
Ni–N1	1.91	2.1	1.97	2.1
Ni–N2	2.00	2.1	2.00	2.1
Ni–N3	1.95	2.1	1.92	2.1

<sup>a</sup> Bond distances are in Å, and bond angles are in degrees.

illustrated in the rightmost part of Figure S4 (Supporting Information). Ni(II) can also form 5- and 6-coordinate complexes as illustrated in the central and left and panels of Figure S4, respectively. If the separation between the  $e_g$  and  $t_{2g}$  orbitals is small, the energy gained from forming a triplet state from additional ligands compensates orbital destabilization penalties, leading to 6-coordinate paramagnetic complexes. In 5-coordinated  $C_{4v}$  systems the ( $x^2 - y^2$ ) and ( $z^2$ ) orbitals are no longer degenerate; however due to the electron–electron repulsion contribution, the triplet state will become the ground state, such that the complex is also paramagnetic. Accordingly, both the 5- and 6-coordinate geometries were treated as paramagnetic species in the QM calculations; attempts to optimize both complexes as singlets lead to dissociation of the axial ligands.


**Description of Geometries and Electronic Structure.** The computed DFT geometries are summarized in Table 2 and illustrated in Figures 4 and S5 (Supporting Information). Figure 4 shows details of the coordination around the Ni centers. Relevant hydrogen-bond interactions involving Ni ligands and neighboring groups are included, while atoms comprising the backbone atoms included in the model are omitted for clarity. These atoms are shown in Figure S5 (Supporting Information), which also compares the X-ray experimental and the optimized model cluster structures. A ribbon representation of the local protein backbone is included to help demonstrate how the model clusters were defined. In all cases the coordination is well maintained, although shifts in the coordination spheres, presumably associated with the omission of much of the protein, as described below, are evident. While differences in the structures do exist, as elaborated below, the ability of the DFT calculations to yield coordination geometries consistent with those observed experimentally further validates our crystallographic results.

Optimization of the 6-coordinate large cluster resulted in geometries that differ from the experimental structure (Table 2). Six-coordination around Ni was only possible when the complexes were optimized in their corresponding triplet spin-

(57) Fauquant, C.; Diederix, R. E.; Rodrigue, A.; Dian, C.; Kapp, U.; Terradot, L.; Mandrand-Berthelot, M. A.; Michaud-Soret, I. *Biochimie* **2006**, *88*, 1693–1705.


(58) Li, Y.; Zamble, D. *Biochemistry* **2009**, *48*, 2486–2496.

(59) Greenwood, N. N.; Earnshaw, A. *Chemistry of the Elements*, 2nd ed.; Butterworth-Heinemann; Woburn, MA, 1997.



**Figure 4.** Optimized coordination geometries with hexa, penta and tetra coordination extracted from the large cluster optimized structures. (A) The 6-coordinated large cluster (Ni3). (B) The 5-coordinated large cluster (Ni4); (C and D) The 4-coordinated large clusters, Ni2 and Ni1, respectively. Numbers are used throughout the text for referencing relevant atoms.

states. The Ni–O bond distances are consistently longer in the optimized model complex compared to the experimental data, and no trends among the Ni–O distances are observed in either structure. The differences between the computed and experimental structures can be attributed to the omission of electronic effects of neighboring moieties on the water ligands associated with interactions not accounted for in the computational model including (1) the omission of side chains and (2) the Ni(II) sites being located on the protein surface. A more complete study of these effects is beyond the scope of the present work. On the other hand the Ni–N distances show better agreement, and the trend among the Ni–N distances observed experimentally are reproduced in the computational model. This better agreement may result, in part, from the geometrical constraints imposed by the protein. Other structural features are similar for the calculated and experimental geometries in both the 5- and 6-coordinate complexes. For example, the  $O_{ax}-Ni-O_{ax,2}$  angle




**Figure 6.** Spin-density of the small model complexes: (A) is for the 6-coordinated complex, and (B) is for the 5-coordinated complex.

deviates from  $180^\circ$ , being approximately  $172^\circ$  and  $175^\circ$ , for the computed and experimental structures.

Frontier orbitals that include metallic character for the optimized small model complexes for the 5- and 6-coordinate geometries are shown in Figure 5. For the 6-coordinate complex the frontier orbitals are from the optimized small complex where a plane of symmetry was maintained to mimic the large cluster. The computed frontier orbitals closely resemble the qualitative MO interpretation for a 6-coordinate nickel site (Figure S4, Supporting Information). Some orbital mixing is visible, particularly for the orbitals (e.g.,  $1a'$  and  $2a'$ ) corresponding to the  $t_{2g}$  set of an ideal complex with s-donor ligands only. The mixing occurs in both the  $\alpha$  and  $\beta$  sets, and involves mostly the  $xz$  orbital of Ni and a p orbital of the equatorial water oxygen. Orbitals  $1a'$  and  $2a'$  are the resulting bonding and antibonding combinations. Orbitals mix more in the  $\beta$  set but the pattern is similar to that of the  $\alpha$  set. The 6-coordinate Ni(II) complexes are stable, despite the  $d^8$  configuration, because the splitting of the  $e_g$  and  $t_{2g}$  orbitals is relatively small; the energy difference between the  $3a'$  and  $4a'$  ( $e_g$ ) orbitals in Figure 5A and the highest energy orbital of the  $t_{2g}$  set,  $2a'$ , being 1.3 eV. Inspection of the plot of the spin density for the 6-coordinated complex in Figure 6A confirms single occupation of the  $x^2 - y^2$  and  $z^2$  orbitals.

The minimized structure of the 5-coordinated large cluster shows good agreement between the computed and crystal structures around Ni(II) (Table 2). The Ni–N and Ni–O bond



**Figure 5.** Illustration of the frontier orbitals of the 6- (A) and 5-coordinated (B) small complexes of Ni. A mirror plane was used during the optimization of the 6-coordinated model, and no symmetry was used in the optimization of the 5-coordinated model.



separations follow the same trend in both structures, the biggest difference being 0.04 Å. In the QM model there are significant out-of-plane distortions as measured by the O, Ni, and the equatorial ligand angles (Table 2), with those distortions being more pronounced in the experimental structure. The positioning of the oxygen atoms' participation in the hydrogen-bonding network with the equatorial water differs significantly in the two structures.

Interpretation of the electronic structure of the 5-coordinated model (Figure 5B) closely follows the arguments used in the case of 6-coordinated complexes. Interestingly, symmetry could not be used during optimization of the small model complex. All attempts resulted in a 4-coordinated complex of pseudotetrahedral geometry and loss of one of the water ligands. The consequence of the lack of symmetry is more orbital mixing. In Figure 5B, only MOs with a clear metallic character are shown. Increasing in energy are  $z^2$  and  $x^2 - y^2$  in both the  $\alpha$  and  $\beta$  sets and  $xz$ ,  $xy$  in the  $\alpha$  set alone. All others are heavily mixed and cannot be easily assigned. Interesting is the nature of the two topmost orbitals of the  $\alpha$  set, 7a and 8a. They are derived from the  $z^2$  and  $x^2 - y^2$  orbitals of Ni and correspond to the 3a' and 4a' of the 6-coordinated model. It is relevant to note that orbital 7a of the square base pyramid complex has a lower energy than the equivalent orbital of the octahedral complex, 3a', their respective energies being  $-15.63$  and  $-14.55$  eV. The energy of the  $x^2 - y^2$  derived orbital is almost the same,  $-13.79$  and  $-13.47$  eV for the 5- and 6-coordinated complexes. This agrees with the qualitative MO picture of Figure S4 (Supporting Information). The spin density shown in Figure 6B confirms single occupation of the  $x^2 - y^2$  and  $z^2$  orbitals.

Geometry optimizations of both 4-coordinated Ni centers were performed by enforcing a singlet ground state. The optimized geometries maintained the square-planar geometry around both Ni sites (Figure 4C and D). Distortion of the optimized clusters relative to the experimental geometries is smaller than with the 5- and 6-coordinated centers (Figure S5 (Supporting Information)). The geometrical parameters shown in Table 2 are very similar for both Ni centers. Compared to the higher coordination complexes, Ni–N bonds are considerably shorter. There is also a clear trend with Ni–N2 (opposite to the Ni–S bond) which is considerably longer than Ni–N1 and Ni–N3.


**Mechanistic Implications.** HpNikR recognizes multiple DNA sequences. Half of the target DNA sequences are recognized with high affinity ( $K_d = \text{nM}$ ) and the other half with low affinity ( $K_d = \mu\text{M}$ );<sup>16</sup> and a key, unanswered question is how does HpNikR distinguish between its target promoter sequences? We hypothesize that the 5-/6-coordinate site plays a role in DNA discrimination. Molecular dynamics studies of HpNikR support a model in which the DBD are highly flexible until DNA is introduced.<sup>60</sup> We speculate that the presence of both 4-coordinate and 5-/6-coordinate sites promote conformational diversity and contribute to recognition of multiple DNA sequences. Another possible role for the 5-/6-coordinate site is to link nickel occupancy to DNA binding. In this scenario, the overall conformation of HpNikR may be altered as a function of nickel: protein stoichiometry. As a result, differential DNA recognition occurs as a function of nickel occupancy. Such a role would correlate well with *in vivo* studies that have demonstrated that the function of HpNikR is related to nickel availability.<sup>20</sup> A final scenario involves the nickel ions switching between the

4-coordinate and 5-/6-coordinate sites upon DNA-binding. The thermodynamic cost of this switching would serve to control the protein/DNA interaction. Inspection of the crystal structure identifies two histidine residues (His 75 and His 105) located between the two sites which may serve to traffic nickel between the two sites. The exact role of the 5-/6-coordinate site in DNA recognition remains to be determined; however, we propose that this site is an important player in the complex mechanism of DNA recognition by HpNikR.

**Comparison to Other NikR Structures.** A key structural difference between HpNikR, EcNikR, and PhNikR is that the nickel-binding domains are asymmetric in HpNikR and symmetric in both EcNikR and PhNikR. The structures were attained under different pH conditions (EcNikR and PhNikR at pH 7.5–8.5, HpNikR at pH 5.6). By comparing all three structures, we can identify certain residues that may explain why holo-Ni(II)HpNikR binds nickel at a 5-/6-coordinate site in the absence of DNA. Among the five residues involved in coordinating nickel in the two distinct classes of-binding sites, four belong to the central  $\beta$ -sheet sandwich of the NikR fold while one, His74, is located on a loop between a helix and a  $\beta$ -strand. We identified the conformational change that would occur if nickel were to move from a 5-/6-coordinate to a 4-coordinate geometry by aligning the MBD of HpNikR with its homologues. Pairs of MBDs forming the interface containing both nickel-binding sites were superimposed based on the  $C_\alpha$  atoms using PyMOL, resulting in overall RMSD of 1.4/1.9 Å relative to EcNikR and PhNikR, respectively. Interestingly, the corresponding RMSD values are reduced to 0.7/0.9 Å when only the  $\beta$ -sheets are included in superposition, while the RMSD increases dramatically (4.2/3.1 Å) when only the  $\alpha$ -helices are aligned. In addition, alignment of the EcNikR and PhNikR MBDs yields RMSD values of 0.5 Å for the  $\beta$ -sheets and 1.0 Å for the  $\alpha$ -helices, showing the greater similarity of these homologues versus HpNikR. This analysis suggests that the conformational changes required to accommodate all square-planar sites in HpNikR could be described as positional shifts of  $\alpha$ -helices.

Figure 7A shows the alignment of HpNikR and EcNikR based on only the  $\beta$ -sheet  $C_\alpha$  atoms. To shift the Ni(II) at the 5-/6-coordinate geometry to a 4-coordinate geometry, the second  $\alpha$ -helix of the HpNikR MBD would need to shift such that there would be steric clash at the N-terminus of the helix. Sequence alignment indicates that HpNikR contains an NQ-motif (Asn80-Gln81) in this region that is absent in the other homologues. These two residues form the basis of the extended hydrogen-bonding network (Figure 7B) which appears to be crucial in determining the conformation of the loop that contains His74. On the “4-coordinate side” (lower region of Figure 7B), the side chain of Gln81 makes a hydrogen bond to the backbone of His74', potentially making it more difficult for that residue to ligate the nickel ion. On the “5-/6-coordinate side” (upper region of Figure 8b), Asn80 makes a hydrogen bond to Tyr72 which leads to a shift in the position of Gln81', such that it no longer interacts with His74, but rather with the backbone of Glu78 and stabilizes the salt bridge between Arg77 and Asp85. We propose that these hydrogen-bonding interactions help stabilize the 5-/6-coordinate geometries relative to the 4-coordinate geometry by allowing His74 to assume the proper orientation necessary to ligate the nickel ion. In both EcNikR and NikR, the NQ motif is replaced by less polar residues, and an analogous hydrogen-bonding network is absent (Figures S6 and S7 (Supporting Information)).

(60) Sindhikara, D. J.; Roitberg, A. E.; Merz, K. M., Jr. *Biochemistry* **2009**, *48*, 12024–12033.



**Figure 7.** (A) MBD of holo-Ni(II)*HpNikR* ( $\alpha$ -helices are colored green) overlaid with MBD of holo-Ni(II)*EcNikR* ( $\alpha$ -helices are colored red).  $\beta$ -Sheets from both homologues are colored yellow. Structural alignment was based on the  $C_{\alpha}$  atoms in the  $\beta$ -sheets. (B) The extended hydrogen-bonding network at the N-terminus of neighboring  $\alpha$ -helices of the MBD of holo-Ni(II)*HpNikR*. Amino acid side chains are shown as ball and stick. Dashed lines indicate the hydrogen bonds.

Recently, Michaud-Soret and co-workers reported the structural characterization of several mutants of *HpNikR* to assess the role of the external site observed in their nickel-soaked apo-*HpNikR* structures.<sup>61</sup> This site is unoccupied in our holo-

Ni(II)*HpNikR* structure. The mutant designed to address the role of the external site is a double histidine mutant H74G/H75G. This mutant also disrupts the 5-/6-coordinate site so it is not specific for the external site. As with wt-*HpNikR*, the published structure was generated by soaking crystals of apo-H74G/H75G with excess nickel. In the nickel-bound structure of the mutant, only two square-planar sites occupied by nickel ions are present which adds further support to the idea that holo-Ni(II)*HpNikR* only has two square-planar nickel sites.

The H74G/H75G mutant exhibits decreased DNA-binding affinity compared to wild type and has no *in vivo* activity.<sup>61</sup> On the basis of these results it was concluded that the external site is required for DNA-binding because the His74G/His75G mutation has disrupted this site. However, His74 is one of the ligands in the 5-/6-coordinate site in addition to participating in the external site, and the mutation may disrupt the 5-/6-coordinate site as well as the external site. Thus, an alternate interpretation is that the decrease in DNA-binding coupled with lack of *in vivo* activity observed for the mutant may be due to disruption of the 5-/6-coordinate site, an interpretation which is consistent with our hypothesis that the 5-/6-coordinate site is essential for *HpNikR* function. Further studies are required to address these issues.

## Conclusion

Here, we report the first crystal structure of holo-Ni(II)*HpNikR*. This structure reveals that holo-Ni(II)*HpNikR* contains two distinct classes of high affinity nickel sites: two 5-/6-coordinate sites and two 4-coordinate sites. DFT calculations show that the 5-/6-coordinate sites are stabilized by key amino acid side chains from the protein. Spectroscopic analysis of nickel-binding to *HpNikR* at a range of pH values suggest that occupation of the nickel sites is pH independent. On the basis of these observations we hypothesize that the 5-/6-coordinate site plays a role in *HpNikR*'s ability to specifically and selectively bind to a series of promoter sequences.

**Acknowledgment.** We thank the Department of Pharmaceutical Sciences at the University of Maryland-Baltimore and NIH GM051501 to A.D.M. for financial support for this research.

**Supporting Information Available:** Movie demonstrating the differences in the ligand geometries for the 4- and 5-/6-coordinate sites, additional DFT and structural figures, pH-dependent Ni titrations, and complete reference 38. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JA104118R

(61) Bahlawane, C.; Dian, C.; Muller, C.; Round, A.; Fauquant, C.; Schauer, K.; de Reuse, H.; Terradot, L.; Michaud-Soret, I. *Nucleic Acids. Res.* **2010**, *38*, 3105–3118.

(62) Weiss, M. S. *J. Appl. Crystallogr.* **2001**, *34*, 130–135.

(63) Weiss, M. S.; Hilgenfeld, R. *J. Appl. Crystallogr.* **1997**, *30*, 203–205.