

Side-Chain Cross-Linked Short α -Helices That Behave like Original Proteins in Biomacromolecular Interactions

Masaaki Kajino, Kazuhisa Fujimoto,* and Masahiko Inouye*

Graduate School of Pharmaceutical Sciences, University of Toyama, Sugitani 2630, Toyama 930-0194, Japan

S Supporting Information

ABSTRACT: We explored the effect of α -helical stabilization upon the binding of short peptides to DNAs. The short peptides were designed according to the binding domains of DNA-binding proteins and were cross-linked between their side chains with diacetylenic or isophthalic cross-linking agents to keep stable α -helices. The binding abilities of the peptides to DNAs were evaluated by fluorescence resonance energy transfer analysis. When a cross-linked peptide based on the homeodomain of the transcription factor was titrated with a target DNA duplex, its dissociation constant (K_d) was calculated to be ~ 0.5 nM. This value was the double-digit smaller than that of the corresponding non-cross-linked peptide. The cross-linked peptide showed high substrate specificity for DNAs at the same level as the original DNA-binding protein.

Protein–biomacromolecule (protein and DNA/RNA) interactions play a central role in numerous biological functions in organisms. Therefore, powerful inhibitors for the interactions are expected to bring breakthroughs in the resolution of intractable diseases and to be next-generation drugs. Such inhibitors are required to cover large contact areas upon the binding sites of the target macromolecules, and it is hard for conventional inhibitors with small molecular weights to accomplish this mission. Although protein-based inhibitors promise high inhibition efficiency, there are serious problems with their use as practical inhibitors with respect to molecular size, cell permeability, chemical modification, folding, large-scale synthesis, etc.

In view of these difficulties, artificial helical foldamers or stable α -helices of short peptides are considered to be potent candidates for the above mission because of the importance of α -helices in protein–biomacromolecule interactions.¹ Several artificial helical foldamers have been developed on the basis of, for example, pyridyl pyridinone, benzourea, and β -peptide skeletons.^{2,3} However, their unnatural bodies not only need many synthetic efforts but also complicate the faithful replication of α -helical architectures. Indeed, the β -peptide-based foldamers tend to adopt 14-helices, implying that the relative positions and directions of their side chains differ from those of the α -helices seen in their original proteins. On the other hand, stable α -helices of short peptides are divided into three classes: miniature proteins,⁴ hydrogen-bonding surrogates (HBSs),⁵ and side-chain cross-linked helices.⁶ Miniature proteins consist of only natural amino acids, including avian pancreatic polypeptide or Trp-cage as a helix stabilizer.⁴ Although the stabilization method can be applied to a variety of native proteins, miniature proteins readily suffer from degradation with protease. HBS α -helices show

relatively high binding affinities for target macromolecules and resistance to protease. Recently, the inhibition of a transcription coactivator interaction by using an HBS α -helix has been reported.^{5a} However, the HBS helices need much higher binding affinity to mature into practical inhibitors. We previously reported a general method for stabilizing α -helices of short peptides by using a variety of cross-linking agents.^{6a} This methodology can copy α -helix domains out as faithfully as possible and result in high α -helicities of the “isolated” peptides. Therefore, such copies are expected to be practical inhibitors for biomacromolecular interactions. In the present work, we synthesized short helical peptides based on the binding domains of DNA-binding proteins and examined their strong affinities to the target DNAs.

We chose homeodomains of transcription factors in eukaryotic genes as a helix-domain source among various DNA-binding proteins because the homeodomain and its variants have been comprehensively studied for their binding properties.⁷ One of the variants, QKS0, has been reported to strongly bind to a specific DNA duplex, QRE, with a dissociation constant (K_d) of ~ 0.01 nM.^{7c} We designed the sequence of peptide **qk-1** according to the crystal structure of the complex of QKS0 and QRE (Figure 1a). The original sequence of the binding domain in QKS0 is peptide **qk-2**. Two ornithine (Orn) residues were placed at the *i* and *i* + 11 positions on **qk-1** for a cross-linking reaction in place of two Ile residues on **qk-2**. On the basis of the crystal structure of the complex of QKS0 and QRE, their side chains should point in the direction opposite to those of the binding residues. Therefore, the resulting cross-linkers were expected not only to reproduce the original bindings but also to have little direct influence upon the bindings. Furthermore, three Ala residues were introduced into **qk-1** instead of Trp, Phe, and Lys to simplify the sequence because their bulky side chains make little contribution to the association with QRE. For fluorescence resonance energy transfer (FRET) analysis, the amino termini of the peptides were modified with a Tokyo Green (TG) derivative,⁸ while one strand of the duplex QRE was labeled with TAMRA at its 5' end; “A” is used as the abbreviation of the labeled duplex (Figure 1b,c). The emission property of the TG fluorophore is similar to that of fluorescein, a FRET donor against TAMRA. Among a series of cross-linking agents previously reported by us, the diacetylenic molecule **1**⁶ was used to afford a cross-linked peptide **qk-1**[^]**1** because of its high ability for stabilizing the α -helices of short peptides at the *i* and *i* + 11 cross-linking positions (Figure 1d).

The helical contents of the peptide **qk-1** and the cross-linked peptide **qk-1**[^]**1** were calculated to be ~ 20 and 60%, respectively,

Received: July 30, 2010

Published: December 21, 2010




Figure 1. (a) Sequences of peptides qk-1, qk-2, and hnf-1 and of HNF-3 γ . The arrows indicate residues changed from the original sequences. (b) Sequences of DNAs A–C. (c) Chemical structures of Tokyo Green (TG) and tetramethylcarboxyrhodamine (TAMRA). (d) The diacetylenic and isophthalic cross-linking agents 1 and 2.

from the mean residue ellipticity at 222 nm (Figure S1 in the Supporting Information).^{6c,9} This relatively high value for qk-1^{^1} reveals that the TG modification scarcely disturbed the α -helical structure.^{6c} This finding suggests that a variety of functional groups can be introduced at the N-termini of cross-linked peptides without damaging the α -helical stabilization.

Cross-linked qk-1^{^1} was titrated with duplex A (QRE) at 25 °C by fluorescence spectroscopy. When duplex A was added to a solution of qk-1^{^1} under irradiation with excitation light at 500 nm, the fluorescence intensity from the TG fluorophore at 520 nm decreased (Figure 2a).¹⁰ Although the fluorescence decrease was also observed from the titration experiment involving the combination of non-cross-linked qk-1 and duplex A (Figure S2), the tendency was much more sluggish than that observed in the case of qk-1^{^1}. This finding indicates that the helical qk-1^{^1} binds to duplex A much more strongly than qk-1 does. To quantitatively compare the binding affinities of qk-1^{^1}, qk-1, and the original qk-2 for duplex A, their K_d values were estimated from their binding isotherms according to a 1:1 binding mode by nonlinear analysis (Figure 2b), and the values are presented in Table 1. The K_d values for qk-1^{^1}, qk-1, and qk-2 were <1.0, 30 \pm 5, and 23 \pm 4 nM, respectively. Unfortunately, it was difficult to estimate exactly the binding affinity of qk-1^{^1} for duplex A at a NaCl concentration of 150 mM because of the low limit of detection. Thus, additional titration experiments were performed at NaCl concentrations of 300 and 500 mM, where the binding affinity was reduced as a result of electrostatic interactions. In the presence of 300 (500) mM NaCl, the K_d values of qk-1^{^1} and qk-1 changed to 3 \pm 1 (87 \pm 5) and 165 \pm 13 (>2500) nM, respectively, and $\Delta G_{\text{qk-1}^1} - \Delta G_{\text{qk-1}} \approx 4$ kJ/mol at both NaCl concentrations (Figure S3). Since the difference in ΔG values could be considered independent of the salt concentration, the K_d value of qk-1^{^1} in the presence of 150 mM




Figure 2. (a) Fluorescence spectra of qk-1^{^1} (10 nM) titrated with duplex A (QRE) (0–160 nM) in a 10 mM Tris-HCl buffer containing 150 mM NaCl and 6 mM MgCl₂. (b) Isotherm curves for the fluorescence titrations of qk-1^{^1} (red) and qk-1 (black) in the presence of 150 mM NaCl. The observation wavelengths were 520 and 518 nm for qk-1^{^1} and qk-1, respectively.

Table 1. Dissociation Constants (in units of nM) for Peptides with the DNAs QRE and ERE at various NaCl concentrations (in mM) Obtained from the FRET Analysis

	QRE			ERE
	150	300	500	150
qk-1 ^{^1}	~0.5	3 \pm 1	87 \pm 5	11 \pm 3
qk-1	30 \pm 5	165 \pm 13	>2500	34 \pm 4
qk-2	23 \pm 4	–	–	–

NaCl could be recalculated as ~0.5 nM. This value is much smaller than the values of 7.5 and 17 nM in the systems of Woolley's side-chain cross-linked α -helix^{7a} and Schepartz's miniature protein,^{7b} respectively.

The K_d values were also determined by surface plasmon resonance (SPR) analysis (Figure S4).¹¹ The K_d values of nonlabeled qk-1^{^1}, qk-1, and qk-2 for nonlabeled duplex A were calculated to be ~70, 6000, and 1000 nM, respectively, from their sensorgrams. Although the order of the binding affinities, qk-1^{^1} > qk-2 > qk-1, was the same as that observed in the FRET analysis, the K_d values were ~100 times as large as those in the FRET analysis at [NaCl] = 150 mM. The high values might have large errors because the molecular weights of 2100–2400 made it hard to obtain strong signals and good signal-to-noise ratios in the SPR analysis. On the other hand, the K_d values were close to those (87 and >2500 nM for qk-1^{^1} and

qk-1, respectively) at the NaCl concentration of 500 mM in the FRET analysis. As shown in Figure S3b, the isotherm curves accompanying small error bars afforded the relatively accurate K_d values at the NaCl concentration of 500 mM. Thus, the K_d value of ~ 0.5 nM for the cross-linked qk-1^{^1} in the FRET analysis at [NaCl] = 150 mM expresses its real binding affinity.

To support the effect of the cross-linkage on the binding affinity, the melting temperatures (T_m) of nonlabeled duplex A (QRE) (1 μ M) were measured in the absence and presence of nonlabeled qk-1^{^1} and qk-1. The T_m value was 69.0 °C in the absence of the peptides. The value increased to 70.8 °C in the presence of 3 equiv of qk-1^{^1} but scarcely changed upon addition of qk-1. This T_m increase also certifies the effect of the α -helical stabilization upon the interaction between the cross-linked peptide and the duplex.

To examine the specificity of the cross-linked qk-1^{^1}, fluorescence titration experiments were carried out with another DNA duplex having a different sequence, B (ERE), which is known to bind QK50 more weakly than QRE (Figure 1b).^{7c} The K_d of qk-1^{^1} for duplex B was ~ 11 nM (Figure S5), and the $K_{d(A:QRE)}/K_{d(B:ERE)}$ ratio was 22, which is close to that for the original protein QK50 (34).^{7c} On the other hand, the non-cross-linked qk-1 exhibited no DNA specificity; the K_d for duplex B was ~ 34 nM.

To show the generality of our cross-linking method, the DNA-binding protein HNF-3 γ was selected as another example.¹² As shown in Figure 1a, the sequence of a peptide hnf-1 was arranged on the basis of the original sequence of HNF-3 γ . Four Ala residues were introduced into hnf-1 for the same reason as for qk-1 mentioned above. Two Orn residues were positioned at the i and $i + 7$ on hnf-1 for the cross-linking reaction in place of the Gln and Leu residues in the original sequence. The isophthalic cross-linker 2 can effectively stabilize α -helices of short peptides at the i and $i + 7$ cross-linking positions.^{6a} Furthermore, the sequence of hnf-1 includes an extra Arg-repeated moiety at the C-terminus. The original HNF-3 γ was reported to bind to a target DNA duplex through the α -helical and the Arg-repeated moieties; the duplex labeled with TAMRA is abbreviated as "C" (Figure 1b). Therefore, the Arg-repeated sequence was added to the C-terminus of hnf-1. The helical contents of the peptide hnf-1 and the cross-linked hnf-1^{^2} were calculated to be ~ 20 and 70%, respectively (Figure S6).^{6c,9} In the titration experiments involving the peptides and duplex C, a binding enhancement by the α -helical stabilization similar to that seen in the above case was observed. The K_d value changed from 570 nM for the non-cross-linked hnf-1 to 4 nM for the cross-linked hnf-1^{^2} (Figure S7). Thus, our strategy proved to be applicable for emulating various DNA-binding proteins.

Finally, we explored the resistance property of the cross-linked peptides against digestive enzymes. The cross-linked qk-1^{^1} and the non-cross-linked qk-1 (50 μ M) were digested with trypsin (0.5 μ M), and the digested peptides were analyzed by HPLC (Figure S8). The half-life for digestion ($\tau_{1/2}$) of qk-1^{^1} was 65 min, while the $\tau_{1/2}$ value for qk-1 was 15 min. Thus, the cross-linked peptides could somewhat obtain the resistance property against digestive enzymes because of the α -helical stabilization and/or the presence of the cross-linking moiety. This finding shows the possibility for future applications of the cross-linked peptides such as to in vivo use.

In conclusion, cross-linked peptides with stable α -helices were found to work well, like the original proteins. Their binding affinities for DNAs were strong (sub-nanomolar level), and their substrate specificities were comparable to those of the original

proteins. Furthermore, in the presence of digestive enzymes, the cross-linked peptides survived much longer than native peptides. These results have encouraged us to extend our strategy to protein–protein interactions related to intractable diseases.

■ ASSOCIATED CONTENT

Supporting Information. Figures S1–S8 and all experimental procedures. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

fujimoto@pha.u-toyama.ac.jp; inouye@pha.u-toyama.ac.jp

■ ACKNOWLEDGMENT

This work was partially supported by Grants-in-Aid for Scientific Research in Priority Areas "New Frontiers in Photochromism (471) and Young Scientists (B) (21750167) from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT) of Japan. We thank Professor Ikuo Fujii (Osaka Prefecture University) and Professor Mineyuki Mizuguchi (University of Toyama) for measurements of SPR sensorgrams.

■ REFERENCES

- (1) (a) Arora, P. S.; Ansari, A. Z. *Nature* **2009**, *462*, 171–173. (b) Guarracino, D. A.; Arora, P. S. *Chem. Biol.* **2009**, *16*, 919–920. (c) Henchey, L. K.; Jochim, A. L.; Arora, P. S. *Curr. Opin. Chem. Biol.* **2008**, *12*, 692–697.
- (2) (a) Cummings, C. G.; Hamilton, A. D. *Curr. Opin. Chem. Biol.* **2010**, *14*, 341–346. (b) Davis, J. M.; Tsou, L. K.; Hamilton, A. D. *Chem. Soc. Rev.* **2007**, *36*, 326–334.
- (3) For recent papers on β -peptides, see: (a) Bautista, A. D.; Appelbaum, J. S.; Craig, C. J.; Michel, J.; Schepartz, A. *J. Am. Chem. Soc.* **2010**, *132*, 2904–2906. (b) Harker, E. A.; Schepartz, A. *ChemBioChem* **2009**, *10*, 990–993. (c) Michel, J.; Harker, E. A.; Tirado-Rives, J.; Jorgensen, W. L.; Schepartz, A. *J. Am. Chem. Soc.* **2009**, *131*, 6356–6357.
- (4) For recent papers, see: (a) Smith, B. A.; Daniels, D. S.; Coplin, A. E.; Jordan, G. E.; McGregor, L. M.; Schepartz, A. *J. Am. Chem. Soc.* **2008**, *130*, 2948–2949. (b) Daniels, D. S.; Schepartz, A. *J. Am. Chem. Soc.* **2007**, *129*, 14578–14579. (c) Harman, R. E.; Badders, D.; Fuller, M.; Makienko, E. G.; Houston, M. E.; Quay, S. C.; Johnson, P. H. *J. Biol. Chem.* **2007**, *282*, 9813–9824.
- (5) For recent papers, see: (a) Henchey, L. K.; Kushal, S.; Dubey, R.; Chapman, R. N.; Olenyuk, B. Z.; Arora, P. S. *J. Am. Chem. Soc.* **2010**, *132*, 941–943. (b) Bao, J.; Dong, X. Y.; Zhang, J. Z. H.; Arora, P. S. *J. Phys. Chem. B* **2009**, *113*, 3565–3571. (c) Wang, D.; Lu, M.; Arora, P. S. *Angew. Chem., Int. Ed.* **2008**, *47*, 1879–1882.
- (6) For recent papers, see: (a) Fujimoto, K.; Kajino, M.; Inouye, M. *Chem.—Eur. J.* **2008**, *14*, 857–863. (b) Zhang, F.; Sadvoski, O.; Xin, S. J.; Woolley, G. A. *J. Am. Chem. Soc.* **2007**, *129*, 14154–14155. (c) Bernal, F.; Tyler, A. F.; Korsmeyer, S. J.; Walensky, L. D.; Verdine, G. L. *J. Am. Chem. Soc.* **2007**, *129*, 2456–2457.
- (7) For example, see: (a) Guerrero, L.; Smart, O. S.; Woolley, G. A.; Allemann, R. K. *J. Am. Chem. Soc.* **2005**, *127*, 15624–15629. (b) Montclare, J. K.; Schepartz, A. *J. Am. Chem. Soc.* **2003**, *125*, 3416–3417. (c) Ades, S. E.; Sauer, R. T. *Biochemistry* **1995**, *34*, 14601–14608.
- (8) Mineno, T.; Ueno, T.; Urano, Y.; Kojima, H.; Nagano, T. *Org. Lett.* **2006**, *8*, 5963–5966.
- (9) Fujimoto, K.; Oimoto, N.; Katsuno, K.; Inouye, M. *Chem. Commun.* **2004**, 1280–1281.
- (10) The fluorescence intensity was hardly changed in a similar titration experiment using duplex A (QRE) without the TAMRA

labeling. The decrease in the fluorescence intensity resulting from the addition of the labeled A was due not to quenching by nucleobase residues such as guanine but to FRET from TG to TAMRA.

(11) Karlsson, R. *Anal. Biochem.* **1994**, *221*, 142–151.

(12) Clark, K. L.; Halay, E. D.; Lai, E.; Burley, S. K. *Nature* **1993**, *364*, 412–420.