

Integrated Electrochemical–Chemical Oxidation Mediated by Alkoxyulfonium Ions

Yosuke Ashikari, Toshiki Nokami, and Jun-ichi Yoshida*

Department of Synthetic Chemistry and Biological Chemistry, Graduate School of Engineering, Kyoto University, Nishikyo-ku, Kyoto 615-8510, Japan

S Supporting Information

ABSTRACT: Generation of carbocations by the “cation pool” method followed by reaction with dimethyl sulfoxide (DMSO) gave the corresponding alkoxyulfonium ions. Alkoxyulfonium ions could also be generated by in situ DMSO trapping of electrochemically generated carbocations. The resulting alkoxyulfonium ions were transformed into carbonyl compounds by treatment with triethylamine. The present integrated electrochemical–chemical oxidation can be applied to the oxidation of diarylmethanes to diaryl ketones, toluenes to benzaldehydes, and aryl-substituted alkenes to 1,2-diketones. Moreover, the oxidation of unsaturated compounds bearing a nucleophilic group in an appropriate position gives cyclized carbonyl compounds.


Organic synthesis has been developed mainly on the basis of step-by-step synthesis. However, combining multiple steps without isolating intermediates is strongly needed to enhance the power and efficiency of organic synthesis.^{1,2} To this end, chemists have been interested in the integration of chemical reactions to develop synthetic transformations that would be otherwise impossible, such as the generation of unstable highly reactive species that are swiftly utilized for a further reaction before they decompose.³

In addition to chemical reactions using chemical reagents and/or catalysts, electrochemical reactions^{4,5} using electron transfer on the surface of the electrode serve as a powerful means of generating highly reactive species. Electrochemistry allows for the selective introduction and removal of electrons under mild conditions, whereas this process usually requires harsh conditions when the chemical method is employed.

Among a variety of reactions used in organic synthesis, selective oxidation of organic compounds still remains as a major challenge.⁶ We envisaged that integration of electrochemical oxidation and chemical oxidation would be a nice approach to this challenge, and we report here an example of such integration that is mediated by an unstable reactive intermediate.

We chose to use alkoxyulfonium ions as key intermediates because treatment of an alkoxyulfonium ion with a base such as triethylamine eventually gives the corresponding carbonyl compound; this serves as one of the mildest methods for oxidation of organic compounds [dimethyl sulfoxide (DMSO) oxidation] (Scheme 1).^{7,8} Although alkoxyulfonium ions can be generated by several chemical methods, the oxidation state of the precursors ($RR'CHX$; $X = \text{heteroatom}$) and the resulting alkoxyulfonium ions ($RR'CH-OS^+Me_2$) are the same, and the overall

Scheme 1. Chemical Oxidation Mediated by Alkoxyulfonium Ions




reaction to give $RR'C=O$ is a two-electron oxidation. If substrates of lower oxidation state, for example $RR'CH_2$, could be used to generate alkoxyulfonium ions, the overall reaction would be a four-electron oxidation, and the method would serve as a highly useful way of oxidizing organic compounds.

It is well-known that electrochemical oxidation serves as a powerful method for generating carbocations and onium ions under mild conditions.⁴ In fact, various types of electrochemical oxidation reactions proceed by a mechanism involving carbocations, though carbocations are usually trapped spontaneously by nucleophiles that are present in the reaction medium. However, we have developed an effective method for generating and accumulating carbocations or onium ions by low-temperature electrochemical oxidation. The method, which is called the “cation pool” method, has been successfully applied to *N*-acyliminium ions,⁹ alkoxy-carbenium ions,¹⁰ and diaryl-carbenium ions.¹¹ On the basis of these backgrounds, we envisaged that carbocations generated by the cation pool method could be used for a subsequent reaction with DMSO to generate alkoxyulfonium ions, which would then react with an amine base to give the corresponding carbonyl compounds (Scheme 2). In fact, Mayr and co-workers observed in their mechanistic studies that diaryl-carbenium ions generated by chemical S_N1 reaction reacted with DMSO and that subsequent treatment with triethylamine gave the diaryl ketones.¹² It should be noted that unlike the chemical methods, the oxidation state of the precursor in the electrochemical method is lower than that of the resulting alkoxyulfonium ion. Therefore, we initiated our studies of the integration of the electrochemical method and the chemical method mediated by alkoxyulfonium ions.

To test the feasibility of the present concept, we first examined the reaction of an electrochemically generated diaryl-carbenium ion with DMSO. Thus, diaryl-carbenium ion **2** was generated and

Received: March 30, 2011

Published: July 11, 2011

Scheme 2. Integrated Electrochemical–Chemical Oxidation Mediated by Alkoxysulfonium Ions


accumulated by the electrochemical oxidation of diarylmethane **1** in CD_2Cl_2 using Bu_4NBF_4 as a supporting electrolyte at -78°C according to the “cation pool” method (Figure 1, method A).¹¹ The addition of DMSO to **2** gave alkoxysulfonium ion **3**, which was well-characterized by NMR analysis at -78°C . Treatment of the solution of **3** with triethylamine gave the corresponding diaryl ketone **4** in 91% yield. This proof-of-principle experiment revealed that the electrochemical reaction and the chemical reaction could be integrated effectively through the intermediacy of a carbocation and an alkoxysulfonium ion. The method was also applicable to diarylmethane **5** to produce the corresponding diaryl ketone **6** (Table 1).

In-situ trapping of carbocations by DMSO is also effective when the substrate oxidation potential is lower than that of DMSO (1.76 V vs SCE). For example, the electrochemical oxidation of 4-methoxytoluene (**7**) (1.38 V vs SCE) in 1:2 $\text{DMSO}-d_6/\text{CD}_2\text{Cl}_2$ (Figure 1, method B) gave alkoxysulfonium ion **8**, which was characterized by NMR analysis. In this case, the electrolysis could be carried out at 24°C because the alkoxysulfonium ion is more stable than the corresponding carbocation. Treatment of **8** with triethylamine gave aldehyde **9** in 86% yield. It is noteworthy that the cation pool method cannot be used in this case because 4-methoxybenzyl cation is not stable enough to accumulate in the solution even at -78°C . It is also noteworthy that the success of the in situ method is attributed to the fact that **7** is more easily oxidized than DMSO. In contrast, the oxidation potential of diarylmethane **1** (1.96 V vs SCE) is higher than that of DMSO, and therefore, the electrochemical oxidation of **1** should be carried out in the absence of DMSO. However, a wide range of organic compounds can serve as substrates for the in situ method. For example, xanthene (**10**) was oxidized effectively by method B to give 9-xanthenone (**11**) (Table 1).

The present method could be extended to the oxidation of aryl-substituted alkenes. For example, 4,4'-dichloro-*trans*-stilbene (**12**) was electrochemically oxidized in the presence of $\text{DMSO}-d_6$ at 0°C (Figure 2). The NMR study indicated the formation of dialkoxysulfonium ion **13** in the solution. Treatment of **13** with triethylamine gave the corresponding 1,2-diketone **14** in 83% yield. In contrast, chemical oxidation of *trans*-stilbene using $\text{RuCl}_3/\text{NaIO}_4$ gives the corresponding 1,2-diketone (benzil) in only 5% yield and the carbon–carbon bond cleavage product (benzoic acid) as the major product.¹³ In addition, the electrochemical method also often suffers from overoxidation. For example, anodic oxidation of alkenes¹⁴ and of 1,2-diols¹⁵ and their derivatives often leads to carbon–carbon bond cleavage. The present method solves this otherwise formidable problem by integration with an extremely mild chemical method such as DMSO oxidation. Therefore, the present method serves as a powerful and highly selective method for the conversion of alkenes to 1,2-dicarbonyl compounds (Table 1).

To demonstrate the utility and versatility of the present method, oxidation reactions of alkenes bearing a nucleophilic group in an



Method A: trapping of a “cation pool” with DMSO

Method B: trapping of a carbocation with DMSO in-situ


Figure 1. Typical examples of the integrated electrochemical–chemical oxidation and ^1H NMR spectra of the cationic intermediates **2**, **3**, and **8**.

appropriate position that can participate in the electrochemical oxidation were examined.^{16,17} The electrochemical oxidation of **23** took place in the presence of $\text{DMSO}-d_6$ at 0°C (Figure 3). The formation of cyclized alkoxysulfonium ion **24** was confirmed by NMR analysis. Treatment of **24** with triethylamine gave the corresponding cyclized carbonyl compound **25**.

The alkene substrate scope was also investigated (Table 2). Alkenes bearing a nucleophilic moiety such as a tosylamide (**23** and **26**), a hydroxyl group (**28**), or an aromatic ring (**30**) gave the corresponding cyclized ketones in moderate to high yields. Moreover, the electrolysis of 1,6-dienes **32**, **34**, **36**, and **38** gave the corresponding cyclized products **33**, **35**, **37**, and **39**, respectively.

In order to gain deeper insight into the mechanism, we measured the oxidation potentials of the substrates (Table 2) and compared them with those of olefinic compounds without nucleophilic functional groups. The oxidation potentials of **23**, **26**, **28**, and **30** were less positive than that of β -methylstyrene (1.42 V vs SCE), indicating that the radical cation generated by one-electron oxidation undergoes rapid cyclization. Also, the oxidation potential of **32** (1.16 V vs SCE) was less positive than that of **40** (1.54 V vs SCE), suggesting that the radical cation rapidly reacts with the other carbon–carbon double bond,¹⁸ as shown in Figure 4.

Table 1. Integrated Electrochemical–Chemical Oxidation of Organic Compounds^a

substrate	oxidation potential (V) ^b	method	electricity (F/mol)	product	yield (%) ^c
	1.96	A	2.5		91
	1.75	A	2.5		62
	1.38	B	2.5		86 ^d
	1.36	B	4.0		84
	1.36	B	2.1		83
	1.31	B	2.1		70 (71) ^e
	1.44	B	2.1		50 (74) ^f
	1.20	B	2.1		71
	1.09	B	2.5		55

^a The reactions were carried out on a 0.25 mmol scale. ^b Determined by rotating disk electrode (RDE) voltammetry in 0.1 M LiClO₄/CH₃CN using SCE as a reference electrode. ^c Isolated yields are given, unless otherwise stated. ^d GC yield. ^e The reaction was carried out on a 1.3 mmol scale. ^f NMR yield.




Figure 2. Integrated electrochemical–chemical oxidation of 4,4'-dichloro-*trans*-stilbene and the ¹H NMR spectrum of the cationic intermediate 13.

The electrochemical method is often cited as being environmentally favorable because it enables straightforward




Figure 3. Integrated electrochemical–chemical oxidation of alkene 23 and the ¹H NMR spectrum of the cyclized cationic intermediate 24.




Figure 4. Comparison of oxidation potentials and a proposed mechanism involving the rapid cation radical cyclization.

Table 2. Integrated Electrochemical–Chemical Oxidation Involving Cyclization^a

substrate	oxidation potential (V) ^b	electricity (F/mol)	product	yield (%) ^c
	1.34	2.1		86 ^d
	1.31	2.5		85
	1.20	3.0		29 52 ^d
	1.21	2.1		31 73
	1.16	2.2		33 72 ^e
	1.23	2.1		35 90 ^e
	1.17	2.1		37 78
	1.24	2.1		39 76

^a The reactions were carried out on a 0.25 or 0.13 mmol scale. ^b Determined by RDE voltammetry in 0.1 M LiClO₄/CH₃CN using SCE as a reference electrode. ^c Isolated yields. ^d 1:2 DMSO/CH₂Cl₂ was used. ^e Bu₄NB(C₆F₅)₄ was used as a supporting electrolyte.

transformations under mild conditions without the use of hazardous strong chemical reagents.¹⁹ The present method based on integration of the electrochemical method with an extremely mild, environmentally benign chemical method opens a new aspect of the chemistry of oxidation reactions. Further

work to explore the full range of the scope of the present approach and its application to the synthesis of molecules with interesting biological activities and functions is in progress.

■ ASSOCIATED CONTENT

S Supporting Information. Experimental procedures, spectroscopic data for new compounds, and crystallographic data (CIF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

yoshida@sbchem.kyoto-u.ac.jp

■ ACKNOWLEDGMENT

This work was supported by a Grant-in-Aid for Scientific Research and a Grant-in-Aid for the Global COE program from MEXT, Japan.

■ REFERENCES

- (1) For reviews, see: (a) Tietze, L. F. *Chem. Rev.* **1996**, *96*, 115. (b) Ryu, I.; Sonoda, N.; Curran, D. P. *Chem. Rev.* **1996**, *96*, 177. (c) Parsons, P. J.; Penkett, C. S.; Shell, A. J. *Chem. Rev.* **1996**, *96*, 195. (d) Louie, J.; Bielawski, C. W.; Grubbs, R. H. *J. Am. Chem. Soc.* **2001**, *123*, 11312. (e) Ugi, I. *Pure Appl. Chem.* **2001**, *73*, 187. (f) Fogg, D. E.; dos Santos, E. N. *Coord. Chem. Rev.* **2004**, *248*, 2365. (g) de Meijere, A.; von Zezschwitz, P.; Bräse, S. *Acc. Chem. Res.* **2005**, *38*, 413. (h) Wasilke, J.-C.; Obrey, S. J.; Baker, R. T.; Bazan, G. C. *Chem. Rev.* **2005**, *105*, 1001. (i) Schmidt, B. *Pure Appl. Chem.* **2006**, *78*, 469. (j) Enders, D.; Hüttl, M. R. M.; Grondal, C.; Raabe, G. *Nature* **2006**, *441*, 861.
- (2) For recent examples, see: (a) Orita, A.; Yaruva, J.; Otera, J. *Angew. Chem., Int. Ed.* **1999**, *38*, 2267. (b) Orita, A.; Yoshioka, N.; Struwe, P.; Braier, A.; Beckmann, A.; Otera, J. *Chem.—Eur. J.* **1999**, *5*, 1355. (c) Dömling, A.; Ugi, I. *Angew. Chem., Int. Ed.* **2000**, *39*, 3168. (d) Bienaymé, H.; Hulme, C.; Odon, O.; Schmitt, P. *Chem.—Eur. J.* **2000**, *6*, 3321. (e) Balme, G.; Bossharth, E.; Monteiro, N. *Eur. J. Org. Chem.* **2003**, 4101. (f) Ajamian, A.; Gleason, J. L. *Angew. Chem., Int. Ed.* **2004**, *43*, 3754. (g) Shimizu, M.; Shimon, K.; Hiyama, T. *Chem. Lett.* **2006**, *35*, 838. (h) Clarke, P. A.; Santos, S.; Martin, W. H. C. *Green Chem.* **2007**, *9*, 438. (i) Bremner, W. S.; Organ, M. G. *J. Comb. Chem.* **2007**, *9*, 14. (j) Fuwa, H.; Tako, T.; Ebine, M.; Sasaki, M. *Chem. Lett.* **2008**, *37*, 904. (k) Nokami, T.; Tsuyama, H.; Shibuya, A.; Nakatsutsumi, T.; Yoshida, J. *Chem. Lett.* **2008**, *37*, 942. (l) Ikeda, S.; Shibuya, M.; Kanoh, N.; Iwabuchi, Y. *Chem. Lett.* **2008**, *37*, 962. (m) Numata, Y.; Kawashima, J.; Hara, T.; Tajima, Y. *Chem. Lett.* **2008**, *37*, 1018. (n) Yamaguchi, K.; Kotani, M.; Kamata, K.; Mizuno, N. *Chem. Lett.* **2008**, *37*, 1258. (o) Babu, G.; Orita, A.; Otera, J. *Chem. Lett.* **2008**, *37*, 1296. (p) Cernak, T. A.; Lambert, T. H. *J. Am. Chem. Soc.* **2009**, *131*, 3124. (q) Hardee, D. J.; Lambert, T. H. *J. Am. Chem. Soc.* **2009**, *131*, 7536.
- (3) (a) Suga, S.; Yamada, D.; Yoshida, J. *Chem. Lett.* **2010**, *39*, 404. (b) Yoshida, J.; Saito, K.; Nokami, T.; Nagaki, A. *Synlett* **2011**, 1189.
- (4) For recent reviews of organic electrochemistry, see: (a) Moeller, K. D. *Tetrahedron* **2000**, *56*, 9527. (b) Lund, H. J. *Electrochem. Soc.* **2002**, *149*, S21. (c) Sperry, J. B.; Wright, D. L. *Chem. Soc. Rev.* **2006**, *35*, 605. (d) Yoshida, J.; Kataoka, K.; Horcajada, R.; Nagaki, A. *Chem. Rev.* **2008**, *108*, 2265.
- (5) For some recent examples, see: (a) Hasegawa, M.; Ishii, H.; Cao, Y.; Fuchigami, T. *J. Electrochem. Soc.* **2006**, *153*, D162. (b) Hayashi, K.; Kim, S.; Chiba, K. *Electrochemistry* **2006**, *74*, 621. (c) Wu, X.; Dube, M. A.; Fry, A. J. *Tetrahedron Lett.* **2006**, *47*, 7667. (d) Tajima, T.; Kurihara, H.; Fuchigami, T. *J. Am. Chem. Soc.* **2007**, *129*, 6680. (e) Mitsudo, K.; Kaide, T.; Nakamoto, E.; Yoshida, K.; Tanaka, H. *J. Am. Chem. Soc.* **2007**, *129*, 2246. (f) Horii, D.; Fuchigami, T.; Atobe, M. *J. Am. Chem. Soc.* **2007**, *129*, 11692. (g) Park, Y. S.; Little, R. D. *Electrochim. Acta* **2009**, *54*, 5077. (h) Kakiuchi, F.; Kochi, T.; Mutsutani, H.; Kobayashi, N.; Urano, S.; Sato, M.; Nishiyama, S.; Tanabe, T. *J. Am. Chem. Soc.* **2009**, *131*, 11310. (i) Kirste, A.; Schnakenburg, G.; Stecker, F.; Fischer, A.; Waldvogel, S. R. *Angew. Chem., Int. Ed.* **2010**, *49*, 971.
- (6) For example, see: Lenoir, D. *Angew. Chem., Int. Ed.* **2006**, *45*, 3206.
- (7) For Swern–Moffatt-type oxidation, see: Mancuso, A. J.; Swern, D. *Synthesis* **1981**, 165.
- (8) For Kornblum oxidation, see: Kornblum, N.; Powers, J. W.; Anderson, G. J.; Jones, W. J.; Larson, H. O.; Levand, O.; Weaver, W. M. *J. Am. Chem. Soc.* **1957**, *79*, 6562.
- (9) For N-acyliminium ion pools, see: (a) Yoshida, J.; Suga, S.; Suzuki, S.; Kinomura, N.; Yamamoto, A.; Fujiwara, K. *J. Am. Chem. Soc.* **1999**, *121*, 9546. (b) Suga, S.; Okajima, M.; Fujiwara, K.; Yoshida, J. *J. Am. Chem. Soc.* **2001**, *123*, 7941. (c) Suga, S.; Watanabe, M.; Yoshida, J. *J. Am. Chem. Soc.* **2002**, *124*, 14824. (d) Yoshida, J.; Suga, S. *Chem.—Eur. J.* **2002**, *8*, 2651. (e) Maruyama, T.; Mizuno, Y.; Shimizu, I.; Suga, S.; Yoshida, J. *J. Am. Chem. Soc.* **2007**, *129*, 1902.
- (10) For alkoxy-carbenium ion pools, see: (a) Suga, S.; Suzuki, S.; Yamamoto, A.; Yoshida, J. *J. Am. Chem. Soc.* **2000**, *122*, 10244. (b) Okajima, M.; Suga, S.; Itami, K.; Yoshida, J. *J. Am. Chem. Soc.* **2005**, *127*, 6930. (c) Suga, S.; Matsumoto, K.; Ueoka, K.; Yoshida, J. *J. Am. Chem. Soc.* **2006**, *128*, 7710.
- (11) For diarylcarbenium ion pools, see: (a) Okajima, M.; Soga, K.; Nokami, T.; Suga, S.; Yoshida, J. *Org. Lett.* **2006**, *8*, 5005. (b) Nokami, T.; Ohata, K.; Inoue, M.; Tsuyama, H.; Shibuya, A.; Soga, K.; Okajima, M.; Suga, S.; Yoshida, J. *J. Am. Chem. Soc.* **2008**, *130*, 10864. (c) Okajima, M.; Soga, K.; Watanabe, T.; Terao, K.; Nokami, T.; Suga, S.; Yoshida, J. *Bull. Chem. Soc. Jpn.* **2009**, *82*, S94. (d) Terao, K.; Watanabe, T.; Suehiro, T.; Nokami, T.; Yoshida, J. *Tetrahedron Lett.* **2010**, *51*, 4107. (e) Nokami, T.; Watanabe, T.; Musya, N.; Morofuji, T.; Tahara, K.; Tobe, Y.; Yoshida, J. *Chem. Commun.* **2011**, *47*, 5575. (f) Nokami, T.; Watanabe, T.; Musya, N.; Suehiro, T.; Morofuji, T.; Yoshida, J. *Tetrahedron* **2011**, *67*, 4464.
- (12) (a) Phan, T. B.; Nolte, C.; Kobayashi, S.; Ofial, A. R.; Mayr, H. *J. Am. Chem. Soc.* **2009**, *131*, 11392. A reaction of benzyl cations with DMSO has also been reported in the literature. See: (b) Scheit, K. H.; Kampe, W. *Angew. Chem., Int. Ed. Engl.* **1965**, *4*, 787.
- (13) Tabatabaeian, K.; Mamaghani, M.; Mahmoodi, N. O.; Khorshidi, A. *Catal. Commun.* **2008**, *9*, 416. For other methods of oxidizing alkenes to 1,2-diketones, see: Manandhar, S.; Singh, R. P.; Eggers, G. V.; Shreeve, J. M. *J. Org. Chem.* **2002**, *67*, 6415.
- (14) (a) Halas, S. M.; Okyne, K.; Fry, A. J. *Electrochim. Acta* **2003**, *48*, 1837. (b) Bäumer, U.-St.; Schäfer, H. J. *J. Appl. Electrochem.* **2005**, *35*, 1283.
- (15) Shono, T.; Matsumura, Y.; Hashimoto, T.; Hibino, K.; Hamaguchi, H.; Aoki, T. *J. Am. Chem. Soc.* **1975**, *97*, 2546.
- (16) Electrochemical dimethoxylation of styrenes and related compounds has been reported. See: (a) Raoult, E.; Sarrazin, J.; Tallec, A. *J. Appl. Electrochem.* **1984**, *14*, 639. (b) Ogibin, Y. N.; Ilovaiskii, A. I.; Nikishin, G. I. *Russ. Chem. Bull.* **1994**, *43*, 1536.
- (17) For electron-transfer-mediated cyclization reactions, see: (a) Tokuda, M.; Miyamoto, T.; Fujita, H.; Sugimoto, H. *Tetrahedron* **1991**, *47*, 747. (b) Tokuda, M.; Fujita, H.; Miyamoto, T.; Sugimoto, H. *Tetrahedron* **1993**, *49*, 2413. (c) New, D. G.; Tesfai, Z.; Moeller, K. D. *J. Org. Chem.* **1996**, *61*, 1578. (d) Xu, H.-C.; Moeller, K. D. *J. Am. Chem. Soc.* **2008**, *130*, 13542. (e) Liu, L.; Floreancig, P. E. *Angew. Chem., Int. Ed.* **2010**, *49*, 3069. (f) Xu, H.-C.; Moeller, K. D. *Angew. Chem., Int. Ed.* **2010**, *49*, 8004.
- (18) (a) Moeller, K. D.; Tino, L. V. *J. Am. Chem. Soc.* **1992**, *114*, 1033. (b) Reddy, S. H. K.; Chiba, K.; Sun, Y.; Moeller, K. D. *Tetrahedron* **2001**, *57*, 5183. (c) Huang, K. D.; Moeller, K. D. *Tetrahedron* **2006**, *62*, 6536.
- (19) (a) Matthews, M. A. *Pure Appl. Chem.* **2001**, *73*, 1305. (b) Rusling, J. F. *Pure Appl. Chem.* **2001**, *73*, 1895. (c) Mentel, M.; Beier, M. J.; Breinbauer, R. *Synthesis* **2009**, 1463 and references cited therein.