

# Comparison of Two Yeast MnSODs: Mitochondrial *Saccharomyces cerevisiae* versus Cytosolic *Candida albicans*

Yuewei Sheng,<sup>†</sup> Troy A. Stich,<sup>§</sup> Kevin Barnese,<sup>†</sup> Edith B. Gralla,<sup>†</sup> Duilio Cascio,<sup>‡</sup> R. David Britt,<sup>§</sup> Diane E. Cabelli,<sup>\*||</sup> and Joan Selverstone Valentine<sup>\*†,⊥</sup>

<sup>†</sup>Department of Chemistry and Biochemistry, and <sup>‡</sup>Department of Energy-Institute for Genomics and Proteomics, University of California Los Angeles, 420 Westwood Plaza, Los Angeles, California 90095, United States

<sup>§</sup>Department of Chemistry, University of California Davis, One Shields Avenue, Davis, California 95616, United States


<sup>||</sup>Chemistry Department, Brookhaven National Laboratory, Upton, New York 11973, United States

<sup>⊥</sup>Department of Bioinspired Sciences, Ewha Womans University, Seoul 120-750, Republic of Korea

**S** Supporting Information

**ABSTRACT:** Human MnSOD is significantly more product-inhibited than bacterial MnSODs at high concentrations of superoxide ( $O_2^{\cdot-}$ ). This behavior limits the amount of  $H_2O_2$  produced at high  $[O_2^{\cdot-}]$ ; its desirability can be explained by the multiple roles of  $H_2O_2$  in mammalian cells, particularly its role in signaling. To investigate the mechanism of product inhibition in MnSOD, two yeast MnSODs, one from *Saccharomyces cerevisiae* mitochondria (ScMnSOD) and the other from *Candida albicans* cytosol (CaMnSODc), were isolated and characterized. ScMnSOD and CaMnSODc are similar in catalytic

kinetics, spectroscopy, and redox chemistry, and they both rest predominantly in the reduced state (unlike most other MnSODs). At high  $[O_2^{\cdot-}]$ , the dismutation efficiencies of the yeast MnSODs surpass those of human and bacterial MnSODs, due to very low level of product inhibition. Optical and parallel-mode electron paramagnetic resonance (EPR) spectra suggest the presence of two  $Mn^{3+}$  species in yeast  $Mn^{3+}$ SODs, including the well-characterized 5-coordinate  $Mn^{3+}$  species and a 6-coordinate L– $Mn^{3+}$  species with hydroxide as the putative sixth ligand (L). The first and second coordination spheres of ScMnSOD are more similar to bacterial than to human MnSOD. Gln154, an H-bond donor to the Mn-coordinated solvent molecule, is slightly further away from Mn in yeast MnSODs, which may result in their unusual resting state. Mechanistically, the high efficiency of yeast MnSODs could be ascribed to putative translocation of an outer-sphere solvent molecule, which could destabilize the inhibited complex and enhance proton transfer from protein to peroxide. Our studies on yeast MnSODs indicate the unique nature of human MnSOD in that it predominantly undergoes the inhibited pathway at high  $[O_2^{\cdot-}]$ .




## INTRODUCTION

Superoxide dismutases (SODs) are enzymes that function to convert the superoxide ( $O_2^{\cdot-}$ ) anion to dioxygen and hydrogen peroxide ( $H_2O_2$ ) catalytically via sequential reduction and oxidation of a metal cofactor at virtually diffusion-controlled rate constants (Scheme 1).

Manganese-containing superoxide dismutases (MnSODs) are found in a wide variety of species, and, in these species, it can be localized either in the cytosol or in the mitochondria or in both. In bacteria, MnSOD is cytosolic, while in yeast both mitochondrial and cytosolic forms exist. In mammalian cells, it is mitochondrial. MnSOD catalyzes the disproportionation of  $O_2^{\cdot-}$  by a mechanism that distinguishes it from the other SODs in that it becomes a less effective antioxidant when  $O_2^{\cdot-}$  levels are high relative to enzyme concentration.<sup>1</sup> Specifically, human MnSOD<sup>2</sup> has been shown to be very inefficient at removal of high concentrations of  $O_2^{\cdot-}$ , whereas the MnSOD from bacteria (*Escherichia coli*<sup>2</sup> and *Deinococcus radiodurans*<sup>3</sup>) and yeast

## Scheme 1. The Dismutase Mechanism







(*Saccharomyces cerevisiae*)<sup>4</sup> are more efficient under the same conditions. It is this feature of differential efficiency at  $O_2^{\cdot-}$  removal that led to our interest in exploring the activity of MnSODs from different cellular sources and locations.

Efficient removal of high concentrations of  $O_2^{\cdot-}$  will, by necessity, produce high local concentrations of  $H_2O_2$ .  $H_2O_2$  is a biologically important class of reactive oxygen species (ROS). It can be reduced to hydroxyl radical ( $\cdot OH$ ) through a Fenton-type reaction with a reduced metal ( $M^{(n-1)+} + H_2O_2 \rightarrow M^{n+} + OH^- + \cdot OH$ ), whose reaction with cellular components causes severe

Received: August 16, 2011

Published: November 12, 2011

## Scheme 2. Catalytic Mechanism of MnSOD



damage to lipids, proteins, and nucleotides. The oxidative damage caused both directly and indirectly by  $\text{H}_2\text{O}_2$  is implicated in cancer and aging.<sup>5,6</sup> Besides its toxic effects, the roles of  $\text{H}_2\text{O}_2$  in aiding phagocytes in the killing of ingested microorganisms as well as in some oxidative biosynthesis are well-known.<sup>7</sup> In addition, in recent years there is rising evidence of  $\text{H}_2\text{O}_2$  functioning as a signaling agent in higher organisms.<sup>7–11</sup>

The mechanism by which MnSOD removes high  $\text{O}_2^-$  concentrations less efficiently than other SODs involves product inhibition. The overall reduction of  $\text{O}_2^-$  by an oxidized metal (Scheme 2), when carried out by  $\text{Mn}^{2+}\text{SOD}$ , occurs either through a pathway (reaction 2) where delivery of protons to the bound peroxo moiety is effectively instantaneous (an “outer-sphere” pathway) or through a pathway (reaction 3) where an identifiable transient is formed (an “inner-sphere” pathway), defined here as a “product-inhibited” complex that has been described as a side-on Mn(III)–peroxo species.<sup>12</sup> The inner-sphere pathway can become significant at high  $\text{O}_2^-$  concentrations, and, if the dissociation of peroxide ( $k_4$ ) is relatively slow, this can lead to reduced catalytic efficiency of MnSOD. The level of product inhibition is described by  $k_2/k_3$ , the gating ratio of the two pathways.

*S. cerevisiae* is an important single-cell model for studying human and other eukaryotic cells. Like human cells, it expresses a homotetrameric MnSOD in its mitochondria (ScMnSOD), which shares sequence similarity with human MnSOD. To our surprise, our previous study showed ScMnSOD to be the most efficient at removing high  $\text{O}_2^-$  concentrations, among MnSODs that have been characterized to date.<sup>4</sup> However, the origins of the low level of product inhibition are unclear.

*S. cerevisiae* and humans have a common cellular localization of their SODs. They both have MnSODs only in their mitochondrion and copper–zinc SODs (CuZnSOD) in their cytosol. In contrast, a closely related yeast *Candida albicans* expresses a MnSOD that lacks the mitochondrial leader sequence and is active in the cytosol (CaMnSODc) in addition to its mitochondrial CaMnSOD and cytosolic CaCuZnSOD. CaMnSODc shares 70% sequence similarity with ScMnSOD and has been proposed to be important for survival of *C. albicans* in nutrient-limiting conditions.<sup>13</sup> (We have chosen for the sake of convenience to distinguish between cytosolic and mitochondrial MnSODs by adding a “c” to the MnSOD to indicate that it is a cytosolic enzyme. This distinction becomes important in those organisms (*C. albicans*) where MnSOD is found in both locations. This distinction is not relevant to human and *S. cerevisiae* as all of the MnSOD is mitochondrial.) Overexpression of CaMnSODc has been shown to rescue the phenotypes of *S. cerevisiae* strain lacking the cytosolic CuZnSOD.<sup>13</sup>

To understand further the mechanism of product inhibition in MnSOD and its implication in  $\text{H}_2\text{O}_2$  signaling in higher organisms, we isolated and characterized ScMnSOD and CaMnSODc. Herein, we report the first comparison of the catalytic activity of

the two yeast MnSODs, the crystal structures of their metal-binding sites, and their spectroscopic characterization. We discuss how these factors relate to the function of yeast MnSODs; we further compare human, bacterial, and yeast MnSODs; and we explore the implications with respect to the evolution of  $\text{H}_2\text{O}_2$  signaling.

## MATERIALS AND METHODS

**DNA Manipulation of CaMnSODc and Transformation into *S. cerevisiae*.** The pVT102U-CaMnSODc (with *URA3* and *AMP* marker) vector containing the complete coding sequence of CaMnSODc was generously provided by Prof. Bourbonnais.<sup>13</sup> Sequencing showed that it is an I169V mutant of the wild-type protein accidentally created during cloning. The primers 5'-GTTCCAAT-TATTGCCATCGATGCTTGGGAA-C-3' and 5'-GTTCCCAAGCA-TCGATGGCAATAATTGGA-AC-3' were used to create the wild-type cDNA (mutations are in bold type). The PCR products were transformed into *E. coli* DH5 $\alpha$  strain and screened by ampicillin selection. The purified pVT102U-CaMnSODc vector was transformed into *S. cerevisiae* *sod2* $\Delta$  strain (EG110; *Mat-alpha*, *ura3*, *leu2*, *trp1*, *his3*, *sod2::TRP1*).

**Expression and Purification of ScMnSOD and CaMnSODc.** Yeast cells carrying Yep352-ScMnSOD<sup>4</sup> were grown in YPEG media (1% yeast extract, 2% peptone, 3% glycerol, 2% ethanol, pH 4) supplemented with 0.5 mM Mn(II) sulfate at 30 °C to OD > 20, and then harvested by centrifugation at 12 000g for 10 min. ScMnSOD was isolated as previously described.<sup>4</sup>

Yeast cells carrying pVT102U-CaMnSODc were grown in YPD (1% yeast extract, 2% peptone, 2% dextrose, pH 4) media supplemented with 0.5 mM Mn(II) sulfate at 30 °C to OD > 10, and then harvested by centrifugation at 12 000g for 10 min. The procedures for isolation of CaMnSODc and ScMnSOD were the same<sup>4</sup> except that the procedures of the DE52 column were different. CaMnSODc does not bind tightly to the DE52 resin, while most other proteins in the crude extract do. The active fractions from the hydrophobic interaction column were pooled and dialyzed against water and then 5 mM potassium phosphate (pH 8) at 4 °C. The protein then was loaded on a DE52 column, and washed with 2 column volumes of 5 mM potassium phosphate (pH 8). CaMnSODc fell into the load and wash during the chromatography. The load and wash were combined, concentrated, and loaded onto a G300 gel filtration column.

**Pulse Radiolysis.** The pulse radiolysis experiments were carried out using the 2 MeV Van de Graaff accelerator at Brookhaven National Laboratory. Upon irradiation of water with a pulse of energetic electrons, hydrated electrons ( $e_{\text{aq}}^-$ ), hydroxyl radicals ( $\cdot\text{OH}$ ), and, in lesser yield, hydrogen atoms ( $\text{H}\cdot$ ) are the primary radicals produced. Superoxide radical is then generated in air-saturated aqueous solution containing sodium formate through the following reactions:  $\cdot\text{OH} + \text{HCO}_2^- \rightarrow \text{H}_2\text{O} + \text{CO}_2^{\cdot-}$ ,  $\text{O}_2 + \text{CO}_2^{\cdot-} \rightarrow \text{O}_2^{\cdot-} + \text{CO}_2$ ,  $e_{\text{aq}}^- + \text{O}_2 \rightarrow \text{O}_2^{\cdot-}$ ,  $\text{H}\cdot + \text{O}_2 \rightarrow \text{HO}_2\cdot$ .

The experiments to measure catalytic rates were carried out using two methodologies. One set of studies involved oxidizing the resting enzyme with substoichiometric quantities of  $\text{O}_2^-$  and following the appearance of  $\text{Mn}^{3+}\text{SOD}$ , which has an absorbance band at around 480 nm (extinction coefficient  $\approx 800 \text{ M}^{-1} \text{ cm}^{-1}$ ). The rate of reaction 2 ( $k_2$ ) was determined by fitting the resultant time trace of  $\text{Mn}^{3+}$  growth to a first-order reaction. In this set of experiments, 2:1 [ $\text{H}_2\text{O}_2$ ]:[MnSOD] was added as previous studies had shown that  $\text{H}_2\text{O}_2$  effectively reduces  $\text{Mn}^{3+}\text{SOD}$  to  $\text{Mn}^{2+}\text{SOD}$ . The other method involved following the decay of various concentrations of  $\text{O}_2^-$  at 260 nm using a 1:1 to 1:50 ratio of [MnSOD]:[ $\text{O}_2^-$ ]. Here, all of the rate constants except  $k_2$  were calculated by fitting the data obtained to a mechanism described in Scheme 2 using the Chemical Kinetics program in PRWIN.<sup>14</sup>

All pulse radiolysis samples were prepared in 10 mM potassium phosphate, 10 mM sodium formate, and 10 or 100  $\mu$ M EDTA at 25 °C. All MnSOD concentrations were taken as the ICP-measured concentration of manganese in the sample. As all of the manganese concentrations were less than the corresponding protein concentrations, we assumed that all of the manganese was bound. This had the advantage of allowing direct comparison of activity of a tetrameric and a dimeric enzyme as everything is normalized to a monomeric unit. However, it also assumes no cooperativity between metal sites. The pH of the buffer was adjusted using ultrapure (Baker Ultrex) sodium hydroxide and sulphuric acid as needed.

**EPR Spectroscopy.** X-band (9.47 GHz) perpendicular-polarization ( $B_0 \perp B_1$ ) continuous-wave electron paramagnetic resonance (CW EPR) spectra were recorded using a Bruker ECS106 spectrometer equipped with a TE102 cavity (ER3902ST). Parallel-polarization ( $B_0 \parallel B_1$ ) data were acquired using a dual-mode cavity (ER 4116DM) operating at 9.39 GHz. Cryogenic temperatures were achieved and controlled using an Oxford Instruments ESR900 liquid helium cryostat in conjunction with an Oxford Instruments ITC503 temperature and gas flow controller. Spectral simulations were performed in MatLab using EasySpin 3.1.<sup>15,16</sup>

**Crystallization of ScMnSOD and CaMnSODc.** The apoprotein of ScMnSOD aggregates after the native protein is heated at 70 °C in MOPS, while the metalated ScMnSOD stays folded in solution. Fully metalated ScMnSOD was obtained by heating as-isolated protein (20 mM MOPS, pH 7) at 70 °C for 1 h and removing aggregated protein by centrifugation. The reductive methylation of lysine residues of ScMnSOD was carried out as described, to improve the diffraction of protein crystals.<sup>17</sup> The activities of native and methylated ScMnSOD are similar in the SOD activity assay.<sup>18</sup> All of the free amino groups of lysine residues and the N-terminal were methylated as confirmed by mass spectrometry (Figure S1).


Methylated ScMnSOD was crystallized by hanging-drop vapor diffusion at 4 °C against a well solution of 0.2 M sodium malonate (pH 7) in 20% (w/v) polyethylene glycol 3350 with a protein concentration of 7 mg/mL. Native CaMnSODc was crystallized by hanging-drop vapor diffusion at 4 °C against a solution of 0.1 M magnesium chloride, 0.1 M sodium chloride, and 0.1 M trisodium citrate (pH 5.5) in 30% (w/v) polyethylene glycol 400 with a protein concentration of 7 mg/mL. ScMnSOD and CaMnSODc crystals were cryo-protected in mother liquor solution containing 30% glycerol and flash frozen in liquid nitrogen prior to data collection.

**Crystallography: Data Collection and Refinement.** All data were collected at 100 K at the UCLA X-ray diffraction facility, using a Rigaku FRE+ generator and a Rigaku HTC detector. All data were processed using DENZO and SCALEPACK.<sup>19</sup> ScMnSOD was phased by molecular replacement using human MnSOD Q143N mutant (1QNM). CaMnSODc was phased by molecular replacement using ScMnSOD (3LSU). All of the molecular replacement was done using PHASER.<sup>20</sup> The models were built using COOT.<sup>21</sup> All model refinement was done using REFMAC<sup>22</sup> and PHENIX.<sup>23</sup>

Coordinates and structure factors have been deposited in the PDB with accession numbers 3LSU for the ScMnSOD structure and 3QVN for the CaMnSODc structure.

**Miscellaneous Methods.** The metal contents of the purified proteins were determined by ICP-MS (Agilent 7500 series). UV-visible absorption spectra were collected on a Shimadzu UV-2501 PC spectrophotometer.

The mass weight of the protein subunit was determined by electrospray ionization mass spectrometry (ESI-MS) via a triple quadrupole instrument (API III, Applied Biosystems). An  $m/z$  600–2200 range was scanned (positive mode, orifice 90 V, 0.3 Da step size, 5.61 s/scan). Data were processed and analyzed using MacSpec 3.3, Hypermass, and BioMultiview 1.3.1 software (Applied Biosystems).



**Figure 1.** The HPLC-SEC profiles of CaMnSODc (a) and ScMnSOD (b). Inset: The plot of the molecular weight of the five standards (●), ScMnSOD (green ▲), and CaMnSODc (red ■) versus their retention time. The concentration of proteins loaded into the column is 6 mg/mL. The column was calibrated using five standards: (1) bovine thyroglobulin (670 kDa), (2) bovine  $\gamma$ -globulin (158 kDa), (3) ovalbumin (44 kDa), (4) horse myoglobin (17 kDa), and (5) vitamin B12 (1.35 kDa). The buffer contained 50 mM sodium phosphate (pH 6.7) and 50 mM sodium chloride.


The mass weight of the native protein was determined by HPLC (Agilent 1200 series) using a size exclusion column (Tosoh Bioscience, TSK gel G2000SW 7.5 mm ID X 30 cm) at a flow rate of 0.5 mL/min. The column was calibrated using five standards: bovine thyroglobulin (670 kDa), bovine  $\gamma$ -globulin (158 kDa), ovalbumin (44 kDa), horse myoglobin (17 kDa), and vitamin B12 (1.35 kDa).

## RESULTS

**Tetramer versus Dimer.** CaMnSODc was overexpressed and purified from *S. cerevisiae*, with 0.6 equiv of manganese per subunit as determined by ICP-MS. The subunit mass weight is 22.7 kDa as determined by ESI-MS. The purity was confirmed by SDS-PAGE gel (Figure S2). CaMnSODc has been reported to be a homotetramer,<sup>13</sup> but our investigation of the native protein using gel filtration chromatography indicates that its molecular weight in solution under our conditions is  $\sim$ 46 kDa, corresponding to a homodimer (Figure 1). It should be noted, however, that we found that CaMnSODc crystallized as a tetramer under other conditions (see below). ScMnSOD, in contrast to CaMnSODc and analogous to the human MnSOD, is a homotetramer under our conditions, although the two yeast enzymes share great sequence similarity.

**Kinetic Studies.** Solutions containing 1  $\mu$ M MnSOD at pH 7 were pulse irradiated, and the disappearance of  $O_2^-$  was followed at 260 nm; see Figure 2. The traces we collected for both CaMnSODc and ScMnSOD were plotted along with previous data collected under identical conditions for MnSOD from human, *E. coli* and *D. radiodurans*.<sup>2,3</sup> When the  $O_2^-$  concentration was in 7-fold excess over the enzyme, human MnSOD already showed a depressed activity, yet the  $O_2^-$  disappearance as a function of time was similar for the other four enzymes and could be fit to a first-order process (Figure 2A). When the ratio of  $[O_2^-]:MnSOD$  increased to 41, the five MnSODs displayed






**Figure 2.** Decay of 7  $\mu\text{M}$  (A) and 41  $\mu\text{M}$  (B)  $\text{O}_2^-$  catalyzed by 1  $\mu\text{M}$  (in Mn) human (a, black), *E. coli* (b, red), *D. radiodurans* (c, blue) MnSOD, ScMnSOD (d, green), and CaMnSODc (e, pink). The sample for pulse radiolysis contained 10 mM potassium phosphate (pH 7), 10 mM sodium formate, and 10  $\mu\text{M}$  EDTA. The  $\text{O}_2^-$  concentration in these figures is calculated from the absorbance at 260 nm.

dramatically different efficiencies at removing  $\text{O}_2^-$  (Figure 2B), which scale as CaMnSODc > ScMnSOD > *D. radiodurans* (*Dr*MnSOD) > *E. coli* (*Ec*MnSOD) > human MnSOD. For human and *E. coli* MnSOD, the  $\text{O}_2^-$  decay was characterized by a fast decrease, followed by a slow almost zero-order phase. It is the biphasic nature of the trace that suggests dismutation proceeded by a fast catalytic pathway and a slower, concomitant product-inhibited pathway. The gating values for these two enzymes are 1 and 5, respectively.<sup>2</sup> The trace obtained for the *Dr*MnSOD showed a much less distinct plateau region, and fitting the data yielded a gating value of 12.<sup>3</sup> Our earlier study of the reaction of ScMnSOD with high  $\text{O}_2^-$  concentrations led to a calculated gating value  $k_2/k_3$  of 16–20, indicating that the fast protonation pathway dominates.<sup>4</sup>

CaMnSODc removes high concentrations of  $\text{O}_2^-$  even more efficiently than ScMnSOD. Here,  $k_2$  of CaMnSODc was determined to be  $1.0 \times 10^9 \text{ M s}^{-1}$  by following the appearance of  $\text{Mn}^{3+}$  SOD ( $\lambda_{\text{max}} = 480 \text{ nm}$ ; see method, Figure S3) as CaMn<sup>2+</sup>SODc was oxidized by substoichiometric amounts of  $\text{O}_2^-$ . However, at physiological pH (6.5–7.5), CaMnSODc removed various concentrations of  $\text{O}_2^-$  ( $[\text{O}_2^-]:[\text{MnSOD}]$  from 2–48:1) at similar




**Figure 3.** The effects of pH on the overall activity of MnSODs. The rate constant was determined by fitting the disproportionation of low doses of  $\text{O}_2^-$  ( $[\text{O}_2^-]:\text{MnSOD}$  from 1–3) to a first-order process. Under these  $[\text{O}_2^-]:\text{MnSOD}$  ratios, the reaction of  $\text{Mn}^{2+}$ SOD with  $\text{O}_2^-$  proceeds predominantly through the outer-sphere pathway. This rate constant reflects a combination of  $k_1$  and  $k_2$ . MnSOD is from human ( $\blacklozenge$ ), *E. coli* (red  $\blacksquare$ ), *S. cerevisiae* (green  $\bullet$ ), and *C. albicans* (pink  $\blacktriangledown$ ). The data points marked with an arrow correspond to the rate constants measured for the yeast enzymes after adjusting the pH of the sample solution from 9.5 to neutral.

rates, showing that only minimal product inhibition occurred under these conditions. Even at  $[\text{O}_2^-]:\text{MnSOD}$  ratios as high as 48, the activity of CaMnSODc was comparable to that of *S. cerevisiae* CuZnSOD, an enzyme that does not exhibit product inhibition (Figure S4). The disappearance of  $\text{O}_2^-$  catalyzed by CaMnSODc could be fitted to a first-order process.

To achieve such a high efficiency, CaMnSODc would either carry such a high gating ratio ( $k_2/k_3$ ) that the reaction of  $\text{Mn}^{2+}$  SOD with  $\text{O}_2^-$  proceeds almost completely via the fast protonation (reaction 2), or it has such a high dissociation rate ( $k_4$ ) that the product-inhibited complex decomposes before it is detected during pulse radiolysis (reaction 4). Assuming that  $k_4$  is  $50 \text{ s}^{-1}$ , which is close to that of bacterial dimeric MnSODs, the calculated  $k_2/k_3$  is 65. Assuming that  $k_2/k_3$  is 20, which is comparable to that for ScMnSOD,  $k_4$  is calculated to be  $210 \text{ s}^{-1}$ . The latter scenario is more likely, because a dissociation rate of  $210 \text{ s}^{-1}$  is not dramatically different from a  $k_4$  of  $130 \text{ s}^{-1}$  for human MnSOD and 90–140  $\text{s}^{-1}$  for ScMnSOD. In addition, dissociation rates as large as  $1000 \text{ s}^{-1}$  have been reported in human Y34 mutants.<sup>24</sup> A  $k_2/k_3$  ratio of 65, however, is 3 times as large as the gating ratio of ScMnSOD, the least inhibited MnSOD that has been characterized to date. Therefore, the high efficiency of CaMnSODc is more likely due to the fast dissociation of the inhibited complex from the metal center.

Previous studies have shown that the rate of enzymatic catalysis as a function of pH is different for the human and bacterial MnSODs. Given that a major difference between the human and bacterial MnSODs is that the former is a tetramer while the latter are dimers, we compare here two yeast MnSODs, where one is a dimer and the other is a tetramer. In these studies, we were initially interested only in the change in the fast catalytic cycle ( $k_1$  and  $k_2$ ), which was measured by following the disappearance of  $\text{O}_2^-$  in the presence of ScMnSOD or CaMnSODc under conditions where the  $[\text{O}_2^-]:\text{MnSOD}$  ratio was between 1 and 3. The overall activity of both ScMnSOD and CaMnSODc decreased considerably as pH was increased (Figure 3).



**Figure 4.** Dependence of the level of product inhibition of ScMnSOD on pH. Decay of  $48 \mu M O_2^-$  catalyzed by  $1 \mu M$  (in Mn) ScMnSOD at pH 7.5 (a) and 8.9 (b). Inset: Dependence of the values of  $k_2/k_3$  (●) and  $k_4$  (▼) of ScMnSOD on pH. The sample for pulse radiolysis contained 10 mM potassium phosphate (pH 7), 10 mM sodium formate, and  $10 \mu M$  EDTA.

Both yeast enzymes resemble *EcMnSOD*, with a relatively constant activity up to pH 9, but are distinct from human MnSOD, where catalysis is relatively unchanged even up to pH 11. It is surprising that ScMnSOD and CaMnSODc are different in their quaternary structure, yet they exhibit such similar pH dependence. It is notable that the yeast MnSODs are even more sensitive to the change of pH than the bacterial MnSODs. The  $pK$  is  $\sim 8.5$  in two yeast MnSODs,  $\sim 9.5$  in *EcMnSOD*, and  $\sim 10.5$  in human MnSOD. Adjusting the pH of yeast MnSOD sample solutions from 9.5 back to 7 resulted in the restoration of their activity to  $\sim 50\%$  of that of the intact enzymes (Figure 3).

pH also has an impact on the level of product inhibition exhibited by ScMnSOD. The slow almost zero-order phase, which is a sign of the saturation of MnSOD by  $O_2^-$ , was discernible from the reaction of high concentrations of  $O_2^-$  ( $[O_2^-]:MnSOD = 48:1$ ) with ScMnSOD at pH 8.9 (Figure 4). This feature was barely seen under the same conditions at pH 7.5. The gating ratio  $k_2/k_3$  and the dissociation rate  $k_4$  were calculated by fitting the  $O_2^-$  decay into the four-reaction mechanism (Scheme 2; Figure 4, inset). The calculated  $k_2/k_3$  is a constant value over the pH range of 7–9.5. The dissociation rate  $k_4$ , however, dramatically decreases when pH is raised to 8 and reaches a plateau at pH  $\sim 8.5$  with a  $pK$  of  $\sim 8$ . Because of a depressed  $k_4$ , the dissociation of  $H_2O_2$  from the active site may become the rate-limiting step at high pH, resulting in a more inhibited ScMnSOD. The values of  $k_2/k_3$  and  $k_4$  of ScMnSOD were difficult to calculate at pH  $> 9.5$ , at which point  $k_1$  and  $k_2$  corresponded to the rate-limiting steps of the reaction. In the case that  $k_4$  stays constant at the plateau while  $k_1$ ,  $k_2$ , and  $k_3$  continue decreasing at pH  $> 9.5$  the dissociation of the inhibited complex may occur faster than its formation, leading to a decay of  $O_2^-$  at pH  $> 9.5$  that appeared to be noninhibited.

CaMnSODc, in contrast, did not show a pronounced occurrence of the slow zero-order phase even at elevated pH (Figure S5). The decay of  $O_2^-$  over the pH range of 7–9.5 could still be fit to a first-order process ( $R^2 = 1.5$ ). In this case, it is difficult to calculate either the ratio of  $k_2/k_3$  or the value of  $k_4$  based on fitting.


**Oxidation State.** Our previous work showed that tetrameric ScMnSOD, unlike all other wild-type MnSODs studied thus far,

is isolated predominantly in the 2+ oxidation state.<sup>4</sup> Dimeric CaMnSODc, structurally similar to bacterial MnSODs, is also isolated in the  $Mn^{2+}$  state (Figure S6–A). The two enzymes from yeast are also very similar with respect to the thermodynamics of oxidation and reduction. From this point on, if the results are very similar for the two yeast MnSODs, data from only one enzyme are shown.

The optical absorption spectrum of as-isolated ScMnSOD has a very weak visible absorption band around 480 nm (Figure 5A) that suggests that a small amount of  $Mn^{3+}$ SOD is present along with a significantly larger amount of  $Mn^{2+}$ SOD. The addition of sodium hyposulfite fully reduced the enzyme and completely quenched this absorption intensity (Figure 5A). On the basis of the difference in absorption between the fully reduced and the as-isolated enzyme (Figure 5A, inset) and using the extinction coefficient for  $Mn^{3+}$ SOD obtained from pulse radiolysis ( $\epsilon_{475} = 780 M^{-1} cm^{-1}$ ),  $\sim 10\%$  of the as-isolated enzyme is oxidized. The oxidized enzymes found in both as-isolated ScMnSOD and CaMnSODc have spectra that differ from those of most reported  $Mn^{3+}$ SODs in that additional absorption intensity at  $\sim 390$  nm is visible (Figure 5A, inset; Figure S6–A, inset). They are, however, similar to the spectrum of ScMn<sup>3+</sup>SOD that was reported previously.<sup>25</sup>

Oxidation of as-isolated ScMn<sup>2+</sup>SOD or CaMn<sup>2+</sup>SODc by 0.75 equiv of potassium permanganate rapidly gave absorption spectra characteristic of 5-coordinate  $Mn^{3+}$ SOD with a maximum at 480 nm (Figure 5B, S6–B) with no residual absorbance of permanganate. Both yeast MnSODs remained oxidized in solution and retained full activity as determined by pulse radiolysis. Interestingly, an absorption intensity around 390 nm increased slowly over time (Figure 5B, S6–B). On the basis of the difference between the immediately scanned and the final spectrum (Figure S7), a  $Mn^{3+}$ -containing species different from the 5-coordinate  $Mn^{3+}$ SOD was formed over time. Further addition of 1 equiv of sodium ascorbate selectively bleached the absorption of ScMn<sup>3+</sup>SOD or CaMn<sup>3+</sup>SODc around 390 nm, yielding the characteristic 5-coordinate  $Mn^{3+}$ SOD spectra (Figure 5C, S6–C), which has been reported previously.<sup>25</sup> Treatment of the yeast enzymes with 10-fold molar excess of ascorbate yielded the completely reduced enzymes. Therefore, the unknown species is kinetically more prone to reduction by ascorbate than the well-characterized 5-coordinate species. In contrast, 0.5 equiv of  $H_2O_2$  reduced the 5-coordinate  $Mn^{3+}$  and the unknown species simultaneously in 1 equiv of yeast  $Mn^{3+}$ SODs, giving optical spectra of fully reduced MnSOD (data not shown). The slow formation of the 6-coordinate adduct and its susceptibility to reduction by ascorbate are very puzzling, and we will continue to explore its origin.

One species known to absorb around 390 nm is the low-temperature *EcMn*<sup>3+</sup>SOD:azide adduct. At room temperature, the 5-coordinate *EcMn*<sup>3+</sup>SOD binds azide in such a fashion that the coordination number of the Mn ion does not change. At lower temperatures ( $T_m = 200$  K), this adduct transforms into an inner-sphere 6-coordinate azide– $Mn^{3+}$ SOD complex.<sup>26</sup> The room-temperature spectrum of azide-treated ScMn<sup>3+</sup>SOD shows a shift of the absorption band similar to that observed in other azide– $Mn^{3+}$ SODs<sup>26</sup> (Figure S8). However, there remains additional intensity at 390 nm that is reminiscent of the features present in the spectrum of the low-temperature azide adduct of *EcMn*<sup>3+</sup>SOD where azide is directly bound to the Mn ion. This may indicate that the unknown species in ScMn<sup>3+</sup>SOD is also 6-coordinate (L– $Mn^{3+}$ SOD, L is the putative sixth ligand).



**Figure 5.** The oxidation state of ScMnSOD. (A) Optical spectra of as-isolated ScMnSOD (—) and ScMnSOD reduced by sodium hydrosulfite (---). Inset: The difference spectrum between as-isolated and reduced ScMnSOD. (B) Optical spectra of ScMnSOD oxidized by potassium permanganate ( $[\text{KMnO}_4]:\text{MnSOD} = 0.75:1$ ) at pH 7.4 as measured over time. Time 0 refers to the start of the first scan immediately after mixing  $\text{KMnO}_4$  with the enzyme. (C) Reduction of  $\text{ScMn}^{3+}\text{SOD}$  (—, oxidized by 0.75 equiv of  $\text{KMnO}_4$  and allowed to equilibrate at room temperature for 2 h) by 1 equiv of sodium ascorbate ( $\cdots$ ). (D) Optical spectra of  $\text{ScMn}^{3+}\text{SOD}$  at different pH. The sample solutions contained  $190 \mu\text{M}$  (in Mn) enzyme in 25 mM potassium phosphate (pH 7.4).


The measurement of the redox potential of MnSOD is notoriously difficult. Our electrochemical studies on ScMnSOD and CaMnSODc failed to achieve efficient electron transfer between the electrode and the enzymes and thus only yielded an approximation of their redox potential as  $\sim 0.3$  V. This value is comparable to those determined for *E. coli* (0.3 V)<sup>27</sup> and human (0.4 V)<sup>28</sup> MnSOD, which are a reflection of the redox potential of the 5-coordinate  $\text{Mn}^{3+}\text{SOD}$  species. The redox potentials of human and *E. coli* MnSOD are close to that of ascorbate $^{\cdot-}$ ,  $\text{H}^+$ /ascorbate $\text{H}^-$  ( $E^\circ = 282$  mV), at neutral pH.

The  $\text{Mn}^{3+}\text{SOD}$  spectra obtained from pulse radiolysis, in which ScMn $^{2+}\text{SOD}$  or CaMn $^{2+}\text{SODc}$  was oxidized by substoichiometric amount of  $\text{O}_2^-$  (see method), merely showed formation of the well-characterized 5-coordinate  $\text{Mn}^{3+}\text{SOD}^4$  and were comparable to those of other  $\text{Mn}^{3+}\text{SODs}$ <sup>3,29</sup> (Figure 6). However, the time scale of each single measurement at each wavelength was shorter than 1 ms. Over 10 ms the reduction of yeast  $\text{Mn}^{3+}\text{SODs}$  by excess  $\text{H}_2\text{O}_2$  was discernible. Therefore, the

generation of the putative L– $\text{Mn}^{3+}\text{SOD}$  species likely occurred more slowly under pulse radiolysis conditions than the reduction of the 5-coordinate  $\text{Mn}^{3+}$  center by  $\text{H}_2\text{O}_2$ . Indeed the shifting of the spectrum around 390 nm was discernible in a much longer time scale ( $>10$  min) from the oxidation of as-isolated ScMnSOD using  $^{60}\text{Co}$  radiation as the source of  $\text{O}_2^-$  (Figure S9).

MnSODs that are isolated in the  $\text{Mn}^{3+}$  state possess absorption spectra that change modestly as pH is raised. For example, at pH  $> 9$  the 478 nm absorption feature corresponding to EcMn $^{3+}\text{SOD}$  begins to diminish slightly in intensity and shifts to shorter wavelength. This pK event has been ascribed to deprotonation of Tyr34 (numbering in ScMnSOD).<sup>30</sup> However, for ScMnSOD and CaMnSODc, the absorption intensity around 390 nm increased as pH was increased (Figure 5D, S6–D). ScMnSOD and CaMnSODc may not have the same pK as other  $\text{Mn}^{3+}\text{SODs}$ , or the spectral change due to the pK may have been obscured by the absorption shifting around 390 nm. The  $\text{Mn}^{3+}$  spectra of ScMnSOD obtained from pulse radiolysis at pH 7 and






**Figure 6.** The  $\text{Mn}^{3+}$ SOD spectrum of  $\text{CaMnSODc}$  obtained from pulse radiolysis. The pulse radiolysis sample contained 60  $\mu\text{M}$  (in Mn)  $\text{CaMnSODc}$  in 10 mM potassium phosphate (pH 7), 10 mM sodium formate, and 100  $\mu\text{M}$  EDTA. The enzyme was reduced prior to each pulse with 120  $\mu\text{M}$   $\text{H}_2\text{O}_2$ .

10 were compared, and a roughly 20% decrease in intensity but no shift in peak for the  $\text{Mn}^{3+}$ SOD absorbance at 480 nm were observed (Figure S10). The spectral shift over pH indicates that the sixth ligand is likely a hydroxide that with a higher concentration binds to  $\text{Mn}^{3+}$  more readily at high pH.

**EPR Spectra.** In agreement with the electronic absorption data shown above, results from perpendicular-mode X-band EPR experiments confirm that  $\text{CaMnSODc}$  is isolated predominately in the  $\text{Mn}^{2+}$  oxidation state (Figure S11-A). These spectra are similar to those obtained for  $\text{EcMn}^{2+}$ SOD with five separate electronic spin transitions observed due to zero-field splitting of the manifold of spin levels of the high-spin ( $S = 5/2$ )  $\text{Mn}^{2+}$  ion. These resonances are further split by the hyperfine interaction of the electron spin with the  $^{55}\text{Mn}$  nuclear spin ( $I = 5/2$ ). The corresponding parallel-mode EPR spectrum for as-isolated  $\text{CaMnSODc}$  possesses little intensity (Figure S11-B). In contrast, the spectra of the pH 7.65 permanganate-oxidized samples have at least eight intense transitions centered at 80 mT ( $g = 8.4$ ) (Figure 7A) that are somewhat similar to the six parallel-mode EPR transitions separated by  $\sim 10$  mT observed previously for  $\text{EcMn}^{3+}$ SOD.<sup>31</sup> Campbell et al.<sup>31</sup> showed that these signals arise from transitions within the  $m_S = |\pm 2\rangle$  manifolds of a high-spin ( $S = 2$ ) Mn(III) system. Analysis of the temperature dependence of the integrated signal intensity led to an evaluation of the magnitude and, importantly, sign of the zero-field splitting parameters,  $D = +2.10 \text{ cm}^{-1}$  and  $E = 0.24 \text{ cm}^{-1}$ . (Note: These values are consistent with those determined by analysis of variable-temperature, variable-field magnetic circular dichroism saturation behavior.<sup>32</sup>) That  $D > 0$  indicates that the lowest unoccupied metal-centered molecular orbital (MO) is  $3d_z^2$ -based.<sup>33</sup> These conclusions are borne out by the trigonal bipyramidal coordination geometry of the  $\text{Mn}^{3+}$  site determined by X-ray crystallography as well as results of numerous electronic structure calculations.<sup>34</sup>

In the present case of  $\text{ScMn}^{3+}$ SOD and  $\text{CaMn}^{3+}$ SODc, more than six parallel-mode resonances are observed, indicating that there are at least two  $\text{Mn}^{3+}$ -containing species present in the sample. Subtraction of the EPR spectrum of  $\text{EcMn}^{3+}$ SOD from that of  $\text{ScMn}^{3+}$ SOD reveals an underlying sextet with interpeak spacings of approximately 4.5–5.0 mT (Figure 7B). Such a small effective  $^{55}\text{Mn}$  hyperfine coupling parameter ( $A_{\parallel}$ ) has been




**Figure 7.** (A) Parallel-mode CW EPR spectrum of  $\text{CaMn}^{3+}$ SODc and  $\text{ScMn}^{3+}$ SOD oxidized by potassium permanganate ( $[\text{ScMnSOD}]:\text{KMnO}_4 = 1:0.75$ ) and of  $\text{ScMn}^{3+}$ SOD treated with 1 equiv of sodium ascorbate. (B) The deconvolution of the two spectra (top two spectra in panel A) of the permanganate-oxidized species shows two components, one assigned to a  $\text{Mn}^{3+}$  center with a  $^5A_{1g}$  ground state and one with a  $^5B_{1g}$  ground state. Instrument settings:  $\nu = 9.3959 \text{ GHz}$ ; microwave power = 32 mW; sweep rate = 1.81 mT/s; temp = 4.9 K.

observed previously for several 6-coordinate Mn(III) centers.<sup>33,35,36</sup> There has also been a preliminary report that showed a fraction of 5-coordinate  $\text{EcMn}^{3+}$ SOD converts to a new species at pH 11.54 that is characterized by a parallel-mode EPR spectrum with  $A_{\parallel}(^{55}\text{Mn}) = 3.3 \text{ mT}$ .<sup>31</sup>

The dramatic difference in  $A_{\parallel}$  for the two spectroscopically distinct  $\text{ScMn}^{3+}$ SOD species likely resulted from a change in the electronic ground state for the  $\text{Mn}^{3+}$  ion, from  $^5A_{1g}$  to  $^5B_{1g}$ .<sup>33</sup> Respectively, these states correspond to d-electron configurations wherein the  $3d_{z^2}$ -based and  $3d_{x^2-y^2}$ -based MOs are empty. A simple explanation for this change would be a new, strong ligand binding in the equatorial plane to the  $\text{Mn}^{3+}$  ion. This event would raise the energy of the  $3d_{x^2-y^2}$ -based MO allowing the fourth d-electron to occupy the  $3d_{z^2}$  orbital instead.

The EPR transition observed for both electron configurations is between the  $m_S = |\pm 2\rangle$  spin levels. For the  $^5B_{1g}$  state, the axial zero-field splitting interaction ( $D$ ) will be negatively signed, putting the  $|\pm 2\rangle$  manifold lowest in energy. Thus, this manifold will be populated at all temperatures, and if the energy splitting ( $\Delta$ ) of the  $m_S = |\pm 2\rangle$  spin levels at zero field is less than the energy of the incident microwave radiation ( $h\nu = 0.319 \text{ cm}^{-1}$  for the 9.396 GHz radiation employed here), a parallel-mode EPR transition will be observed. However, for the  $^5A_{1g}$  state, the sign of  $D$  is positive and the  $|\pm 2\rangle$  manifold will not be populated at low temperatures. For  $\text{CaMn}^{3+}$ SODc, increasing the temperature thermally populates this manifold, and the EPR transition becomes apparent. By monitoring the temperature-dependent intensity of the peak at 58 mT, we have determined that



**Figure 8.** Active site structure of ScMnSOD, showing the coordination sphere of manganese (chain A). Some second coordination sphere residues are also shown. Coordination and hydrogen bonds are indicated as solid and dashed lines, respectively.

$D = +1.90 \text{ cm}^{-1}$  and  $E = 0.20 \text{ cm}^{-1}$ , values similar to those found for the 5-coordinate form of *EcMn*<sup>3+</sup>SOD (see Figure S12).

Upon treatment of ScMn<sup>3+</sup>SOD with 1 equiv of ascorbate, the <sup>5</sup>B<sub>1g</sub> contribution to the parallel-mode EPR spectrum completely disappears (compare second and third traces in Figure 7A). This spectral change coincides with bleaching of the 390 nm feature in the electronic absorption spectrum (Figure 5C) upon addition of ascorbate, suggesting the narrowly split EPR signal and the 390 nm absorption intensity are spectral signatures of the same species.

**Crystal Structure.** The active site structure of the two yeast MnSODs was investigated by crystallography (Table S1). The crystals of both yeast MnSODs are colorless. The X-ray crystallography structure of ScMnSOD is always a tetramer (Figure S13–A). To our surprise, the quaternary structure of CaMnSODc varied with crystallization conditions. It is either a dimer (data not shown) or a tetramer (Figure S13–B). We believe that the quaternary structure of CaMnSODc depends on buffer condition (ion strength and viscosity) and protein concentration, and we will explore the factors that control its quaternary structure in a future publication.

ScMnSOD and CaMnSODc both maintain the 5-coordinate sphere of manganese conserved in all other MnSODs (Figure 8, S14).<sup>34,37–40</sup> Their active sites are superimposable, giving a root-mean-square deviation (rmsd) of 0.20 Å (Figure S14). For both enzymes, the observed manganese bond lengths are ~2.2 Å to each His NE2, 2.0 Å to Asp OD2, and 2.3 Å to OH2 (Table S2). These are only slightly longer than the corresponding distances in MnSODs that rest predominantly in the Mn<sup>3+</sup> state. Spectroscopic and computational evidence suggests that in the reduced state Mn is bound by a water molecule and in the oxidized state it is bound by a hydroxide.<sup>41</sup> The distance of the Mn–O bond should be farther in the bound water as it is not as strong a bond and closer in the bound hydroxide. The long Mn–O (solvent molecule) distances in both yeast MnSODs are consistent with Mn<sup>2+</sup>–OH<sub>2</sub> moieties. Although these shifts are small and within possible experimental error, the observed longer

coordination bond distances could reflect the 2+ oxidation state of ScMnSOD and CaMnSODc.

During X-ray crystallography data collections, metalloenzyme crystals are likely to be reduced by photoelectrons generated from the excitation of protein and solvent atoms by the incident X-ray beam.<sup>42</sup> As ScMnSOD and CaMnSODc are both isolated in the reduced form as indicated by their optical and EPR spectra (Figures 5, S6, and S11), the X-ray structures are certainly those of the reduced (Mn<sup>2+</sup>SOD) enzymes. For other MnSODs that are isolated in the Mn<sup>3+</sup> oxidation state, active site bond distances measured in their X-ray structures (Table S2) would suggest that the human and *Thermus thermophilus* crystals contained primarily Mn<sup>3+</sup>SOD and the remaining MnSODs were likely primarily reduced during the data collection.

## DISCUSSION


**The Relationship of Product Inhibition to Active Site Structure.** On the basis of the structural and sequence similarity shared between ScMnSOD and human MnSOD, ScMnSOD was expected to exhibit a level of product inhibition similar to human MnSOD. However, we found that ScMnSOD is dramatically gated toward the fast protonation pathway. CaMnSODc resembles ScMnSOD in its efficiency of removing O<sub>2</sub><sup>•−</sup> at high O<sub>2</sub><sup>•−</sup> doses. The yeast MnSODs are more like bacterial MnSODs in terms of function. Our studies on yeast MnSODs place human MnSOD in a unique class that predominantly undergoes the product-inhibited pathway at high levels of O<sub>2</sub><sup>•−</sup> among MnSODs that have been characterized to date.

Numerous studies on the structure–function relationship of MnSOD have been focused on the structure of second coordination sphere structure,<sup>24,43–47</sup> which is highly conserved in MnSOD from different organisms.<sup>34,37–39</sup> Superimposition of all active site residues (within 6 Å of Mn) of ScMnSOD on those of human or *E. coli* MnSOD gives rmsd values of 0.25 and 0.18 Å, respectively (Figure 9), indicating that in terms of second coordination sphere, ScMnSOD resembles *EcMnSOD* more than it does human MnSOD, despite the fact that ScMnSOD and human MnSOD are both tetramers and share greater sequence similarity. Of particular significance are highly conserved residues Tyr34 and His30, mutations of which usually cause changes in product inhibition.<sup>24,48</sup> Their positions in ScMnSOD overlap more with those in *EcMnSOD* than with those in human MnSOD (Figure 9). One would expect subtle shifts of these residues in the second coordination sphere to cause dramatic differences in the degree of product inhibition among MnSOD from different species. The factors that control the level of product inhibition will be subjects of future investigations.

**The Unusual Resting State of Yeast MnSODs and the Role of Gln154.** In contrast to human and bacterial MnSODs that rest predominantly as Mn<sup>3+</sup>SOD, the as-isolated proteins of yeast MnSODs are both ~90% reduced. Our preliminary electrochemical results indicate that the redox potentials of yeast MnSODs are comparable to those determined for human and bacterial MnSODs. However, previous studies on *EcMnSOD* have shown spontaneous oxidation of *EcMn*<sup>2+</sup>SOD in air,<sup>29</sup> which does not occur in yeast MnSODs. Therefore, the unusual resting state of the yeast enzymes may result from thermodynamically driven shifts in oxidation–reduction equilibria.

The fact that yeast MnSODs rest as Mn<sup>2+</sup>SOD could be linked with the positioning of an outer-sphere residue, Gln154 (the numbering in MnSODs: ScMnSOD, Gln154; CaMnSODc,





**Figure 9.** Superimposition of the first and second coordination spheres of ScMnSOD (green) to that of human MnSOD (pink, A) and EcMnSOD (yellow, B).

Gln155; human, Gln143; *E. coli*, Gln146). Conserved in all Mn and FeSODs, Gln154 donates an H-bond to the metal-bound solvent molecule and is the only contact of the metal-bound solvent molecule to the protein beyond the active site. Gln154 makes three hydrogen bonds to three highly conserved residues, Tyr34, Asn80, and Trp133 (numbering in ScMnSOD). Its hydrogen-bonding potential is fully utilized, and Gln154 seems firmly fixed in place. Gln154 together with Asp168 that is an H-bond acceptor from the metal-coordinated H<sub>2</sub>O or OH<sup>-</sup> determine the position of the coordinated solvent molecule. Mutagenesis on Gln154 often causes dramatic depression of reactivity,<sup>43,44,46</sup> indicating the key role of the NE2(Gln154)···O(solv) bond.


The active site reduction midpoint potential ( $E_m$ ) contributes directly to the driving forces of the two half reactions of O<sub>2</sub><sup>-</sup> dismutation. In Mn and FeSOD (structural homologues), Gln154 is critical in modulating  $E_m$  and thus contributes to the ion specificity as well as determining the resting state of Mn and FeSOD.<sup>12,49</sup> The second coordination sphere of Mn and FeSOD is similar except that the Gln154 amide nitrogen is ~0.7 Å further from the metal in FeSOD. DFT calculations on the structure of Mn and FeSOD suggest that reason for MnSOD resting

predominantly in the Mn<sup>3+</sup> state is the destabilization of coordinated H<sub>2</sub>O versus OH<sup>-</sup> by the closely placed glutamine side chain.<sup>12,50</sup> In Mn-substituted EcFeSOD in which Mn is placed away from Gln69, the  $E_m$  is elevated by 670 mV, and the enzyme completely rests as the reduced form.<sup>51</sup> Mutation of Gln143 to asparagine in human MnSOD also causes the enzyme to rest as Mn<sup>2+</sup>SOD.<sup>43</sup> Either EcMn(Fe)SOD or human Q143 mutants is only able to catalyze one half reaction, the oxidation of O<sub>2</sub><sup>-</sup>.<sup>43,51</sup>

Most MnSODs with solved structure conserve a NE2-(Gln154)···O(solv) H-bond of 2.85–2.95 Å (Table S2), and, as indicated by previous studies, the further the Gln154 amide nitrogen is away from Mn, the more the 2+ state is favored by the enzyme. Thus, the observation that the distance of Gln154 to Mn is slightly longer in the two yeast MnSODs (4.79 and 4.73 Å in ScMnSOD and CaMnSODc, respectively) than in the MnSODs that predominantly rest in the oxidized form (human, 4.65 Å; *E. coli*, 4.64 Å; *D. radiodurans*, 4.54 Å; *Caenorhabditis elegans*, 4.54 Å) may be related to the fact that the yeast MnSODs rest in the reduced state.

**The Relationship of Product Inhibition of Human MnSOD to H<sub>2</sub>O<sub>2</sub> Signaling in Mammals.** Human MnSOD is much less efficient at turning O<sub>2</sub><sup>-</sup> into H<sub>2</sub>O<sub>2</sub> than ScMnSOD when O<sub>2</sub><sup>-</sup> levels are high, although the two enzymes share great sequence similarity and are both located in mitochondria. Mitochondria are recognized as the source of O<sub>2</sub><sup>-</sup> with the production occurring through respiratory chain. H<sub>2</sub>O<sub>2</sub> is generated in mitochondrial matrix via the dismutation reaction of O<sub>2</sub><sup>-</sup> catalyzed by MnSOD. The produced H<sub>2</sub>O<sub>2</sub> then diffused into cytosol. The roles of H<sub>2</sub>O<sub>2</sub> are complex in higher organisms. In mammalian cells, the upregulated expression of antioxidant enzymes is not a universal response to supraphysiologic levels of H<sub>2</sub>O<sub>2</sub>.<sup>7</sup> Instead, physiologic levels of H<sub>2</sub>O<sub>2</sub> stimulate cell division, transformation, migration, or apoptosis.<sup>7,8,10,11</sup> This along with the fact that mammalian cells generate H<sub>2</sub>O<sub>2</sub> in response to various stimuli supports that H<sub>2</sub>O<sub>2</sub> is utilized as a signaling agent in numerous mammalian signaling pathways.<sup>7–11,52,53</sup> A well-established H<sub>2</sub>O<sub>2</sub>-induced signaling is the modulation of the tyrosine phosphorylation of proteins by causing the oxidative inhibition of tyrosine phosphatases.<sup>9</sup> Reversible inactivation of tyrosine phosphatases has been demonstrated to play a critical role in PTP and insulin signaling.<sup>54,55</sup> Several mitochondrial signaling pathways are stimulated by increased H<sub>2</sub>O<sub>2</sub>, including tumor necrosis factor (TNF)- $\alpha$ -induced apoptosis<sup>56</sup> and c-Jun NH<sub>2</sub>-terminal kinase (JNK)-induced apoptosis.<sup>52,57</sup> The participation of H<sub>2</sub>O<sub>2</sub> in signaling cascades may require tight regulation of its production in mitochondria.

A recent study demonstrates the production of large bursts of O<sub>2</sub><sup>-</sup>, termed “superoxide flashes”, in human mitochondria.<sup>58</sup> The events of superoxide flashes occur randomly in space and time, last for 20–30 s, and are confined to tiny elliptical areas.<sup>58</sup> Under the conditions that were tested, human MnSOD in mitochondrial matrix was expected to eliminate superoxide flashes in a shorter time scale, considering that the reactivity of human MnSOD was thought to be diffusion controlled ( $k = 2.3 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ ) and its intramitochondrial concentration is 10–40  $\mu\text{M}$ .<sup>52,59</sup> The observation of superoxide flashes indicates that human MnSOD is unable to maintain a steady-state concentration of O<sub>2</sub><sup>-</sup> in tiny elliptical areas in mitochondria. The most likely explanation for this phenomenon is that human MnSOD is significantly inhibited at these high concentrations of O<sub>2</sub><sup>-</sup>. Under these conditions, bursts of O<sub>2</sub><sup>-</sup> are not translated into bursts of

Scheme 3. Proposed Mechanism of the Reaction of Yeast Mn<sup>2+</sup>SOD with O<sub>2</sub><sup>-</sup>

H<sub>2</sub>O<sub>2</sub> that could result in aberrant oxidant-driven signaling in mitochondria.

Mitochondria are the main source of H<sub>2</sub>O<sub>2</sub>, and its decomposition in the mitochondrial matrix is catalyzed by glutathione peroxidase and catalase. The rate constants of mammalian glutathione peroxidase and catalase are  $5 \times 10^7$  and  $4.6 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ , respectively, and their intramitochondrial concentration is  $\sim 10\%$  that of MnSOD.<sup>59</sup> These antioxidants maintain a steady-state intramitochondrial H<sub>2</sub>O<sub>2</sub> concentration of  $\sim 0.5 \times 10^{-8} \text{ M}$ .<sup>59</sup> As no comparison of H<sub>2</sub>O<sub>2</sub> concentration in areas with or without superoxide flashes has been reported, it is unclear whether superoxide flashes cause “H<sub>2</sub>O<sub>2</sub> flashes” in tiny confined areas in mitochondria.

In contrast, neither superoxide flashes nor evidence of H<sub>2</sub>O<sub>2</sub> signaling in yeast have been reported. The different levels of product inhibition found in human and ScMnSOD may well be explained by the necessity for constrained generation of H<sub>2</sub>O<sub>2</sub> in human mitochondria, especially under the conditions associated with “superoxide flashes”.

**Binding of a Sixth Ligand and the Proposed Mechanism of Yeast MnSODs.** Native ScMn<sup>3+</sup>SOD and CaMn<sup>3+</sup>SODc display optical absorption spectra that are different from other Mn<sup>3+</sup>SODs. When the yeast enzymes were chemically oxidized, the spectra changed over time from characteristic 5-coordinate Mn<sup>3+</sup> spectra to spectra with additional absorption intensity around 390 nm. The growth of this absorption intensity can be ascribed to the binding of a sixth ligand, L, leading to formation of an L–Mn<sup>3+</sup>SOD complex, and because its formation is favored at high pH the sixth ligand is highly likely to be a hydroxide. EPR spectra suggest the presence of at least two Mn<sup>3+</sup>-containing species with distinct electronic ground states in yeast Mn<sup>3+</sup>SODs. One of them is the well-characterized 5-coordinate Mn<sup>3+</sup>SOD, and the other could be an L–Mn<sup>3+</sup>SOD species with the sixth ligand bound in the equatorial plane.

A 6-coordinate MnSOD that binds two H<sub>2</sub>O/OH<sup>-</sup> has been obtained in the past at cryogenic conditions or at high pH. The structure of cyro-trapped EcMnSOD crystallized at pH 8.5 shows a H<sub>2</sub>O/OH<sup>-</sup> bound to the equatorial plane of Mn, giving an octahedral coordination center.<sup>60</sup> Probing the Mn<sup>2+</sup> center of EcMnSOD using high-field EPR indicates transfer of a water molecule to the Mn center at 240–268 K.<sup>61</sup> EcMn<sup>3+</sup>SOD at pH 11.65 forms a putative 6-coordinate species, giving a heterogeneous EPR spectrum that is comparable to those reported here for ScMn<sup>3+</sup>SOD and CaMn<sup>3+</sup>SODc at pH 7.65.<sup>31</sup> Recently, Porta et al. have reported a 6-coordinate structure of EcMnSOD with peroxide bound as a side-on ligand to active site manganese.<sup>62</sup> However, the existence of a 6-coordinate MnSOD binding two H<sub>2</sub>O/OH<sup>-</sup> that is stable in solution at room temperature and neutral pH is novel.

Examination of the second coordination sphere of ScMnSOD shows two outer-sphere solvent molecules within 6 Å of the Mn

center, in addition to the Mn-coordinated water. As discussed by Tabares et al.,<sup>61</sup> the sixth ligand is most likely the water that is hydrogen bonded to Tyr34 and His30 (Figure 8). Tyr34 and His30 are highly conserved residues that are part of the proposed hydrogen-bonding network, which pumps protons from protein surface to active site.<sup>1,40,45,46</sup> They are also involved in the gating between fast and slow protonation pathways as well as the dissociation of H<sub>2</sub>O<sub>2</sub> from the active site.<sup>24,48</sup> This water, however, only occupies 50% of the subunits in the structure of ScMnSOD or EcMnSOD and cannot be seen in the structure of human MnSOD or CaMnSODc.

The binding site of O<sub>2</sub><sup>-</sup> in the MnSOD active site has been investigated using small anions like fluoride and azide as O<sub>2</sub><sup>-</sup> analogues. Azide binds to the sixth ligand site in proximity of Tyr34 in the plane of His26, His81, and Asp168 at low temperatures.<sup>26</sup> Fluoride also sits close to the same binding site and is likely hydrogen bonded to Tyr34.<sup>63</sup> Therefore, to react with the Mn ion, O<sub>2</sub><sup>-</sup> is highly likely bound in the sixth ligand site in the vicinity to Tyr34. O<sub>2</sub><sup>-</sup> has also been proposed to form hydrogen bonds with Tyr34 and the Mn-coordinated H<sub>2</sub>O but does not directly bind Mn.<sup>63</sup> In both cases, the outer-sphere water, which has been proposed to become the sixth ligand, is located in the vicinity to the O<sub>2</sub><sup>-</sup> binding site.

Herein, we propose a mechanism based on the tendency of yeast Mn<sup>3+</sup>SODs to bind two solvent molecules, which may explain their high efficiency at high O<sub>2</sub><sup>-</sup> doses (Scheme 3). When O<sub>2</sub><sup>-</sup> binds to (or merely approaches) the sixth ligand binding site close to Tyr34, Mn<sup>2+</sup> transfers an electron to O<sub>2</sub><sup>-</sup> (which becomes peroxide) and is oxidized to Mn<sup>3+</sup>. Because Mn<sup>3+</sup>SOD tends to bind two solvent molecules, the water molecule that is hydrogen bonded to Tyr34 and His30 may travel to the Mn<sup>3+</sup> center to compete with the peroxide for the sixth ligand site (Scheme 3). This could cause steric effects at the metal binding site, which may destabilize the inner-sphere peroxo–Mn<sup>3+</sup>SOD complex. This is consistent with the fact that ScMnSOD and CaMnSODc are significantly gated toward the outer-sphere fast protonation pathway, in which formation of the peroxo–Mn<sup>3+</sup>SOD species is disfavored. The outer-sphere water may not necessarily bind Mn<sup>3+</sup> ion, which could be a slow process, but merely fluctuate between the Mn center and Tyr34/His30 to create steric effects at the active site. Deviations in the positioning of outer-sphere solvent molecules have been reported to be important in enhancing proton transfer in other metalloenzymes.<sup>64</sup> In human carbonic anhydrase II (HACII), the positioning and structuring of water molecules in the active site cavity vary with the conformation of an outer-sphere residue, His64.<sup>64</sup>

The putative translocation of the outer-sphere solvent molecule in yeast MnSODs may also facilitate dissociation of the bound peroxide by enhancing proton transfer. Numerous previous studies show that MnSODs from various organisms exhibit

a similar hydrogen isotope effect of  $\sim 2$ , suggesting that proton transfer is involved.<sup>65</sup> Dissociation of  $\text{H}_2\text{O}_2$  from MnSOD active site requires two protons transferred from the protein. Rapid proton transfer from the axial solvent ligand to peroxide is expected to occur in all MnSOD. In yeast MnSODs, the other proton is likely donated by the outer-sphere  $\text{H}_2\text{O}$  that sits between Tyr34 and His30 upon translocation, which may highly enhance the proton transfer and facilitate the removal of the peroxide from the active site.

Indeed, the rate of peroxide release from active site ( $k_4$ ) in ScMnSOD is dramatically depressed at increased pH with a  $\text{pK}$  of  $\sim 8$ , which causes the level of product inhibition to increase (Figure 4). This is significantly different from human MnSOD, which maintain a constant  $k_4$  over a wide range of pH (6–10).<sup>65</sup> The depression of  $k_4$ , which is likely caused by slower proton transfer, could result from either deprotonation of the axial  $\text{H}_2\text{O}$  or the outer-sphere  $\text{H}_2\text{O}$  that is hydrogen bonded to Tyr34 and His30 or from deprotonation of Tyr34 as proposed to be responsible for the spectroscopic  $\text{pK}$  of EcMnSOD.<sup>30</sup> In the latter scenario, deprotonation of Tyr34 may destabilize the potential sixth ligand by eliminating its hydrogen bonding with Tyr34. Considering that the  $\text{pK}$  of  $k_4$  for ScMnSOD is as low as 8 (Figure 4) while the  $\text{pK}$  of Tyr34 is 11.5, deprotonation of the solvent ligands is more likely. This may support the hypothesis that the outer-sphere solvent molecule approaches the sixth ligand site of the  $\text{Mn}^{3+}$  ion and transfers a proton to the peroxide to facilitate its leaving.

However, the binding of two  $\text{H}_2\text{O}/\text{OH}^-$  in ScMn<sup>3+</sup>SOD and CaMn<sup>3+</sup>SODc may not be relevant to their fast catalytic kinetics and low level of product inhibition, but merely a coincidence. We will therefore continue investigating the correlation between product inhibition and the L–Mn<sup>3+</sup>SOD species by studying yeast MnSOD with mutations at certain second coordination sphere residues.

In conclusion, MnSODs from *S. cerevisiae* mitochondria and *C. albicans* cytosol both exhibit higher efficiencies of  $\text{O}_2^-$  dismutation at high  $[\text{O}_2^-]$  than do human and bacterial MnSODs. The unusual resting state of yeast MnSODs could be attributed to the slightly longer distance between Gln154 and Mn. Both yeast Mn<sup>3+</sup>SODs form a 6-coordinate species. The potential sixth ligand could be an outer-sphere water, and its translocation to Mn ion in the catalytic cycle may facilitate the reaction of Mn<sup>2+</sup>SOD with  $\text{O}_2^-$  through the noninhibited pathway. Among MnSODs that have been characterized to date, human MnSOD is the only enzyme that significantly undergoes the product-inhibited pathway at high  $[\text{O}_2^-]$ , which can be ascribed to complicated roles of  $\text{H}_2\text{O}_2$  in mammalian cells, especially as a signaling agent.

## ■ ASSOCIATED CONTENT

**S Supporting Information.** Crystallization conditions, pulse radiolysis, optical and EPR spectra, crystallography information, and active site bond distances. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## ■ AUTHOR INFORMATION

### Corresponding Author

[jsv@chem.ucla.edu](mailto:jsv@chem.ucla.edu); [cabelli@bnl.gov](mailto:cabelli@bnl.gov)

## ■ ACKNOWLEDGMENT

This work was supported by Grant DK46828, KOSEF/MEST through WCU project (R31-2008-000-10010-0) to J.S.V. and

National Institutes of Health and Grant GM48242 to R.D.B. Radiolysis studies were carried out at the Center for Radiation Chemistry Research at BNL, which is funded under Contract DE-AC02-98CH10886 with the U.S. Department of Energy and supported by its Division of Chemical Sciences, Geosciences, and Biosciences, Office of Basic Energy Sciences. We acknowledge Minglei Zhao (Department of Chemistry and Biochemistry, UCLA) for assisting us in the X-ray structural determination. We acknowledge Professor Michael Hill (Occidental College) for assisting us in the redox potential measurements.

## ■ REFERENCES

- (1) Abreu, I. A.; Cabelli, D. E. *Biochim. Biophys. Acta* **2010**, *1804*, 263.
- (2) Zheng, J.; Domsic, J. F.; Cabelli, D.; McKenna, R.; Silverman, D. N. *Biochemistry* **2007**, *46*, 14830.
- (3) Abreu, I. A.; Hearn, A.; An, H.; Nick, H. S.; Silverman, D. N.; Cabelli, D. E. *Biochemistry* **2008**, *47*, 2350.
- (4) Barnese, K.; Sheng, Y.; Stich, T. A.; Gralla, E. B.; Britt, R. D.; Cabelli, D. E.; Valentine, J. S. *J. Am. Chem. Soc.* **2010**, *132*, 12525.
- (5) Valko, M.; Rhodes, C. J.; Moncol, J.; Izakovic, M.; Mazur, M. *Chem.-Biol. Interact* **2006**, *160*, 1.
- (6) Giorgio, M.; Trinei, M.; Migliaccio, E.; Pelicci, P. G. *Nat. Rev. Mol. Cell Biol.* **2007**, *8*, 722.
- (7) Veal, E. A.; Day, A. M.; Morgan, B. A. *Mol. Cell* **2007**, *26*, 1.
- (8) Stone, J. R.; Yang, S. *Antioxid. Redox Signaling* **2006**, *8*, 243.
- (9) Rhee, S. G. *Science* **2006**, *312*, 1882.
- (10) Finkel, T.; Holbrook, N. J. *Nature* **2000**, *408*, 239.
- (11) Finkel, T. *Curr. Opin. Cell Biol.* **2003**, *15*, 247.
- (12) Jackson, T. A.; Brunold, T. C. *Acc. Chem. Res.* **2004**, *37*, 461.
- (13) Lamarre, C.; LeMay, J. D.; Deslauriers, N.; Bourbonnais, Y. *J. Biol. Chem.* **2001**, *276*, 43784.
- (14) Schwarz, H. BNL Pulse Radiolysis Program. Brookhaven National Laboratory.
- (15) Stoll, S.; Britt, R. D. *Phys. Chem. Chem. Phys.* **2009**, *11*, 6614.
- (16) Stoll, S.; Schweiger, A. *J. Magn. Reson.* **2006**, *178*, 42.
- (17) Walter, T. S.; Meier, C.; Assenberg, R.; Au, K. F.; Ren, J. S.; Verma, A.; Nettleship, J. E.; Owens, R. J.; Stuart, D. I.; Grimes, J. M. *Structure* **2006**, *14*, 1617.
- (18) Sutherland, M. W.; Learmonth, B. A. *Free Radical Res.* **1997**, *27*, 283.
- (19) Otwinowski, Z.; Minor, W. *Methods Enzymol.* **1997**, *276*, 307.
- (20) McCoy, A. J.; Grosse-Kunstleve, R. W.; Adams, P. D.; Winn, M. D.; Storoni, L. C.; Read, R. J. *J. Appl. Crystallogr.* **2007**, *40*, 658.
- (21) Emsley, P.; Cowtan, K. *Acta Crystallogr., Sect. D* **2004**, *60*, 2126.
- (22) Murshudov, G. N.; Vagin, A. A.; Dodson, E. J. *Acta Crystallogr., Sect. D* **1997**, *53*, 240.
- (23) Adams, P. D.; Grosse-Kunstleve, R. W.; Hung, L. W.; Ioerger, T. R.; McCoy, A. J.; Moriarty, N. W.; Read, R. J.; Sacchettini, J. C.; Sauter, N. K.; Terwilliger, T. C. *Acta Crystallogr., Sect. D: Biol. Crystallogr.* **2002**, *58*, 1948.
- (24) Perry, J. J.; Hearn, A. S.; Cabelli, D. E.; Nick, H. S.; Tainer, J. A.; Silverman, D. N. *Biochemistry* **2009**, *48*, 3417.
- (25) Ravindranath, S. D.; Fridovich, I. *J. Biol. Chem.* **1975**, *250*, 6107.
- (26) Whittaker, M. M.; Whittaker, J. W. *Biochemistry* **1996**, *35*, 6762.
- (27) Vance, C. K.; Miller, A. F. *Biochemistry* **2001**, *40*, 13079.
- (28) Leveque, V. J.; Vance, C. K.; Nick, H. S.; Silverman, D. N. *Biochemistry* **2001**, *40*, 10586.
- (29) Whittaker, J. W.; Whittaker, M. M. *J. Am. Chem. Soc.* **1991**, *113*, 5528.
- (30) Maliekal, J.; Karapetian, A.; Vance, C.; Yikilmaz, E.; Wu, Q.; Jackson, T.; Brunold, T. C.; Spiro, T. G.; Miller, A. F. *J. Am. Chem. Soc.* **2002**, *124*, 15064.
- (31) Campbell, K. A. Dissertation Thesis, University of California Davis, 1999.
- (32) Jackson, T. A.; Karapetian, A.; Miller, A. F.; Brunold, T. C. *Biochemistry* **2005**, *44*, 1504.



- (33) Campbell, K. A.; Force, D. A.; Nixon, P. J.; Dole, F.; Diner, B. A.; Britt, R. D. *J. Am. Chem. Soc.* **2000**, *122*, 3754.
- (34) Edwards, R. A.; Baker, H. M.; Whittaker, M. M.; Whittaker, J. W.; Jameson, G. B.; Baker, E. N. *J. Biol. Inorg. Chem.* **1998**, *3*, 161.
- (35) Krivokapic, I.; Noble, C.; Klitgaard, S.; Tregenna-Piggott, P.; Weihe, H.; Barra, A. L. *Angew. Chem., Int. Ed.* **2005**, *44*, 3613.
- (36) Campbell, K. A.; Lashley, M. R.; Wyatt, J. K.; Nantz, M. H.; Britt, R. D. *J. Am. Chem. Soc.* **2001**, *123*, 5710.
- (37) Borgstahl, G. E.; Parge, H. E.; Hickey, M. J.; Beyer, W. F., Jr.; Hallewell, R. A.; Tainer, J. A. *Cell* **1992**, *71*, 107.
- (38) Fluckiger, S.; Mittl, P. R.; Scapozza, L.; Fijten, H.; Folkers, G.; Grutter, M. G.; Blaser, K.; Cramer, R. *J. Immunol.* **2002**, *168*, 1267.
- (39) Trinh, C. H.; Hunter, T.; Stewart, E. E.; Phillips, S. E.; Hunter, G. J. *Acta Crystallogr., Sect. F: Struct. Biol. Cryst. Commun.* **2008**, *64*, 1110.
- (40) Perry, J. J.; Shin, D. S.; Getzoff, E. D.; Tainer, J. A. *Biochim. Biophys. Acta* **2010**, *1804*, 245.
- (41) Jackson, T. A.; Xie, J.; Yikilmaz, E.; Miller, A. F.; Brunold, T. C. *J. Am. Chem. Soc.* **2002**, *124*, 10833.
- (42) Carugo, O.; Djinovic Carugo, K. *Trends Biochem. Sci.* **2005**, *30*, 213.
- (43) Hsieh, Y. S.; Guan, Y.; Tu, C. K.; Bratt, P. J.; Angerhofer, A.; Lepock, J. R.; Hickey, M. J.; Tainer, J. A.; Nick, H. S.; Silverman, D. N. *Biochemistry* **1998**, *37*, 4731.
- (44) Leveque, V. J.; Stroupe, M. E.; Lepock, J. R.; Cabelli, D. E.; Tainer, J. A.; Nick, H. S.; Silverman, D. N. *Biochemistry* **2000**, *39*, 7131.
- (45) Whittaker, M. M.; Whittaker, J. W. *Biochemistry* **1997**, *36*, 8923.
- (46) Edwards, R. A.; Whittaker, M. M.; Whittaker, J. W.; Baker, E. N.; Jameson, G. B. *Biochemistry* **2001**, *40*, 15.
- (47) Guan, Y.; Hickey, M. J.; Borgstahl, G. E.; Hallewell, R. A.; Lepock, J. R.; O'Connor, D.; Hsieh, Y.; Nick, H. S.; Silverman, D. N.; Tainer, J. A. *Biochemistry* **1998**, *37*, 4722.
- (48) Davis, C. A.; Hearn, A. S.; Fletcher, B.; Bickford, J.; Garcia, J. E.; Leveque, V.; Melendez, J. A.; Silverman, D. N.; Zucali, J.; Agarwal, A.; Nick, H. S. *J. Biol. Chem.* **2004**, *279*, 12769.
- (49) Yikilmaz, E.; Porta, J.; Grove, L. E.; Vahedi-Faridi, A.; Bronshteyn, Y.; Brunold, T. C.; Borgstahl, G. E. O.; Miller, A. F. *J. Am. Chem. Soc.* **2007**, *129*, 9927.
- (50) Yikilmaz, E.; Xie, J.; Brunold, T. C.; Miller, A. F. *J. Am. Chem. Soc.* **2002**, *124*, 3482.
- (51) Miller, A. F.; Vance, C. K. *Biochemistry* **2001**, *40*, 13079.
- (52) Cadenas, E. *Mol. Aspects Med.* **2004**, *25*, 17.
- (53) Droge, W. *Physiol. Rev.* **2002**, *82*, 47.
- (54) Mahadev, K.; Wu, X.; Zilbering, A.; Zhu, L.; Lawrence, J. T.; Goldstein, B. J. *J. Biol. Chem.* **2001**, *276*, 48662.
- (55) Tonks, N. K. *Cell* **2005**, *121*, 667.
- (56) Melendez, J. A.; Dasgupta, J.; Subbaram, S.; Connor, K. M.; Rodriguez, A. M.; Tirosh, O.; Beckman, J. S.; JourD'Heuil, D. *Antioxid. Redox Signaling* **2006**, *8*, 1295.
- (57) Chen, K.; Thomas, S. R.; Albano, A.; Murphy, M. P.; Keaney, J. F., Jr. *J. Biol. Chem.* **2004**, *279*, 35079.
- (58) Wang, W.; et al. *Cell* **2008**, *134*, 279.
- (59) Cadenas, E.; Davies, K. J. *Free Radical Biol. Med.* **2000**, *29*, 222.
- (60) Borgstahl, G. E. O.; Pokross, M.; Chehab, R.; Sekher, A.; Snell, E. H. *J. Mol. Biol.* **2000**, *296*, 951.
- (61) Tabares, L. C.; Cortez, N.; Agalidis, I.; Un, S. *J. Am. Chem. Soc.* **2005**, *127*, 6039.
- (62) Porta, J.; Vahedi-Faridi, A.; Borgstahl, G. E. *J. Mol. Biol.* **2010**, *399*, 377.
- (63) Tabares, L. C.; Cortez, N.; Hiraoka, B. Y.; Yamakura, F.; Un, S. *Biochemistry* **2006**, *45*, 1919.
- (64) Merz, K. M.; Toba, S.; Colombo, G. *J. Am. Chem. Soc.* **1999**, *121*, 2290.
- (65) Hearn, A. S.; Stroupe, M. E.; Cabelli, D. E.; Lepock, J. R.; Tainer, J. A.; Nick, H. S.; Silverman, D. N. *Biochemistry* **2001**, *40*, 12051.