

Observing a Model Ion Channel Gating Action in Model Cell Membranes in Real Time in Situ: Membrane Potential Change Induced Alamethicin Orientation Change


Shuji Ye,^{*,†} Hongchun Li,[†] Feng Wei,[†] Joshua Jasensky,[‡] Andrew P. Boughton,[§] Pei Yang,[§] and Zhan Chen^{*,‡,§}

[†]Hefei National Laboratory for Physical Sciences at Microscale, and Department of Chemical Physics, University of Science and Technology of China, Hefei, Anhui 230026, People's Republic of China

[‡]Department of Biophysics, and [§]Department of Chemistry, University of Michigan, Ann Arbor, Michigan 48109, United States

S Supporting Information

ABSTRACT: Ion channels play crucial roles in transport and regulatory functions of living cells. Understanding the gating mechanisms of these channels is important to understanding and treating diseases that have been linked to ion channels. One potential model peptide for studying the mechanism of ion channel gating is alamethicin, which adopts a split $\alpha/3_{10}$ -helix structure and responds to changes in electric potential. In this study, sum frequency generation vibrational spectroscopy (SFG-VS), supplemented by attenuated total reflectance Fourier transform infrared spectroscopy (ATR-FTIR), has been applied to characterize interactions between alamethicin (a model for larger channel proteins) and 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine (POPC) lipid bilayers in the presence of an electric potential across the membrane. The membrane potential difference was controlled by changing the pH of the solution in contact with the bilayer and was measured using fluorescence spectroscopy. The orientation angle of alamethicin in POPC lipid bilayers was then determined at different pH values using polarized SFG amide I spectra. Assuming that all molecules adopt the same orientation (a δ distribution), at pH = 6.7 the α -helix at the N-terminus and the 3_{10} -helix at the C-terminus tilt at about 72° (θ_1) and 50° (θ_2) versus the surface normal, respectively. When pH increases to 11.9, θ_1 and θ_2 decrease to 56.5° and 45° , respectively. The δ distribution assumption was verified using a combination of SFG and ATR-FTIR measurements, which showed a quite narrow distribution in the angle of θ_1 for both pH conditions. This indicates that all alamethicin molecules at the surface adopt a nearly identical orientation in POPC lipid bilayers. The localized pH change in proximity to the bilayer modulates the membrane potential and thus induces a decrease in both the tilt and the bend angles of the two helices in alamethicin. This is the first reported application of SFG to the study of model ion channel gating mechanisms in model cell membranes.



1. INTRODUCTION

Ion channel proteins are transmembrane proteins that regulate ionic permeability in cell membranes. They are key elements in signaling and sensing pathways and provide selective transport in and out of cells.¹ Defective ion channels may be a factor in diseases such as cystic fibrosis, cardiac arrhythmias, and Parkinson's disease.^{2–9} The study of ion channel structures is important to understanding disease mechanisms and developing effective treatments.

Voltage-gated ion channels are a subset of transmembrane ion channels, activated by changes in electrical potential across the cell membrane.^{10,11} Much research has been done in obtaining ion channel crystal structures and understanding how these channels sense and respond to membrane potential changes.^{12–24} Although these studies have substantially improved our understanding of ion channel voltage sensing, the detailed mechanism for this process is still an open question. Currently, three models are under debate: the transporter model, the helical screw model, and the paddle model.^{22,23} We believe that direct, in situ structural measure-

ments of ion channels and related membrane proteins will elucidate the precise mechanism of ion channel gating.

Alamethicin, frequently used as a model for larger channel proteins,^{25–35} is a 20-residue peptide extracted from the fungus *Trichoderma viride*. Alamethicin molecules can form voltage-gated ion channels in membranes.^{25–38} The molecular structure and conformational features of alamethicin have been studied extensively.^{25–42} The X-ray crystal structure of alamethicin crystallized from methanol was determined to be predominantly helical with an N-terminal α -helix and a C-terminal domain containing a 3_{10} -helical element.³⁹ The Pro14 residue separating the two domains induces a 20° – 35° bend in alamethicin.³⁹

The mechanisms of alamethicin action within cell membranes have been widely studied.^{25–38} It is currently believed that alamethicin interacts with cell membranes through the barrel-stave mode, in which conducting pores in the membrane

Received: November 25, 2011

Published: March 15, 2012

are formed by parallel bundles of 3–12 helical alamethicin monomers surrounding a central, water-filled pore.^{27,28,37–39,43–45} However, the molecular-level details of the mechanism of alamethicin channel formation remain unclear,^{46,47} with contradictory results regarding its orientation in cell membranes in the absence of membrane potential. It has been variously suggested that alamethicin adopts a trans-membrane orientation,^{40–42,48–52} lies on the membrane surface,^{53–55} or both.^{56,57} A continuous distribution of orientations has also been proposed.⁵⁸ Recently, we applied sum frequency generation vibrational spectroscopy (SFG-VS) to investigate the interactions between alamethicin and different lipid bilayers without the presence of a membrane potential.⁵⁹ SFG-VS is a nonlinear optical spectroscopic technique which provides vibrational spectra of surfaces and interfaces.^{59–71} Our previous studies have indicated that alamethicin inserts into fluid-phase lipid bilayers, but lies down and/or aggregates on gel-phase bilayer surfaces.⁵⁹

To understand the gating mechanism of ion channels, it is important to investigate these biological molecules in the presence of a membrane potential. In this study, for the first time, we investigate the interactions between alamethicin and model membranes in situ in the presence of a transmembrane potential.

Membrane potential (e.g., lipid membrane dipole potential) plays important roles in protein–membrane interactions.^{72–74} In addition to an external electric field, protons and hydroxides (or pH variations) can also effectively modulate lipid membrane potential by altering surface charge and dipole potential.^{75–79} Upon exposing a lipid membrane to aqueous solutions of varying pH, the phosphate and choline groups of lipid molecules may change charge distribution at the membrane interface, resulting in alterations of their Debye length, membrane surface charge density, and zeta potential.^{75–79} This can lead to a change in membrane potential and an increase in channel protein conductance.⁷⁹ Because pH plays a crucial role in modulating the electric characteristics of zwitterionic-based lipid membranes, changes in pH have been used to modulate the transport properties of alamethicin oligomers inserted in zwitterionic-based model lipid membranes.⁷⁵ In addition, earlier studies have indicated that high pH values increased the open-state probability of the channel of *Entamoeba histolytica*, which has a mechanism of action similar to alamethicin.⁸⁰ With this in mind, we modulated the membrane potential by changing the pH of the subphase in contact with the lipid bilayer. We then investigated how changes in subphase pH affected the orientation of alamethicin in membranes using SFG-VS and attenuated total reflectance-Fourier transform infrared spectroscopy (ATR-FTIR). Results from this in situ study will help to understand ion channel proteins in action with the presence of a membrane potential.

2. EXPERIMENTAL SECTION

2.1. Materials and Sample Preparations. Alamethicin from *Trichoderma viride* was purchased from Sigma-Aldrich (St. Louis, MO) with a minimum purity of 90%. The lipids 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine (POPC) and 1,2-dimyristoyl-*sn*-glycero-3-phosphoethanolamine-*N*-(lissamine rhodamine B sulfonyl) ammonium salt (Rhod-DMPE) were purchased from Avanti Polar Lipids (Alabaster, AL). Potassium chloride (KCl), potassium phosphate (K_3PO_4), and deuterated water (D_2O) were ordered from Aldrich (Milwaukee, WI) with a purity of >99.0% and were used as received. The stock salt solution (1.0 M) of pH 6.7 or 11.9 was prepared by dissolving KCl or K_3PO_4 into ultrapure water from a Millipore system

(Millipore, Bedford, MA). Right-angle CaF_2 prisms were purchased from Altos (Trabuco Canyon, CA). They were thoroughly cleaned as described elsewhere⁵⁹ and were verified using SFG to be free of contamination before lipid deposition.

Single lipid bilayers were prepared on CaF_2 substrates using Langmuir–Blodgett and Langmuir–Schaefer (LB/LS) methods with a KSV2000 LB system.^{59,69,71} Lipid bilayers were immersed in water inside a 1.6 mL reservoir throughout the entire experiment. No SFG C–H stretching signal was detected after depositing a deuterated lipid monolayer or bilayer on a CaF_2 surface, indicating that there was no contamination on the CaF_2 surface when constructing the lipid monolayer or bilayer. For alamethicin–bilayer interaction experiments, $\sim 15 \mu L$ of alamethicin solution (in methanol with a concentration of 2.5 mg/mL) was injected into the reservoir, resulting in a final alamethicin concentration of 23 $\mu g/mL$. A magnetic micro stirrer was used to ensure a homogeneous concentration of peptide molecules in the subphase below the bilayer.

2.2. Polarized ATR-FTIR and SFG-VS Experiments. Details regarding SFG theories and instruments have been reported previously.^{59–71} The SFG and ATR-FTIR experimental details and data analysis methods used in this research are presented in the Supporting Information.

2.3. Estimation of Membrane Potential Difference. It has been demonstrated that membrane potentials can be measured directly using fluorescence spectroscopy.^{81–83} In this study, we estimated the membrane potential change based on the fluorescent intensity change of (10%Rhod-DMPE+90%POPC (in mass ratio))/POPC and POPC/(10%Rhod-DMPE+90%POPC) bilayers using our SFG experimental geometry (shown in the Supporting Information). The potential difference across the POPC/POPC bilayer was estimated to be -16.2 mV when the subphase pH was changed from 7.3 to 11.8. This value agrees with the results of phospholipid 1-stearoyl-2-oleoyl-phosphatidyl-choline reported by Zhou et al.⁷⁶

3. RESULTS AND DISCUSSION

3.1. SFG Results at pH = 6.7. SFG ssp and ppp spectra of alamethicin in a POPC/POPC bilayer at pH = 6.7 are shown in Figure 1. Spectra were collected after an injection of a 15 μL alamethicin/methanol solution into the subphase (~ 1.6 mL) of a POPC/POPC bilayer at pH 6.7. The resulting concentration of alamethicin in the subphase is 23 $\mu g/mL$. We observed three peaks in these spectra: a 1670 cm^{-1} peak originating from a




Figure 1. SFG spectra of alamethicin in a POPC/POPC bilayer at pH = 6.7.

helical structure dominated by α -helix with minor contribution from a 3_{10} -helix,⁵⁹ a 1635 cm^{-1} peak due to the 3_{10} -helical structure,⁵⁹ and a 1720 cm^{-1} signal generated by the carbonyl groups of the lipid bilayer. More details on the amide I peak assignments can be found in ref 59 and the references therein.

Alamethicin consists of two helical segments due to the presence of the helix-breaking Pro14 residue.³⁹ Here, we define the tilt angles between the principal axis of the α -helix (residues 1–13) or 3_{10} -helix (residues 14–20) and the POPC/POPC bilayer surface normal to be θ_1 and θ_2 , respectively. Molecular orientation information can be obtained by relating SFG susceptibility tensor elements χ_{ijk} ($i,j,k = x,y,z$) to the SFG molecular hyperpolarizability tensor elements β_{lmn} ($l,m,n = a,b,c$).^{59–71} For α -helices and 3_{10} -helices, the details of orientation determination using polarized SFG spectra have been published previously.⁶⁹ Using the relationship between the measured ppp and ssp spectral intensity ratios of the peaks at 1670 and 1635 cm^{-1} , it is possible to determine the orientation angles θ_1 and θ_2 .⁵⁹ Here, the experimentally measured $\chi_{\text{ppp}}/\chi_{\text{ssp}}$ ratios of the peaks at 1670 and 1635 cm^{-1} for alamethicin in the POPC/POPC bilayer at pH = 6.7 are 2.65 and 2.18. The resulting orientation angles θ_1 and θ_2 are calculated to be about 72° and 50° , assuming a δ -orientation distribution.

Our previous SFG studies indicated that alamethicin molecules also adopt a tilted orientation in a DMPC/DMPC bilayer with θ_1 and θ_2 calculated to be 63° and 43° .⁵⁹ In the POPC/POPC bilayer used in this work, the alamethicin molecules are tilted more toward the bilayer surface. “Hydrophobic matching” between the hydrophobic region of a peptide and the hydrocarbon region of a lipid bilayer is the major requirement for the stable transmembrane insertion of peptaibols.^{42,50–52} However, factors such as the hydrophobic moment of the peptide, the lipid saturated/unsaturated fatty acyl chain compositions, and the lipid bilayer phase transition temperature are also important.^{31,50–52} Although the hydrophobic thickness of a POPC bilayer is larger than that of a DMPC bilayer, the phase transition temperature of POPC is much lower than that of DMPC because of the unsaturated chain in POPC. This may lead to the difference in peptide tilt angle observed.

3.2. SFG Results at pH = 11.9. After all measurements were performed at pH = 6.7, the subphase pH was raised to 11.9 by adding $100\ \mu\text{L}$ of K_3PO_4 (in 1 M solution) into the subphase ($\sim 1.6\text{ mL}$ of DI water), which resulted in a 0.06 M K_3PO_4 solution. ATR-FTIR experiments confirmed that the POPC/POPC bilayer is stable at pH 11.9 in the absence of peptide based on the ATR-FTIR signals observed in the C–H stretching frequency region (not shown). Figure 2 shows the time-dependent intensity change of the SFG amide I peptide signals (1665 cm^{-1}) in the ppp polarization after changing the subphase pH from 6.7 to 11.9. The intensity increased immediately after adjusting the pH, indicating that the pH change rapidly affected the alamethicin interaction with the bilayer. Figure 3 presents the SFG spectra of alamethicin in a POPC/POPC bilayer at pH = 11.9, which shows that the SFG amide I intensity from alamethicin increased substantially in the POPC/POPC bilayer.

After adjusting the pH, the fitted $\chi_{\text{ppp}}/\chi_{\text{ssp}}$ ratios for the peaks shown in Figure 3 changed to 1.93 and 2.34. From these ratios, the orientation angles were determined to be $\theta_1 = 56.5^\circ$ and $\theta_2 = 45^\circ$. We compared the fitted χ_{ppp} and χ_{ssp} of the 1670 cm^{-1} peak at different pH values and found that $\chi_{\text{ppp}}(\text{pH} = 11.9)/$




Figure 2. The time-dependent intensity change of ppp SFG spectra at 1665 cm^{-1} in the alamethicin bound POPC/POPC bilayer after a change in pH from 6.7 to 11.9.




Figure 3. SFG ssp and ppp spectra of alamethicin in a POPC/POPC bilayer at pH = 11.9.

$\chi_{\text{ppp}}(\text{pH} = 6.7) = 3.63$ and $\chi_{\text{ssp}}(\text{pH} = 11.9)/\chi_{\text{ssp}}(\text{pH} = 6.7) = 4.11$. It is well-known that SFG signal intensity is related to the number of molecules detected and their orientations. Because the orientation was deduced above from the $\chi_{\text{ppp}}/\chi_{\text{ssp}}$ ratio, the effect of a change in the number of molecules can be isolated. Thus, by comparing spectral intensities, we quantified that the number of the molecules that inserted into the lipid bilayer increased about 2 times after adjusting the pH from 6.7 to 11.9.

To verify that the observed effect arose from the pH change alone instead of a change in the ionic strength of the subphase, we also investigated the effect of salt on the SFG signals from alamethicin in a POPC/POPC bilayer at pH = 6.7. After $300\ \mu\text{L}$ of KCl solution (1.0 M) (pH = 6.7) was added into the aqueous subphase (resulting in a 0.16 M KCl solution), alamethicin amide I intensity decreased (data shown in the Supporting Information). After the addition of KCl, the fitted χ_{ssp} and χ_{ppp} values for the alamethicin amide I signal decreased by a factor of 1.5 and 1.7, respectively. Overall, the ratio $\chi_{\text{ppp}}/\chi_{\text{ssp}}$ decreased slightly in this case, which would indicate that alamethicin stood up relative to the bilayer.^{59,69} In that case, an increase in signal intensity would be expected if the number of peptide molecules in the lipid bilayer remained the same.

Therefore, the observed decrease in SFG amide I intensity must come from the “salting out” effect, which results in fewer peptides in the bilayer. Such an effect has been observed in an earlier study of the effect of NaCl addition on the incorporation of alamethicin to a dioleoylphosphatidylcholine (DOPC) bilayer.⁸⁴ The salt experiment presented here confirms that the pH change induced more alamethicin to insert into the lipid bilayer. This change in intensity and the number of peptide molecules present was found not to be fully reversible when the pH of the subphase was returned to 7. After the SFG spectra in Figure 3 were collected, about 40 μL of H_3PO_4 (in 1.0 M stock solution) solution was injected into the subphase, and SFG spectra were collected after equilibration (while the subphase pH is ~ 8.0). After that, an additional 20 μL of H_3PO_4 (1.0 M) solution was injected into the subphase, and SFG spectra were again collected after further equilibration (at a subphase pH ~ 7.0). The SFG spectra and the time-dependent SFG signal change are shown in the Supporting Information. With the addition of H_3PO_4 , the alamethicin SFG amide I intensity decreased quickly. The final SFG amide I signal intensity of alamethicin after the pH returned to ~ 7 is lower than the initial SFG intensity, which may be due to the combination of pH and salt effects. Another possible reason may be due to the slightly altered lipid bilayer structure induced by the inserted alamethicin. As shown in Figure 1, a peak at $1710\text{--}1720\text{ cm}^{-1}$ generated by the carbonyl groups of the lipid bilayer was observed, while this peak disappeared in Figure 3. On the basis of this, we believe that the pH effect is not fully reversible.

3.3. ATR-FTIR Results. ATR-FTIR was used as a supplemental technique to substantiate SFG results. Polarized ATR-FTIR spectra of alamethicin in a POPC/POPC bilayer are displayed in Figure 4. According to our previous results,⁸⁵




Figure 4. Polarized ATR-FTIR spectra of alamethicin in a POPC/POPC bilayer at pH = 6.7 and 11.9.

spectra were fitted using three peaks centered at 1623, 1635, and 1660 cm^{-1} . The intensity ratio (R) of the signal measured using the p- versus s-polarized beam for the 1660 cm^{-1} peak is 1.15 at pH = 6.7 and 1.50 at pH = 11.9. Assuming a δ orientation distribution, the orientation angle is determined to be $\theta_1 = 74^\circ$ at pH = 6.7 and $\theta_1 = 58^\circ$ at pH = 11.9. The results from these ATR-FTIR experiments are comparable to the SFG

results of $\theta_1 = 72^\circ$ at pH = 6.7 and $\theta_1 = 56.5^\circ$ at pH = 11.9 presented above. In addition, the fitted absorbance area of the 1660 cm^{-1} peak increased 1.6 and 2.1 times for s and p polarizations, respectively. On the basis of the changes in absorbance area and dichroic ratio, we can deduce that the number of molecules inserted into the lipid bilayer increased by $\sim 2\times$ after adjusting the pH from 6.7 to 11.9, in agreement with our SFG result.

3.4. Combined SFG and ATR-FTIR Studies. In sections 3.1 and 3.2, we assumed that all molecules adopted the same orientation (a δ distribution). This is supported by the good agreement between results from SFG and ATR-FTIR (which measure different orientational parameters). By combining these results, it is possible to characterize a Gaussian distribution directly and measure both the average orientation angle θ and the distribution width σ simultaneously. As discussed previously,⁸⁶ SFG can measure $\langle \cos \theta \rangle$ and $\langle \cos^3 \theta \rangle$ (where “ $\langle \rangle$ ” means average). Here, the $\chi_{\text{ppp}}/\chi_{\text{ssp}}$ ratio measures $\langle \cos \theta \rangle / \langle \cos^3 \theta \rangle$. ATR-FTIR measures $\langle \cos^2 \theta \rangle$. Table 1 lists

Table 1. Deduced Orientation Angle and Angle Distribution Widths of θ_1 from SFG or ATR-FTIR Measurements, When a Gaussian Distribution Is Used

pH	method	Gaussian width			
		$\sigma = 0$	$\sigma = 5$	$\sigma = 8$	$\sigma = 10$
6.7	SFG	72	75	79	85
6.7	ATR-FTIR	74	75	77	78
11.9	SFG	56.5	57.5	58.5	60
11.9	ATR-FTIR	58	58.5	58.5	59

the deduced combinations of the average angle and angle distribution from measured SFG and ATR-FTIR data when a Gaussian distribution is used. Combining SFG and ATR-FTIR measurements, at pH = 6.7, $\theta_1 = 75^\circ$, $\sigma = 5^\circ$. At pH = 11.9, $\theta_1 = 58.5^\circ$, $\sigma = 8^\circ$. Nevertheless, the average angles are similar to those deduced by the delta orientation distribution, and the angle distribution widths are quite narrow. We believe that the narrow orientation distribution width is due to the specific interactions between lipid bilayer and alamethicin. This is different from protein-polymer interactions, where many different interactions can be involved, leading to broad orientation distributions for proteins adsorbed onto polymers.⁸⁷

3.5. Further Discussion. Surface charge-induced $\chi^{(3)}$ contributions have been extensively discussed in second harmonic generation spectroscopy.^{88–90} We have considered the possibility that the surface charge (or surface electric field) could induce $\chi^{(3)}$ contributions to SFG amide I signals.⁹¹ Such a method was also reported in a review article by Space and his colleagues.⁹² We found that for a static field of $2 \times 10^8\text{ V/m}$, the $\chi^{(3)}$ contribution is at least 1 order of magnitude smaller than the $\chi^{(2)}$ contribution. In the experiments presented in this work, our measured membrane potential is much smaller: only -16.2 mV surface potential difference for two different pH conditions. If we take the lipid bilayer thickness to be 5 nm, the calculated field is $3 \times 10^6\text{ V/m}$, which is almost 2 orders of magnitude smaller than the case discussed previously.⁹¹ Therefore, we expect the contribution from the $\chi^{(3)}$ term to be negligible as compared to the SFG signals ($\chi^{(2)}$ term).

Because orientation analysis depends on a known secondary structure, we next considered the possibility that the change in pH would affect protein conformation. In general, when alamethicin is denatured, a coiled-coil structure is formed,⁹³

which shifts the peak centers of the SFG amide I signal and requires additional spectral fitting using new peak centers. Both ssp and ppp SFG spectra collected from alamethicin at pH = 6.7 and pH = 11.9 can be fitted using the same two amide I peaks between 1600 and 1700 cm^{-1} . Previous studies have also indicated that alamethicin is stable above its pK_a (5.2–5.8).⁹⁴ Moreover, we performed unpolarized ATR-FTIR measurements and CD experiments on alamethicin in the POPC/POPC bilayer at pH 6.7 and pH 11.9 (Supporting Information), and no substantial secondary structural changes were observed. Together, these findings indicate that no denaturation occurred for alamethicin at pH = 11.9.

Our SFG and ATR-FTIR studies both indicate that increasing the pH alters not only the orientation of alamethicin, but also the number of alamethicin molecules present in the lipid bilayer.

It is interesting to observe that the bend angle between the two helical components in alamethicin ($\theta_1 - \theta_2$) is about 22° at pH = 6.7 and 11.5° at pH = 11.9 when we assume the plane containing both helical components is perpendicular to the membrane surface. These results suggest that an increase in subphase pH changes not only the peptide orientation (θ_1, θ_2), but also the angle of the bend between helical segments ($\phi = \theta_1 - \theta_2$) (as shown in Figure 5). The change in pH can alter the




Figure 5. (A) The definition of tilt angle and bend angle of alamethicin in POPC/POPC bilayer. (B) A schematic to show the pH-dependent channel gating action of alamethicin.

membrane surface charge density and membrane dipole potential. In conjunction with possible lateral pressure effects within the lipid membrane,^{75–79} these changes in the membrane environment can induce alamethicin to stand up more and unbend as the pH is increased. The observed pH-dependent alamethicin action is quite similar to the models proposed by Stockner et al.⁹⁵ to explain the effect of lateral pressure upon conformational transition of an ion channel from the closed to the open state. They built these ion channel models based on bacterial homologues of the nAChR channel with different structural properties:^{96,97} (i) In the asymmetric tilted helix channel model, the orientation angle becomes more narrow in the open state. (ii) In the symmetric bent helix model, the bend angle between the two helical components becomes smaller in the active state. Stockner's simulation results support both models, and a change in lateral pressure led to a more pronounced effect for the bent-helix model.⁹⁵ Our experimental results presented above also support both models.

Generally, the ion channel will not completely open unless the potential difference is larger than 100 mV. The potential

difference used in this study was only ~ -16.2 mV, which was not enough to open the channel completely. However, this potential difference was enough to lead to a detectable change in alamethicin orientation, which will provide molecular information on the voltage dependence of the alamethicin channels formed in membranes.

Although crystal structures are an excellent source of high-resolution structural information, proteins are far more dynamic and can interact quite differently in their native environments (membrane or otherwise) than in the crystal lattice. Although crystal structures reveal little direct information on how these proteins interact with membranes, they are certainly valuable in guiding interpretations. In this study, we successfully applied SFG-VS and ATR-FTIR to study alamethicin in lipid bilayer in situ, using the alamethicin crystal structure as a starting point to interpret our observations. We found that upon membrane potential change, alamethicin changes its membrane orientation. Even though we could not directly relate our observations to interpret gating mechanisms for larger ion channel proteins, in the future, the application of similar experimental approaches on more complex systems, for example, larger ion channel proteins, with the help of newly developed data analysis software,⁹⁸ will provide further insight into different gating mechanisms. Recently, we developed a software package that can read crystal structures from the protein data bank and calculate SFG amide I signals of helical structures in the protein with different polarization combinations as a function of protein orientation (or orientations of various helical structures within the protein). In the literature, it is stated that the models for the modes of voltage sensing under debate are a transporter model, helical screw model, and paddle model. These models were proposed for complex ion channel protein systems, examples of which include the voltage-gated potassium channel protein KvAP, which contains six hydrophobic segments per subunit, S1–S6.^{15–17} The SFG spectral responses due to these three models should be different. In the transporter model, orientation of S4 helix changes; in the helical screw model, none of the helices changes orientation; while in the paddle model, substantial orientation changes of both S3 and S4 helices occur.⁹⁹ When studying large ion channel proteins in the future, we will be able to use the methodology developed in this study, along with the software package mentioned above, to investigate orientation changes of helices in situ upon a change in membrane potential. This will provide direct evidence on which model best interprets the ion channel working mechanisms.

4. CONCLUSION

SFG has developed into a powerful tool to study peptide and protein structures at interfaces.^{100–115} We have applied SFG to characterize interactions between the model ion-channel peptide alamethicin and POPC lipid bilayers, in situ in real time with the presence of a membrane potential and without the need for exogenous labels. The membrane potential was controlled by altering the pH of the bilayer subphase. The orientation of alamethicin in POPC/POPC lipid bilayers was determined by deducing the orientations of the α - and 3_{10} -helical structural segments using polarized SFG amide I spectra and was substantiated by polarized ATR-FTIR studies. We found that at pH = 6.7, the N-terminal α -helix tilts at about 72° versus the surface normal in POPC bilayer (θ_1), and the C-terminal 3_{10} -helix (beyond the Pro14 residue) tilts at about 50° versus the surface normal (θ_2). When pH was increased to 11.9,

the orientation angles θ_1 and θ_2 decreased to 56.5° and 45° , respectively, indicating a change in peptide orientation as well as in the bend between helical segments. The change in membrane potential caused by the change in pH also induced the immediate insertion of more alamethicin molecules into the membrane. Our results from SFG and ATR-FTIR were in good agreement, showing that the orientation distribution of alamethicin in the POPC bilayer is quite narrow. This work provides a basis for future elucidation of gating mechanisms of larger ion channel proteins using SFG in physiologically relevant environments in situ.

■ ASSOCIATED CONTENT

Supporting Information

Details about ATR-FTIR, SFG, fluorescence, and CD experiments, as well as data analysis procedures. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

shujiye@ustc.edu.cn; zhanc@umich.edu

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

S.Y. gratefully acknowledges support from the National Natural Science Foundation of China (21073175, 91127042), National Basic Research Program of China (2010CB923300), and the Fundamental Research Funds for the Central Universities. Z.C. is thankful for support from the U.S. National Institutes of Health (1R01GM081655-01A2).

■ REFERENCES

- (1) Kelkar, D. A.; Chattopadhyay, A. *Biochim. Biophys. Acta* **2007**, *1768*, 2011–2025.
- (2) Cooper, E. C.; Jan, L. Y. *Proc. Natl. Acad. Sci. U.S.A.* **1999**, *96*, 4759–4766.
- (3) Jentsch, T. J.; Hübner, C. A.; Fuhrmann, J. C. *Nat. Cell Biol.* **2004**, *6*, 1039–1047.
- (4) Verkman, A. S.; Galiotta, L. J. V. *Nat. Rev. Drug Discovery* **2009**, *8*, 153–171.
- (5) Berger, A. L.; Ikuma, M.; Welsh, M. J. *Proc. Natl. Acad. Sci. U.S.A.* **2005**, *102*, 455–460.
- (6) Gadsby, D. C.; Vergani, P.; Csanady, L. *Nature* **2006**, *440*, 477–483.
- (7) Tan, H. L.; Bink-Boelkens, M. T. E.; Bezzina, C. R.; Viswanathan, P. C.; Beaufort-Krol, G. C. M.; van Tintelen, P. J.; van den Berg, M. P.; Wilde, A. A. M.; Balsem, J. R. *Nature* **2001**, *409*, 1043–1047.
- (8) Armstrong, N.; Sun, Y.; Chen, G. Q.; Gouaux, E. *Nature* **1998**, *395*, 913–917.
- (9) Stutts, M. J.; Canessa, C. M.; Olsen, J. C.; Hamrick, M.; Cohn, J. A.; Rossier, B. C.; Boucher, R. C. *Science* **1995**, *269*, 847–850.
- (10) Mortimer, J. T.; Bhadra, N. Fundamentals of electrical stimulation. In *Neuromodulation*; Krames, E., Peckham, P. H., Reza, A. R., Eds.; 2009; Vol. 2, p 111.
- (11) Armstrong, C. M.; Hille, B. *Neuron* **1998**, *20*, 371–380.
- (12) Gribkoff, V. K.; Leonard, K. K., Eds. *Structure, Function and Modulation of Neuronal Voltage-Gated Ion Channels*; John Wiley & Sons: New York, 2009.
- (13) Lorincz, A.; Nusser, Z. *Science* **2010**, *328*, 906–909.
- (14) Sasaki, M.; Takagi, M.; Okamura, Y. *Science* **2006**, *312*, 589–592.
- (15) Doyle, D. A.; Cabral, J. M.; Pfuetzner, R. A.; Kuo, A. L.; Gulbis, J. M.; Cohen, S. L.; Chait, B. T.; MacKinnon, R. *Science* **1998**, *280*, 69–77.
- (16) Long, S. B.; Campbell, E. B.; MacKinnon, R. *Science* **2005**, *309*, 897–903.
- (17) Long, S. B.; Tao, X.; Campbell, E. B.; MacKinnon, R. *Nature* **2007**, *450*, 376–382.
- (18) Wu, Y. K.; Yang, Y.; Ye, S.; Jiang, Y. X. *Nature* **2010**, *466*, 393–U148.
- (19) Swartz, K. J. *Nature* **2008**, *456*, 891–897.
- (20) Green, D. R.; Reed, J. C. *Science* **1998**, *281*, 1309–1312.
- (21) Giancotti, F. G.; Ruoslahti, E. *Science* **1999**, *285*, 1028–1032.
- (22) MacKinnon, R. *Science* **2004**, *306*, 1304–1305.
- (23) Sands, Z.; Grottesi, A.; Sansom, M. S. *Curr. Biol.* **2005**, *15*, R44–47.
- (24) Vinothkumar, K. R.; Henderson, R. Q. *Rev. Biophys.* **2010**, *43*, 65–158.
- (25) Sansom, M. S. P. *Prog. Biophys. Mol. Biol.* **1991**, *55*, 139–235.
- (26) Woolley, G. A.; Wallace, B. A. *J. Membr. Biol.* **1992**, *129*, 109–136.
- (27) Sansom, M. S. P. *Eur. Biophys. J.* **1993**, *22*, 105–124.
- (28) Sansom, M. S. P. *Q. Rev. Biophys.* **1993**, *26*, 365–421.
- (29) Cafiso, D. S. *Annu. Rev. Biophys. Biomol. Struct.* **1994**, *23*, 141–165.
- (30) Marsh, D. *Biochem. J.* **1994**, *23*, 345–361.
- (31) Bechinger, B. *J. Membr. Biol.* **1997**, *156*, 197–211.
- (32) Duclouhier, H.; Wroblewski, H. *J. Membr. Biol.* **2001**, *184*, 1–12.
- (33) Nagaraj, R.; Balaram, P. *Acc. Chem. Res.* **1981**, *14*, 356–362.
- (34) Kessel, A.; Cafiso, D. S.; Ben-Tal, N. *Biophys. J.* **2000**, *78*, 571–583.
- (35) Mathew, M. K.; Balaram, P. *Mol. Cell. Biochem.* **1983**, *50*, 47–64.
- (36) Hall, J. E.; Vodyanoy, I.; Balasubramanian, T. M.; Marshall, G. R. *Biophys. J.* **1984**, *45*, 233–247.
- (37) Mathew, M. K.; Balaram, P. *FEBS Lett.* **1983**, *157*, 1–5.
- (38) Leitgeb, B.; Szekeres, A.; Manczinger, L.; Vagvolgyi, C.; Kredics, L. *Chem. Biodiversity* **2007**, *4*, 1027–1051.
- (39) Fox, R. O.; Richards, F. M. *Nature* **1982**, *300*, 325–330.
- (40) North, C. L.; Barranger-Mathys, M.; Cafiso, D. S. *Biophys. J.* **1995**, *69*, 2392–2397.
- (41) Bak, M.; Baywater, R. P.; Hohwy, M.; Thomsen, J. K.; Adelhorst, K.; Jakobsen, H. J.; Sorensen, O. W.; Nielsen, N. C. *Biophys. J.* **2001**, *81*, 1684–1698.
- (42) Salnikov, E. S.; Friedrich, H.; Li, X.; Bertani, P.; Reissmann, S.; Hertweck, C.; O’Neil, J. D. J.; Raap, J.; Bechinger, B. *Biophys. J.* **2009**, *96*, 86–100.
- (43) Duclouhier, H. *Eur. Biophys. J.* **2004**, *33*, 169–174.
- (44) Laver, D. R. *Biophys. J.* **1994**, *66*, 355–359.
- (45) Vodyanoy, I.; Hall, J. E.; Balasubramanian, T. M. *Biophys. J.* **1983**, *42*, 71–82.
- (46) Milov, A. D.; Samoilova, R. I.; Tsvetkov, Y. D.; Zotti, M. D.; Toniolo, C.; Raap, J. *J. Phys. Chem. B* **2008**, *112*, 13469–13472.
- (47) Stella, L.; Burattini, M.; Mazzuca, C.; Palleschi, A.; Venanzi, M.; Coin, I.; Peggion, C.; Toniolo, C.; Pispisa, B. *Chem. Biodiversity* **2007**, *4*, 1299–1312.
- (48) Kessel, A.; Cafiso, D. S.; Ben-Tal, N. *Biophys. J.* **2000**, *78*, 571–583.
- (49) Marsh, D.; Jost, M.; Peggion, C.; Toniolo, C. *Biophys. J.* **2007**, *92*, 473–481.
- (50) Marsh, D.; Jost, M.; Peggion, C.; Toniolo, C. *Biophys. J.* **2007**, *92*, 4002–4011.
- (51) Marsh, D. *Biochemistry* **2009**, *48*, 729–737.
- (52) Salnikov, E. S.; Zotti, M. D.; Formaggio, F.; Li, X.; Toniolo, C.; O’Neil, J. D. J.; Raap, J.; Dzuba, S. A.; Bechinger, B. *J. Phys. Chem. B* **2009**, *113*, 3034–3042.
- (53) Banerjee, U.; Zidovetzki, R.; Birge, R. R.; Chan, S. I. *Biochemistry* **1985**, *24*, 7621–7627.
- (54) Ionov, R.; El-Abed, A.; Angelova, A.; Goldmann, M.; Peretti, P. *Biophys. J.* **2000**, *78*, 3026–3035.
- (55) Mottamal, M.; Lazaridis, T. *Biophys. Chem.* **2006**, *122*, 50–57.
- (56) Chen, F. Y.; Lee, M. T.; Huang, H. W. *Biophys. J.* **2002**, *82*, 908–914.
- (57) Huang, H. W.; Wu, Y. *Biophys. J.* **1991**, *60*, 1079–1087.

- (58) Spaar, A.; Munster, C.; Salditt, T. *Biophys. J.* **2004**, *87*, 396–407.
- (59) Ye, S. J.; Nguyen, K. T.; Chen, Z. *J. Phys. Chem. B* **2010**, *114*, 3334–3340.
- (60) Miranda, P. B.; Shen, Y. R. *J. Phys. Chem. B* **1999**, *103*, 3292–3307.
- (61) Kim, J.; Somorjai, G. A. *J. Am. Chem. Soc.* **2003**, *125*, 3150–3158.
- (62) Kim, J.; Cremer, P. S. *ChemPhysChem* **2001**, *2*, 543–546.
- (63) Li, G.; Ye, S.; Morita, S.; Nishida, T.; Osawa, M. *J. Am. Chem. Soc.* **2004**, *126*, 12198–12199.
- (64) Voges, A. B.; Al-Abadleh, H. A.; Musorriti, M. J.; Bertin, P. A.; Nguyen, S. T.; Geiger, F. M. *J. Phys. Chem. B* **2004**, *108*, 18675–18682.
- (65) Li, Q. F.; Hua, R.; Chea, I. J.; Chou, K. C. *J. Phys. Chem. B* **2008**, *112*, 694–697.
- (66) Ye, H. K.; Gu, Z. Y.; Gracias, D. H. *Langmuir* **2006**, *22*, 1863–1868.
- (67) Yatawara, A. K.; Tiruchinapally, G.; Bordenyuk, A. N.; Andreana, P. R.; Benderskii, A. V. *Langmuir* **2009**, *25*, 1901–1904.
- (68) Liu, J.; Conboy, J. C. *Biophys. J.* **2005**, *89*, 2522–2532.
- (69) Nguyen, K. T.; Le Clair, S.; Ye, S. J.; Chen, Z. *J. Phys. Chem. B* **2009**, *113*, 12169–12180.
- (70) Ye, S. J.; Nguyen, K. T.; Le Clair, S.; Chen, Z. *J. Struct. Biol.* **2009**, *168*, 61–77.
- (71) Chen, X.; Wang, J.; Boughton, A. P.; Kristalyn, C. B.; Chen, Z. *J. Am. Chem. Soc.* **2007**, *129*, 1420–1427.
- (72) Zheng, C.; Vanderkooi, G. *Biophys. J.* **1992**, *63*, 935–941.
- (73) Cladera, J.; O'Shea, P. *Biophys. J.* **1998**, *74*, 2434–2442.
- (74) Luchian, T.; Mereuta, L. *Langmuir* **2006**, *22*, 8452–8457.
- (75) Chiriac, R.; Luchian, T. *Biophys. Chem.* **2007**, *130*, 139–147.
- (76) Zhou, Y.; Raphael, R. M. *Biophys. J.* **2007**, *92*, 2451–2462.
- (77) Capone, S.; Walde, P.; Seebach, D.; Ishikawa, T.; Caputo, R. *Chem. Biodiversity* **2008**, *5*, 16–30.
- (78) Garcia-Manyes, S.; Oncins, G.; Sanz, F. *Electrochim. Acta* **2006**, *51*, 5029–5036.
- (79) Boggara, M. B.; Faraone, A.; Krishnamoorti, R. *J. Phys. Chem. B* **2010**, *114*, 8061–8066.
- (80) Keller, F.; Hanke, W.; Trissl, D.; Bakkergrunwald, T. *Biochim. Biophys. Acta* **1989**, *982*, 89–93.
- (81) Shapiro, H. M. *Methods* **2000**, *21*, 271–279.
- (82) Cohen, L. B.; Salzberg, B. M. *Rev. Physiol. Biochem. Pharmacol.* **1978**, *83*, 35–88.
- (83) Pfligek, J.; Sigler, K. J. *Photochem. Photobiol.*, **B** **1996**, *33*, 101–124.
- (84) Stankowski, S.; Schwarz, U. D.; Schwarz, G. *Biochim. Biophys. Acta* **1988**, *941*, 11–18.
- (85) Haris, P. I.; Chapman, D. *Biochim. Biophys. Acta* **1988**, *943*, 375–380.
- (86) Wang, J.; Paszti, Z.; Clarke, M. L.; Chen, X.; Chen, Z. *J. Phys. Chem. B* **2007**, *111*, 6088–6095.
- (87) Wang, J.; Lee, S.-H.; Chen, Z. *J. Phys. Chem. B* **2008**, *112*, 2281–2290.
- (88) Zhao, X. L.; Ong, S.; Eienthal, K. B. *Chem. Phys. Lett.* **1993**, *202*, 513–520.
- (89) Ong, S.; Zhao, X. L.; Eienthal, K. B. *Chem. Phys. Lett.* **1992**, *191*, 327–335.
- (90) Geiger, F. M. *Annu. Rev. Phys. Chem.* **2009**, *60*, 61–83.
- (91) Chen, X.; Wang, J.; Paszti, Z.; Wang, F.; Schraub, J. N.; Tarabara, V. V.; Schmaier, A. H.; Chen, Z. *Anal. Bioanal. Chem.* **2007**, *388*, 65–72.
- (92) Perry, A.; Neipert, C.; Space, B.; Moore, P. B. *Chem. Rev.* **2006**, *106*, 1234–1258.
- (93) Woolley, G. A.; Wallace, B. A. *Biochemistry* **1993**, *32*, 9819–9825.
- (94) Payne, J. W.; Jakes, R.; Hartley, B. S. *Biochem. J.* **1970**, *117*, 757–766.
- (95) Jerabek, H.; Pabst, G.; Rappolt, M.; Stockner, T. *J. Am. Chem. Soc.* **2010**, *132*, 7990–7997.
- (96) Hilf, R. J. C.; Dutzler, R. *Nature* **2009**, *457*, 115–118.
- (97) Hilf, R. J. C.; Dutzler, R. *Nature* **2008**, *452*, 375–379.
- (98) Boughton, A. P.; Yang, P.; Tesmer, V. M.; Ding, B.; Tesmer, J. J.; Chen, Z. *Proc. Natl. Acad. Sci. U.S.A.* **2011**, *108*, E667–E673.
- (99) Tombola, F.; Pathak, M. M.; Isacoff, E. Y. *Annu. Rev. Cell Dev. Biol.* **2006**, *22*, 23–52.
- (100) Mermut, O.; Phillips, D. C.; York, R. L.; McCrea, K. R.; Ward, R. S.; Somorjai, G. A. *J. Am. Chem. Soc.* **2006**, *128*, 3598–3607.
- (101) Phillips, D. C.; York, R. L.; Mermut, O.; McCrea, K. R.; Ward, R. S.; Somorjai, G. A. *J. Phys. Chem. C* **2007**, *111*, 255–261.
- (102) Jung, S. Y.; Lim, S. M.; Albertorio, F.; Kim, G.; Gurau, M. C.; Yang, R. D.; Holden, M. A.; Cremer, P. S. *J. Am. Chem. Soc.* **2003**, *125*, 12782–12786.
- (103) Kim, G.; Gurau, M.; Kim, J.; Cremer, P. S. *Langmuir* **2002**, *18*, 2807–2811.
- (104) Kim, G.; Gurau, M. C.; Lim, S. -M.; Cremer, P. S. *J. Phys. Chem. B* **2003**, *107*, 1403–1409.
- (105) Evans-Nguyen, K. M.; Fuierer, R. R.; Fitchett, B. D.; Tolles, L. R.; Conboy, J. C.; Schoenfish, M. H. *Langmuir* **2006**, *22*, 5115–5121.
- (106) Weidner, T.; Apte, J. S.; Gamble, L. J.; Castner, D. G. *Langmuir* **2009**, *26*, 3433–3440.
- (107) Weidner, T.; Samuel, N. T.; McCrea, K.; Gamble, L. J.; Ward, R. S.; Castner, D. G. *Biointerphases* **2010**, *5*, 9–16.
- (108) Weidner, T.; Breen, N. F.; Li, K.; Drohny, G. P.; Castner, D. G. *Proc. Natl. Acad. Sci. U.S.A.* **2010**, *107*, 13288–13293.
- (109) Fu, L.; Ma, G.; Yan, E. C. *J. Am. Chem. Soc.* **2010**, *132*, 5405–5412.
- (110) Chen, X.; Wang, J.; Sniadecki, J. J.; Even, M. A.; Chen, Z. *Langmuir* **2005**, *21*, 2662.
- (111) Clarke, M. L.; Wang, J.; Chen, Z. *J. Phys. Chem. B* **2005**, *109*, 22027–22035.
- (112) Chen, X.; Chen, Z. *Biochim. Biophys. Acta* **2006**, *1758*, 1257–1273.
- (113) Doyle, A. W.; Fick, J.; Himmelhaus, M.; Eck, W.; Graziani, I.; Prudovsky, I.; Grunze, M.; Maciag, T.; Neivandt, D. J. *Langmuir* **2004**, *20*, 8961–8965.
- (114) Nguyen, K. T.; Soong, R.; Im, S.; Waskell, L.; Ramamoorthy, A.; Chen, Z. *J. Am. Chem. Soc.* **2010**, *132*, 15112–15115.
- (115) Nguyen, K. T.; King, J. T.; Chen, Z. *J. Phys. Chem. B* **2010**, *114*, 8291–8300.