

Circularly Polarized Luminescence of Curium: A New Characterization of the 5f Actinide Complexes

Ga-Lai Law,^{†,‡} Christopher M. Andolina,^{†,‡} Jide Xu,[‡] Vinh Luu,^{||} Philip X. Rutkowski,[§] Gilles Muller,^{||} David K. Shuh,[§] John K. Gibson,[§] and Kenneth N. Raymond^{*,‡,§}


[†]Department of Chemistry, University of California, Berkeley, California 94720-1460, United States

[§]Chemical Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, United States

^{||}Department of Chemistry, San José State University, One Washington Square, San José, California, 95192-0101

S Supporting Information

ABSTRACT: A key distinction between the lanthanide (4f) and the actinide (5f) transition elements is the increased role of f-orbital covalent bonding in the latter. Circularly polarized luminescence (CPL) is an uncommon but powerful spectroscopy which probes the electronic structure of chiral, luminescent complexes or molecules. While there are many examples of CPL spectra for the lanthanides, this report is the first for an actinide. Two chiral, octadentate chelating ligands based on orthoamide phenol (IAM) were used to complex curium(III). While the radioactivity kept the amount of material limited to micromole amounts, spectra of the highly luminescent complexes showed significant emission peak shifts between the different complexes, consistent with ligand field effects previously observed in luminescence spectra.



■ INTRODUCTION

Due to their radioactivity, consequent rarity, and difficulty of handling, the actinides (particularly those beyond uranium in the periodic table) have electronic levels which are less understood than other d or f transition metals. Although the optical properties of Cm(III) were the subject of an excellent recent review,¹ less is known at the molecular level concerning the electronic and structural properties of actinides in excited states,^{1–4} in contrast to the 4f lanthanides. The latter are therefore often used as surrogates for the actinides. The significant electronic and chemical differences between the 4f and the 5f elements require direct investigation to differentiate.^{5–8}

Recent advances in computational modeling for these relativistic and many-electron elements have rekindled the interest of experimentalists in the molecular chemistry and physics of the actinides.^{5,9–11} This effort is significant, since most of the prior actinide photoluminescence studies have been in the solid state and based on direct excitation of the metal center. While antennae sensitization has been commonly used for indirect enhancement of the luminescence of lanthanides (even in commercial use), there are very few reports of the sensitization of actinides. In this study a new window into the electronic structure of actinides is opened through the use of circularly polarized luminescence (CPL) spectroscopy.^{12,13} Compared to standard photoluminescence spectroscopy, this spectroscopic technique is rarely reported.^{14–18} The only example of CPL from an actinide complex is a study of the charge transfer band of uranyl complexes, which is not an f–f transition.^{19,20}

CPL is the emission analog of circular dichroism (CD). Similar to CD, CPL can also discriminate between structurally similar luminescent chiral complexes/molecules.^{21–25} In this work, octadentate chiral phenolate ligands, H(2,2)BnMe IAMS(–)/R(+) chiral ligands (both enantiomers denoted as L1), as well as a newly synthesized chiral cage, EtH(2,2)-BIAM (L2), were used to form Cm(III) complexes (Scheme 1).^{24,26} Ligands L1 and L2 were selected based on the photophysical properties of the L1 terbium(III) complexes.²⁴ The sensitizer/chelator group 2-hydroxyisophthalamide (IAM) was thought likely to provide a favorable pairing of electronic states for sensitization of Cm(III) (Scheme 1).^{27,28} As previously shown for the lanthanides, L1 and L2 are both strongly coordinating and strongly sensitizing ligands. Given the relative energy levels (see Figure S1, Supporting Information) we anticipated efficient luminescence, a large quantum yield, and perhaps strong CPL response for the Cm(III) complexes. Although Am lies below Eu and so might be considered the most promising 5f luminescent ion, relativistic effects and spin–orbit coupling³ make Cm (which lies below Gd) the best choice, as shown in Scheme 1.


■ EXPERIMENTAL SECTION

Synthesis of L1 has been previously reported,²⁴ and L2 preparation will be reported in another pending manuscript. The integrity of these ligands was verified by elemental analysis, proton NMR, and ESI/MS, and for L1, ligand characterization agreed with reported values.²⁴

Received: July 11, 2012

Published: August 25, 2012

Scheme 1. (Top) Structures of the Chiral Ligands Used in This Study;^a (Bottom) Cartoon of CmL1 Structure (left) and Partial Energy Diagram Generally Describing the Energy Transfer from the Chromophore (IAM) to Cm(III) and Tb(III) (right)



^aAsterisk (*) indicates the chiral center for L1 and L2.

Complexes were prepared in situ by dissolving the appropriate amount of the ligand L1 (H₄R (+) BnMeH(2,2)IAM or L2, H₄EtBH(2,2)IAM) to yield either a 10 or a 200 μM solution in HPLC-grade methanol dried with molecular sieves. Curium(III) solutions were prepared in dry methanol at the same concentrations as the ligand solution (10 or 200 μM) from the Cm(III) acidified (HCl) stock. Complexes were prepared by mixing 1.00 mL of the ligand solution with 1.00 mL of the Cm(III) methanol solution in a 1:1 stoichiometric ratio and thoroughly mixed by gentle pipetting. A base, 5 μL of dry pyridine, was added in excess to ensure deprotonation of the IAM moieties. Samples were incubated for at least 48 h to allow for equilibration. As the complexes were formed in situ, characterization of the complexes was done by ESI/MS. For the photoluminescence measurements, the stock solution was diluted to approximately 5, 4, 3, 2, and 1 μM in methanolic solution.

ESI/MS. All ESI/MS experiments were performed using an Agilent 6340 QIT-MS in the Heavy Elements Research Laboratory (HERL) at the Lawrence Berkeley National Laboratory (LBNL). The instrument has a detection range of 50–2200 *m/z* with a resolution of ~0.25 *m/z*. Mass spectra were recorded using the negative-ion mode. Solutions were injected into an electrospray needle via a syringe pump and nebulized using nitrogen gas upon exiting the needle, resulting in creation of small droplets and eventually gaseous ions. The gas-phase ions entered a charged capillary, where they were further desolvated by dry heated nitrogen from a liquid nitrogen dewar, which also supplied the nebulizing gas. The ions then passed through an ion-focusing skimmer and into two focusing octopoles for transfer into the ion trap. Mass spectra were acquired using the following instrumental parameters: Solution flow rate, 60 μL/h; nebulizer gas pressure, 15 psi; dry gas flow rate, 5 L/min; dry gas temperature, 325 °C; capillary voltage, 4500 V; capillary exit voltage, -147.3 V; skimmer voltage, -40.0 V; octopole 1 and 2 dc voltage, -12.00 and -3.67 V; octopole rf amplitude, 200.0 Vpp; lens 1 and 2 voltages, 5.0 and 60.0 V; trap drive, 150.5.

CmL2 [Cm(III)(EtBH(2,2)IAM)]⁻: ESI-MS negative mode (M⁻) *m/z* expected 1405.32 (found 1405.0).

CmL1R [Cm(III)R(+)/BnMeH22IAM]⁻: ESI-MS negative mode (M⁻) *m/z* expected 1544.46 (found 1544.5).

CmL1S [Cm(III)S(-)/BnMeH22IAM]⁻: ESI-MS negative mode (M⁻) *m/z* expected 1544.46 (found 1544.5).

Photophysics. UV-vis absorbance measurements were collected using the Cary 6000i UV-vis in the HERL. Solution spectra of the Cm(III) complexes were recorded at 25.0 °C. Emission spectra were acquired on a HORIBA Jobin Yvon IBH FluoroLog-3 spectrofluorometer equipped with a 450 W xenon lamp and a temperature controller operating at 25.0 °C. This instrument is described in detail elsewhere.^{21–26} Quantum yields were determined by the optically dilute method using eq 1.

$$\frac{\Phi_x}{\Phi_r} = \left[\frac{A_r(\lambda_r)}{A_x(\lambda_x)} \right] \left[\frac{I(\lambda_r)}{I(\lambda_x)} \right] \left[\frac{n_x^2}{n_r^2} \right] \left[\frac{D_x}{D_r} \right] \quad (1)$$

A quantum yield standard, quinine sulfate, in 0.1 M H₂SO₄ (Φ_r = 0.577) was used as a reference and prepared according to the literature.^{9,27} Absorbance at the excitation wavelength was varied from approximately 0.02 to 0.13 for both the curium complexes and the quinine sulfate reference. A total of 5 CmL samples and quinine sulfate reference solutions were prepared for each quantum yield determination at different absorbances within the previously given range. The following refractive indices were used for the reference and the sample *n*_{H₂O} = 1.333 and *n*_{MEOH} = 1.329, respectively. Quantum yields are reported as the average of three independent measurements unless stated otherwise. Samples and references were excited at 350 nm with a 3 nm band pass. Emission was collected in 1.0 or 0.1 nm increments, 0.2 s integration times, and with a 1.0 nm band pass.

For time-resolved luminescent lifetime measurements, the sub-microsecond xenon flash lamp (Jobin Yvon, S000XeF) unit of the Fluorolog-3 was used as the light source also described in detail elsewhere.^{21–26} Samples in methanol were excited at 350 nm; the emission slit was set to the maximum intensity determined from the emission spectra for each sample. Excitation and emission slits were adjusted to prevent saturation of the detector electronics utilizing ratiometers included in the time-resolved luminescence software

package. The goodness of fit was assessed by minimizing the reduced chi squared function, χ^2 , and a visual inspection of the weighted residuals. Each trace contained at least 10 000 points, and the reported lifetime values result from at least three independent measurements.

Circularly polarized luminescence and total luminescence spectra were recorded on an instrument described previously,^{21–25} operating in a differential photon-counting mode. The light source for indirect excitation was a continuous wave 1000 W xenon arc lamp from a Spex FluoroLog-2 spectrofluorometer, equipped with excitation and emission monochromators with dispersions of 4 nm/mm (SPEX, 1681B). The optical detection system consisted of a focusing lens, long-pass filter, and 0.22 m monochromator. Emitted light was detected by a cooled EMI-9558B photomultiplier tube operating in photon-counting mode. All measurements were performed in quartz cuvettes with a path length of 0.4 or 1.0 cm.

RESULTS AND DISCUSSION

The two ligands (**L1** and **L2**) differ by the presence of an additional capping moiety and incorporation of the chiral centers. The open-ended octadentate ligands, **L1R** and **L1S** (collectively **L1**, see Scheme 1), were synthesized in enantiopure forms.²¹ These molecules are chiral due to insertion of chiral centers on the opposite end of the IAM chelator to the capping group referred to as “H(2,2)”. The **L2** macrocyclic ligand is a derivative of **L1** and is more synthetically challenging. It is a macrocyclic octadentate ligand in which the four chelating chromophores are tethered together at both ends, and it was designed to enhance kinetic and thermodynamic stability. Rigidity generally improves the photoluminescent properties such as quantum yield and excited state lifetime (relative to **L1**), however, not necessarily the CPL response. Since the **L2** ligand is capped at either end of the IAM chromophore, chirality is introduced into the molecule by addition of ethyl groups on one of the capping scaffolds H(2,2) (Scheme 1).

The CPL spectra of **CmL1R/S** and **CmL2** are shown in Figures 1 and 2, respectively. The CPL activity of the Cm(III) complexes in methanol solutions was determined using a custom-built instrument that measures left and right CPL and is quantified as g_{lum} , the luminescence dissymmetry factor (eq 2).^{21–24} Here, I_L is the intensity of left CPL and I_R is the intensity of right CPL.

$$g_{\text{lum}} = \frac{\Delta I}{I} = \frac{I_L - I_R}{\frac{1}{2}(I_L + I_R)} \quad (2)$$




Figure 1. CPL spectra of **CmL1** (S[−]) (top) and **CmL1** (R⁺) (bottom), $\lambda_{\text{EX}} = 350$ nm at 25.0 °C in MeOH.




Figure 2. CPL spectra of the **CmL2** (top) and emission spectra (bottom) ($\lambda_{\text{EX}} = 350$ nm at 25.0 °C in MeOH), 4 μM concentration.

The CPL response is typically of greater magnitude at transitions corresponding to magnetic dipole rather than electric dipole allowed transitions.^{12,13} In this case Cm(III) emission is observed at $\sim 611\text{--}3$ nm for the complexes. g_{lum} values for the **R** and **S** isomers were -0.050 and $+0.048$ at an emission wavelength of 611 nm, respectively (Table 1). Since the complexes are enantiomers these spectra are mirror images (Figure 1).

Table 1. CPL Properties of Cm(III) Chiral Complexes (4 μM in MeOH at 25.0 °C)

ligand	g_{lum}	λ_{EM} (nm)
L1R	-0.050	611
L1S	$+0.048$	611
L2	$+0.0018$	600
	-0.0014	609

Comparison of the CPL of **CmL1** and **CmL2** shows that the more rigid **CmL2** has a smaller g_{lum} than the **CmL1** complex. The CPL spectrum of **CmL2** is characterized by a negative g_{lum} value of -0.0014 (a greater degree of right CPL) at an emission wavelength of ~ 609 nm (Table 1) with another peak seen at 600 nm with a g_{lum} value of $+0.0018$. In comparison to the lanthanides, these values for the Cm(III) complexes are not as large as the g_{lum} values for **[EuL1R/S]**, previously reported with values of $+0.298$ and -0.294 (at an emission wavelength of 596 nm),²⁴ or with any other lanthanide complexes we reported.^{22–25} g_{lum} values for the **CmL1R/S** complexes were similar to the Tb(III) values determined with the same ligand **L1R/S** ($+0.044$ and -0.048 , respectively). It is important to note that although Tb(III) and Cm(III) have some similarities with respect to excited state energies, it is unlikely that these similarities result in similar g_{lum} , since other parameters (including coordination geometry and inner-sphere solvent coordination as well as vibronic relaxations) play an influential role. In the case of **L2**, capping of the ligand has enhanced the photoluminescent properties through the increased rigidity of the ligand.

We suggest that the energy transfer mechanism to Cm(III) occurs via the “antennae effect”; absorbed light excites the IAM

ligand to a singlet, which decays to a triplet state followed by excitation of the accepting state of the metal ion. The lowest triplet state of the IAM chromophore has been previously determined as $23\,300\text{ cm}^{-1}$, as shown in the Jablonski diagram (Scheme 1).^{28,29} The intramolecular energy transfer from the triplet state of the chromophore is believed to proceed to a high-energy Cm(III) excited state (above the lowest lying excited state), which then cascades to the lowest emitting state ($^5D'_{7/2}$) before relaxing to the ground state ($^8S'_{7/2}$) by radiative (photon emission) and nonradiative (such as vibronic coupling) relaxation processes (Scheme 1).

Photophysical data of the Cm(III) complexes are reported in Table 2. All measurements were performed in methanolic

Table 2. Photophysical Properties of the Cm(III) Chiral Complexes in MeOH at 25.0 °C

complex	ϵ ($M^{-1}\text{ cm}^{-1}$)	λ_{EM} (nm)	Φ (%)	τ_{MeOH} (μs)	q_{MeOH}
CmL1R	25 600	609	0.38	471	0.3, ^a 0.2 ^b
CmL1S	25 100	609	0.35	502	0.3, ^a 0.2 ^b
CmL2	19 300	606	0.50	353	0.5, ^a 0.3 ^b

^aCalculated with eq 2a. ^bCalculated using eq 2b.

solutions at 25.0 °C (Supporting Information). Sensitization of Cm(III) by IAM is surprisingly efficient. This is essential since such low concentrations of the complexes necessarily were used for the photoluminescent (and CPL) measurements ($4\ \mu\text{M}$). The UV absorption maximum is observed at 335–345 nm for both complexes, with a large molar absorptivity from the π – π^* transitions. This is characteristic of the IAM chromophore, as seen in our previous lanthanide studies.^{26,28,30} The primary emission band, observed at 610 nm, is assigned to an f – f transition ($^5D'_{7/2} \rightarrow ^8S'_{7/2}$) of Cm(III). The emission spectra and photophysical properties for the enantiomers (CmL1 R/S) are nearly identical, as expected (Figure 3). The photoluminescence emission spectra comparing the two different ligands are shown in Figure 3.

The coordination differences of the ligands (L1 and L2) are more apparent upon comparison of the photophysical properties of the complexes (Table 2 and Figure 3). CmL2, the bicapped macrocycle, has enhanced photoluminescent properties relative to CmL1, improved quantum yield efficiency, and




Figure 3. Normalized emission spectra of CmL2, CmL1R, and CmL1S in methanol $\lambda_{EX} = 350 \pm 4\text{ nm}$ at $5\ \mu\text{M}$ and 25.0 °C.

brightness. The quantum yield of the CmL2 complex is significantly larger ($\Phi = 0.50$ versus 0.38–0.35) when compared to the CmL1 isomers. The values of the molar absorptivity ($19\,300\text{ M}^{-1}\text{ cm}^{-1}$) and observed lifetime ($353\ \mu\text{s}$) are slightly smaller and shorter for CmL2 compared to CmL1, respectively. We suggest that the increased rigidity, stability, and reduction of external quenching by solvent molecules, due to more effective encapsulation of the CmL2 complex, accounts for the enhancement of these photophysical properties. Solvation numbers, q_{MeOH} , of the complexes were estimated using a modified Cm(III) equation originally developed by Kimura³¹ and then by Tian³² (eqs 2a and 2b, respectively).³³ To obtain q_{MeOH} , the numerical values of both original equations were taken as 1/2 to approximate the quenching of a methanol molecule since one methanol contains a single O–H oscillator whereas water is composed of two oscillators.³³ However, it is important to note that these two equations fail to account for many nonradiative decay processes such as back energy transfer to a chromophore or quenching by other nearby oscillators.¹

$$q_{\text{H}_2\text{O}} = 0.65xk_{\text{obs}} - 0.88 \quad (2a)$$

$$q_{\text{H}_2\text{O}} = 0.63xk_{\text{obs}} - 0.29 \quad (2b)$$

The q_{MeOH} number determined for the CmL1 complexes is estimated to be 0.2–0.3 and 0.5–0.3 for CmL2, although other quenching processes may be occurring and unaccounted for in both systems. Alternatively, partial dissociation of the chelator allowing for a small fraction of CmL species with an inner-sphere water cannot be definitively ruled out and may influence a nonzero q_{MeOH} number. However, the large quantum yields and monoexponential time-resolved photoluminescent data together are suggestive of a single species present in solution with no solvent molecules directly coordinated to either complex.

Inspection of the photoluminescence emission spectra from the two different octadentate ligands (L1 and L2) shows the characteristic Cm(III) emission band, which is observed between 590 and 614 nm (Figure 3). The smaller bands observed in the emission spectra (570 and 590 nm) are likely a result of the crystal field splitting of the lowest lying excited state ($^5D'_{7/2}$).³ The time-resolved photoluminescence lifetime decays were fitted to a monoexponential decay function, consistent with one species in solution for all Cm(III) samples discussed here (Table 2). In addition, only one major peak for the predicted Cm(III) complexes was observed in the ESI/MS spectra (Figure S2, Supporting Information).

The bathochromic shift (red shift) of the emission wavelength maxima between CmL1 and CmL2 has been discussed for a range of Cm(III) complexes.³ Comparison of either complex to directly excited methanolic solutions of Cm(III) ($\lambda_{EX} = 396\text{ nm}$) with spectra peak maxima at 600 nm shows a larger red shift. This red shift of 8–10 nm is much larger than those observed for Ln(III) complexes and free cations (0–3 nm).^{34,35} For comparison, aqueous Cm(III) has reported emission maxima at 593.8 nm.^{1,32,36} The nephelauxetic effect seen in the luminescence spectra of a range has been previously explained (see Table 4 of ref 1).

CONCLUSIONS

We report the first actinide CPL spectra of an f – f transition. Highly luminescent Cm(III) complexes of antenna ligands

show significant emission peak shifts between the two complexes studied. The apparent ligand field effects are consistent with the significant differences in electronic structure between 5f Cm(III) and 4f analogs.

■ ASSOCIATED CONTENT

📄 Supporting Information

Energy level diagram depicting luminescent actinides in relation to the IAM chromophore and ESI/MS spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

E-mail raymond@socrates.berkeley.edu

Author Contributions

[†]These authors contributed equally to this manuscript.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Some of this research was supported by NIH grant HL069832. Work at LBNL was supported by the Director, Office of Science, Office of Basic Energy Sciences, and the Division of Chemical Sciences, Geosciences, and Biosciences of the U.S. Department of Energy at LBNL under Contract No. DE-AC02-05CH11231. G.M. thanks the National Institutes of Health, Minority Biomedical Research support (1 SC3 GM089589-03), and the Henry Dreyfus Teacher-Scholar Award for financial support. G.M. also thanks Mr. Victor Maraschin from the San José State University Nuclear Science facility for his technical help in handling the samples.

■ REFERENCES

- (1) Edelstein, N. M.; Klenze, R.; Fanghänel, T.; Hubert, S. *Coord. Chem. Rev.* **2006**, *250*, 948.
- (2) Billard, I.; Geipel, G. *Springer Ser. Fluoresc.* **2008**, *5*, 465.
- (3) Carnall, W. T. *J. Phys. Chem.* **1992**, *96*, 8713.
- (4) Gorden, A. E. V.; Xu, J.; Szigethy, G.; Oliver, A.; Shuh, D. K.; Raymond, K. N. *J. Am. Chem. Soc.* **2007**, *129*, 6674.
- (5) Polly, R.; Schimmelpfennig, B.; Rabung, T.; Flörsheimer, M.; Klenze, R.; Geckeis, H. *Radiochim. Acta* **2010**, *98*, 627.
- (6) Elgazzar, S.; Ruzs, J.; Oppeneer, P. M.; Colineau, E.; Griveau, J. C.; Magnani, N.; Rebizant, J.; Caciuffo, R. *Phys. Rev. B: Condens. Matter* **2010**, *81*, 235117.
- (7) Söderlind, P.; Gonis, A. *Phys. Rev. B: Condens. Matter* **2010**, *82*, 033102.
- (8) Odoh, S. O.; Schreckenbach, G. *J. Phys. Chem. A* **2010**, *114*, 1957.
- (9) Marie, C.; Miguiditchian, M.; Guillaumont, D.; Tosseng, A.; Berthon, C.; Guilbaud, P.; Duvail, M.; Bisson, J.; Guillauneux, D.; Pipelier, M.; Dubreuil, D. *Inorg. Chem.* **2011**, *50*, 6557.
- (10) Pérez-Villa, A.; David, J.; Fuentealba, P.; Restrepo, A. *Chem. Phys. Lett.* **2011**, *507*, 57.
- (11) Tecmer, P.; Gomes, A. S. P.; Ekström, U.; Visscher, L. *Phys. Chem. Chem. Phys.* **2011**, *13*, 6249.
- (12) Martin, B.; Richardson, F. S. *Q. Rev. Biophys.* **1979**, *12*, 181.
- (13) Riehl, J. P.; Richardson, F. S. *Chem. Rev.* **1986**, *86*, 1.
- (14) Panak, P.; Klenze, R.; Kim, J. I.; Wimmer, H. *J. Alloys Compd.* **1995**, *225*, 261.
- (15) Klenze, R.; Panak, P.; Kim, J. I. *J. Alloys Compd.* **1998**, *271–273*, 746.
- (16) Nugent, L. J.; Burnett, J. L.; Baybarz, R. D.; Werner, G. K.; Tanner, S. P.; Tarrant, J. R.; O. L. Keller, J. *J. Phys. Chem.* **1969**, *73*, 1540.
- (17) Dem'yanova, T. A.; Stepanov, A. V.; Babaev, A. S.; Aleksandruk, V. M. *Radiokhimiya* **1986**, *28*, 494.

- (18) Yusov, A. B. *J. Radioanal. Nucl. Chem.* **1990**, *143*, 287.
- (19) Samoilov, B. N. *J. Exp. Theor. Phys.* **1948**, *18*, 1030.
- (20) Murata, K.; Yamazaki, Y.; Morita, M. *J. Lumin.* **1979**, *18–19* (Part 1), 407.
- (21) Seitz, M.; Moore, E. G.; Ingram, A. J.; Muller, G.; Raymond, K. N. *J. Am. Chem. Soc.* **2007**, *129*, 15468.
- (22) Seitz, M.; Do, K.; Ingram, A. J.; Moore, E. G.; Muller, G.; Raymond, K. N. *Inorg. Chem.* **2009**, *48*, 8469.
- (23) Samuel, A. P. S.; Lunkley, J. L.; Muller, G.; Raymond, K. N. *Eur. J. Inorg. Chem.* **2010**, *2010*, 3343.
- (24) Petoud, S.; Muller, G.; Moore, E. G.; Xu, J.; Sokolnicki, J.; Riehl, J. P.; Le, U. N.; Cohen, S. M.; Raymond, K. N. *J. Am. Chem. Soc.* **2007**, *129*, 77.
- (25) D'Aleo, A.; Xu, J.; Do, K.; Muller, G.; Raymond, K. N. *Helv. Chim. Acta* **2009**, *92*, 2439.
- (26) Xu, J.; Corneillie, T. M.; Moore, E. G.; Law, G.-L.; Butlin, N. G.; Raymond, K. N. *J. Am. Chem. Soc.* **2011**, *133*, 19900.
- (27) Eastman, J. W. *Photochem. Photobiol.* **1967**, *6*, 55.
- (28) Samuel, A. P. S.; Xu, J.; Raymond, K. N. *Inorg. Chem.* **2008**, *48*, 687.
- (29) Moore, E. G.; Samuel, A. P. S.; Raymond, K. N. *Acc. Chem. Res.* **2009**, *42*, 542.
- (30) Petoud, S.; Cohen, S. M.; Bünzli, J.-C. G.; Raymond, K. N. *J. Am. Chem. Soc.* **2003**, *125*, 13324.
- (31) Kimura, T.; Nagaishi, R.; Kato, Y.; Yoshida, Z. *Radiochim. Acta* **2001**, *89*, 125.
- (32) Tian, G.; Edelstein, N. M.; Rao, L. *J. Phys. Chem. A* **2011**, *115*, 1933.
- (33) Horrocks, W. D., Jr.; Sudnick, D. R. *Science* **1979**, *206*, 1194.
- (34) Andolina, C. M.; Mathews, R. A.; Morrow, J. R. *Helv. Chim. Acta* **2009**, *92*, 2330.
- (35) Andolina, C. M.; Holthoff, W. G.; Page, P. M.; Mathews, R. A.; Morrow, J. R.; Bright, F. V. *Appl. Spectrosc.* **2009**, *63*, 483.
- (36) Rao, L.; Tian, G. *Dalton Trans.* **2011**, *40*, 914.