

Correction to Palladium-Catalyzed Cross-Coupling of Aryl Chlorides and Triflates with Sodium Cyanate: A Practical Synthesis of Unsymmetrical Ureas

Ekaterina V. Vinogradova, Brett P. Fors, and Stephen L. Buchwald*

J. Am. Chem. Soc. **2012**, *134*, 11132–11135. DOI: 10.1021/ja305212v

Page 11134. Reaction conditions in footnotes corresponding to Tables 2 and 3 were switched. PhOH (2 mmol) should be present in the reaction conditions used for Table 3. The conditions in the footnotes should read:

Table 2: Step 1: ArX (1 mmol), NaOCN (2 mmol), Pd₂(dba)₃ (0.5 mol %), L1 (1.2 mol %), NEt₃ (25 mol %), toluene (2 mL). Pd₂(dba)₃ and L1 were preheated in toluene (2 mL) at 120 °C for 3 min. Step 2: ArNH₂ (1.2 mmol), toluene (2 mL).

Table 3: Step 1: ArX (1 mmol), NaOCN (2 mmol), PhOH (2 mmol), Pd₂(dba)₃ (0.5 mol %), L1 (1.2 mol %), NEt₃ (25 mol %), toluene (2 mL). Pd₂(dba)₃ and L1 were preheated in toluene (2 mL) at 120 °C for 3 min. Step 2: ArNH₂ (1.2 mmol), toluene (2 mL).