

Estimation of Free Gossypol in Cottonseed Meal and Cottonseed Meats: Modified Method

AN EARLIER MODIFICATION (3) of a spectrophotometric method (2) for the determination of free gossypol, as the dianilino derivative, in cottonseed meal and meats has been simplified. Ethanol, water and ethyl ether are combined into a single solvent mixture for the extraction of the gossypol in a homogenizer which is cooled by immersion in a water bath. The modified method has the precision and accuracy of the original.

Reagents are as follows: (a) Solvent mixture: 715 ml 95% ethanol, 285 ml of distilled water, 200 ml ethyl ether (peroxide free) and 0.2 ml glacial acetic acid. Ether is tested as follows: a mixture of 0.1 g of vanadic oxide and 2 ml concentrated sulfuric acid is heated on a steam bath for 15 min, cooled and diluted to 50 ml with water (1). A pink color is obtained when 2 ml of reagent is shaken with 10 ml of ether containing peroxides. (b) Aniline: Freshly distilled from approximately 1 g of 30 mesh granular zinc. The distillate should be colorless. (c) Acid washed Hyflo Super-Cel: Prepared as previously described (3,4).

The procedure is as follows: Add 60 ml of the solvent mixture to either 1 g of finely ground cottonseed meal or 0.250 g of cottonseed meats placed in a 200 ml chamber of a Sorvall omni-mixer or similar comminution apparatus. Homogenize for 5 min after

surrounding the mixing chamber with water at 27–30 C. Filter the homogenate under vacuum through a filter tube (Corning 9480) previously prepared by inserting a porcelain plate over which is formed a layer of asbestos followed by a 1/8 in. mat of Hyflo Super-Cel. The filtrate and washings are received in a 100 ml volumetric flask containing 5 ml of ether, placed under a bell jar. Wash the equipment and residue with the ethanolic solvent mixture dispensed from a fine-tipped wash bottle. Dilute the extract to 100 ml, mix and transfer 5 ml aliquots to two 25 ml volumetric flasks. Dilute one of the aliquots to 25 ml with the solvent mixture, mix and reserve as the reference solution. Add 0.5 ml of freshly distilled aniline to the other aliquot and heat gently either on a steam bath or in a water bath at 70–75 C for 40 min. After cooling dilute the sample to 25 ml with the ethanol-water-ether solvent mixture, mix and determine the absorbance at 445 m μ using the aliquot without aniline as the reference.

The gossypol content of the samples may be calculated from the extinction coefficient or determined graphically from the absorbance-concentration curve, as previously described (4,5).

Typical results for two samples of cottonseed meal by the present and the earlier modifications are shown in Table I and are in good agreement.

F. H. SMITH
Nutritional Biochemistry Section
Animal Science Department
North Carolina State University
Raleigh, North Carolina 27607

REFERENCES

1. Baskerville, C., and W. A. Hamor, *Ind. Eng. Chem. Anal. Ed.* **3**, 387 (1911).
2. Lyman, C. M., B. R. Holland and F. Hale, *Ibid.* **15**, 489–491 (1943).
3. Smith, F. H., *Ibid.* **18**, 43–45 (1946).
4. Smith, F. H., *JAOCS* **42**, 145–147 (1965).
5. Smith, F. H., *Ibid.* **44**, 267–269 (1967).

[Received July 8, 1968]

TABLE I

Free Gossypol Values for Cottonseed Meals by Modified Methods^a

Sample	Number	Determination	Modification	
			Present %	Earlier %
1		a	0.328	0.320
		b	0.334	0.328
		c	0.324	0.325
		d	0.320	0.321
		Average	0.327	0.324
2		a	0.026	0.027
		b	0.029	0.027
		c	0.029	0.029
		Average	0.028	0.028

^a The present modification is compared with an earlier one (3).

Identification of Arachidic Acid in Porcine Glycerides

LITERATURE REPORTS INDICATE that porcine glycerides contain about 1% linolenic acid, but there is little support for the presence of arachidic acid in such amounts. Magidman et al. (1) reported that lard contains 0.5% linolenic acid and 0.3–0.4% arachidic acid. Gas-liquid chromatography of the methyl esters of porcine glycerides, using an F & M Model 720 dual column gas chromatograph with thermal conductivity detectors and 10 ft \times 1/4 in.

OD stainless steel columns packed with 20% diethylene glycol succinate on 60/80 mesh firebrick, shows a peak with a retention time between methyl linoleate and methyl eicosanoate (Table I), which could represent the methyl ester of either linolenic acid or arachidic acid or both. Chromatographic conditions were as follows: column temperature, 230 C; injection temperature, 275 C; detector temperature, 240 C; and flow rate of helium gas, 50 ml/min.

Standard samples of methyl linolenate and methyl eicosanoate, when chromatographed under the above conditions revealed the same retention time as the unknown peak. Semilogarithmic plots of relative

¹ Present address: School of Household Economics, University of Alberta, Edmonton, Alberta.

² Present address: Department of Food Science, University of Guelph, Guelph, Ontario, Canada.