

expansion of the structural information normally obtainable through NMR spectroscopy of lipid derivatives. Further details, including the scope and limitations of the use of csr, will be presented at the American Oil Chemists' Society Meeting in Atlantic City, October 1971, and in papers to be submitted later to the Journal.

DANIEL SWERN
JOHN P. WINEBURG
Fels Research Institute
Department of Chemistry
Temple University
Philadelphia, Pennsylvania 19122

ACKNOWLEDGMENT

Work supported in part by the National Cancer Institute of the

U.S. Public Health Service.

REFERENCES

1. Hopkins, C.Y., in "Nuclear Magnetic Resonance of Fatty Acids and Glycerides," Vol. 8, Edited by R.T. Holman, Pergamon Press, New York, 1966.
2. Sanders, J.K.M., and D.H. Williams, Chem. Commun. 1970:422.
3. Sanders, J.K.M., and D.H. Williams, J. Amer. Chem. Soc. 93:641 (1971).
4. Briggs, J., G.H. Frost, F.A. Hart, G.P. Moss and M.L. Staniforth, Chem. Commun. 1970:749.
5. DeMarco, P.V., T.K. Elzey, R.P. Burton and E. Wenkert, J. Amer. Chem. Soc. 92:5734 (1970).
6. Rondeau, R.E., and R.E. Sievers, Ibid. 93:1522 (1971).

[Received May 7, 1971]

ERRATUM

JAOCS 48:140-142 (1971), S.P. Fore, E.T. Rayner and H.P. Dupuy: "Determination of Residual Solvent in Oilseed Meals and Flours: III. Isopropanol." On page 142, Reference 1 should read: Dupuy, H.P., and S.P. Fore, JAOCS 47:17 (1970).
