

List of Physico-chemical Symbols adopted by the Chemical Society.
 [See J.C.S., 1921, 119, 502—512.]

1. Mathematical Symbols.

| | Usual symbol. | Alternative symbol. |
|--|---------------|---------------------|
| Base of natural (Napierian) logarithms | e | |
| Diameter | d | |
| Radius | r | |
| Ratio of circumference to diameter | π | |
| Summation | Σ | |
| Variation | δ | |
| Total differential | dx | |
| Partial differential | ∂x | |

2. Universal Constants.

| | | |
|---|-----|--|
| Acceleration due to gravity | g | |
| Mechanical equivalent of heat | J | |
| Avogadro's constant [number of molecules in 1 gram-molecule (mole)] | N | |
| Gas constant per mole | R | |
| Faraday's constant (number of coulombs per gram-equivalent of an ion) | F | |
| Charge on an electron | e | |

3. General Physics and Chemistry.

| | | |
|--|--|-----|
| Length | l | |
| Height | h | |
| Mass | m | |
| Time | t | |
| Volume | v, V | |
| Density (mass per unit volume) | d | |
| Pressure | p, P | |
| Concentration | c, C | |
| Mole fraction | x | |
| Critical constants : pressure, volume, temperature (centigrade), temperature (absolute), density | $\left\{ \begin{array}{l} p_c, v_c \\ t_c, T_c \\ d_c \end{array} \right.$ | D |
| Reduced quantities : pressure, volume, temperature, density | $\left\{ \begin{array}{l} p_r, v_r \\ t_r, T_r, d_r \end{array} \right.$ | |
| van der Waals's constants | a, b | |
| Fluidity | ϕ | |
| Viscosity | η | |
| Surface tension | γ | |
| Diffusion coefficient | Δ | |
| Atomic weight | A | |
| Molecular weight | M | |
| Velocity coefficient of reaction | k | |
| Equilibrium constant | $K, (K_e, K_p)$ | |
| van't Hoff coefficient | i | |
| Degree of dissociation (electrolytic, thermal, etc.) | α | |

LIST OF PHYSICO-CHEMICAL SYMBOLS.

4. Heat and Thermodynamics.

| | Usual symbol. | Alternative symbol. |
|--|------------------|---------------------|
| Temperature (centigrade) | t | |
| Temperature (absolute) | T | θ |
| Critical temperature | t_c, T_c | |
| Reduced temperature | t_r, T_r | |
| Critical solution temperature | t_{cs}, T_{cs} | |
| Quantity of heat | Q | |
| Entropy | S | |
| Specific heat | c | |
| Specific heat at constant pressure | c_p | |
| Specific heat at constant volume | c_v | |
| Ratio of specific heats, $c_p : c_v$ | γ | |
| Molecular heat | C | |
| Molecular heat at constant pressure | C_p | |
| Molecular heat at constant volume | C_v | |
| Latent heat per gram | l | |
| Latent heat per mole | L | |
| Maximum work (diminution of free energy) | A | |

5. Optics.

| | | |
|--|------------------------|-------|
| Wave-length of light | λ | |
| Refractive index | n | n_r |
| Specific refractive power (Gladstone and Dale) | $r_g, [r_g]_\lambda^o$ | |
| Specific refractive power (Lorentz and Lorenz) | $r_L, [r_L]_\lambda^o$ | |
| Molecular refractive power | R_g, R_L | |
| Angle of optical rotation | a | |
| Specific rotatory power | $[a]$ | |
| Molecular rotatory power | $M[a]$ | |
| Specific magnetic rotation | $[\omega]$ | |
| Molecular magnetic rotation | $M[\omega]$ | |

6. Electricity and Magnetism.

| | | |
|---|---|--------------------|
| Quantity of electricity | Q | |
| Current intensity | I | |
| Resistance | R | |
| Electromotive force | E | |
| Electrode potential, or discharge potential of an ion | E | ϵ |
| Electrode potential referred to the normal hydrogen or normal calomel electrode respectively, the potential of which is taken as zero | E_h, E_c | e_h, e |
| Normal potential, i.e., the electrode potential referred to the normal hydrogen or normal calomel electrode respectively, when the solution is molecular-normal in respect of all participating substances and ions of variable concentration | ${}_0 E_h, {}_0 E_c$ | ${}_0 e_h, {}_0 e$ |
| Dielectric constant | ϵ | |
| Conductivity (specific conductance) | κ | |
| Equivalent conductivity | Λ | |
| Equivalent conductivity at different dilutions—volumes in litres containing 1 gram-equivalent | $\Lambda_{10}, \Lambda_v, \Lambda_\infty$ | |

6. Electricity and Magnetism—(continued).

| | Usual symbol. | Alternative symbol. |
|---|-------------------------|---------------------|
| Equivalent conductivity of cation and of anion | Δ_k, Δ_a | |
| Equivalent conductivity of specified ions... | Δ_K, Δ_{Cl} | |
| Molecular conductivity | μ | |
| Velocity of cation and of anion in cm./ sec. when the potential gradient is 1 volt per cm. | U_k, U_a | |
| Transport number of cation and of anion ... | n_k, n_a | |
| Magnetic permeability | μ | |
| Magnetic susceptibility | κ | |

List of Symbols, Arranged Alphabetically.

| Symbol. | Name of quantity. |
|-------------------------------------|---|
| <i>A</i> | Atomic weight; maximum work. |
| <i>a</i> | Van der Waals's constant. |
| <i>b</i> | Van der Waals's constant. |
| <i>C</i> | Concentration; molecular heat. |
| <i>c</i> | Concentration; specific heat. |
| <i>C_p, C_v</i> | Molecular heat at constant pressure, and at constant volume. |
| <i>c_p, c_v</i> | Specific heat at constant pressure, and at constant volume. |
| <i>D</i> | Alternative symbol for density. |
| <i>d</i> | Diameter; total differential; density. |
| <i>d_c</i> | Critical density. |
| <i>d</i> | Reduced density. |
| <i>E</i> | Electromotive force; electrode potential. |
| <i>e</i> | Base of Napierian logarithms; charge on an electron. |
| <i>E_h, E_o</i> | Electrode potential referred to the normal hydrogen or the normal calomel electrode, respectively, the potential of which is taken as zero. |
| <i>E_h, E_o</i> | Normal potential, that is, the electrode potential referred to the normal hydrogen or the normal calomel electrode respectively, when the solution is molecular-normal in respect of all participating substances and ions of variable concentration. |
| <i>F</i> | Faraday's constant (number of coulombs per gram-equivalent of an ion). |
| <i>g</i> | Acceleration due to gravity. |
| <i>h</i> | Height. |
| <i>I</i> | Current. |
| <i>i</i> | Van't Hoff's coefficient. |
| <i>J</i> | Mechanical equivalent of heat. |
| <i>K</i> | Equilibrium constant. |
| <i>K_c, K_p</i> | Equilibrium constant, when molar concentrations and partial pressures respectively are employed. |
| <i>k</i> | Velocity coefficient of reaction. |
| <i>L</i> | Latent heat per mole. |
| <i>l</i> | Length; latent heat per gram. |
| <i>M</i> | Molecular weight. |
| <i>M^[α]</i> | Molecular rotatory power. |
| <i>M^[ω]</i> | Molecular magnetic rotatory power. |
| <i>m</i> | Mass. |
| <i>N</i> | Avogadro's constant (Loschmidt's number) or number of molecules in 1 gram-molecule. |
| <i>n</i> | Refractive index. |

List of Symbols, Arranged Alphabetically—(continued).

| Symbol. | Name of quantity. |
|---|---|
| n_k, n_a | Transport number of kation and of anion. |
| n | Refractive index (alternative symbol). |
| P | Pressure. |
| p | Pressure. |
| p_c, p_r | Critical pressure : reduced pressure. |
| Q | Quantity of heat; quantity of electricity. |
| R | Gas constant per mole; electrical resistance. |
| R_g, R_L | Molecular refractive power, according to Gladstone and Dale, and to Lorentz and Lorenz respectively. |
| r | Radius. |
| r_g, r_L | Specific refractive power according to Gladstone and Dale, and to Lorentz and Lorenz respectively. |
| S | Entropy. |
| T | Absolute temperature. |
| T_c | Critical temperature (on the absolute scale). |
| T_r | Reduced temperature (absolute). |
| T_{cs} | Critical solution temperature (absolute). |
| t | Time; temperature (centigrade). |
| t_c | Critical temperature (centigrade). |
| t_{cs} | Critical solution temperature (centigrade). |
| t_r | Reduced temperature (centigrade). |
| U_k, U_a | Velocity of kation and of anion in cm./sec. when the potential gradient is 1 volt per cm. |
| V | Volume. |
| v | Volume. |
| v_c, v_r | Critical volume : reduced volume. |
| W | Electrical resistance (alternative symbol). |
| x | Mole fraction. |
| α | Degree of dissociation (electrolytic, thermal, etc.); angle of optical rotation. |
| $[a]$ | Specific rotatory power. |
| γ | Surface tension; ratio of specific heats. |
| Δ | Diffusion coefficient. |
| δ | Variation. |
| ∂ | Partial differential. |
| ϵ | Electrode potential (alternative symbol); dielectric constant. |
| ϵ_h, ϵ | Electrode potential referred to the normal hydrogen or the normal calomel electrode respectively, the potential of which is taken as zero (alternative symbols). |
| $\epsilon^{e_h}, \epsilon^e$ | Normal potential, that is, the electrode potential referred to the normal hydrogen or the normal calomel electrode respectively, when the solution is molecular-normal in respect of all participating substances and ions of variable concentration (alternative symbols). |
| η | Viscosity. |
| θ | Temperature (centigrade), (alternative symbol). |
| κ | Specific conductance (conductivity); magnetic susceptibility. |
| Λ | Equivalent conductivity. |
| $\Lambda_{10}, \Lambda_v, \Lambda_\infty$ | Equivalent conductivity at different dilutions (volumes in litres containing 1 gram-equivalent). |
| Λ_k, Λ_a | Equivalent conductivity of kation and of anion. |
| λ | Wave-length of light. |
| μ | Molecular conductivity; magnetic permeability. |
| π | Ratio of circumference to diameter. |
| Σ | Summation. |
| σ | Surface tension (alternative symbol). |
| ϕ | Fluidity. |
| $[\omega]$ | Specific magnetic rotation. |