

Gas-phase Thermolyses. Part 3.¹ Gas-phase Thermolysis of Silylated Thionocarboxylic Acid Derivatives: a Route to Thioketens?


By Lars Carlsen* and Helge Egsgaard, Chemistry Department, Risø National Laboratory, DK-4000 Roskilde, Denmark

Ernst Schaumann,* Herbert Mrotzek, and Wolf-Rüdiger Klein, Institut für Organische Chemie und Biochemie der Universität Hamburg, D-2000 Hamburg 13, Martin-Luther-King-Platz 6, Federal Republic of Germany

The unimolecular gas-phase thermolytic decomposition of three silylated thionocarboxylic acid derivatives (2b), (3), and (8) have been studied by the flash vacuum thermolysis-field ionization mass spectrometry technique in the temperature range from 783 to 1 404 K in order to elucidate its possible applicability as a route to thioketens. Only very minor amounts of the expected thioketens were found, whereas the corresponding ketens were obtained as the major products. A possible mechanism for keten formation is discussed.

THE β -elimination of hydrogen chloride from acyl chlorides containing an α -hydrogen atom is perhaps the most important synthetic pathway to ketens.² The possible corresponding formation of thioketens from thioacyl chlorides can, however, be applied only in exceptional cases.^{3,4} Alternatively, thionocarboxylic

carboxylic acid *silyl* esters are found to be *ca.* 62 kJ mol⁻¹ more stable than the corresponding thio compounds.⁷ Hence, the thermodynamically unfavourable thiono to thio rearrangement together with the known ready elimination of silyl ethers from silylated carboxylic acid derivatives,⁸ suggest the use of silylated



SCHEME 1 Reagents: i, Prⁱ₂NLi; ii, Me₃SiCl; iii, B₂H₆; iv, H₂O₂; v, Cr^{VI} or Mn^{VIII}; vi, SOCl₂; vii, MeC=NSiMe₃

esters may be expected to yield thioketens upon thermolysis. However, this method may be of limited value, since a competing reaction analogous to the Chapman rearrangement is expected to lead to thiolocarboxylate esters⁵ rather than alcohol elimination. Thus, to our knowledge only one example of thioketen formation from a thionoester has been reported,⁶ the thioketen being isolated as its dimer.

In contrast to *alkyl* thionocarboxylates thiono-

thiocarboxylic acid derivatives as thioketen precursors. In this paper we report on the gas-phase thermolyses of three silylated thionocarboxylic acid derivatives in order to study the possible formation of thioketens. The starting materials (2b), (3), and (8) (see Scheme 1) have been chosen so as to provide sterically stabilized products, which would facilitate subsequent preparative operations.

The thionoester (2a) was synthesized by reaction of the

acyl chloride (1a) with a dithiocarbamate followed by silylation of the intermediate by chlorotrimethylsilane as reported by Kricheldorf and Leppert.⁹ The sterically highly hindered ester (2b) was prepared by a reaction of (1b) with the anion of *N*-trimethylsilylthioacetamide, adopting the method of Lücke.¹⁰ The thionoester (2a)




FIGURE 1 Field ionization mass spectra obtained after thermolysis of (3) at 1 043 K (a) and 1 404 K (b)

could be reacted further with lithium di-isopropylamine and chlorotrimethylsilane to the keten OS-acetal (3). A similar conversion was not observed in the case of (2b).

The precursor (7) to the thionoester (8) was prepared from homofenchene (4) by a procedure similar to that reported for the synthesis of 2-*t*-butyl-3,3-dimethylbutyryl chloride.¹¹ By using KMnO_4 in the presence of tetra-*n*-butylammonium bromide¹² in the oxidation of (5) to (6) instead of chromium trioxide the yield could be raised from 43 to 75%. The final silylation of (7) into (8) was performed as mentioned above using the metalated *N*-trimethylsilylthioacetamide reagent.

Compounds (5)–(8) can exist as *exo*- and/or *endo*-conformers. The primary alcohol (5) exhibits two sets of three methyl ^1H n.m.r. signals, which strongly suggests the presence of both possible configurations

at C-2. Similarly, more than three methyl signals and two signals for 2-H are observed for compounds (7) and (8). The carboxylic acid (6) apparently exists exclusively as a single conformer. However, the very small difference expected in the 4J coupling between *exo*-2-H and *exo*-6-H, and *endo*-2-H and *anti*-7-H, respectively (see *exo*- and *endo*-2-fenchol¹³), unfortunately affords no possibilities for the determination of the actual configuration.

Gas-phase Thermolyses of (2b), (3), and (8).—The unimolecular gas-phase thermolytic decomposition of the silylated thionocarboxylic acid derivatives (2b), (3), and (8) were studied by the flash vacuum thermolysis-field ionization mass spectrometry (f.v.t.-f.i.m.s.) technique^{14,15} (see Experimental section) in the temperature range from 783 to 1 404 K. Field ionization is known to give intense molecular ions and little fragmentation.¹⁴

In Figures 1a and 2a the f.i.m.s. spectra obtained from the thermolysis at 1 043 K of compounds (3) and (8), respectively, are shown. At this temperature the compounds are almost completely decomposed. Lowering the thermolysis temperature to 783 K did not change





FIGURE 2 Field-ionization mass spectra obtained after thermolysis of (8) at 1 043 K (a) and 1 404 K (b)

the composition of the product mixture, but only the relative yields. Also the absolute yields are changed, as *ca.* 95% of the starting materials were recovered.

The spectrum depicted in Figure 1a exhibits two

peaks at m/e 178 and 98. The compound responsible for the latter peak evidently does not contain sulphur, since only an isotopic cluster corresponding to ^{13}C and ^{18}O is seen, whereas the characteristic ^{34}S isotopic peak is lacking. The isotopic cluster surrounding the peak




of m/e 178 is consistent with the presence of two trimethylsilyl groups and one sulphur atom, as the relative intensities of the peaks of m/e 178/179/180/181 were found and calculated to be 1.00/0.19/0.12/0.02 and 1.00/0.18/0.13/0.02, respectively. On this basis the thermolysis of (3) is then formulated as in Scheme 2.


An analogous analysis of the spectrum depicted in Figure 2a leads to the conclusion that compound (8) upon thermolysis decomposes into the keten (11) and trimethylsilanethiol (12). Additionally a product exhibiting a molecular ion at m/e 136, which may be assigned to carbene (13) or an isomeric species, is formed (Scheme 3).

In contrast to these very simple fragmentation reactions compound (2b) exhibits a rather complicated

Changing the thermolysis temperature to 1404 K, in the case of (2b), did not give rise to any new product formation, but caused only a change in the relative composition of the product mixture. For compounds (3) and (8), however, small changes are observed by elevating the thermolysis temperature, as new sets of peaks are developed in the f.i.m.s. spectra obtained following thermolysis at 1404 K (Figures 1b and 2b). In both cases appearance of the new peaks can be rationalized as the result of competing thioketen formation. In the case of (8) the higher temperature also



resulted in the appearance of a variety of new peaks of low intensity, caused by more pronounced degradation of the carbon skeleton, probably similar to the thermolytic pathways observed for (2b) (Scheme 5).



decomposition pattern. The product mixture consists of the three possible ketens (14) and the three possible carbenes (15) together with *t*-butyl and *s*-propyl radicals, methylpropene and propene as products of consecutive radical decomposition, and the thiol (12). However, no indication of any thioketen formation was found (Scheme 4).


It should be noted that it is not possible to calculate the yields of the single species directly from the f.i.m.s. spectra, as the single compounds may exhibit rather different field ionization sensitivities.¹⁴ However, as the ketens and thioketens probably exhibit comparable sensitivities,¹⁵ it is obvious that only very minor amounts of the thioketens are formed.

DISCUSSION

The elimination of trimethylsilanethiol (12) from (2b) or (8) and of bistrimethylsilyl sulphide (10) from (3) to give ketens is in apparent contrast to the relative strengths of Si-S and Si-O bonds.⁷ Moreover, (2b)


steric effect may also favour transition states like (21) incorporating sulphur, as the larger size of sulphur compared with oxygen will diminish strain in the four-membered ring.

The reactivity of sulphur in four-centre transition



and (8) obviously cannot furnish ketens *via* a simple one-step process. A rationale for keten formation in these examples can be made by assuming primary enthiolization to give (20a and b), respectively, followed by β -elimination of $\text{Me}_3\text{Si-SH}$ (12), whereas formation of thioketen (18) at the more elevated temperature

states involving silicon is also obvious from the relative stabilities of bis-silylated amides (22) and thioamides (23). Thus compound (22; R = Ph) is gradually cleaved to give benzonitrile only at reflux temperature (*ca.* 470 K),¹⁷ whereas the corresponding thio-compound (23; R = Ph) eliminates $\text{Me}_3\text{Si-S-SiMe}_3$ (10) *in situ* on silylation of *N*-trimethylsilylthiobenzamide at 253 K.¹⁰



probably results from the direct β -elimination of $\text{Me}_3\text{Si-OH}$ (19) from (8). Elimination of $\text{Me}_3\text{Si-S-SiMe}_3$ (10) from (3) to give keten (9) would be in complete analogy to the invoked cleavage of (20).

Previously studied β -eliminations of silanols from *N*-silylcarbamates or imidocarboxylates have been proven to follow intramolecular pathways *via* four-centre transition complexes.¹⁶ This non-ionic mechanism may also operate in the gas phase and thus be applied to the thermolysis of (3) as well as (2b) and (8) *via* (20). This leads to transition state (21a) from (20) and (21b) from (8). The driving force in each case seems to be the pronounced nucleophilicity of sulphur which permits attack at silicon thus overcoming the stability of the Si-O bond. Besides the electronic a

mechanism, a primary thio-thiolo rearrangement followed by rapid decomposition of the latter into the ketens (11) and (14), respectively, and the silanethiol (12), should *a priori* be considered. After thermolyses at 1 043 K only very minor amounts of the starting materials are recovered, which implies that possible primarily generated species are quantitatively decomposed into the observed products before reaching the detection system (ion source). In general it is believed that species with half lives less than *ca.* 10^{-3} s may escape detection.¹⁸ However, assuming $\Delta H_r \simeq \Delta G_r \simeq -62$ kJ mol⁻¹ (*i.e.* $\Delta S_r \simeq 0$)⁷ for the thio-thiolo rearrangement, the equilibrium constants for the thio-thiolo system can, according to the van't Hoff equation, be calculated to be $\log_e K$ (293 K) $\simeq 25$ and $\log_e K$ (1 043 K) $\simeq 7$, respectively, *i.e.* even at the elevated temperature the thio-thiolo rearrangement will be rather favourable. In the present case the latter rearrangement should compete with the possible rethermolytic β -elimination of silanethiol (12) from the thio-species,

the latter totally suppressing the former, corresponding to quantitative rethermolysis. Based on the above figures and the fact that rethermolyses in general may be expected to occur to only a small extent¹⁸ it seems unlikely that a primary thiono-thiolo rearrangement should be expected to play any major role in the thermolyses of (8) and (2b).

In conclusion, it appears that because of the electronic and possibly also steric effect of neighbouring sulphur the Si-O bond is readily cleaved in the silylated thio-carboxylic acid derivatives (2b), (3), and (8) to give ketens, thus excluding this method as a suitable route to thioketens.

EXPERIMENTAL

I.r. spectra were recorded on a Perkin-Elmer 257 spectrograph, and the ¹H n.m.r. spectra (CDCl₃) on Varian T60, Varian EM360, and Perkin-Elmer R32 instruments.

*Flash Vacuum Thermolysis Technique.*¹⁸—The f.v.t. technique used is based on the direct combination of a thermolysis unit with a double focusing mass spectrometer with a field ion source. The thermolysis unit is constructed as a modification of the Pye-Unicam PV4000 pre-column pyrolysis system, which is based on the Curie-point principle, i.e. the high frequency inductive heating in ferromagnetic materials. The thermolysis unit is connected directly to the ion source of the mass spectrometer *via* a heated line-of-sight inlet system.

Samples (*ca.* 50 μg) of the pure compound were introduced (micro-syringe) into the reactor *via* a heated injection block. The contact time in the hot zone has been estimated to be *ca.* 10⁻³–10⁻⁴ s,¹⁸ fulfilling the contact time requirement for f.v.t. equipment. According to the geometry of the system, thermolysis products with half-lives < *ca.* 10⁻³ s are assumed to escape detection.¹⁸ The internal geometry of the reactor (length 40 mm, internal diameter 2 mm) combined with a low actual pressure (*P ca.* 10⁻⁴ Torr) assure a very low frequency of intermolecular collisions relative to the molecular-hot surface collision frequency, i.e. only unimolecular reactions take place.

The mass spectra were recorded on a Varian MAT CH 5D instrument equipped with a combined electron impact ionization-field ionization-field desorption ion source. The field ion emitter was a 10 μm tungsten wire activated in benzonitrile vapour.

Compounds (1a),¹⁹ (1b),²⁰ (4),²¹ and *N*-trimethylsilylthioacetamide (from thioacetamide and hexamethyldisilazane)²² were prepared according to reported procedures.

3,3-Dimethyl(thiobutanoic) Acid O-Trimethylsilyl Ester (2a).—Compound (2a) was prepared by reaction of (1a) with 2-phenylethylamine-carbon disulphide-triethylamine followed by silylation with chlorotrimethylsilane-triethylamine,⁹ yield 49%; b.p. 341–343 K at 11 mbar; i.r. exhibits no absorption in the carbonyl region; δ 0.35 (9 H, SiMe₃), 0.98 (9 H, Bu^t), and 2.64 (2 H, CH₂).

3,3-Dimethyl-2-isopropyl(thiobutanoic) Acid O-Trimethylsilyl Ester (2b).—*N*-Trimethylsilylthioacetamide (12.5 g, 0.086 mol) in hexane (15 ml) was slowly added to BuLi (0.086 mol) in hexane at 195 K. After 1 h (1b) (14.5 g, 0.082 mol) in hexane (20 ml) was slowly added and the solution was allowed to warm to room temperature. The precipitate was removed by filtration (with exclusion of moisture). The filtrate was concentrated *in vacuo* and the

product was isolated by distillation, yield 54%; b.p. 356–358 K at 2 mbar; i.r. exhibits no absorption in the carbonyl region; δ 0.40 (9 H, SiMe₃), 0.97 and 1.09 (6 H, d, *J ca.* 7 Hz, diastereotopic Me of Prⁱ), 1.02 (9 H, s, Bu^t), 2.00 (1 H, m, CH of Prⁱ), and 2.77 (1 H, d, *J* 5 Hz, 2-H).

3,3-Dimethyl-1-trimethylsiloxy-1-trimethylsilylthiobut-1-ene (3).—Compound (2a) (9.4 g, 46 mmol) in dry THF (20 ml) was slowly added to a stirred solution of lithium diisopropylamine (50 mmol) [prepared from di-isopropylamine (50 mmol) and an equimolar amount of BuLi in hexane] at 195 K. The stirring was continued for 3 h followed by dropwise addition of chlorotrimethylsilane (5.4 g, 50 mmol) in dry hexane (10 ml). The mixture was allowed to heat up to room temperature and the precipitated LiCl was removed by filtration after 20 h (exclusion of moisture). The solvent was removed *in vacuo* and the product isolated by distillation, yield 9.0 g (72%); b.p. 348–350 K (at 0.9 mbar); ν_{max.} 1 640 cm⁻¹ (C=C); δ 0.23 and δ 0.27 (both 9 H, SiMe₃), 1.03 (9 H, Bu^t), and 4.70 (1 H, =CH) (Found: C, 52.65; H, 10.35; S, 11.1. C₁₂H₂₈OSSi₂ requires C, 52.15; H, 10.15; S, 11.6%).

1,3,3-Trimethylbicyclo[2.2.1]heptan-2-ylmethanol (5).—Compound (5) was prepared from (4) by reaction with diborane followed by oxidation of the resulting borane with H₂O₂,¹¹ yield 89%; m.p. 334 K; ν_{max.} (KBr) 3 340 cm⁻¹ (OH); [α]_D²⁰ -37.7° (CDCl₃); δ 0.8–1.8 [m, containing δ 0.83, 0.88, 0.96, 1.00, 1.03, and 1.12 (Me)] and 3.68 (d, *J* 7 Hz, CH₂O) (Found: C, 78.25; H, 12.15. C₁₁H₂₀O requires C, 78.5; H, 12.0%).

1,3,3-Trimethylbicyclo[2.2.1]heptane-2-carboxylic Acid (6).—Compound (6) was preferentially prepared by oxidizing (5) according to ref. 12, yield 75%; m.p. 379 K, [α]_D²⁰ -4.4° (CDCl₃); ν_{max.} (KBr) *ca.* 3 000br (OH) and 1 700 cm⁻¹ (C=O); δ 0.9–2.3 [m, containing δ 1.02, 1.14, and 1.17 (Me)], 2.16 (d, *J ca.* 2 Hz, H-2), and 10.7br (OH) (Found: C, 72.5; H, 10.0. C₁₁H₁₈O₂ requires, C, 72.5; H, 9.95%).

1,3,3-Trimethylbicyclo[2.2.1]heptane-2-carbonyl Chloride (7).—Compound (7) was prepared by reaction between (6) and thionyl chloride, yield 60%; b.p. 347 K at 0.95 mbar; ν_{max.} (film) 1 805 cm⁻¹ (C=O); δ 0.9–2.3 [m, containing δ 1.00, 1.09, 1.14, 1.17, and 1.25 (Me)] and 2.61 and 2.70 (both d, *J ca.* 2 Hz, *endo*- and *exo*-2-H) (Found: C, 65.85; H, 8.65. C₁₁H₁₇ClO requires C, 65.8; H, 8.55%).

O-Trimethylsilyl 1,3,3-Trimethylbicyclo[2.2.1]heptane-2-thiocarboxylate (8).—The reaction was analogous to the above preparation of (2b) from (1b), yield 25%; b.p. 383 K at 1.1 mbar; i.r. exhibits no absorption in the carbonyl region; δ 0.41 (9 H, SiMe₃), 0.8–2.2 [m, containing δ 0.98, 1.00, 1.14, and 1.16 (Me)], and 2.76 and 2.87 (both d, *J ca.* 2 Hz, *endo*- and *exo*-2-H).

Compounds (2a), (2b), and (8) did not give satisfactory elemental analyses due to their highly hygroscopic nature.

[9/1860 Received, 23rd November, 1979]

REFERENCES

- Part 2, see ref. 15.
- D. Borrmann, 'Methoden der Organischen Chemie', Houben-Weyl-Müller, Thieme, Stuttgart, 1968, 4 Aufl., Bd. VII/4, p. 94.
- G. Seybold, *Angew. Chem.*, 1975, **87**, 710; *Angew. Chem. Internat. Edn.*, 1975, **14**, 703.
- H. J. Bestmann and G. Schmid, *Angew. Chem.*, 1974, **86**, 274; *Angew. Chem. Internat. Edn.*, 1974, **13**, 273; M. Parmentier, J. Galloy, M. Van Meerssche, and H. G. Viehe, *Angew. Chem.*, 1975, **87**, 33; *Angew. Chem. Internat. Edn.*, 1975, **14**, 53. Regarding

the mesomeric stabilization of the products, see also E. Schaumann, S. Harto, and G. Adiwidjaja, *Chem. Ber.*, 1979, **112**, 2698.

⁵ M. J. Janssen, 'Thiolo, Thiono and Dithio Acids and Esters', in S. Patai, 'The Chemistry of Carboxylic Acids and Esters', Wiley, London, 1969, ch. 15, p. 705.

⁶ A. Schönberg, L. v. Vargha, and H. Kaltschmitt, *Ber. Deutsch. Chem. Gesellschaft*, 1931, **64**, 2582.

⁷ G. A. Gornowicz and J. W. Ryan, *J. Org. Chem.*, 1966, **31**, 3439.

⁸ C. Ainsworth, F. Chen, and Y.-N. Kuo, *J. Organometallic Chem.*, 1972, **42**, 59; C. Ainsworth and Y.-N. Kuo, *ibid.*, p. 73.

⁹ H. R. Kricheldorf and E. Leppert, *Synthesis*, 1971, 435.

¹⁰ H. W. Lüke, Dissertation, Univ. Hamburg, 1978.

¹¹ M. S. Newman, A. Arkell, and T. Fukunaga, *J. Amer. Chem. Soc.*, 1960, **82**, 2498.

¹² A. W. Herriott and D. Picker, *Tetrahedron Letters*, 1974, 1511.

¹³ J. I. Musher, *Mol. Phys.*, 1963, **6**, 93.

¹⁴ H. D. Beckey, 'Field Ionization Mass Spectrometry', Pergamon Press, New York, 1971.

¹⁵ L. Carlsen, H. Egsgaard, and E. Schaumann, *J.C.S. Perkin II*, 1980, 1206.

¹⁶ V. F. Mironov, V. D. Sheludyakov, and V. P. Kozyukov, *Organometal. Chem. Synth.*, 1972, **1**, 329.

¹⁷ J. Pump and E. G. Rochow, *Chem. Ber.*, 1964, **97**, 627.

¹⁸ L. Carlsen and H. Egsgaard, *Thermochim. Acta*, 1980, **38**, 47.

¹⁹ A. H. Homeyer, F. C. Withmore, and V. H. Wallingford, *J. Amer. Chem. Soc.*, 1933, **55**, 4209.

²⁰ A. A. Sacks and J. G. Aston, *J. Amer. Chem. Soc.*, 1951, **73**, 3902.

²¹ O. Wallach, *Annalen*, 1907, **353**, 209.

²² W. Walter, H. W. Lüke, and J. Voss, *Annalen*, 1975, 1808.