

Kinetic Data for Outer-sphere V^{2+} and $[Ru(NH_3)_6]^{2+}$ Reductions of Platinum(IV) Complexes and a Correlation of Rate Constants

By Carmel S. Glennon, Timothy D. Hand, and A. Geoffrey Sykes,* Department of Inorganic and Structural Chemistry, The University, Leeds LS2 9JT

The kinetics of 2 : 1 V^{2+} and $[Ru(NH_3)_6]^{2+}$ reductions of six Pt^{IV} complexes: *trans*- $[Pt(NH_3)_4Cl_2]^{2+}$, *trans*- $[Pt(NH_2CH_3)_4Cl_2]^{2+}$, $[Pt(NH_3)_5Cl]^{3+}$, $[Pt(NH_3)_5Br]^{3+}$, *mer*- $[Pt(NH_3)_3Cl_3]^+$, and $[PtCl_6]^{2-}$ have been studied. Reactions involve one-equivalent changes, $Pt^{IV} \rightarrow Pt^{III} \rightarrow Pt^{II}$ where the first step is rate determining and Pt^{III} is a transient intermediate. A linear correlation of second-order rate constants (15 °C) k_V and k_{Ru} , $\log k_V = 0.89 \log k_{Ru} - 1.68$, is obtained at $I = 0.10M$ ($LiClO_4$), consistent with outer-sphere reactions, and non-utilization (by V^{2+}) of inner-sphere pathways involving Cl^- and Br^- bridges. The occurrence of outer-sphere reactions appears to preclude two-equivalent changes. Rate constants for the V^{2+} reductions of *trans*- $[Pt(NH_3)_4Cl_2]^{2+} > [Pt(NH_3)_5Cl]^{3+}$ differ by a factor of 60, whereas those for corresponding $[Pt(NH_3)_4]^{2+}$ reductions, which involve two-equivalent changes, have previously been shown to differ by a factor 10^4 . Activation parameters have been determined for five of the V^{2+} reductions. Those for the reduction of *trans*- $[Pt(NH_3)_4Cl_2]^{2+}$ and *trans*- $[Pt(NH_2CH_3)_4Cl_2]^{2+}$ differ significantly indicating an influence of the methyl groups.

The present investigations are an extension of earlier studies on the V^{2+} reductions of Pt^{IV} complexes $[PtCl_6]^{2-}$ and $[Pt(NH_3)_5Cl]^{3+}$ [equation (1)] in which outer-sphere



electron transfer in one-equivalent steps was indicated.¹ Evidence for this assignment was based on (a) failure to observe intermediate formation of vanadium(IV) under conditions when this would otherwise have been spectrophotometrically detectable, and (b) the magnitude of rate constants for the V^{2+} reduction of $[PtCl_6]^{2-}$, which were too large to permit substitution into either coordination sphere and formation of an inner-sphere bridged precursor complex. The first step in the $Pt^{IV} \rightarrow Pt^{III} \rightarrow Pt^{II}$ conversion is rate determining and is the focal point of the present investigations. Studies with the one-equivalent substitution-inert reductant $[Ru(NH_3)_6]^{2+}$, complement the V^{2+} studies and help to further substantiate the outer-sphere assignment.

EXPERIMENTAL

Preparation of *trans*- $[Pt(NH_3)_4Cl_2]Cl_2$.—The procedure for preparing this compound has been described previously.² Chlorine gas was passed through a solution of ca. 1.5 g of the Pt^{II} complex $[Pt(NH_3)_4]^{2+}$ (see below) in a minimum of 6M-HCl for 30 min. Pale yellow *trans*- $[Pt(NH_3)_4Cl_2]Cl_2$ crystals were filtered off, and washed with ethanol and ether {Found: H, 3.1; Cl, 35.3; N, 13.7. Calc. for $[Pt(NH_3)_4Cl_2]Cl_2$: H, 2.96; Cl, 35.1; N, 13.8%}. The spectrum in H_2O gave λ_{max} 335 nm (ϵ 101 l mol⁻¹ cm⁻¹).³

Preparation of *trans*- $[Pt(NH_2CH_3)_4Cl_2]Cl_2$.⁴—Methylamine solution (33%, 6.25 ml) was added to $Na_2[PtCl_4]$ (Johnson Matthey, 1.66 g) in H_2O (60 ml). A colour change, dark red \rightarrow yellow, was observed after the mixture had been warmed on a steam-bath for 15 min. The residue from rotary evaporation to dryness of the mixture (when excess of methylamine was removed) was treated with Cl_2 as above. The yellow product was washed with cold water, ethanol, and ether. Recrystallization was from water by addition of concentrated HCl {Found: C, 10.5; H, 4.45. Calc. for $[Pt(NH_2CH_3)_4Cl_2]Cl_2$: C, 10.4; H,

4.34%}. The spectrum in H_2O gave λ_{max} 340 nm, (ϵ 116.8 l mol⁻¹ cm⁻¹).

Preparation of $[Pt(NH_3)_5Br](ClO_4)_3 \cdot 3H_2O$.—Potassium hexabromoplatinate(IV) (Johnson and Matthey; 2 g), disodium hydrogen phosphate (4 g), and 20% ammonia solution (40 ml) were mixed with H_2O (60 ml). The mixture was heated on a steam-bath at ca. 80 °C for 40 min. Pale yellow crystals began to form, and the solution was cooled in ice. The crystals, the phosphate salt of $[Pt(NH_3)_5Br]^{3+}$, were filtered off and washed with ice-cold ethanol, and then ether. The solid was dissolved in a minimum volume of 0.1M-HClO₄, and the perchlorate salt obtained by addition of small amounts of ice-cold HClO₄ (72%). The spectrum gave λ_{max} 323 nm (ϵ 273 l mol⁻¹ cm⁻¹) {Found: H, 2.7; Br, 11.5; Cl, 14.3; N, 9.2. Calc. for $[Pt(NH_3)_5Br](ClO_4)_3 \cdot 3H_2O$: H, 2.9; Br, 11.1; C, 14.9; N, 9.8%}.

Preparation of *mer*- $[Pt(NH_3)_3Cl_3]Cl \cdot H_2O$.⁵—The Pt^{II} complex (see below) $[Pt(NH_3)_3Cl]Cl$ (1 g) was dissolved in 6M-HCl (250 ml), and a stream of Cl_2 gas was passed through for ca. 40 min. A yellow precipitate of *trans*- $[Pt(NH_3)_4Cl_2]Cl_2$ formed during the oxidation and was discarded. The filtrate was left to evaporate on a steam-bath until yellow crystals were obtained. The mixture was then cooled, and the yellow product filtered off. Recrystallization was from a minimum of H_2O with addition of conc. HCl. The spectrum in H_2O gave λ_{max} 402 nm ($\epsilon = 53$ l mol⁻¹ cm⁻¹); $\lambda(sh)$ 330 (ϵ 220 l mol⁻¹ cm⁻¹). The meridional assignment is consistent with the preparative procedure,⁵ and Pt^{II} products in this study. Calc. for $[Pt(NH_3)_3Cl_3]Cl \cdot H_2O$: H, 2.2; Cl (total), 35.0; Cl (ionic), 8.74; N, 10.3 [Found: H, 2.7; Cl (total), 34.8; Cl (ionic), 9.1; N, 10.2%]. Ionic chloride was determined by a potentiometric method using a Ag/AgCl electrode.⁶

Other Pt^{IV} Complexes.—Samples of ammonium hexachloroplatinate(IV) (Johnson and Matthey), λ_{max} 262 nm (ϵ 2.45×10^4 l mol⁻¹ cm⁻¹) and λ_{max} 454 nm (ϵ 48 l mol⁻¹ cm⁻¹), and chloropenta-ammineplatinum(IV) perchlorate, λ_{max} 286 nm (ϵ 186 l mol⁻¹ cm⁻¹) were as used previously.¹

Preparation of *cis*- $[Pt(NH_3)_2Cl_2]$.—This Pt^{II} complex was prepared by dissolving $K_2[PtCl_4]$ (Johnson and Matthey) in dilute HCl, and treating the solution with NH_3 in the presence of NH_4Cl (buffer control to prevent formation of hydroxo-complexes) as described previously.⁷ Yellow *cis*- $[Pt(NH_3)_2Cl_2]$ was separated from a small amount of in-

soluble Magnus's Green salt $[\text{Pt}(\text{NH}_3)_4][\text{PtCl}_4]$. The spectrum of an unrecrystallized sample gave λ_{max} 302 and 366 nm in satisfactory agreement with literature values.⁸

Preparation of $[\text{Pt}(\text{NH}_3)_3\text{Cl}]\text{Cl}$.⁹—The *cis*- $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$ complex (1 g) and cyanate (0.7 g, KNCO) were dissolved in H_2O (20 ml). After the solution had been boiled for 1 min a bright orange solution was obtained, at which stage concentrated HCl (0.4 ml) was added. Yellow unchanged *cis*- $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$ was filtered off. Addition of an excess of $\text{K}_2[\text{PtCl}_4]$ gave a precipitate of Magnus's Green salt, $[\text{Pt}(\text{NH}_3)_4][\text{PtCl}_4]$, and $[\text{Pt}(\text{NH}_3)_3\text{Cl}][\text{PtCl}_4]$. These were separated (Magnus's Green is the least soluble) by treatment with hot water acidified with a few drops of HCl. The filtrate yielded a flesh-pink solid, which was redissolved (dilute HCl) and treated with $[\text{Pt}(\text{NH}_3)_4]\text{Cl}_2$. The $[\text{PtCl}_4]^{2-}$ component was removed as insoluble Magnus's Green salt, and a solution of $[\text{Pt}(\text{NH}_3)_3\text{Cl}]^+$ and excess of $[\text{Pt}(\text{NH}_3)_4]^{2+}$ remained. The latter crystallized first and was filtered off. A sample of $[\text{Pt}(\text{NH}_3)_3\text{Cl}]\text{Cl}$ was obtained by leaving the solution over P_2O_5 in a desiccator. The spectrum gave λ_{max} 325 (ϵ 34.5 l mol⁻¹ cm⁻¹) and 250 nm (ϵ 122) in good agreement with literature values,⁸ which are quoted for $[\text{Pt}(\text{NH}_3)_3\text{Cl}]\text{SO}_4$.

Preparation of $[\text{Pt}(\text{NH}_3)_4]\text{Cl}_2$.¹⁰—To samples of Magnus's Green $[\text{Pt}(\text{NH}_3)_4][\text{PtCl}_4]$ from the previous preparation (ca. 0.3 g), were added H_2O (25 ml), concentrated HCl (2 ml),




FIGURE 1 U.v. spectra of Pt^{IV} complexes *trans*- $[\text{Pt}(\text{NH}_3)_4\text{Cl}_2]^{2+}$ (\blacktriangledown), *trans*- $[\text{Pt}(\text{NH}_2\text{CH}_3)_4\text{Cl}_2]^{2+}$ (\blacksquare), $[\text{Pt}(\text{NH}_3)_5\text{Cl}]^{2+}$ (\blacktriangle), $[\text{Pt}(\text{NH}_3)_5\text{Br}]^{3+}$ (\times), and *mer*- $[\text{Pt}(\text{NH}_3)_3\text{Cl}_3]^+$ ($+$) in H_2O

and NH_3 (d 0.880, 100 ml). The solution was boiled (ca. 1 h) until the solid had dissolved, the solution being maintained at ca. 100 ml by addition of NH_3 . The solution was concentrated to ca. 50 ml and dilute HCl was added to the filtrate until the solution was just acid to litmus; concentrated HCl (1 ml) was then added. A solid sample was obtained by addition of ($\times 10$ volumes) ethanol-acetone (1:1). The spectrum in H_2O gave λ_{max} 287 nm (ϵ 44 l mol⁻¹ cm⁻¹) (lit.,⁸ ϵ 36 l mol⁻¹ cm⁻¹).

Reductants.—Solutions of hexaquoovanadium(II) perchlorate (hereafter V^{2+}) in dilute perchloric acid were prepared by electrolytic reduction of vanadium(IV) perchlorate. Hexa-ammineruthenium(II) solutions were prepared by Zn/Hg reduction of the chloride salt¹¹ of the Ru^{II} analogue in 0.01M-HClO₄, and standardized as previously described.¹²

Product Analyses.—Detailed spectrophotometric analyses were carried out for the V^{2+} reductions of Pt^{IV} complexes. It has previously been demonstrated that with $[\text{PtCl}_6]^{2-}$ and $[\text{Pt}(\text{NH}_3)_5\text{Cl}]^{3+}$ little or no V^{IV} is formed and that the 2:1 reactions, (1), proceed in consecutive one-equivalent steps. Here we have further confirmed that no V^{IV} is formed in the reaction of V^{2+} with *mer*- $[\text{Pt}(\text{NH}_3)_3\text{Cl}_3]^+$. The identity of the Pt^{II} products is also relevant. The V^{2+} reactant was in ca. 5% excess of the stoichiometric 2:1 ratio. Reactions were at 25 °C, $[\text{H}^+] = 1.0\text{M}$, and $I = 1.0\text{M}$ (perchlorate). Final spectra were corrected for $\text{V}^{2+}/\text{V}^{3+}$ absorbance. Relevant Pt^{IV} and Pt^{II} spectra are shown in Figures 1 and 2.




FIGURE 2 U.v. spectra of *cis*- $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$ (\blacksquare), $[\text{Pt}(\text{NH}_3)_3\text{Cl}]^+$ (\bullet), and $[\text{Pt}(\text{NH}_3)_4]^{2+}$ (\blacktriangle) in H_2O

Spectra of *cis*- and *trans*- $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$ are given in ref. 8. It was found that for the V^{2+} reduction of *trans*- $[\text{Pt}(\text{NH}_3)_4\text{Cl}_2]^{2+}$ the product was at least 75% $[\text{Pt}(\text{NH}_3)_4]^{2+}$. With $[\text{Pt}(\text{NH}_3)_5\text{Cl}]^{3+}$ as oxidant $[\text{Pt}(\text{NH}_3)_4]^{2+}$ is the product, and with *mer*- $[\text{Pt}(\text{NH}_3)_3\text{Cl}_3]^+$ the product is $[\text{Pt}(\text{NH}_3)_3\text{Cl}]^+$. The final spectrum in the latter case gave λ_{max} 325 nm (ϵ 26 l mol⁻¹ cm⁻¹) as compared to λ_{max} 325 nm (ϵ 34 l mol⁻¹ cm⁻¹) for $[\text{Pt}(\text{NH}_3)_3\text{Cl}]^+$. Since *cis*- and *trans*- $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$ have much stronger absorption at λ_{max} 302 (ϵ 128 l mol⁻¹ cm⁻¹) and 315 nm (ϵ 60) peaks respectively,⁸ these products can be excluded.

Kinetics.—Reactions were monitored at Pt^{IV} absorbance peak positions by the stopped-flow method. The V^{2+} reductant was generally in large (>10-fold) excess when plots of absorbance changes $\log(A_t - A_\infty)$ against time (linearity > 80% completion) gave slopes of $2k_V [\text{Pt}^{\text{IV}}]_0 / 2.303$ from which second-order rate constants k_V could be evaluated. Rate constants k_V for V^{2+} reactions with the latter in smaller excess were obtained from the slope ($\times 2.303$) of plots $\log\{1 + ([\text{V}^{2+}]_0 - 2[\text{Pt}^{\text{IV}}]_0)\Delta\epsilon / (A_t - A_\infty)\}$ against time, where $\Delta\epsilon$ (l mol⁻¹ cm⁻¹) = $(\epsilon_{\text{Pt}^{\text{IV}}}/2 + \epsilon_{\text{VII}}) - (\epsilon_{\text{Pt}^{\text{II}}}/2 + \epsilon_{\text{VIII}})$, l is the optical path length (2 cm),

TABLE 1

Rate constants, k_V , for the V^{2+} reduction of $trans$ -[Pt(NH₃)₄Cl₂]²⁺, $I = 1.00M$ (LiClO₄) except as stated, λ 325 nm

$t/^\circ C$	[H ⁺]	10 ³ [V ²⁺]	10 ⁴ [Pt ^{IV}]	k_V
	M	M	M	l mol ⁻¹ s ⁻¹
35.0	0.50	7.4	3.46	95
	0.50	3.7	1.85	107
	0.50	3.7	1.67	107 ^a
	0.10	9.8	2.86	114
25.0	0.10	6.5	2.86	95
	1.00	9.8	4.57	53
	1.00	4.9	4.57	53
	1.00	2.4	4.57	67
	0.50	3.7	1.67	55 ^a
	0.10	9.8	2.86	63
	0.10	6.5	2.86	64
	0.10	6.5	1.33	56
15.0	0.10	3.3	1.33	59
	0.10	9.8	2.86	32
	0.10	6.5	1.33	28
	0.10	3.3	1.33	25
	0.09	5.3	2.30	15.1 ^b

^a [Cl⁻] = 0.01M added initially. ^b $I = 0.10M$ (LiClO₄). Average of 4 runs.

and initial concentrations are indicated. Most [Ru(NH₃)₆]²⁺ reductions were carried out with the Pt^{IV} reactant in excess of the stoichiometric amount, when second-order rate constants were obtained from plots of $\log\{1 + (2[Pt^{IV}]_0 - [Ru^{II}]_0)\Delta\epsilon/(A_t - A_\infty)\}$ against time. For consistency with these equations correction factors are required to rate constants for the V^{2+} reduction of [PtCl₆]²⁻, and the [Ru(NH₃)₆]²⁺ reduction of [Pt(NH₃)₅Cl]³⁺, (all $\times \frac{1}{2}$) reported in the previous paper.¹

Ionic strengths of reactant solutions were made up to $I = 1.00$ and $0.10M$ with V^{2+} , and $0.10M$ with [Ru(NH₃)₆]²⁺ as the reductant using HClO₄/LiClO₄.

Treatment of Data.—Activation parameters for the V^{2+} reductions were obtained using a non-linear least-squares programme¹³ with $1/k^2$ weighting factor. Errors in other cases were estimated using a standard (unweighted) least-squares treatment.

Results. Second-order rate constants, k_V , for the V^{2+} reductions of Pt^{IV} complexes, rate law (2), are listed in

$$-d[Pt^{IV}]/dt = k_V[V^{2+}][Pt^{IV}] \quad (2)$$

Tables 1—4 for $trans$ -[Pt(NH₃)₄Cl₂]²⁺, $trans$ -[Pt(NH₂CH₃)₄Cl₂]²⁺, [Pt(NH₃)₅Br]³⁺, and mer -[Pt(NH₃)₃Cl₃]³⁺ respectively. All rate constants quoted are an average of at least two duplicate runs. Rate constants at $15^\circ C$, $I = 0.10M$ (LiClO₄), were required for the correlation described below.

TABLE 2

Rate constants, k_V , for the V^{2+} reduction of $trans$ -[Pt(NH₂CH₃)₄Cl₂]²⁺, $I = 1.00M$ (LiClO₄) except as stated, λ 340 nm

$t/^\circ C$	[H ⁺]	10 ² [V ²⁺]	10 ³ [Pt ^{IV}]	k_V
	M	M	M	l mol ⁻¹ s ⁻¹
35.0	0.90	4.88	2.03	258
	0.90	0.49	0.24	250
	0.10	0.85	0.45	262 ^a
	0.10	0.85	0.45	184
25.0	0.90	4.88	2.03	195
	0.90	1.69	0.96	192
	0.90	0.49	0.24	192
	0.10	0.85	0.45	184
15.0	0.10	0.42	0.45	199
	0.90	1.69	0.96	130
	0.10	0.42	0.45	138
	0.09	0.53	0.26	57 ^b

^a [Cl⁻] = 0.010M added initially. ^b $I = 0.10M$ (LiClO₄).

No dependence on [H⁺] and [Cl⁻] is observed. Activation parameters were determined for the first three of these reactions, and these are shown alongside values for [Pt(NH₃)₅Cl]³⁺ and [PtCl₆]²⁻ at $I = 1.00M$ (LiClO₄), in Table 5.

Second-order rate constants, k_{Ru} , are defined by a rate

TABLE 3

Rate constants, k_V , for the V^{2+} reduction of [Pt(NH₃)₅Br]³⁺, $I = 1.00M$ (LiClO₄), λ 323 nm

$t/^\circ C$	[H ⁺]	10 ³ [V ²⁺]	10 ³ [Pt ^{IV}]	k_V
	M	M	M	l mol ⁻¹ s ⁻¹
35.0	0.95	0.75	4.81	14.2
	0.90	2.24	11.5	
	0.10	1.23	5.85	14.6
25.0	0.95	1.23	5.92	10.4
	0.95	0.75	4.81	10.9
	0.90	2.24	11.5	10.2
	0.50	0.72	2.89	10.6
	0.10	1.23	5.85	9.5
15.0	0.95	0.75	4.81	5.8
	0.90	2.24	11.5	5.6
	0.50	0.72	2.89	6.5
	0.10	1.23	5.85	6.0

law of the same form as (2). All studies were at $15^\circ C$, $I = 0.10M$ (LiClO₄), Table 6. Difficulties in working with this reductant at higher [H⁺] and [ClO₄⁻], and at higher temperatures, have been referred to previously.¹²

TABLE 4

Rate constants ($15.0^\circ C$), k_V , for the V^{2+} reduction of mer -[Pt(NH₃)₃Cl₃]³⁺, $I = 1.00M$ (LiClO₄) except as stated, λ 400 nm

[H ⁺]	10 ³ [V ²⁺]	10 ⁴ [Pt ^{IV}]	k_V
M	M	M	l mol ⁻¹ s ⁻¹
0.10	4.30	5.35	177
0.30	4.30	4.30	158
0.50	4.30	6.42	141
0.55	2.92	5.35	169
0.75	2.92	4.27	186
1.00	2.92	3.20	140
1.00	2.92	6.42	146
0.09	2.88	5.25	71 [*]

* $I = 0.10M$ (LiClO₄).

Table 7 lists k_V and k_{Ru} values at $15^\circ C$, $I = 0.10M$ (LiClO₄), relevant to the rate constant correlation in Figure 3. Rate constants for the V^{2+} and [Ru(NH₃)₆]²⁺ reductions of [PtCl₆]²⁻, and the V^{2+} reduction of [Pt(NH₃)₅Cl]³⁺ at

TABLE 5

Summary of rate constants ($25^\circ C$) and activation parameters for the V^{2+} reduction of Pt^{IV} complexes, $I = 1.00M$ (LiClO₄)

Oxidant	k_V	ΔH^\ddagger	ΔS^\ddagger
	l mol ⁻¹ s ⁻¹	kcal mol ⁻¹	cal K ⁻¹ mol ⁻¹
$trans$ -[Pt(NH ₃) ₄ Cl ₂] ²⁺	28.3	10.7 ± 0.6	-14.5 ± 2.0
$trans$ -[Pt(NH ₂ CH ₃) ₄ Cl ₂] ²⁺	134	5.1 ± 0.3	-31.0 ± 0.9
[Pt(NH ₃) ₅ Cl] ³⁺ ^a	0.61	9.9 ± 0.5	-25.1 ± 1.7
[Pt(NH ₃) ₅ Br] ³⁺	6.0	7.5 ± 0.5	-28.8 ± 1.6
mer -[Pt(NH ₃) ₃ Cl ₃] ³⁺	159		
[PtCl ₆] ²⁻ ^b	1.23×10^4	4.6 ± 0.2	-23.9 ± 0.7

^a Ref. 1. ^b Ref. 1, rate constants corrected ($\times \frac{1}{2}$) see Experimental section.

$I = 0.10M$ were not previously reported. A linear fit was obtained, see equation (6) below, using all k_{Ru} values in Table 7. Errors for k_V were assumed to be of the same magnitude as observed for the more extensive studies at $I = 1.00M$ (LiClO₄).

TABLE 6

Rate constants (15 °C), k_{Ru} , for the $[\text{Ru}(\text{NH}_3)_6]^{2+}$ reduction of Pt^{IV} complexes, $I = 0.10\text{M}$ (LiClO_4), λ 335 nm

Oxidant	$[\text{H}^+]$	$10^4[\text{Ru}(\text{NH}_3)_6^{2+}]$	$10^4[\text{Pt}^{\text{IV}}]$	$10^3 k_{\text{Ru}}$
	M	M	M	$\text{l mol}^{-1} \text{s}^{-1}$
<i>trans</i> - $[\text{Pt}(\text{NH}_3)_4\text{Cl}_2]^{2+}$	0.10	1.93	6.05	2.07
	0.01	4.61	6.05	2.00
	0.01	1.08	1.01	2.36
<i>trans</i> - $[\text{Pt}(\text{NH}_2\text{CH}_3)_4\text{Cl}_2]^{2+}$	0.10	1.12	2.13	9.3
	0.01	2.45	5.85	6.5
	0.01	0.82	2.10	6.7
$[\text{Pt}(\text{NH}_3)_5\text{Br}]^{3+}$	0.10	0.52	2.29	0.163
	0.01	1.24	2.41	0.198
	0.01	0.20	1.21	0.187
<i>mer</i> - $[\text{Pt}(\text{NH}_3)_3\text{Cl}_3]^+$	0.01	5.8	1.30	16.0
	0.01	5.8	0.73	16.0
	0.02	4.8	0.73	19.5
	0.02	4.8	0.73	17.6
	0.03	3.4	1.09	15.8
	0.04	1.9	0.73	19.8
	0.04	1.9	1.09	16.0
	0.10	6.3	2.43	21.5

DISCUSSION

With other redox partners both the V^{2+} and Pt^{IV} reactants have been shown to be capable of two-equivalent changes. Although in the case of the Cr^{2+} reduction of $[\text{Pt}(\text{NH}_3)_5\text{Cl}]^{3+}$ a considerable barrier is presented in

TABLE 7

Summary of rate constants for the V^{2+} and $[\text{Ru}(\text{NH}_3)_6]^{2+}$ reduction of Pt^{IV} complexes at 15 °C, $I = 0.10\text{M}$ (LiClO_4)

Oxidant	k_{V} $\text{l mol}^{-1} \text{s}^{-1}$	k_{Ru} $\text{l mol}^{-1} \text{s}^{-1}$
<i>trans</i> - $[\text{Pt}(\text{NH}_3)_4\text{Cl}_2]^{2+}$	15.1	2.14×10^3
<i>trans</i> - $[\text{Pt}(\text{NH}_2\text{CH}_3)_4\text{Cl}_2]^{2+}$	57	7.5×10^3
$[\text{Pt}(\text{NH}_3)_5\text{Cl}]^{3+}$	0.26 ^a	17.7 ^b
$[\text{Pt}(\text{NH}_3)_5\text{Br}]^{3+}$		1.8×10^2
<i>mer</i> - $[\text{Pt}(\text{NH}_3)_3\text{Cl}_3]^+$	71	1.78×10^4
$[\text{PtCl}_6]^{2-}$	2.14×10^4 ^c	7.9×10^6 ^d

^a $[\text{V}^{2+}] = 3.0 \times 10^{-3}\text{M}$; $[\text{Pt}^{\text{IV}}] = 1.5 \times 10^{-3}\text{M}$; $[\text{H}^+] = 0.08\text{M}$. ^b Rate constants from ref. 1 corrected ($\times \frac{1}{2}$), see Experimental section. ^c $[\text{V}^{2+}] = 6.5 \times 10^{-4}\text{M}$; $[\text{Pt}^{\text{IV}}] = 7.6 \times 10^{-4}\text{M}$; $[\text{H}^+] = 0.09\text{M}$. ^d Error $\pm 30\%$; $[\text{Ru}(\text{NH}_3)_6^{2+}] = (8.4 - 23.5) \times 10^{-6}\text{M}$; $[\text{Pt}^{\text{IV}}] = 6.8 \times 10^{-6}\text{M}$.

that Cr^{IV} is not easy to generate in aqueous solution, such a two-equivalent inner-sphere pathway appears more favourable than the alternative involving formation of Pt^{III} .¹⁴ Oxidation of V^{2+} to V^{IV} should occur more readily since V^{IV} (as VO^{2+}) is a stable species.¹⁵ However no contributions from the two-electron step, $\text{V}^{\text{II}} + \text{Pt}^{\text{IV}} \rightarrow \text{V}^{\text{IV}} + \text{Pt}^{\text{II}}$, which with excess of V^{2+} is followed by further reactions $\text{V}^{\text{II}} + \text{V}^{\text{IV}} \rightarrow \text{V}^{\text{III}}_2 \rightarrow 2\text{V}^{\text{III}}$, are observed. This alone suggests an outer-sphere assignment, if (as seems likely) the more extensive re-organisation attendant on a two-equivalent change is difficult to achieve in an outer-sphere process.

For the six Pt^{IV} oxidants now investigated all observations are in accordance with a reaction sequence (3)—(4). Further confirmation of the outer-sphere



assignment to reaction (3) was sought by comparison of rate constants k_{V} with k_{Ru} for the corresponding

$[\text{Ru}(\text{NH}_3)_6]^{2+}$ reductions. A plot of $\log k_{\text{V}}$ against $\log k_{\text{Ru}}$ (Figure 3) shows satisfactory linearity and since the slope approaches 1.0 as required by the modified Marcus approach,¹⁶ outer-sphere assignments can be made for both reductants. Initially however the V^{2+} studies were at $I = 1.00\text{M}$ (LiClO_4), whereas the $[\text{Ru}(\text{NH}_3)_6]^{2+}$ studies had to be at a lower ionic strength, $I = 0.10\text{M}$ (LiClO_4). These results (open circles in Figure 3) give a slope similar to that of the broken line, which summarizes existing $\log k_{\text{V}}$ and $\log k_{\text{Ru}}$ data for the reaction of some 24 Co^{III} oxidants [equation (5)].¹⁷ With Pt^{IV} as

$$\log k_{\text{V}} = 0.81 \log k_{\text{Ru}} - 0.64 \quad (5)$$

oxidant a wide range of charges from -2 to $+3$ were investigated, and we therefore saw fit to obtain further




FIGURE 3 Correlation of rate constants for V^{2+} (k_{V}) and $[\text{Ru}(\text{NH}_3)_6]^{2+}$ (k_{Ru}) reductions of Pt^{IV} complexes *trans*- $[\text{Pt}(\text{NH}_3)_4\text{Cl}_2]^{2+}$ (1), *trans*- $[\text{Pt}(\text{NH}_2\text{CH}_3)_4\text{Cl}_2]^{2+}$ (2), $[\text{Pt}(\text{NH}_3)_5\text{Cl}]^{3+}$ (3), $[\text{Pt}(\text{NH}_3)_5\text{Br}]^{3+}$ (4), *mer*- $[\text{Pt}(\text{NH}_3)_3\text{Cl}_3]^+$ (5), and $[\text{PtCl}_6]^{2-}$ (6). Rate constants k_{Ru} at $I = 0.10\text{M}$ (LiClO_4), and k_{V} at $I = 1.00\text{M}$ (LiClO_4) (\circ), and $I = 0.10\text{M}$ (LiClO_4) (\bullet)

data with V^{2+} as reductant at $I = 0.10\text{M}$ (LiClO_4). A significant change in slope is observed, where the shaded

$$\log k_{\text{V}} = (0.89 \pm 0.01) \log k_{\text{Ru}} - (1.68 \pm 0.06) \quad (6)$$

points now give a fit to [equation (6)]. It is difficult to

reappraise the Co^{III} studies making due allowance for effects of I (charges on the Co^{III} complexes vary much less in any case), but we suspect that the slope may also be closer to 1.0,¹⁸ than indicated by equation (5). Correlations involving Cr^{2+} ,^{19,20} $[\text{Cr}(\text{bipy})_3]^{2+}$,^{21,22} Eu^{2+} ,¹⁸ have been shown to be close to 1.0. A notable exception appears to be with $[\text{TiOH}]^{2+}$ as reductant (slope *ca.* 0.55) an effect which as yet is not fully understood.²³ The possibility that this correlation is in fact curved has been suggested but additional data are required.

The correlation in Figure 3 leads to an outer-sphere assignment for the V^{2+} reduction of the complex $[\text{Pt}(\text{NH}_3)_5\text{Br}]^{3+}$. It is concluded that Br^- like Cl^- does not readily function as a lead-in group for inner-sphere V^{2+} reduction, an interesting observation bearing in mind that the rate constants (25 °C) for $[\text{Pt}(\text{NH}_3)_5\text{Br}]^{3+}$ ($6.0 \text{ l mol}^{-1} \text{ s}^{-1}$) and $[\text{Pt}(\text{NH}_3)_5\text{Cl}]^{3+}$ ($0.61 \text{ l mol}^{-1} \text{ s}^{-1}$) do not necessarily exclude an inner-sphere contribution. There are two points to make. One is that V^{2+} has an aversion for the softer halide ions which can be rationalized in terms of hard and soft acid-base theory (V^{2+} is hard).²³ Secondly, assuming that formation of an inner-sphere precursor complex can occur, subsequent electron-transfer might be slow. We believe the first of these is relevant, since the V^{2+} reduction of $[\text{Co}(\text{NH}_3)_5\text{Br}]^{2+}$ ($30 \text{ l mol}^{-1} \text{ s}^{-1}$) and $[\text{Co}(\text{NH}_3)_5\text{Cl}]^{2+}$ ($10.2 \text{ l mol}^{-1} \text{ s}^{-1}$) are also outer-sphere.²² The V^{2+} reduction of $[\text{Co}(\text{NH}_3)_5\text{F}]^{2+}$ on the other hand is inner-sphere,²² and reduction of Pt^{IV} fluoro-complexes [*e.g.* $\text{Pt}(\text{NH}_3)_5\text{F}^{3+}$] would be of interest since they might also proceed by an inner-sphere pathway, and thus provide conditions, appropriate for a two-electron transfer.

Studies on $\text{Pt}^{\text{II}}-\text{Pt}^{\text{IV}}$ exchange reactions have provided evidence for Cl^- bridging and an inner-sphere mechanism.²⁴ A two-equivalent change involving simultaneous or near-simultaneous transfer of two electrons may occur in these reactions. The rate constants for the $[\text{Pt}(\text{NH}_3)_4]^{2+}$ exchange with *trans*- $[\text{Pt}(\text{NH}_3)_4\text{Cl}_2]^{2+}$ is 10^4 greater than for the exchange with $[\text{Pt}(\text{NH}_3)_5\text{Cl}]^{3+}$. This difference is presumed to stem from the presence of a stronger *trans* Pt-NH₃ bond in $[\text{Pt}(\text{NH}_3)_5\text{Cl}]^{3+}$, making the accompanying octahedral-square-planar geometry change (some bond stretching will be required in the activated complex) more difficult to achieve. For the outer-sphere V^{2+} and $[\text{Ru}(\text{NH}_3)_6]^{2+}$ reductions rate constants exhibit much smaller differences of $\times 60$ and $\times 10^2$ respectively, consistent with the less demanding $\text{Pt}^{\text{IV}} \rightarrow \text{Pt}^{\text{III}}$ change. Enthalpies of activation are, moreover, very similar for these two V^{2+} reductions (Table 5) whereas for the $[\text{Pt}(\text{NH}_3)_4]^{2+}$ reductions of $[\text{Pt}(\text{NH}_3)_5\text{Cl}]^{3+}$ ($16.7 \text{ kcal mol}^{-1}$) and *trans*- $[\text{Pt}(\text{en})_2\text{Cl}_2]^{2+}$ ($11.5 \text{ kcal mol}^{-1}$) the differences account for the rate constant pattern. The V^{2+} reduction of $[\text{Pt}(\text{NH}_3)_5\text{Cl}]^{3+}$ might be slower than for *trans*- $[\text{Pt}(\text{NH}_3)_4\text{Cl}_2]^{2+}$ because Pt-N bond vibrations *trans* to the Cl^- are less conducive to electron transfer into the e_g orbital aligned in this direction.²⁵ Rate constants for the reductions of *trans*- $[\text{Pt}(\text{NH}_3)_4\text{Cl}_2]^{2+}$ and *mer*- $[\text{Pt}(\text{NH}_3)_3\text{Cl}_3]^+$ complexes are

very similar, as in the corresponding $[\text{Pt}(\text{NH}_3)_4]^{2+}$ reductions.

Of further interest is the effect of replacing NH_3 by CH_3NH_2 in *trans*- $[\text{Pt}(\text{NH}_3)_4\text{Cl}_2]^{2+}$, when activation parameters (Table 5) change in a way which we do not fully understand, but which could be related to the presence of the hydrophobic methyl groups. The trends are self compensating since rate constants k_V show little variation and are within a factor of $\times 5$.

Platinum(II) reaction products were identified spectrophotometrically, and these indicate that geometry changes accompanying the redox change occur by Cl^- rather than NH_3 loss. Unfortunately our data do not include a study of a *cis*-dichloro-complex. Beattie and Starink²⁶ have shown that for the two-equivalent Cr^{2+} reduction of *cis*- $[\text{Pt}(\text{pn})_2\text{Cl}_2]^{2+}$ (pn = propylenediamine) the Pt^{II} product is $[\text{Pt}(\text{pn})(\text{pnH})\text{Cl}]^+$, whereas for the one-equivalent Eu^{2+} reduction (probably inner-sphere) the final Pt^{II} product is $[\text{Pt}(\text{pn})_2]^{2+}$. For a one-equivalent outer-sphere V^{2+} reduction loss of two *cis*-chloride ligands is to be expected.

C. S. G. and T. D. H. are grateful for S.R.C. research studentships.

[8/2100 Received, 5th December, 1978]

REFERENCES

- 1 A. Bakač, T. D. Hand, and A. G. Sykes, *Inorg. Chem.*, 1975, **14**, 2540.
- 2 'Gmelin' Handbuch der Anorganischen Chemie, 68D Verlag Chemie, Weinheim/Bergstrasse, 1957, p. 491.
- 3 R. R. Rettew and R. C. Johnson, *Inorg. Chem.*, 1965, **4**, 1565.
- 4 L. N. Essen, *Inorg. Synth.*, 1974, **15**, 93.
- 5 Ref. 2, p. 525.
- 6 C. Brown, *J. Amer. Chem. Soc.*, 1934, **56**, 646.
- 7 G. B. Kauffman and D. O. Cowan, *Inorg. Synth.*, 1963, **7**, 241.
- 8 J. Chatt, G. A. Gamlen, and L. E. Orgel, *J. Chem. Soc.*, 1958, 486.
- 9 L. A. Tschugnev, *J. Chem. Soc.*, 1915, 1247.
- 10 R. N. Keller, *Inorg. Synth.*, 1946, **2**, 250.
- 11 J. R. Pladziewicz, T. J. Meyer, J. A. Broomhead, and H. Taube, *Inorg. Chem.*, 1973, **12**, 639.
- 12 C. S. Glennon, J. D. Edwards, and A. G. Sykes, *Inorg. Chem.*, 1978, **17**, 1654.
- 13 R. H. Moore and R. K. Zeigler, Los Alamos Report LA, 2367, 1959, and Addenda.
- 14 J. K. Beattie and F. Basolo, *Inorg. Chem.*, 1971, **10**, 486.
- 15 M. Green and A. G. Sykes, *J. Chem. Soc. (A)*, 1970, 3221 and 1971, 3067.
- 16 D. L. Toppen and R. G. Linck, *Inorg. Chem.*, 1971, **10**, 2635.
- 17 T. D. Hand, Ph.D. Thesis, University of Leeds, 1976, p. 29, and references therein.
- 18 F.-R. F. Fan and E. S. Gould, *Inorg. Chem.*, 1974, **13**, 2647.
- 19 I. Baldea, K. Wiegardt, and A. G. Sykes, *J.C.S. Dalton*, 1977, 80.
- 20 J. C. Chen and E. S. Gould, *J. Amer. Chem. Soc.*, 1973, **95**, 5544.
- 21 A. Zwickel and H. Taube, *Discuss. Faraday Soc.*, 1960, **29**, 42.
- 22 M. R. Hyde, R. S. Taylor, and A. G. Sykes, *J.C.S. Dalton*, 1973, 2730.
- 23 G. A. K. Thompson and A. G. Sykes, *Inorg. Chem.*, 1976, **15**, 638; *ibid.*, 1979, **18**, 2025.
- 24 F. Basolo, M. L. Morris, and R. G. Pearson, *Discuss. Faraday Soc.*, 1960, **29**, 80.
- 25 M. Green, K. Schug, and H. Taube, *Inorg. Chem.*, 1965, **4**, 1184.
- 26 J. K. Beattie and J. Starink, *Inorg. Chem.*, 1975, **14**, 996.