

Additions and Corrections

Reductive Nitrosylation of Tetraoxometalates. Part 15. Generation of $\text{Mn}(\text{NO})^{3+}$, $\text{Mn}(\text{NO})^{2+}$, and $\text{Mn}(\text{NO})_2^{2+}$ Moieties Directly from MnO_4^- . Synthesis, Characterization, and Electrochemistry of Cyanonitrosyl Derivatives of Manganese (1989, 1963)

Ramgopal Bhattacharyya, Nabanita Ghosh, and Gobinda P. Bhattacharjee

Page 1964. In Scheme 1, step (vii) should read 'recrystallization from water-ethanol'.

Page 1965. In Table 1 the experimental %C analysis for compound (**10b**) should be 52.3, not 57.3.

Page 1967. Figure 2 should be replaced by the amended version below.




Figure 2. E.s.r. spectra of (a) $[\text{PPh}_4]_3[\text{Mn}_2(\text{NO})_2(\text{CN})_8] \cdot 3\text{H}_2\text{O}$ (**6b**), polycrystalline, 298 K (----), in frozen CH_3CN (—), and in frozen CH_3CN after coulometry (····), (b) $[\text{PPh}_4]_3[\text{Mn}_2(\text{NO})_2(\text{CN})_8] \cdot 3\text{H}_2\text{O}$ (**6a**), polycrystalline, 298 K (—), in frozen CH_3CN after coulometry (----), and (c) $[\text{NMe}_4]_3[\text{Mn}_2(\text{NO})_2(\text{CN})_8] \cdot 3\text{H}_2\text{O}$ (**7**), polycrystalline, 298 K (····).