

Synthesis of the homoleptic rhodium(III) complex $[\text{Rh}(\text{C}_6\text{Cl}_5)_3]$. Molecular structures of $[\text{Rh}(\text{C}_6\text{Cl}_5)_3]$ and $[\text{Rh}(\text{C}_6\text{Cl}_4\text{-C}_6\text{Cl}_4)(\text{C}_6\text{Cl}_5)_2(\text{SC}_4\text{H}_8)_2]$

María P. García, M. Victoria Jiménez, Fernando J. Lahoz, José A. López and Luis A. Oro

Departamento de Química Inorgánica, Instituto de Ciencia de Materiales de Aragón, Universidad de Zaragoza-C.S.I.C., 50009 Zaragoza, Spain

Received 19th August 1998, Accepted 27th October 1998

The arylation of $[\text{RhCl}_3(\text{SC}_4\text{H}_8)_3]$ (SC_4H_8 = tetrahydrothiophene, tht) with the Grignard reagent $\text{MgCl}(\text{C}_6\text{Cl}_5)$, under different reaction conditions, afforded the neutral rhodium(III) complexes $[\text{Rh}(\text{C}_6\text{Cl}_5)_3]$ **1** and $[\text{Rh}(\text{C}_6\text{Cl}_4\text{-C}_6\text{Cl}_4)(\text{C}_6\text{Cl}_5)_2(\text{tht})_2]$ **2** as well as the paramagnetic rhodium(II) compound $[\text{Rh}(\text{C}_6\text{Cl}_5)_2(\text{tht})_2]$ **3**. The homoleptic complex **1** can also be obtained by treatment of $[\text{P}(\text{CH}_2\text{Ph})\text{Ph}_3][\text{RhCl}(\text{C}_6\text{Cl}_5)_3]$ with AgClO_4 . The complexes are stable to the air and moisture in the solid state and moderately stable in deoxygenated solutions. The crystal structures of compounds **1** and **2** have been determined. They display distorted octahedral environments. All C_6Cl_5 groups in **1** or one C_6Cl_5 group in **2** act as chelating ligands being bonded through the *ipso*-C atom and one of the *ortho*-Cl atoms. Complex **2** additionally incorporates two tetrahydrothiophene ligands and a perchloro-2,2'-biphenylene ligand, formed by coupling of two pentachlorophenyl groups.

Introduction

The chemistry of perhalogenoaryl derivatives of rhodium and iridium has been a subject of our attention in recent years. In this area we have reported a variety of neutral, anionic, mono- and di-nuclear pentafluorophenyl complexes of rhodium and iridium in oxidation states III¹⁻⁵ and I.⁶ Interestingly, the related, but less frequently used, pentachlorophenyl ligand has allowed the stabilisation of mononuclear rhodium and iridium complexes^{7,8} in the rare oxidation state II, including the preparation of novel homoleptic complexes of formula $[\text{M}^{\text{II}}(\text{C}_6\text{Cl}_5)_4]^{2-}$ (M = Rh or Ir).^{9,10} A relevant feature, recently observed for some rhodium pentachlorophenyl complexes such as $[\text{Rh}^{\text{III}}(\text{C}_6\text{Cl}_5)_4]^-$, is the capability of the pentachlorophenyl group to present two co-ordination modes, the conventional σ bond and the unusual chelating bonding through the C_{ipso} atom and one of the *o*-Cl atoms.¹⁰

On the other hand, the preparation of aryl derivatives can be achieved by the reaction of the appropriate halogen starting metal complex with aryl-lithium or -magnesium compounds, but these reactions are very sensitive to the chosen conditions. For instance, the nature and yield of the products of the reactions of $[\text{RhCl}_3(\text{tht})_3]$ (tht = tetrahydrothiophene) with lithium derivatives depends on the experimental conditions and on the steric properties of the aryl or alkyl group used; so an excess of LiC_6Cl_5 leads¹⁰ to the preparation of the arylated rhodium(II) dianion $[\text{Rh}(\text{C}_6\text{Cl}_5)_4]^{2-}$ and with $\text{Li}(2,4,6\text{-}i\text{-Pr}_3\text{C}_6\text{H}_2)$ yields¹¹ the neutral square-planar complex $[\text{Rh}(2,4,6\text{-}i\text{-Pr}_3\text{C}_6\text{H}_2)_2(\text{tht})_2]$. During these arylating reactions a one-electron reduction process has taken place. However, with LiC_6F_5 it gives¹ the square-pyramidal rhodium(III) dianion $[\text{Rh}(\text{C}_6\text{F}_5)_3]^{2-}$, while LiMe affords¹² the octahedral $[\text{RhMe}_6]^{3-}$. Quite likely, in these reactions radical processes are involved.

In general, organomagnesium compounds are less reactive than organolithium compounds, but they are also less basic and poorer one-electron reducing agents, and so less prone to side-reactions arising from these properties. For these reasons, and following our interest in the chemistry of the pentachlorophenyl ligand, we have studied the arylation of $[\text{RhCl}_3(\text{tht})_3]$ with pentachlorophenylmagnesium chloride. We report here the synthesis and full characterisation of the new homoleptic $[\text{Rh}^{\text{III}}(\text{C}_6\text{Cl}_5)_3]$ compound, which constitutes the first metal complex

with three pentachlorophenyl groups acting as chelating ligands in a "RhC₃Cl₃" co-ordination environment, together with a related complex of formula $[\text{Rh}^{\text{III}}(\text{C}_6\text{Cl}_4\text{-C}_6\text{Cl}_4)(\text{C}_6\text{Cl}_5)_2(\text{tht})_2]$. The latter complex is the result of the unusual coupling of two pentachlorophenyl groups.

Results and discussion

The previously reported arylation of $[\text{RhCl}_3(\text{tht})_3]$ with the lithium derivative LiC_6Cl_5 afforded¹⁰ the paramagnetic rhodium(II) dianion $[\text{Rh}(\text{C}_6\text{Cl}_5)_4]^{2-}$. In contrast, the treatment of $[\text{RhCl}_3(\text{tht})_3]$ in THF with the Grignard reagent $\text{MgCl}(\text{C}_6\text{Cl}_5)$ (1:6 ratio, 4 h reflux), work-up and column chromatography of the reaction product allow the preparation of the neutral homoleptic rhodium(III) complex $[\text{Rh}(\text{C}_6\text{Cl}_5)_3]$ **1** (31% yield). The MS (FAB) of **1** shows the parent peak at m/z 851, together with the peak at m/z 601 corresponding to the $[\text{Rh}(\text{C}_6\text{Cl}_5)_2]$ fragment. The IR spectrum exhibits only absorptions due to the C_6Cl_5 groups.¹³ Its molecular geometry was established by an X-ray structural determination.

Trying to improve the yield in complex **1** in the Grignard reaction, the mixture of $[\text{RhCl}_3(\text{tht})_3]$ and $\text{MgCl}(\text{C}_6\text{Cl}_5)$ was left at reflux temperature for 18 h. After work-up, in similar way to that before, the yield in complex **1** was smaller (around 10%) but a new fraction was collected from the column containing the complex $[\text{Rh}(\text{C}_6\text{Cl}_4\text{-C}_6\text{Cl}_4)(\text{C}_6\text{Cl}_5)_2(\text{tht})_2]$ **2** (15% yield). So the coupling of two C_6Cl_5 groups has taken place with the formation of a new C-C σ bond.¹⁴ A similar process takes place when MCl_4 (M = Si or Ge) reacts with $\text{MgCl}(\text{C}_6\text{Cl}_5)$.¹⁵

A coproduct in the above syntheses is the paramagnetic mononuclear rhodium(II) complex $[\text{Rh}(\text{C}_6\text{Cl}_5)_2(\text{tht})_2]$ **3**, formed in relatively small yield (about 20%) and easily separated from the column by elution with CH_2Cl_2 . Complex **3** is paramagnetic and its X-band EPR spectrum at room temperature is analogous to those found for the neutral rhodium(II) complexes $[\text{Rh}(\text{C}_6\text{Cl}_5)_2\text{L}_2]$ [L_2 = cycloocta-1,5-diene (cod), $\{\text{P}(\text{O}^i\text{Pr})_3\}_2$, $\{\text{P}(\text{OMe})_3\}_2$, $\{\text{PPh}_3\}_2$, py_2 , 1,2-bis(diphenylphosphino)ethane (dppe), bis(diphenylphosphino)methane (dppm)].⁸ The values $g_x = 2.73$, $g_y = 2.60$ and $g_z = 1.95$ can be understood if the unpaired electron is mainly in a d_z orbital with the z axis perpendicular to the first co-ordination rhodium plane. No evidence of hyperfine structure has been found.




Fig. 1 Molecular diagram of the complex $[\text{Rh}(\text{C}_6\text{Cl}_5)_3]$ **1**, together with the numbering scheme used.




Fig. 2 Molecular diagram of the complex $[\text{Rh}(\text{C}_6\text{Cl}_4\text{-C}_6\text{Cl}_4)(\text{C}_6\text{Cl}_5)(\text{tht})_2]$ **2**, together with the numbering scheme used.

Complex **1** can also be obtained by reaction of $[\text{P}(\text{CH}_2\text{Ph})\text{Ph}_3][\text{RhCl}(\text{C}_6\text{Cl}_5)_3]$ with AgClO_4 ; AgCl and $[\text{P}(\text{CH}_2\text{Ph})\text{Ph}_3]\text{ClO}_4$ are separated and the solution provides $[\text{Rh}(\text{C}_6\text{Cl}_5)_3]$ (72% yield).

Molecular structures of $[\text{Rh}(\text{C}_6\text{Cl}_5)_3]$ **1** and $[\text{Rh}(\text{C}_6\text{Cl}_4\text{-C}_6\text{Cl}_4)(\text{C}_6\text{Cl}_5)(\text{tht})_2]$ **2**

The crystal structures of complexes $\mathbf{1} \cdot \frac{1}{2}\text{CH}_2\text{Cl}_2$ and **2** have been solved by single-crystal X-ray crystallography. The respective molecular structures of the complexes, including a conventional image of the displacement parameters, are presented in Figs. 1 and 2, together with the atomic labelling schemes used. Selected bond distances and angles are collected in Tables 1 and 2.

Although the complex **1** has only three ligands, the rhodium atom is in a distorted-octahedral environment formed by three Rh–C σ bonds from the three C_6Cl_5 ligands (in a *fac* disposition), and three Rh–Cl secondary bonds, each involving one *o*-chlorine of different pentachlorophenyl groups (Fig. 1). This arrangement of the perhalogenophenyl groups results in a helix structure with a pseudo- C_3 axis, making this complex chiral, but with both enantiomers present in the crystal struc-

Table 1 Selected bond lengths (\AA) and angles ($^\circ$) for $[\text{Rh}(\text{C}_6\text{Cl}_5)_3]$ **1**

Rh–Cl(2)	2.6171(16)	Rh–C(1)	1.986(6)
Rh–Cl(8)	2.5889(17)	Rh–C(7)	1.993(7)
Rh–Cl(14)	2.6020(18)	Rh–C(13)	1.994(6)
Cl(2)–C(2)	1.764(6)	Cl(8)–C(8)	1.763(7)
Cl(14)–C(14)	1.764(6)	mean Cl–C ^a	1.722(3)
Cl(2)–Rh–Cl(8)	99.76(5)	C(1)–Rh–C(7)	100.4(3)
Cl(2)–Rh–Cl(14)	107.70(6)	C(1)–Rh–C(13)	100.8(2)
Cl(8)–Rh–Cl(14)	96.99(6)	C(7)–Rh–C(13)	100.8(3)
Cl(2)–Rh–C(1)	67.84(18)	Cl(8)–Rh–C(1)	164.56(19)
Cl(2)–Rh–C(7)	86.31(18)	Cl(8)–Rh–C(7)	68.64(19)
Cl(2)–Rh–C(13)	167.73(18)	Cl(8)–Rh–C(13)	92.17(18)
Cl(14)–Rh–C(1)	95.66(18)	Rh–Cl(2)–C(2)	73.2(2)
Cl(14)–Rh–C(7)	161.84(19)	Rh–Cl(8)–C(8)	73.3(2)
Cl(14)–Rh–C(13)	67.6(2)	Rh–Cl(14)–C(14)	73.4(2)
Rh–C(1)–C(2)	105.8(4)	Rh–C(1)–C(6)	138.0(5)
Rh–C(7)–C(8)	104.3(5)	Rh–C(7)–C(12)	137.9(5)
Rh–C(13)–C(14)	105.7(4)	Rh–C(13)–C(18)	136.3(5)

^a Mean bond length of the non-co-ordinated chlorines.

Table 2 Selected bond lengths (\AA) and angles ($^\circ$) for $[\text{Rh}(\text{C}_6\text{Cl}_4\text{-C}_6\text{Cl}_4)(\text{C}_6\text{Cl}_5)(\text{tht})_2]$ **2**

Rh–S(1)	2.4275(12)	Rh–S(2)	2.4997(11)
Rh–Cl(2)	2.8682(14)	Rh–C(1)	2.046(4)
Rh–C(7)	1.990(4)	Rh–C(18)	2.020(4)
C(2)–Cl(2)	1.748(5)	mean Cl–C ^a	1.728(2)
C(12)–C(13)	1.472(5)		
S(1)–Rh–S(2)	82.11(4)	S(1)–Rh–Cl(2)	104.93(4)
S(1)–Rh–C(1)	167.74(14)	S(1)–Rh–C(7)	94.18(12)
S(1)–Rh–C(18)	90.86(11)	S(2)–Rh–Cl(2)	77.64(4)
S(2)–Rh–C(1)	97.47(12)	S(2)–Rh–C(7)	97.42(12)
S(2)–Rh–C(18)	172.76(11)	Cl(2)–Rh–C(1)	63.19(14)
Cl(2)–Rh–C(7)	159.24(12)	Cl(2)–Rh–C(18)	106.01(12)
C(1)–Rh–C(7)	98.02(18)	C(1)–Rh–C(18)	89.77(16)
C(7)–Rh–C(18)	81.31(16)	Rh–Cl(2)–C(2)	71.02(16)
Rh–C(1)–C(2)	109.5(3)	Rh–C(1)–C(6)	135.7(4)

^a Mean bond length of the non-co-ordinated chlorines.

Table 3 Metal complexes containing chelating pentachlorophenyl ligands

Complex	Number of C_6Cl_5 ligands			Ref.
	Total	Chelating	σ -Bonded	
$[\text{Rh}(\text{C}_6\text{Cl}_5)_3]$	3	3	0	This work
$[\text{Rh}(\text{C}_6\text{Cl}_4\text{-C}_6\text{Cl}_4)(\text{C}_6\text{Cl}_5)(\text{tht})_2]$	1	1	0	This work
$[\text{Rh}(\text{C}_6\text{Cl}_5)_4]$	4	2	2	10
$[\text{Rh}(\text{C}_6\text{Cl}_5)_3\text{Cl}]^-$	3	2	1	10
$[\text{Rh}\{\text{C}(\text{O})\text{C}_6\text{Cl}_5\}_2(\text{C}_6\text{Cl}_5)\text{Cl}]^-$	1	1	0	10
$[\text{Pt}(\text{C}_6\text{Cl}_5)_4]$	4	2	2	16
$[\text{Cr}(\text{C}_6\text{Cl}_5)_4]^-$	4	2	2	17

ture. This chelating co-ordination mode of the pentachlorophenyl groups has previously been reported for the anionic rhodium complexes¹⁰ $[\text{Rh}(\text{C}_6\text{Cl}_5)_4]^-$, $[\text{Rh}(\text{C}_6\text{Cl}_5)_3\text{Cl}]^-$ and $[\text{Rh}\{\text{C}(\text{O})\text{C}_6\text{Cl}_5\}_2(\text{C}_6\text{Cl}_5)\text{Cl}]^-$, for the platinum complex¹⁶ $[\text{Pt}(\text{C}_6\text{Cl}_5)_4]$ and, very recently, also for a chromium(III) complex¹⁷ $[\text{Cr}(\text{C}_6\text{Cl}_5)_4]^-$, but the present case is the first homoleptic one with all the C_6Cl_5 groups acting as chelate ligands (Table 3).

The deviation from a perfect octahedral co-ordination arises fundamentally from the chelating co-ordination of the pentachlorophenyl ligands, which form strained four-membered metallacycles. This distortion of the metal environment can be visualised in the Cl–Rh–C bite angles, 67.84(18), 68.64(19) and 67.6(2) $^\circ$, and is the origin of an asymmetric co-ordination of the phenyl rings, which are tilted to permit the approach of the *o*-chlorine atoms to the metal centre. This tilting can be charac-

terised with the angles around the C_{ipso} carbons (see Table 1), or in terms of the value of the angle between the Rh–C bond and the C_{ipso} – C_{para} direction (the pseudo-2-fold axis of the ligand), 162.9, 160.9 and 162.9(4)° around C(1), C(7) and C(13), respectively.

The lengths observed for these Rh–*o*-Cl bonds [2.6171(16), 2.5889(17) and 2.6020(18) Å for Cl(2), Cl(8) and Cl(14), respectively] are, as expected, significantly longer than the usual Rh^{III} –Cl distances (mean value 2.374 Å),¹⁸ but short enough to be considered as secondary bonds.¹⁰ Furthermore, they are shorter than the secondary bonds described in the previously mentioned rhodium complexes [range 2.6277(11)–2.8863(9) Å]. Interestingly, in compound **1** all the chelating chlorines are *trans* disposed to the C_{ipso} atom of another C_6Cl_5 ring, as is also described for $[Rh(C_6Cl_5)_3Cl]^-$ and for the homoleptic complexes $[M(C_6Cl_5)_4]^-$ (M = Rh or Cr) and $[Pt(C_6Cl_5)_4]$.

In compound **2** the metal centre also exhibits a distorted octahedral environment, completing the co-ordination sphere with three different type of ligands: two tetrahydrothiophene ligands, an octachlorobiphenylene chelating ligand and a C_6Cl_5 ring linked to the metal through a Rh–C σ bond and through a Rh–Cl secondary bond. In this complex the distortion is also due to the small bite angle of the C_6Cl_5 chelating group [63.19(14)°] and, to a lesser extent, of the perhalogenobiphenylene ligand [81.31(16)°]. The tht ligands are in a *cis* disposition, the Rh–*o*-Cl bond being *trans* to a carbon atom of the perhalogenobiphenylene ligand.

The perchloro-2,2'-biphenylene group presents a considerable twist between the two phenyl rings [torsion angle C(11)–C(12)–C(13)–C(14) 36.3(7)°] due to the steric hindrance of Cl(11) and Cl(14). This conformation is similar to that reported for the compounds $[M(C_6Cl_4-C_6Cl_4)(C_6Cl_5)_2]$ (M = Si or Ge)¹⁵ with torsion angles of about 34°. The two Rh–C bond lengths in this ligand are slightly different [1.990(4) and 2.020(4) Å], probably due to the different *trans* atoms; the long one corresponds to the bond *trans* to the sulfur atom and is very similar to the Rh–C distance for the C_6Cl_5 group [2.046(4) Å], also *trans* to a tht ligand. The short distance, *trans* to an *o*-Cl, is equal to those calculated in **1** for Rh– C_6Cl_5 bonds, also *trans* to an *o*-chlorine.

The Rh–Cl bond [2.8682(14) Å] is longer than those observed for **1** [2.6171(16), 2.5889(17) and 2.6020(18) Å], but similar to those reported for $[Rh\{C(O)C_6Cl_5\}_2(C_6Cl_5)Cl]^-$ [2.8863(9) Å]. In fact there seems to be a correlation between the number of chelating C_6Cl_5 groups and the length of these Rh–*o*-Cl bonds. The shorter distances correspond to the compound with three chelating C_6Cl_5 rings (**1**, mean 2.603 Å), there are intermediate values in the complexes with two chelating C_6Cl_5 ligands $\{[Rh(C_6Cl_5)_4]^-$, 2.6277 and 2.7521(11) Å; $[Rh(C_6Cl_5)_3Cl]^-$, 2.656 and 2.678(4) Å; mean value 2.688 Å} and longer distances in the compounds with one chelating C_6Cl_5 ligand (**2**, 2.8682(14) Å; $[Rh\{C(O)C_6Cl_5\}_2(C_6Cl_5)Cl]^-$, 2.8863(9) Å; mean value 2.881 Å).

Experimental

General

Analyses (C, H and S) and IR spectra (in Nujol suspension between polyethylene sheets) were obtained as described elsewhere.⁸ Mass spectra were measured in a VG Autospec double-focusing mass spectrometer operating in the positive mode; ions were produced with the standard Cs^+ gun at ca. 30 kV; 3-nitrobenzyl alcohol was used as matrix; high-resolution FAB spectra are in accordance with the simulated isotopic pattern distribution. The starting rhodium(III) compound $[RhCl_3(tht)_3]$ was prepared as described in the literature.¹⁹ The compound $MgCl(C_6Cl_5)$ was obtained by treating C_6Cl_6 with Mg, in tetrahydrofuran, using I_2 as the initiator, and keeping the mixture stirred at reflux temperature for 3 h; it was directly used with-

out further isolation. Diethyl ether, THF, CH_2Cl_2 and hexane were distilled under nitrogen from the appropriate drying agents.

Preparations

[Rh(C_6Cl_5)₃] **1**. *Method a*. Solid $[RhCl_3(tht)_3]$ (0.500 g, 1.05 mmol) was added to a tetrahydrofuran solution (30 mL) of $MgCl(C_6Cl_5)$ (6.33 mmol) and stirred for 4 h at reflux temperature. The mixture was slowly allowed to reach room temperature; the excess of Grignard compound was hydrolysed with aqueous diethyl ether and then evaporated till dryness. The residue was extracted with CH_2Cl_2 (50 mL), filtered and the filtrate concentrated to a few millilitres. Addition of diethyl ether caused the precipitation of a brown solid. This solid was chromatographed on a silica gel column (2.5 × 25 cm) using hexane as stationary phase. Elution with hexane afforded a yellow band; the hexane was evaporated till dryness and the residue dissolved in CH_2Cl_2 ; addition of methanol gave the yellow microcrystalline solid $[Rh(C_6Cl_5)_3]$ **1** (31% yield). Elution with CH_2Cl_2 yield another yellow band, subsequently identified as $[Rh(C_6Cl_5)_2(tht)_2]$ **3**.

Method b. To a CH_2Cl_2 solution of $[P(CH_2Ph)Ph_3][RhCl(C_6Cl_5)_3]$ (0.100 g, 0.08 mmol) was added solid $AgClO_4$ (16.6 mg, 0.08 mmol). The solution was stirred for 1 h under a nitrogen atmosphere and with exclusion of light to yield a white solid ($AgCl$ and $[P(CH_2Ph)Ph_3]ClO_4$) which was removed by filtration through Kieselguhr. Concentration of the resulting solution under reduced pressure to a volume of 1 mL and addition of methanol (10 mL) afforded **1** as a yellow solid which was filtered off, washed with methanol and dried under vacuum. The compound was recrystallised from CH_2Cl_2 –methanol (72% yield) (Found: C, 25.37. Calc. for $C_{18}Cl_{15}Rh$: C, 25.40%). IR (Nujol; cm^{-1}): $\nu(C_6Cl_5)$ 1345, 1313, 1279, 831 (X-sensitive), 679, 637 [$\nu(Rh-C)$]. MS [FAB; m/z (%): 851 (39) [M^+ , $Rh(C_6Cl_5)_3$], 601 (51) [$Rh(C_6Cl_5)_2$].

[Rh($C_6Cl_4-C_6Cl_4$)(C_6Cl_5)(tht)₂] **2**. The preparation and isolation of complex **2** was similar to that described for **1**, in Method a, but keeping the Grignard reaction at reflux temperature for 18 h. The silica gel column chromatography (2.5 × 25 cm) of the reaction product using hexane as stationary phase gave two yellow bands. Concentration of the hexane solutions provided complexes **1** and **2**. Data for **2**: yield 15% (Found: C, 32.38; H, 1.62; S, 6.40. Calc. for $C_{26}H_{16}Cl_{13}RhS_2$: C, 32.65; H, 1.69; S, 6.70%). IR (Nujol; cm^{-1}) $\nu(C_6Cl_5)$ 1307, 1298, 1289, 1280, 795 and 674; MS [FAB; m/z (%)] 781 (30) [$Rh(C_6Cl_4-C_6Cl_4)(C_6Cl_5)$].

[Rh(C_6Cl_5)₂(tht)₂] **3**. Elution of the column of the previous synthesis with CH_2Cl_2 gave an orange band; concentration of the CH_2Cl_2 solution and addition of methanol gave an orange solid identified as $[Rh(C_6Cl_5)_2(tht)_2]$ **3**. Yield: 20% (Found: C, 30.95; H, 1.97; S, 8.20. Calc. for $C_{20}H_{16}Cl_{10}RhS_2$: C, 30.88; H, 2.07; S, 8.24%). IR (Nujol; cm^{-1}) $\nu(C_6Cl_5)$ 1310, 1290, 820, 665 and 602.

Crystal structure determination of $[Rh(C_6Cl_5)_3] \cdot \frac{1}{2}CH_2Cl_2$, $1 \cdot \frac{1}{2}CH_2Cl_2$ and $[Rh(C_6Cl_4-C_6Cl_4)(C_6Cl_5)(tht)_2]$ **2**

A summary of crystal data and refinement parameters is reported in Table 4. Data were collected on a Siemens-Stoe AED-2 diffractometer with graphite-monochromated Mo- $K\alpha$ radiation, using the ω – 2θ scan method. Three standard reflections were monitored every 55 min throughout data collection; no important variations were observed. All data were corrected for Lorentz-polarisation effects, and for absorption using a semiempirical method (ψ scan);²⁰ minimum and maximum transmission factors are listed in Table 4. All structures were solved by direct methods (SIR 92)²¹ and Fourier techniques, and refined by full-matrix least squares on F^2 (SHELXL 97).²²

Table 4 Crystallographic data for $[\text{Rh}(\text{C}_6\text{Cl}_5)_3]_2\cdot\frac{1}{2}\text{CH}_2\text{Cl}_2$ **1**· $\frac{1}{2}\text{CH}_2\text{Cl}_2$ and $[\text{Rh}(\text{C}_6\text{Cl}_4\text{-C}_6\text{Cl}_4)(\text{C}_6\text{Cl}_5)(\text{tht})_2]$ **2**

	1 · $\frac{1}{2}\text{CH}_2\text{Cl}_2$	2
Crystal color and habit	Yellow, prismatic block	Orange, prismatic block
Crystal size/mm	0.14 × 0.21 × 0.35	0.16 × 0.21 × 0.45
Formula	$\text{C}_{18.5}\text{HCl}_{16}\text{Rh}$	$\text{C}_{26}\text{H}_{16}\text{Cl}_{13}\text{RhS}_2$
<i>M</i>	893.30	956.27
Crystal system	Triclinic	Monoclinic
Space group	$P\bar{1}$ (no. 2)	$P2_1/n$ (no. 14)
<i>T</i> /K	233(2)	295(2)
<i>a</i> /Å	10.8207(11)	9.2192(5)
<i>b</i> /Å	10.8539(15)	32.660(2)
<i>c</i> /Å	14.0152(15)	11.3747(7)
<i>a</i> °	108.608(5)	—
<i>β</i> °	102.214(4)	105.805(6)
<i>γ</i> °	106.976(6)	—
<i>V</i> /Å ³	1405.2(3)	3295.4(3)
<i>Z</i>	2	4
<i>μ</i> /mm ⁻¹	2.142	1.721
<i>θ</i> range/°	1.6–23.0	1.9–25.0
Collected reflections	4173	6264
Unique reflections	3925 (<i>R</i> _{int} = 0.020)	5795 (<i>R</i> _{int} = 0.018)
Minimum, maximum transmission factors	0.8146, 1.0000	0.5147, 0.5625
Data/restraints/parameters	3925/7/329	5795/28/399
<i>R</i> (<i>F</i>) [<i>F</i> ² > 2σ(<i>F</i> ²)] ^a	0.0472	0.0364
<i>wR</i> (<i>F</i> ²) [all data] ^b	0.1306	0.0913

^a $R(F) = \sum ||F_o| - |F_c|| / \sum |F_o|$, for 3261 and 4613 observed reflections, respectively. ^b $wR(F^2) = [\sum w(F_o^2 - F_c^2)^2 / \sum w(F_o^2)^2]^{1/2}$.

The function minimised was $\sum [w(F_o^2 - F_c^2)^2]$ with the weight defined as $w^{-1} = [\sigma^2(F_o^2) + (xP)^2 + yP]$ where $P = (F_o^2 + 2F_c^2)/3$. Atomic scattering factors, corrected for anomalous dispersion, were used as implemented in the refinement program.²²

Complex 1· $\frac{1}{2}\text{CH}_2\text{Cl}_2$. Yellow prismatic crystals suitable for the X-ray diffraction study were obtained from a CH_2Cl_2 –methanol solution of complex **1**. Anisotropic displacement parameters were used in the last cycles of refinement for all atoms, except for the dichloromethane solvent (one half per asymmetric unit), which was two-fold disordered and refined with geometrical restraints, free but complementary occupancy factors [0.277 and 0.224(5)], and two refined isotropic displacement parameters. The refinement converged to $R(F) = 0.0472$ [$F^2 > 2\sigma(F^2)$] and $wR(F^2) = 0.1306$ (all data), with weight parameters $x = 0.0787$ and $y = 3.6004$. Residual peaks in the final difference map were 1.74 and $-0.99 \text{ e } \text{Å}^{-3}$.

Complex 2. Suitable crystals were obtained from a CH_2Cl_2 –methanol solution. After refinement of all the non-hydrogen atoms with anisotropic displacement parameters, the hydrogen atoms of one tht ligand were introduced in calculated positions and refined riding on the corresponding carbon atoms; the hydrogens of the other tht ligand were not introduced because two carbons of this group were two-fold disordered [occupancy factors 0.53 and 0.47(3)]. The refinement converged to $R(F) = 0.0364$ and $wR(F^2) = 0.0913$, with weight parameters $x = 0.0334$ and $y = 4.3269$. Residual peaks in the final difference map were 0.45 and $-0.34 \text{ e } \text{Å}^{-3}$.

CCDC reference number 186/1218.

See <http://www.rsc.org/suppdata/dt/1998/4211/> for crystallographic files in .cif format.

Acknowledgements

We thank Professor Dr P. J. Alonso for the measurement of the EPR spectrum and the DGICYT (Dirección General de Investigación Científica y Técnica) for financial support (project PB95-0318).

References

- M. P. García, L. A. Oro and F. J. Lahoz, *Angew. Chem.*, 1988, **100**, 1766.

- M. P. García, M. V. Jiménez, F. J. Lahoz, L. A. Oro, A. Tiripicchio and J. A. López, *J. Chem. Soc., Dalton Trans.*, 1990, 1503.
- M. P. García, M. V. Jiménez, F. J. Lahoz and L. A. Oro, *J. Chem. Soc., Dalton Trans.*, 1995, 917.
- M. P. García, M. V. Jiménez and L. A. Oro, *J. Organomet. Chem.*, 1992, **438**, 229.
- M. P. García, M. V. Jiménez, T. Luengo and L. A. Oro, *J. Organomet. Chem.*, 1996, **510**, 189.
- M. P. García, M. V. Jiménez, F. J. Lahoz and L. A. Oro, *Inorg. Chem.*, 1995, **34**, 2153.
- M. P. García, M. V. Jiménez, F. J. Lahoz, L. A. Oro and P. J. Alonso, *Angew. Chem.*, 1992, **104**, 1512.
- M. P. García, M. V. Jiménez, F. J. Lahoz, L. A. Oro, J. M. Casas and P. J. Alonso, *Organometallics*, 1993, **12**, 3257.
- M. P. García, M. V. Jiménez, L. A. Oro, F. J. Lahoz, M. Camellini Tiripicchio and A. Tiripicchio, *Organometallics*, 1993, **12**, 4660.
- M. P. García, M. V. Jiménez, A. Cuesta, C. Siurana, L. A. Oro, F. J. Lahoz, J. A. López, M. P. Catalán, A. Tiripicchio and M. Lanfranchi, *Organometallics*, 1997, **10**, 1026.
- R. S. Hay-Motherwell, S. U. Koschmieder, G. Wilkinson, B. Hussain-Bates and M. B. Hursthouse, *J. Chem. Soc., Dalton Trans.*, 1991, 2821.
- R. S. Hay-Motherwell, G. Wilkinson, B. Hussain-Bates and M. B. Hursthouse, *J. Chem. Soc., Chem Commun.*, 1989, 1436; *Polyhedron*, 1990, 2071.
- E. Maslowsky, *Vibrational Spectra of Organometallics Compounds*, Wiley, New York, 1977, p. 437.
- B. J. Wakefield, *Organomagnesium Methods in Organic Synthesis*, Academic Press, London, 1995, ch. 8, pp. 151–160.
- L. Fajari, L. Juliá, J. Riera, E. Molins and C. Miravittles, *J. Organomet. Chem.*, 1990, **381**, 321.
- J. Forniés, B. Menjón, R. M. Sanz-Carrillo, M. Tomás, N. G. Connelly, J. G. Crossley and A. G. Orpen, *J. Am. Chem. Soc.*, 1995, **117**, 4295.
- P. J. Alonso, L. R. Falvello, J. Forniés, M. A. García-Monforte, A. Martín, B. Menjón and G. Rodríguez, *Chem. Commun.*, 1998, 1721.
- A. G. Orpen, L. Brammer, F. H. Allen, O. Kennard, D. G. Watson and R. Taylor, in *International Tables for Crystallography*, Kluwer, Dordrecht, 1992, vol. C, p. 707.
- E. A. Allen and G. Wilkinson, *J. Chem. Soc., Dalton Trans.*, 1972, 613.
- A. C. T. North, D. C. Phillips and F. S. Mathews, *Acta Crystallogr., Sect. A*, 1968, **24**, 351.
- A. Altomare, G. Cascarano, C. Giacovazzo and A. Guagliardi, *J. Appl. Crystallogr.*, 1994, **27**, 435.
- G. M. Sheldrick, SHELXL 97; University of Göttingen, 1997.