

This article was downloaded by:

On: 25 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Liquid Crystals

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713926090>

Photochromic chiral liquid crystalline systems containing spiro-oxazine with a chiral substituent I. Synthesis and characterization of compounds

Hideshi Hattori; Toshiyuki Uryu

Online publication date: 06 August 2010

To cite this Article Hattori, Hideshi and Uryu, Toshiyuki(2011) 'Photochromic chiral liquid crystalline systems containing spiro-oxazine with a chiral substituent I. Synthesis and characterization of compounds', *Liquid Crystals*, 28: 5, 809

To link to this Article: DOI: 10.1080/02678290110039462

URL: <http://dx.doi.org/10.1080/02678290110039462>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Corrigendum

Photochromic chiral liquid crystalline systems containing spiro-oxazine with a chiral substituent

I. Synthesis and characterization of compounds

HIDESHI HATTORI


Central Research Institute, Dai Nippon Printing Co. Ltd., 250-1 Wakashiba,
Kashiwa, Chiba 277-0871, Japan

and TOSHIYUKI URYU

Department of Environmental and Material Engineering,
Teikyo University of Science and Technology, 2525 Yatsuzawa, Uenohara-cho,
Yamanashi 409-0193, Japan

Scheme 1 in this earlier published paper (2001, *Liq. Cryst.*, **28**, 24–34) contains erroneous structures, as described below:

Each oxazine ring in the (*R*)-spiro-oxazine racemic mixture, and in the (*R*, *X**)-spiro-oxazine optically active mixture, erroneously depicts [spiro-carbon]–N=CH– bonds. These should have been drawn as [spiro-carbon]–CH=N– bonds.



Scheme 1. Schematic representation of the concept.