

On the Sum of the Series $\sum_{\nu=0}^{\infty} (t^{\nu}/[u\nu + m]!)$

By Bogumił Leszczyński

An estimation of error in summing the series

$$(1) \quad \sum_{\nu=0}^{\infty} \frac{t^{\nu}}{\Gamma(\beta\nu + 2)}, \quad 0 < \beta < 1,$$

obtained by a solution of Volterra's equation

$$(2) \quad \phi(x) = ax + b \int_0^x (x-z)^{\beta-1} \phi(z) dz, \quad a = \text{const}, b = \text{const},$$

leads to series of the form

$$(3) \quad S_{u,m}(t) = \sum_{\nu=0}^{\infty} \frac{t^{\nu}}{[u\nu + m]},$$

where $[x]$ denotes the largest integer no greater than the real number x , m a non-negative integer, and u real, positive and rational.

The series (3) is absolutely and uniformly convergent for all real values of t and therefore not dependent on the sequence of terms. Changing the sequence of terms in (3) we obtain

$$(4) \quad S_{u,m}(t) = \sum_{j=0}^{q-1} \sigma_{u,m}^j(t),$$

where

$$(5) \quad \sigma_{u,m}^j(t) = \sum_{i=0}^{\infty} \frac{t^{qi+j}}{[uj + m + pi]},$$

and

$$(6) \quad u = \frac{p}{q},$$

where p and q are positive integers with greatest common divisor equal to 1. Substituting in (5)

$$(7) \quad z^p = t^q, *$$

we have

$$(8) \quad \sigma_{u,m}^j(t) = t^{j-[uj+m]/u} s_{u,m}^j(z),$$

where

$$(9) \quad s_{u,m}^j(z) = \sum_{i=0}^{\infty} \frac{z^{[uj+m+pi]}}{[uj + m + pi]}.$$

Differentiating (9) $[uj + m]$ times with respect to z we obtain

$$(10) \quad s(z) = \frac{d^{[uj+m]} s_{u,m}^j(z)}{dz^{[uj+m]}} = \sum_{i=0}^{\infty} \frac{z^{pi}}{(pi)!}.$$

Received December 21, 1964.

* z is an imaginary number when q is odd and p even.

But (10) is the solution of the p th order linear differential equation

$$(11) \quad s^{(p)}(z) = s(z)$$

with initial conditions

$$(12) \quad s(0) = 1, \quad s^{(k)}(0) = 0 \quad (k = 1, 2, \dots, p-1).$$

The characteristic equation of (12) is

$$(13) \quad r^p - 1 = 0.$$

If r denotes a primitive root of p th order of unity, the general solution of (11) has the form

$$(14) \quad s(z) = \sum_{l=1}^p A_l \exp r^l z,$$

and taking (12) into consideration we have

$$(15) \quad \begin{aligned} \sum_{l=1}^p A_l &= 1, & A_l &= \text{const}, \\ \sum_{l=1}^p A_l r^{lk} &= 0 & (k &= 1, 2, \dots, p-1). \end{aligned}$$

For finding A_l we use Cramer's rule, and since there are Vandermonde's determinants in our case

$$(16) \quad A_l = \frac{\prod_{k=2}^p \prod_{n=1}^{k-1} (b_k - b_n)}{\prod_{k=2}^p \prod_{n=1}^{k-1} (r^k - r^n)}, \quad \text{where } b_n = \begin{cases} r^n & \text{when } n \neq l, \\ 0 & \text{when } n = l, \end{cases}$$

or

$$(17) \quad A_l = \prod_{k=1; k \neq l}^p \frac{1}{1 - r^{l-k}} = \prod_{k=1}^{p-1} \frac{1}{1 - r^k},$$

it follows that all A_l are equal, and, thus, from (15),

$$(18) \quad A_l = \frac{1}{p}.$$

Putting (18) into (14) we have

$$(19) \quad s(z) = \frac{1}{p} \sum_{l=1}^p \exp(r^l z).$$

Integrating both sides of (19) $[uj + m]$ times in the interval $(0, z)$ we obtain

$$(20) \quad s_{u,m}^j(z) = \frac{1}{p} \sum_{l=1}^p \exp(r^l z) r^{-l[uj+m]} - B,$$

where

$$(21) \quad B = \frac{1}{p} \sum_{k=1}^{[uj+m]} \frac{z^{[uj+m]-k}}{[uj+m-k]!} \sum_{l=1}^p r^{-lk}.$$

But

$$(22) \quad \sum_{l=1}^p r^{-lk} = \sum_{l=1}^p r^{lk} = \begin{cases} 0 & \text{when } k \text{ is divisible by } p, \\ p & \text{when } k \text{ is not divisible by } p, \end{cases}$$

and therefore

$$(23) \quad B = \sum_{k=1}^{[(uj+m)/p]} \frac{z^{[uj+m]-pk}}{[uj + m - pk]!}.$$

Returning to the variable t we obtain finally

$$(24) \quad S_{u,m}(t) = \sum_{j=0}^{q-1} t^{j-[uj+m]/u} \cdot \left\{ \frac{1}{p} \sum_{l=1}^p \exp(r^l t^u) r^{-l[uj+m]} - \sum_{k=1}^{[(uj+m)/p]} \frac{t^{[uj+m-pk]/u}}{[uj + m - pk]!} \right\}.$$

Warsaw Agricultural University
 Chair of Mathematics
 ul. Rakowiecka 8
 Warsaw, Poland