

# Runge-Kutta Methods with Constrained Minimum Error Bounds\*

By Richard King

**Abstract.** Optimum Runge-Kutta methods of orders  $m = 2, 3$ , and 4 are developed for the differential equation  $y' = f(x, y)$  under Lotkin's conditions on the bounds for  $f$  and its partial derivatives, and with the constraint that the coefficient of  $\partial^m f / \partial x^m$  in the leading error term be zero. The methods then attain higher order when it happens that  $f$  is independent of  $y$ .

**1. Introduction.** Anthony Ralston in [3] developed optimum Runge-Kutta methods of orders two, three, and four for a single first-order differential equation  $y' = f(x, y)$ . They are best in the sense that in each case the sum of the magnitudes of the coefficients in the leading truncation error term assumes a minimum under the following conditions: in the region of interest,

$$(1.1) \quad |f(x, y)| < M \quad \text{and} \quad \left| \frac{\partial^{i+j} f}{\partial x^i \partial y^j} \right| < L^{i+j} / M^{j-1},$$

where  $M$  and  $L$  are constants and  $i + j \leq m$ . These are the conditions used by Lotkin in [2]. Here, using Ralston's notation, the solution is to be advanced from  $x_0$  to  $x_1, x_2, \dots$  by the  $m$ th-order Runge-Kutta approximation

$$(1.2) \quad y_{n+1} = y_n + \sum_{i=1}^m w_i k_i,$$

where  $y_n = y(x_n)$ , the  $w_i$  are constants,

$$(1.3) \quad k_i = hf \left( x_n + \alpha_i h, y_n + \sum_{j=1}^{i-1} \beta_{ij} k_j \right),$$

and  $h = x_{n+1} - x_n$ . For each such approximation, it turns out that  $\alpha_1 = 0$  and

$$(1.4) \quad \alpha_i = \sum_{j=1}^{i-1} \beta_{ij}, \quad i = 2, 3, \dots, m.$$

The leading truncation error term,  $Ek^{m+1}$ , then satisfies

$$(1.5) \quad |Ek^{m+1}| < cML^m h^{m+1};$$

Ralston minimized  $c$  as a function of the parameters to be determined. Other measures of the truncation error have been considered by Hull and Johnston [1].

Our purpose is to find optimum methods of orders  $m = 2, 3$ , and 4 which will attain higher order when it happens that  $f$  is independent of  $y$ . This requires that in each case the coefficient of  $\partial^m f / \partial x^m$  (and, in fact, of  $D^m f$ ) in the leading error term be zero; here  $D$  is defined as

$$(1.6) \quad D = \partial / \partial x + f_n \partial / \partial y, \quad f_n = f(x_n, y_n),$$

and

---

Received November 17, 1965.

\* Work performed under the auspices of the U.S. Atomic Energy Commission.

$$(1.7) \quad D^s = \sum_{k=0}^s \binom{s}{k} f^k \partial^s / \partial x^{s-k} \partial y^k.$$

Furthermore, the vanishing of the term involving  $D^m f$  implies that conditions (1.1) need only be satisfied for  $i + j \leq m - 1$ .

**2. Second-order Methods.** The coefficient of  $h^3$  in the error function is

$$(2.1) \quad E = [\frac{1}{6} - (\alpha_2^2 w_2 / 2)] D^2 f + [\frac{1}{6}] f_y D f.$$

Equating the coefficient of  $D^2 f$  to zero yields

$$(2.2) \quad \alpha_2 = \frac{2}{3}, \quad \beta_{21} = \frac{2}{3}, \quad w_1 = \frac{1}{4}, \quad w_2 = \frac{3}{4},$$

so that the procedure becomes

$$(2.3) \quad y_{n+1} - y_n = (\frac{1}{4}) h f(x_n, y_n) + (\frac{3}{4}) h f(x_n + (\frac{2}{3}) h, y_n + (\frac{2}{3}) h f_n).$$

This is the same as Ralston's second-order method, and the truncation error is

$$(2.4) \quad | E h^3 | < (\frac{1}{3}) M L^2 h^3.$$

In this instance, no minimization problem appears. For  $f$  independent of  $y$  the procedure becomes Radau quadrature of order 3.

**3. Third-order Methods.** Here the coefficient of  $h^4$  in the error function is given by

$$(3.1) \quad E = a_1 D^3 f + a_2 f_y D^2 f + a_3 D f D f_y + a_4 f_y^2 D f,$$

where

$$(3.2) \quad \begin{aligned} a_1 &= \frac{1}{4!} - \frac{1}{3!} (\alpha_2^3 w_2 + \alpha_3^3 w_3), \\ a_2 &= \frac{1}{4!} - \frac{1}{2!} \alpha_2^2 \beta_{32} w_3, \\ a_3 &= \frac{3}{4!} - \alpha_2 \alpha_3 \beta_{32} w_3, \\ a_4 &= \frac{1}{4!}. \end{aligned}$$

But the vanishing of  $a_1$  implies that

$$(3.3) \quad 6\alpha_2\alpha_3 - 4(\alpha_2 + \alpha_3) + 3 = 0.$$

Along this hyperbola the error is bounded as follows:

$$(3.4) \quad | E | < [ | a_2 | + | 2a_2 + a_3 | + | a_2 + a_3 | + 2 | a_3 | + 2 | a_4 | ] M L^3,$$

with

$$(3.5) \quad \begin{aligned} a_2 &= \frac{1}{24} - \frac{\alpha_2}{12}, \\ a_3 &= \frac{1}{8} - \frac{\alpha_3}{6}, \\ a_4 &= \frac{1}{24}. \end{aligned}$$

If we now substitute (3.3) into (3.4) and minimize the right-hand side of (3.4) (as a function of  $\alpha_2$  or of  $\alpha_3$ ), we get  $\alpha_2 = \frac{1}{3}$ ,  $\alpha_3 = \frac{5}{6}$ . With these parameter values the suggested procedure becomes

$$(3.6) \quad y_{n+1} - y_n = \frac{1}{10}k_1 + \frac{1}{2}k_2 + \frac{2}{5}k_3,$$

where

$$(3.7) \quad \begin{aligned} k_1 &= hf(x_n, y_n), \\ k_2 &= hf(x_n + \frac{1}{3}h, y_n + \frac{1}{3}k_1), \\ k_3 &= hf(x_n + \frac{5}{6}h, y_n - \frac{5}{12}k_1 + \frac{5}{4}k_2). \end{aligned}$$

The resulting bound on  $Ek^4$  is

$$(3.8) \quad |Ek^4| < .1389ML^3h^4,$$

compared with  $.1111ML^3h^4$ , in Ralston's third-order procedure.

But if  $f$  is independent of  $y$ , then the procedure is fourth-order instead of third and the error bound is

$$(3.9) \quad |Ek^5| < 3.858 \times 10^{-5}ML^4h^5.$$

**4. Fourth-order Methods.** If we set to zero the coefficient

$$(4.1) \quad b_1 = \frac{1}{120} - \frac{1}{24}(\alpha_2^4 w_2 + \alpha_3^4 w_3 + w_4)$$

of  $D^4f$  in the leading error term, we again get an hyperbola in  $\alpha_2$  and  $\alpha_3$ :

$$(4.2) \quad 10\alpha_2\alpha_3 - 5(\alpha_2 + \alpha_3) + 3 = 0.$$

Along this curve  $b_3 = -4b_1$  vanishes also. The elements in

$$(4.3) \quad \begin{aligned} |E| < [8|b_2| + 8|b_4| + |b_5| + |2b_5 + b_7| + |b_5 + b_6 + b_7| \\ + |b_6| + |2b_6 + b_7| + |b_7| + 2|b_8|]ML^4 \end{aligned}$$

then become (see [1], p. 307)

$$(4.4) \quad \begin{aligned} b_2 &= \frac{5\alpha_3 - 3}{240}, & b_4 &= \frac{\alpha_3 - 1}{240(2\alpha_3 - 1)}, & b_5 &= -b_4 = \frac{1 - \alpha_3}{240(2\alpha_3 - 1)}, \\ b_6 &= \frac{(250\alpha_3^4 - 300\alpha_3^3 + 10\alpha_3^2 + 93\alpha_3 - 27)}{[(240)(10\alpha_3^2 - 12\alpha_3 + 3)]}, \\ b_7 &= \frac{2 - 5\alpha_3}{120}, & b_8 &= \frac{1}{120}. \end{aligned}$$

Minimizing the right-hand side of (4.3) along the hyperbola (4.2), we get

$$(4.5a) \quad \alpha_2 = \frac{4 - (6)^{1/2}}{10} = .1550510257, \quad \alpha_3 = \frac{4 + (6)^{1/2}}{10} = .6449489743,$$

so that

$$\begin{aligned}
 (4.5b) \quad & w_1 = 0, \quad w_2 = \frac{16 - (6)^{1/2}}{36}, \quad w_3 = \frac{16 + (6)^{1/2}}{36}, \quad w_4 = \frac{1}{9}, \\
 & \alpha_4 = 1, \quad \beta_{21} = \frac{4 - (6)^{1/2}}{10}, \quad \beta_{31} = -\left(\frac{11 + 4(6)^{1/2}}{25}\right), \\
 & \beta_{32} = \frac{42 + 13(6)^{1/2}}{50}, \quad \beta_{41} = \frac{1 + 5(6)^{1/2}}{4}, \\
 & \beta_{42} = -\left(\frac{3 + 2(6)^{1/2}}{2}\right), \quad \beta_{43} = \frac{9 - (6)^{1/2}}{4}.
 \end{aligned}$$

This defines the following Runge-Kutta scheme:

$$(4.6) \quad y_{n+1} - y_n = .3764030627k_2 + .5124858262k_3 + .1111111111k_4,$$

with

$$\begin{aligned}
 (4.7) \quad & k_1 = hf(x_n, y_n), \\
 & k_2 = hf(x_n + .1550510257h, y_n + .1550510257k_1), \\
 & k_3 = hf(x_n + .6449489743h, y_n - .8319183588k_1 + 1.476867333k_2), \\
 & k_4 = hf(x_n + h, y_n + 3.311862178k_1 - 3.949489743k_2 + 1.637627564k_3).
 \end{aligned}$$

The error bound is

$$(4.8) \quad |Eh^5| < \left(\frac{11 + 14(6)^{1/2}}{480}\right) ML^4h^5 = .0944ML^4h^5,$$

as compared with

$$(4.9) \quad |Eh^5| < .0546ML^4h^5$$

for Ralston's fourth-order procedure.

In this case the method becomes fifth order when  $f$  is independent of  $y$ , with error bound

$$(4.10) \quad |Eh^6| < 1.389 \times 10^{-5} ML^5h^6.$$

**5. An Additional Constraint.** Now suppose we consider the second error term—that involving  $h^{m+2}$ . In this term, setting the coefficient of  $D^{m+1}f$  to zero leads to

$$(5.1) \quad 10\alpha_2\alpha_3 - 5(\alpha_2 + \alpha_3) + 3 = 0$$

for third-order methods and to

$$(5.2) \quad 2(\alpha_2^2\alpha_3 + \alpha_2\alpha_3^2) - (\alpha_2^2 - \alpha_2\alpha_3 + \alpha_3^2) - (\alpha_2 + \alpha_3) + 1 = 0$$

for fourth-order methods. The intersection of (5.1) and (3.3) is the point

$$(\alpha_2, \alpha_3) = \left(\frac{6 - (6)^{1/2}}{10}, \frac{6 + (6)^{1/2}}{10}\right),$$

which determines the parameters

$$(5.3) \quad \begin{aligned} w_1 &= \frac{1}{9}, & w_2 &= \frac{16 + (6)^{1/2}}{36}, & w_3 &= \frac{16 - (6)^{1/2}}{36}, \\ \beta_{21} &= \frac{6 - (6)^{1/2}}{10}, & \beta_{31} &= -\left(\frac{54 + 19(6)^{1/2}}{250}\right), & \beta_{32} &= \frac{102 + 22(6)^{1/2}}{125} \end{aligned}$$

and thus defines the third-order procedure

$$(5.4) \quad y_{n+1} - y_n = .1111111111k_1 + .5124858262k_2 + .3764030627k_3,$$

where

$$(5.5) \quad \begin{aligned} k_1 &= hf(x_n, y_n), \\ k_2 &= hf(x_n + .3550510257h, y_n + .3550510257k_1), \\ k_3 &= hf(x_n + .8449489743h, y_n - .4021612205k_1 + 1.247110195k_2), \end{aligned}$$

with

$$(5.6) \quad |Eh^4| < .1391ML^3h^4.$$

For derivative functions  $f$  that are independent of  $y$ , this procedure becomes Radau quadrature of order five with leading error term

$$(5.7) \quad |Eh^6| < 1.389 \times 10^{-5}ML^5h^6.$$

Similarly, (5.2) and (4.2) intersect at

$$(\alpha_2, \alpha_3) = \left(\frac{5 - (5)^{1/2}}{10}, \frac{5 + (5)^{1/2}}{10}\right)$$

to yield

$$(5.8) \quad \begin{aligned} w_1 &= \frac{1}{12}, & w_2 &= \frac{5}{12}, & w_3 &= \frac{5}{12}, & w_4 &= \frac{1}{12}, \\ \alpha_4 &= 1, & \beta_{21} &= \frac{5 - (5)^{1/2}}{10}, & \beta_{31} &= -\left(\frac{5 + 3(5)^{1/2}}{20}\right), & \beta_{32} &= \frac{3 + (5)^{1/2}}{4}, \\ \beta_{41} &= \frac{-1 + 5(5)^{1/2}}{4}, & \beta_{42} &= -\left(\frac{5 + 3(5)^{1/2}}{4}\right), & \beta_{43} &= \frac{5 - (5)^{1/2}}{2}. \end{aligned}$$

This is the fourth-order system

$$(5.9) \quad \begin{aligned} y_{n+1} - y_n &= .08333333333k_1 + .4166666667k_2 + .4166666667k_3 \\ &+ .08333333333k_4, \end{aligned}$$

where

$$(5.10) \quad \begin{aligned} k_1 &= hf(x_n, y_n), \\ k_2 &= hf(x_n + .2763932023h, y_n + .2763932023k_1), \\ k_3 &= hf(x_n + .7236067977h, y_n - .5854101966k_1 + 1.309016994k_2), \\ k_4 &= hf(x_n + h, y_n + 2.545084972k_1 - 2.927050983k_2 + 1.381966011k_3), \end{aligned}$$

with

$$(5.11) \quad |Eh^5| < .1218ML^4h^5.$$

For  $f$  independent of  $y$ , (5.9, 10) is Lobatto's sixth-order quadrature formula, with truncation error

$$(5.12) \quad |Eh^7| < 6.614 \times 10^{-7} ML^6 h^7.$$

The restriction that  $\alpha_4 = 1$ , however, precludes having a fourth-order integration scheme corresponding to Radau quadrature, which in this case is of order seven.

**6. Examples.** Both of Ralston's examples and several others have been programmed for a CONTROL DATA 3600 computer, using all of the proposed methods. Results were as good as those for the Ralston schemes. Furthermore, the suggested procedures (3.6, 7), (4.6, 7), (5.4, 5), and (5.9, 10) did indeed produce results of the predicted order of accuracy when the example

$$(6.1) \quad y' = y, \quad y(0) = 1, \quad \text{solution } y(x) = e^x$$

was redone with  $y' = e^x$ . That is, the integration procedures reduce to high-order quadrature formulas and thus could be used to do double duty in a subroutine library.

**7. Acknowledgment.** D. L. Phillips not only gave constant encouragement and advice but also made the observation in the last paragraph of Section 1 and contributed the idea for Section 5.

Argonne National Laboratory  
 Argonne, Illinois.

1. T. E. HULL & R. L. JOHNSTON, "Optimum Runge-Kutta methods," *Math. Comp.*, v. 18, 1964, pp. 306-310. MR **29** #2980.
2. M. LOTKIN, "On the accuracy of Runge-Kutta's method," *MTAC*, v. 5, 1951, pp. 128-133. MR **13**, 286.
3. A. RALSTON, "Runge-Kutta methods with minimum error bounds," *Math. Comp.*, v. 16, 1962, pp. 431-437; Corrigendum, v. 17, 1963, p. 488. MR **27** #940.