

A Note on the Integral

$$\int_0^\infty t^{2\alpha-1}(1+t^2)^{1-\alpha-\beta} J_\nu(x\sqrt{1+t^2}) dt$$

By M. L. Glasser

Abstract. The integral

$$I_\nu^{\alpha,\beta}(x) = \int_0^\infty t^{2\alpha-1}(1+t^2)^{1-\alpha-\beta} J_\nu(x\sqrt{1+t^2}) dt$$

is expressed in terms of Bessel and related functions for various values of the parameters by summing the hypergeometric series representation given by Schmidt.

In a recent paper [1] Schmidt has studied the integral in the title of this note, which he denotes $I_\nu^{\alpha,\beta}(x)$, by expressing it in terms of the hypergeometric series, ${}_1F_2$. In addition, he presented simple expressions for $I_\nu^{1/2,1}$, $I_\nu^{1/2,1/2}$ and $I_\nu^{\alpha,\nu/2+1-\alpha}$ in terms of Bessel functions. The purpose of this note is to augment the results in [1] by a number of simple formulas obtained by summing the hypergeometric series. For brevity the rather messy derivations have been omitted. The notation is that of [2].

$$I_0^{1,1/2}(x) = \frac{1}{x} - J_0(x) + \frac{\pi}{2} [J_0(x)H_1(x) - J_1(x)H_0(x)],$$

$$I_1^{1,1/2}(x) = J_0(x)/x, \quad I_1^{1/2,1}(x) = x^{-1} \sin x, \quad I_0^{1/2,1/2}(x) = x^{-1} \cos x,$$

$$I_{1/2}^{\alpha,5/2-2\alpha}(1) = -\sqrt{\frac{\pi}{8}} [J_{3/2-2\alpha}(1/2)Y_{1/2}(1/2) + Y_{3/2-2\alpha}(1/2)J_{1/2}(1/2)],$$

$$1/2 < \alpha < 1,$$

$$I_0^{1/2,3/2}(1) = -\text{si}(1),$$

$$I_\nu^{(\nu-1/2),(3/2-(1/2)\nu)}(x) = \frac{\Gamma(\nu-1/2)}{2^{\nu+1}\sqrt{\pi}} x^{-\nu} \sin x, \quad \frac{1}{2} < \nu < \frac{5}{2},$$

$$I_\nu^{(\nu+1/2),(1/2-(1/2)\nu)}(x) = \frac{\Gamma(\nu+1/2)}{2^\nu\sqrt{\pi}} x^{-\nu-1} \cos x, \quad -1/2 < \nu < 1/2,$$

$$I_\nu^{(1/2-\nu),((3/2)\nu+1)}(x) = -\frac{\sqrt{\pi}}{2^{\nu+1}} \Gamma(1/2-\nu)x^\nu J_\nu(\frac{1}{2}x) Y_\nu(\frac{1}{2}x), \quad |\nu| < 1/2.$$

$$I_\nu^{1,1/4}(x) = x^{-1/2} \{J_{\nu-1}(x)S_{1/2,\nu}(x) + (\frac{1}{2}-\nu)J_\nu(x)S_{-1/2,\nu-1}(x)\}.$$

Received May 5, 1978.

AMS (MOS) subject classifications (1970). Primary 33A40; Secondary 44A20.

© 1979 American Mathematical Society
 0025-5718/79/0000-0074/\$01.50

This list is not exhaustive and can be extended considerably by use of the relations

$$(x/2\nu)[I_{\nu-1}^{\alpha,\beta}(x) + I_{\nu+1}^{\alpha,\beta}(x)] = I_\nu^{\alpha,\beta+1/2}(x),$$

$$x^{-\nu}[x^\nu I_\nu^{\alpha,\beta}(x)]' = I_{\nu-1}^{\alpha,\beta-1/2}(x),$$

$$x^\nu[x^{-\nu} I_\nu^{\alpha,\beta}(x)]' = -I_{\nu+1}^{\alpha,\beta-1/2}(x).$$

Mathematics and Computer Science Department
Clarkson College of Technology
Potsdam, New York 13676

1. P. W. SCHMIDT, *Math. Comp.*, v. 32, 1978, pp. 265–269.
2. Y. L. LUKE, *The Special Functions and Their Approximations*, Academic Press, New York, 1969.