

The Finite Element Method With Lagrange Multipliers for Domains With Corners*

By Juhani Pitkäranta

Abstract. We study the convergence of the finite element method with Lagrange multipliers for approximately solving the Dirichlet problem for a second-order elliptic equation in a plane domain with piecewise smooth boundary. Assuming mesh refinements around the corners, we construct families of boundary subspaces that are compatible with triangular Lagrange elements in the interior, and we carry out the error analysis of the resulting approximations in weighted Sobolev spaces.

1. Introduction. We consider the model problem

$$(1.1) \quad \begin{aligned} -\Delta u &= f && \text{in } \Omega, \\ u &= g && \text{on } \partial\Omega, \end{aligned}$$

where $\Omega \subset R^2$ is a bounded domain with piecewise smooth boundary $\partial\Omega$, and f and g are given functions defined on Ω . Under sufficient regularity hypotheses on f and g , problem (1.1) admits the following weak formulation: Find a pair (u, ψ) , $u \in H^1(\Omega)$, $\psi \in L_2(\partial\Omega)$, such that

$$(1.2) \quad \begin{aligned} \int_{\Omega} \sum_{i=1}^2 \frac{\partial u}{\partial x_i} \frac{\partial v}{\partial x_i} dx + \int_{\partial\Omega} v\psi dx + \int_{\partial\Omega} u\phi ds \\ = \int_{\Omega} fv dx + \int_{\partial\Omega} g\phi ds \quad \text{for all } (v, \phi) \in H^1(\Omega) \times L_2(\partial\Omega). \end{aligned}$$

For f, g smooth enough, (1.2) has a unique solution (u, ψ) such that u is the weak solution of (1.1) and ψ is defined as

$$(1.3) \quad \psi = -\frac{\partial u}{\partial n} = -n_1 \frac{\partial u}{\partial x_1} - n_2 \frac{\partial u}{\partial x_2}, \quad x \in \partial\Omega,$$

where $n(x) = (n_1(x), n_2(x))$ denotes the unit outward normal vector of $\partial\Omega$ at x .

In the Lagrange multiplier method [2] for approximately solving (1.2) one introduces the finite-dimensional subspaces $M^h \subset H^1(\Omega)$ and $N^h \subset L_2(\partial\Omega)$ and defines the approximate solution as the pair $(u_h, \psi_h) \in M^h \times N^h$ which satisfies

$$(1.4) \quad \begin{aligned} \int_{\Omega} \sum_{i=1}^2 \frac{\partial u_h}{\partial x_i} \frac{\partial v}{\partial x_i} dx + \int_{\partial\Omega} v\psi_h ds + \int_{\partial\Omega} u_h\phi ds \\ = \int_{\Omega} fv dx + \int_{\partial\Omega} g\phi ds \quad \text{for all } (v, \phi) \in M^h \times N^h. \end{aligned}$$

Received September 6, 1979; revised July 9, 1980 and November 24, 1980.

1980 *Mathematics Subject Classification.* Primary 65N30.

* This work was supported by the Finnish National Research Council for Technical Sciences.

Finite element methods based on (1.4) have the benefit that they do not require the fulfillment of the Dirichlet boundary condition in the subspaces. The stability and convergence of such approximations was studied first in [2] (see also [3]). For an abstract Lagrange multiplier method, necessary and sufficient conditions for stability are given in [6].

The actual verification of stability and the construction of the subspaces was considered by the author in [9] for quasi-uniform finite element meshes and in [10] in a more general situation. In [10] a number of local conditions were stated that yield the uniform stability of the Lagrange multiplier method in weighted norms depending on the finite element partitioning of the boundary.

In the present paper we proceed from the ideas of [10] and [4]. We consider the situation where graded meshes are used to handle the singularities arising from the corners of $\partial\Omega$. Assuming mesh refinements of the same type as those considered in [4] for polygonal domains, we construct finite element subspaces that meet with the stability conditions of [10]. We then derive error estimates for the resulting approximations in weighted Sobolev spaces.

The plan of the paper is as follows. In Section 2 we state the basic assumptions and give some properties of weighted Sobolev spaces that are needed in the paper. In Section 3 the stability conditions are restated from [10]. A family of finite element subspaces is introduced in Section 4 and the validity of the stability conditions is verified. Finally, in Section 5, the approximability properties of the subspaces in weighted Sobolev spaces are studied, and a weighted convergence result is derived for the approximate solution.

2. Preliminaries. We consider a bounded, simply connected domain $\Omega \subset \mathbb{R}^2$, whose boundary $\partial\Omega$ consists of a finite number of smooth closed arcs Γ_i , $i = 1, \dots, I$, defined in terms of the smooth mappings $J_i = (J_{i,1}, J_{i,2})$ such that

$$(2.1) \quad \Gamma_i = \{(J_{i,1}(t), J_{i,2}(t)), t \in [0, 1]\},$$

$$[J'_{i,1}(t)]^2 + [J'_{i,2}(t)]^2 \geq C > 0, \quad t \in [0, 1].$$

We will denote the corners, i.e., the endpoints of the arcs Γ_i , by z_i , $i = 1, \dots, I$, numbered in any order. The asymptotic opening angles of the corners into the interior of Ω are denoted by ω_i . We assume that $0 < \omega_i < 2\pi$, $i = 1, \dots, I$, i.e., we exclude needles but include slits.

For $u(x)$ defined on Ω , let $D^k u$ denote the field of all partial derivatives of u of order k . Then the Sobolev space $H^m(\Omega)$, $m \geq 0$, is defined as usual, with the norm

$$\|u\|_{H^m(\Omega)}^2 = \sum_{k=0}^m \int_{\Omega} |D^k u|^2 dx.$$

In what follows, we use the notation

$$(2.2) \quad \varphi_{\beta}(x) = \prod_{i=1}^I |x - z_i|^{\beta_i}, \quad \beta = (\beta_1, \dots, \beta_I) \in \mathbb{R}^I,$$

and $\theta = (1, 1, \dots, 1) \in \mathbb{R}^I$. For $\beta \in \mathbb{R}^I$ and $m \geq 0$ given, we let $W^{m,\beta}(\Omega)$ denote the weighted Sobolev space of functions u , defined on Ω , such that

$$\|u\|_{W^{m,\beta}(\Omega)}^2 = \sum_{k=0}^m \int_{\Omega} \varphi_{\beta+(k-m)\theta} |D^k u|^2 dx < \infty.$$

We state below some inequalities to be needed later on.

LEMMA 2.1. *If $\beta \in R^I$ is such that $0 < \beta_i < 1$, $i = 1, \dots, I$, and $u \in H^1(\Omega)$, then*

$$\int_{\Omega} \varphi_{\beta}^{-2} u^2 dx \leq C \|u\|_{H^1(\Omega)}^2,$$

where C depends on β and Ω .

Proof. Let $\chi_{i,\delta}$ be a smooth function defined on Ω such that $\chi_{i,\delta}(x) \equiv 1$ for $|x - z_i| < \delta$ and $\chi_{i,\delta}(x) \equiv 0$ for $|x - z_i| > 2\delta$. Then, if δ is sufficiently small, a modification of Hardy's inequality gives (see [8]):

$$\int_{\Omega} |x - z_i|^{\alpha} (\chi_{i,\delta} u)^2 dx \leq C \int_{\Omega} |x - z_i|^{\alpha+2} |D^1(\chi_{i,\delta} u)|^2 dx, \quad \alpha > -2.$$

Taking $\alpha = -2\beta_i$, summing over i and using the triangle inequality, the assertion follows after a simple computation. \square

LEMMA 2.2. *If $\beta \in R^I$ is such that $\beta_i < 1$ for some $i \in \{1, \dots, I\}$, then any $u \in W^{2,\beta}(\Omega)$ satisfies*

$$|u(x)| \leq C \|u\|_{W^{2,\beta}(\Omega)}, \quad |x - z_j| > \delta > 0, j \neq i,$$

where C depends on Ω and δ . Moreover, $u(z_i) = 0$.

Proof. See [4]. This result is also implicit in [8].

Consider next functions defined on $\partial\Omega$. We say $\psi \in H^m(\partial\Omega)$, $m \geq 0$, if

$$\|\psi\|_{H^m(\partial\Omega)}^2 = \sum_{i=1}^I \|\psi_i\|_{H^m(0,1)}^2 < \infty,$$

where $\psi_i(t) = \psi(J_i(t))$, $t \in (0, 1)$, $i = 1, \dots, I$. Similarly, $\psi \in H^{m+1/2}(\partial\Omega)$, $m > 0$, if

$$\|\psi\|_{H^{m+1/2}(\partial\Omega)}^2 = \sum_{i=1}^I (\|\psi_i\|_{H^m(0,1)}^2 + |\psi_i^{(m)}|_{1/2}^2) < \infty,$$

where

$$(2.3) \quad |\phi|_{1/2}^2 = \int_0^1 \int_0^1 \frac{[\phi(t) - \phi(t')]^2}{(t - t')^2} dt dt'.$$

We note that $H^{m+1/2}(\partial\Omega)$ can be defined equivalently as the interpolation space [5]

$$H^{m+1/2}(\partial\Omega) = [H^m(\partial\Omega), H^{m+1}(\partial\Omega)]_{1/2,2}.$$

To define weighted Sobolev spaces for functions on $\partial\Omega$, let $\nu = \nu(i)$ and $\mu = \mu(i)$ be integers such that

$$(2.4) \quad J_i(0) = z_{\nu}, \quad J_i(1) = z_{\mu}.$$

With $\beta \in R^I$ and $m \geq 0$, we define $W^{m,\beta}(\Omega)$ as the space of functions ψ such that

$$\|\psi\|_{W^{m,\beta}(\Omega)}^2 = \sum_{i=1}^I \|\psi_i\|_{m,\beta,i}^2 < \infty,$$

where

$$\|\phi\|_{m,\beta,i}^2 = \sum_{j=0}^m \int_0^1 [t^{\beta_i - m + j} (1 - t)^{\beta_i - m + j}]^2 |\phi^{(j)}(t)|^2 dt.$$

Further, we let $W^{m+1/2,\beta}(\partial\Omega)$ denote the space with the norm

$$(2.5) \quad \|\psi\|_{W^{m+1/2,\beta}(\partial\Omega)}^2 = \|\psi\|_{W^{m,\beta-\theta/2}(\partial\Omega)}^2 + \sum_{i=1}^I |t^{\beta_i}(1-t)^{\beta_i}\psi_i^{(m)}|_{1/2}^2,$$

where $|\cdot|_{1/2}$ is defined by (2.3). The following interpolation result is obtained from [11].

LEMMA 2.3. *If $m \geq 0$ and $\beta = \frac{1}{2}(\beta_1 + \beta_2)$, then*

$$W^{m+1/2,\beta}(\partial\Omega) = [W^{m,\beta_1}(\partial\Omega), W^{m+1,\beta_2}(\partial\Omega)]_{1/2,2}.$$

In Section 5 we need the following trace inequality:

LEMMA 2.4. *If $u \in W^{m,\beta}(\Omega)$, $m \geq 1$, then*

$$\|u|_{\partial\Omega}\|_{W^{m-1/2,\beta}(\partial\Omega)} \leq C\|u\|_{W^{m,\beta}(\Omega)}.$$

Proof. In [8] it is proven that if $u \in W^{1,\beta}(\Omega)$, then

$$(2.6) \quad \|u|_{\partial\Omega}\|_{W^{0,\beta-\theta/2}(\partial\Omega)} \leq C\|u\|_{W^{1,\beta}(\Omega)}.$$

From this it is easily concluded that, if $u \in W^{m,\beta}(\Omega)$, $m \geq 1$, then

$$(2.7) \quad \|u|_{\partial\Omega}\|_{W^{m-1,\beta-\theta/2}(\partial\Omega)} \leq C\|u\|_{W^{m,\beta}(\Omega)}.$$

Now let $D_{m-1}u$ be any partial derivative of u of order $m-1$, and let $\psi_i(t) = (D_{m-1}u)(J_i(t))$, $i = 1, \dots, I$. Then

$$(2.8) \quad \sum_{i=1}^I |t^{\beta_i}(1-t)^{\beta_i}\psi_i|_{1/2}^2 \leq C\|\varphi_\beta D_{m-1}u|_{\partial\Omega}\|_{H^{1/2}(\partial\Omega)}^2.$$

Using the trace properties of Sobolev spaces $H^m(\Omega)$ (cf. [1]) the right side can here be further estimated as

$$(2.9) \quad \|\varphi_\beta D_{m-1}u|_{\partial\Omega}\|_{H^{1/2}(\partial\Omega)} \leq C\|\varphi_\beta D_{m-1}u\|_{H^1(\Omega)}.$$

The asserted inequality now follows by comparing (2.5) with (2.7) through (2.9) and noting that

$$\|\varphi_\beta D_{m-1}u\|_{H^1(\Omega)} \leq C\|u\|_{W^{m,\beta}(\Omega)}. \quad \square$$

We state finally a regularity result for the solution of problem (1.1). For the proof, see [8].

THEOREM 2.1. *Let $m \geq 2$ and let $\beta \in R^I$ be such that $\beta_i \geq 0$, $m-1 \geq \beta_i > m-1-\pi/\omega_i$, $i = 1, \dots, I$. Further, let f and g in (1.1) be such that $f = f_1 + f_2$, $f_1 \in H^{m-2}(\Omega)$, $f_2 \in W^{m-2,\beta}(\Omega)$, and $g = g_1 + g_2$, $g_1 \in H^m(\Omega)$, $g_2 \in W^{m,\beta}(\Omega)$. Then problem (1.1) has in $H^1(\Omega)$ a unique weak solution u which satisfies $u = u_1 + u_2$, $u_1 \in H^m(\Omega)$, $u_2 \in W^{m,\beta}(\Omega)$,*

$$\begin{aligned} & \|u_1\|_{H^m(\Omega)} + \|u_2\|_{W^{m,\beta}(\Omega)} \\ & \leq C\{\|f_1\|_{H^{m-2}(\Omega)} + \|f_2\|_{W^{m-2,\beta}(\Omega)} + \|g_1\|_{H^m(\Omega)} + \|g_2\|_{W^{m,\beta}(\Omega)}\}. \end{aligned}$$

3. The Local Stability Conditions. In this section we state a number of conditions that are sufficient for the uniform stability of the Lagrange multiplier method. These were stated already in [10], but as they will be referred to in the sequel, we repeat them here for convenience.

Suppose we are given a partitioning $\tau^h(\partial\Omega)$ of $\partial\Omega$, consisting of connected smooth arcs, a finite-dimensional subspace $N^h \subset L_2(\partial\Omega)$ associated to $\tau^h(\partial\Omega)$ in

the manner described below, and another finite-dimensional space $M^h \subset H^1(\Omega)$. If $S \subset R^n$, we let $d(S)$ denote the diameter of S , and we let K , Λ , and L be some fixed positive parameters, with L an integer. Further, we associate to each pair $\{S_i, S_j\} \subset \tau^h(\partial\Omega)$ a number n_{ij} , defined as the minimum number of subsets $S \in \tau^h(\partial\Omega)$ that have to be crossed when passing from a point in S_i to a point of S_j along $\partial\Omega$.

In the above notation, the sufficient stability conditions are as follows:

A1. To the partitioning $\tau^h(\partial\Omega) = \{S_1, \dots, S_\nu\}$ there corresponds a collection $\{\mathcal{C}_1, \dots, \mathcal{C}_\nu\}$ of discs $\mathcal{C}_j \subset R^2$, such that, for each j , the center of \mathcal{C}_j is in S_j , $d(\mathcal{C}_j) \geq 8Kd(S_j)$, and, for each $\mathcal{C} \in \{\mathcal{C}_j\}$, $\mathcal{C} \cap \mathcal{C}_j \neq \emptyset$ for at most L spheres $\mathcal{C}_j \in \{\mathcal{C}_1, \dots, \mathcal{C}_\nu\}$.

A2. For all pairs $\{S_i, S_j\} \subset \tau^h(\partial\Omega)$,

$$\frac{d(S_i)}{d(S_j)} \leq K(1 + n_{ij})^\Lambda.$$

A3. For each $S_j \in \tau^h(\partial\Omega)$, there exists $\phi_j \in N^h$ such that $\text{supp}\{\phi_j\} \supseteq S_j$, $d(\text{supp}\{\phi_j\}) \leq Kd(S_j)$, and

$$\left| \int_{\partial\Omega} \phi_j ds \right|^2 \geq K^{-1}d(S_j) \int_{\partial\Omega} \phi_j^2 ds.$$

A4. There exists a basis $\{\psi_1, \dots, \psi_\mu\}$ of N^h and a set $\{v_1, \dots, v_\mu\} \subset M^h$ such that

(i) For all i , $1 \leq i \leq \mu$, if $\text{supp}\{\psi_i\} \cap S_j \neq \emptyset$ and $\text{supp}\{\psi_i\} \cap S_k \neq \emptyset$, $S_j, S_k \in \tau^h(\partial\Omega)$, then $n_{j,k} \leq L$.

(ii) For all i , $1 \leq i \leq \mu$, $v_i v_j \neq 0$ for at most L functions v_j , $1 \leq j \leq \mu$.

(iii) If $S_j \in \tau^h(\partial\Omega)$, then

$$\Lambda_j = \{i; \psi_i \neq 0 \text{ on } S_j\} = \{i; v_i \neq 0 \text{ on } S_j\}.$$

(iv) If $S_j \in \tau^h(\partial\Omega)$ and $v_i \neq 0$ on S_j , then

$$K^{-1}d(S_j) \int_{\Omega} |D^1 v_i|^2 dx \leq \|v_i\|_{L_2(S_j)}^2 \leq K \|\psi_i\|_{L_2(S_j)}^2.$$

(v) If $\psi = \sum_1^\mu \beta_i \psi_i$, $v = \sum_1^\mu \beta_i v_i$, $\beta_i \in R^1$, $S_j \in \tau^h(\partial\Omega)$, and Λ_j is as in (iii), then

$$K \sum_{i \in \Lambda_j} \beta_i^2 \|\psi_i\|_{L_2(S_j)}^2 \geq \|\psi\|_{L_2(S_j)}^2 \geq K^{-1} \sum_{i \in \Lambda_j} \beta_i^2 \|\psi_i\|_{L_2(S_j)}^2,$$

$$K \sum_{i \in \Lambda_j} \beta_i^2 \|v_i\|_{L_2(S_j)}^2 \geq \|v\|_{L_2(S_j)}^2 \geq K^{-1} \sum_{i \in \Lambda_j} \beta_i^2 \|v_i\|_{L_2(S_j)}^2,$$

and

$$\int_{S_j} \psi v ds \geq K^{-1} \|\psi\|_{L_2(S_j)}^2.$$

We state below a consequence of the above assumptions, which will be of basic importance in what follows. First, let us define on $[H^1(\Omega) \times L_2(\partial\Omega)]^2$ a bilinear form \mathfrak{B} as

$$(3.1) \quad \mathfrak{B}(u, \psi; v, \phi) = \int_{\Omega} \sum_{i=1}^2 \frac{\partial u}{\partial x_i} \frac{\partial v}{\partial x_i} dx + \int_{\partial\Omega} (u\phi + v\psi) ds,$$

so that (1.4) can be rewritten as

$$\mathfrak{B}(u_h, \psi_h; v, \phi) = \int_{\Omega} f v \, dx + \int_{\partial\Omega} g \phi \, ds, \quad (v, \phi) \in M^h \times N^h.$$

Further, let us introduce on $H^1(\Omega) \times L_2(\partial\Omega)$ the weighted norm

$$\|(u, \psi)\|_{\tau^h}^2 = \int_{\Omega} |D^1 u|^2 \, dx + \sum_j d(S_j)^{-1} \int_{S_j} u^2 \, ds + \sum_j d(S_j) \int_{S_j} \psi^2 \, ds,$$

where the sums are over all $S_j \in \tau^h(\partial\Omega)$. Then we have, by slightly modifying the stability result of [10],

THEOREM 3.1. *Let $\tau^h(\partial\Omega)$, M^h , and N^h be such that the assumptions A1 through A4 are satisfied for some finite K , Λ , and L . Then, if $d(S) < h_0$ for all $S \in \tau^h(\partial\Omega)$, where $h_0 = h_0(\Omega) \in (0, 1)$, we have*

$$\inf_{(u, \psi) \in M^h \times N^h} \sup_{(v, \phi) \in M^h \times N^h} \frac{\mathfrak{B}(u, \psi; v, \phi)}{\|(u, \psi)\|_{\tau^h} \|(v, \phi)\|_{\tau^h}} > C > 0,$$

where C depends on γ , K , Λ , and L .

4. The Finite Element Subspaces. We will now introduce partitionings and subspaces, which meet with the assumptions of Section 3. First, a family of triangulations of Ω is introduced. To this end, let $\beta \in R^I$ be given so that $0 < \beta_i < 1$, $i = 1, \dots, I$, and assume that for each h , $0 < h < 1$, we are given a collection $\tau_{\beta}^h(\Omega)$ of open, disjoint subsets of Ω , such that $\bigcup_{T \in \tau_{\beta}^h(\Omega)} \bar{T} = \bar{\Omega}$ and so that each $T \in \tau_{\beta}^h(\Omega)$ is either a triangle or a triangle with one or two curved sides on $\partial\Omega$. We will further assume that for some positive constant ε , the following conditions hold for all $h \in (0, 1)$.

B. (i) Each $T \in \tau_{\beta}^h(\Omega)$ contains a sphere of radius $\rho > \varepsilon d(T)$.

(ii) If $T \in \tau_{\beta}^h(\Omega)$ and $z_i \notin \bar{T}$, $i = 1, \dots, I$, then

$$\varepsilon h \max_{x \in \bar{T}} \varphi_{\beta}(x) < d(T) < \varepsilon^{-1} h \min_{x \in \bar{T}} \varphi_{\beta}(x).$$

(iii) If $T \in \tau_{\beta}^h(\Omega)$ and $z_i \in \bar{T}$, then

$$\varepsilon h \max_{x \in \bar{T}} \varphi_{\beta}(x) < d(T) < \varepsilon^{-1} h \max_{x \in \bar{T}} \varphi_{\beta}(x).$$

Here φ_{β} refers to (2.1).

For a polygonal domain, triangulations of the above type were discussed previously in [4]. We cite from [4] that the number of triangles in a triangulation $\tau_{\beta}^h(\Omega)$ is bounded by Ch^{-2} , where C depends on Ω and ε , and β . We also point out the following implication of B(iii):

$$(4.1) \quad T \in \tau_{\beta}^h(\Omega) \ \& \ z_i \in \bar{T} \Rightarrow d(T) < Ch^{1/(1-\beta_i)}.$$

To each $\tau_{\beta}^h(\Omega)$ we associate the finite element spaces $M_{\beta, k}^h$, $k = 1, 2, \dots$, defined as subspaces of $H^1(\Omega)$ such that if $u \in M_{\beta, k}^h$ and $T \in \tau_{\beta}^h(\Omega)$, then $u|_T$ is a polynomial of degree at most k .

We let $\tau_{\beta}^h(\partial\Omega)$ denote the boundary partitioning induced by $\tau_{\beta}^h(\Omega)$, i.e., $\tau_{\beta}^h(\partial\Omega)$ consists of the sides on $\partial\Omega$ of the triangles in $\tau_{\beta}^h(\Omega)$. We associate to $\tau_{\beta}^h(\partial\Omega)$ the subspaces $N_{\beta, k}^h \subset L_2(\partial\Omega)$, $k = 1, 2, \dots$. First, assuming the notation

$$(4.2) \quad \tau_{\beta}^h(\partial\Omega) = \{\Gamma_{i,j} \subset \partial\Omega; \Gamma_{i,j} = J_i(I_{i,j}), i = 1, \dots, I, j = 1, \dots, m_i\},$$

where J_i is as in (2.1) and

$$(4.3) \quad I_{i,j} = (t_{j-1}^i, t_j^i); \quad 0 = t_0^i < t_1^i < \dots < t_{m_i}^i = 1,$$

let $N_{\beta,k,i}^h$, $1 \leq i \leq I$, denote the maximal subspace of $C^0[0, 1]$ such that if $\phi \in N_{\beta,k,i}^h$, then (i) if $k = 1$, then $\phi|_{I_{i,j}}$ is a constant for $j = 1$ and $j = m_i$ and a polynomial of degree unity for $2 \leq j \leq m_i - 1$; (ii) if $k > 2$, then $\phi|_{I_{i,j}}$ is a polynomial of degree $k - 1$ for $1 \leq j \leq m_i$. Then define

$$N_{\beta,k}^h = \{ \psi \in L_2(\partial\Omega); \phi(t) = \psi(J_i(t)) \in N_{\beta,k,i}^h, i = 1, \dots, I \}.$$

In order to verify the assumptions A1 through A4 for the above subspaces and partitionings, set $M^h = M_{\beta,k}^h$, $N^h = N_{\beta,k}^h$, and $\tau^h(\partial\Omega) = \{ \partial T \cap \partial\Omega, T \in \tau_{\beta}^h(\Omega), T$ has at least one side on $\partial\Omega \}$. First, we easily conclude from B(ii) that, if $S \in \tau^h(\partial\Omega)$ and $\mathcal{C} \subset R^2$ is any disc of radius $\rho \leq Kd(S)$ containing S , then the number of the sets $S_j \in \tau^h(\partial\Omega)$ that intersect \mathcal{C} is bounded by CK , where, if h is small enough (depending on Ω), C depends only on β and ε . In view of this, there exists for any given β, ε, K , a finite L such that A1 is satisfied for all $h \in (0, h_0)$, $h_0 = h_0(\Omega) \in (0, 1)$.

The verification of A2 is also easy: using B(ii) and (iii), a simple calculation shows that A2 is satisfied if K is large enough (depending on β, ε) and

$$\Lambda = \max_{i=1, \dots, I} \frac{\beta_i}{1 - \beta_i}.$$

To verify A3, let $S \in \tau^h(\partial\Omega)$ be given and let $S_1 \in \tau^h(\partial\Omega)$ be a subset adjacent to S on $\partial\Omega$. Then there exists a nonnegative function $\phi \in N^h$ such that $\text{supp}\{\phi\} \subset S_1 \cup S$. By a scaling argument, we have the inequality $|\int_{\partial\Omega} \phi \, ds|^2 > Cd(S) \int_{\partial\Omega} \phi^2 \, ds$, where $C > 0$ depends on k and on the constant K in A2. Hence, A3 is satisfied.

We come finally to the assumption A4.

LEMMA 4.1. *Let $\beta \in R^I$ be such that $0 \leq \beta_i < 1, i = 1, \dots, I$. Then there exist finite K and L depending on ε and β such that $M^h = M_{\beta,k}^h$ and $N^h = N_{\beta,k}^h$ satisfy A4 for all $h \in (0, h_0)$, $h_0 = h_0(\Omega) \in (0, 1)$.*

Proof. We define first a linear mapping $U: N^h \rightarrow M^h$. To this end, let $\psi \in N^h$ be given and let us associate to each $T \in \tau_{\beta}^h(\Omega)$ a polynomial $p_T = p_T(\psi)$ (on R^2) such that

$$(4.4) \quad (U\psi)(x) = p_T(x), \quad x \in T \in \tau_{\beta}^h(\Omega).$$

We now define the polynomials p_T . First, consider a triangle $T \in \tau_{\beta}^h(\Omega)$ which has a side S on $\partial\Omega$. We choose a cartesian coordinate system $\{x_1, x_2\}$ so that S has the parametrization

$$(4.5) \quad S = \{(x_1, x_2); x_2 = F(x_1), x_1 \in (0, d)\},$$

where F is a smooth function satisfying

$$(4.6) \quad F(0) = F(d) = 0, \quad |F(x_1)| + d|F'(x_1)| \leq Cd^2, \quad x_1 \in (0, d),$$

with C independent of S . Now if S does not touch a corner of $\partial\Omega$, we require that $p_T = p_T(x_1, x_2)$ satisfies

$$p_T(id/k, 0) = \psi(id/k, F(id/k)), \quad i = 0, \dots, k.$$

In the remaining cases, i.e., one endpoint of S is a corner of $\partial\Omega$, we choose the coordinates in (4.5) so that the corner is at $(0, 0)$, and set

$$p_T(0, 0) = 0, \quad p_T(id/k, 0) = (i/k)\psi(id/k, F(id/k)), \quad i = 1, \dots, k.$$

Consider now an arbitrary $T \in \tau_\beta^h(\Omega)$. Recall (cf. [7]) that any polynomial of degree $\leq k$ on R^2 is defined uniquely by its values on a set σ_T given by

$$\sigma_T = \left\{ \frac{1}{k} \sum_{i=1}^3 \nu_i a_i; \nu_i \in \{0, 1, \dots, k\}, \sum_{i=1}^3 \nu_i = k \right\},$$

where a_i are the vertices of T . Now, if T has a side on $\partial\Omega$ and if $x \in \sigma_T$ is located on the line joining the two vertices of T on $\partial\Omega$, then we define $p_T(x)$ as above. At the remaining points $x \in \sigma_T$ we set $p_T(x) = 0$. Then each p_T is uniquely defined. Moreover, it is easy to see that the mapping U defined by (4.4) is linear and into M^h .

We establish next some inequalities for the mapping U . Let $\psi \in N^h$ be given, let $T \in \tau_\beta^h(\Omega)$ have a side S on $\partial\Omega$, parametrized as in (4.5)–(4.6), and let $\phi(x) = \psi(x)$, $x \in S$, if S does not touch a corner of Ω , or $\phi(x) = \phi(x_1, x_2) = (x_1/d)\psi(x_1, x_2)$, $x \in S$, if $(0, 0)$ is a corner of Ω .

Let us first note that, by (4.6),

$$(4.7) \quad \int_0^d [U\psi(x_1, 0) - U\psi(x_1, F(x_1))]^2 dx_1 \\ = \int_0^d \left[\int_0^{F(x_1)} \frac{\partial}{\partial x_2} U\psi(x_1, x_2) dx_2 \right]^2 dx_1 \leq Cd^2 \int_T |\nabla U\psi|^2 dx.$$

Also, since $U\psi(x_1, 0)$ is the Lagrange interpolant of $\phi(x_1, F(x_1))$,

$$(4.8) \quad \int_0^d [U\psi(x_1, 0) - \phi(x_1, F(x_1))]^2 dx_1 \\ \leq Cd^{2k+2} \int_0^d \left[\left(\frac{d}{dx_1} \right)^{k+1} \phi(x_1, F(x_1)) \right]^2 dx_1 \leq C_1 d^2 \int_S \psi^2 ds.$$

Here the last inequality follows from an inverse inequality for the space N^h . (Recall that ψ comes from a polynomial of degree $\leq k-1$ if S touches a corner of Ω , and from a polynomial of degree $\leq k$ otherwise.)

Using (4.7) and (4.8) we obtain

$$\left| \int_S (U\psi - \phi)\psi ds \right| = \left| \int_0^d [(U\psi - \phi)\psi](x_1, F(x_1)) \frac{ds}{dx_1} dx_1 \right| \\ = \left| \int_0^d [U\psi(x_1, F(x_1)) - U\psi(x_1, 0)]\psi(x_1, F(x_1)) \frac{ds}{dx_1} dx_1 \right. \\ \left. + \int_0^d [U\psi(x_1, 0) - \phi(x_1, F(x_1))]\psi(x_1, F(x_1)) \frac{ds}{dx_1} dx_1 \right| \\ \leq Cd \left\{ \int_T |\nabla U\psi|^2 dx \right\}^{1/2} \left\{ \int_S \psi^2 ds \right\}^{1/2} + Cd \int_S \psi^2 ds.$$

On the other hand, by the equivalence of norms in a finite-dimensional space, we have

$$\int_S \psi\phi ds \geq C \int_S \psi^2 ds$$

for d sufficiently small. Combining the last two inequalities we obtain, for d small enough,

$$\int_S (U\psi)\psi \, ds \geq C \int_S \psi^2 \, ds - C_1 d^2 \int_T |\nabla U\psi|^2 \, dx.$$

Now, if S is the only side of T on $\partial\Omega$, then B(i) and (4.8) imply that

$$d(T) \int_T |\nabla U\psi|^2 \, dx \leq C \int_0^d [U\psi(x_1, 0)]^2 \, dx_1 \leq C_1 \int_S \psi^2 \, ds,$$

where C_1 depends on k and ε . More generally, if $T \in \tau_\beta^h(\Omega)$ is a triangle that touches $\partial\Omega$, we conclude from a similar argument that

$$d(T) \int_T |\nabla U\psi|^2 \, dx \leq C \int_S (U\psi)^2 \, ds \leq C_1 \int_S \psi^2 \, ds,$$

where $S = \partial T \cap \partial\Omega$ if T has one or two sides on $\partial\Omega$, and otherwise $S = S_1 \cup S_2$, where $S_i \in \tau^h(\partial\Omega)$ are such that $\bar{S}_1 \cap \bar{S}_2$ is a vertex of T . Combining these inequalities, we see that

$$(4.9) \quad \int_S (U\psi)\psi \, ds \geq (C - C_1 h) \int_S \psi^2 \, ds \geq C_2 \int_S \psi^2 \, ds,$$

h small enough, $S = \partial T \cap \partial\Omega \in \tau^h(\partial\Omega)$, $\psi \in N^h$.

Also, since $U\psi|_T = 0$ if T does not touch $\partial\Omega$, we obtain

$$(4.10) \quad \int_\Omega |\nabla U\psi|^2 \, dx \leq C \sum_{S \in \tau^h(\partial\Omega)} d(S)^{-1} \int_S (U\psi)^2 \, ds$$

$$\leq C_1 \sum_{S \in \tau^h(\partial\Omega)} d(S)^{-1} \int_S \psi^2 \, ds.$$

Using (4.9) and (4.10), it is now easy to complete the proof of Lemma 4.1. First, introduce a basis $\{\psi_j\}$ of N^h such that A4(i) is satisfied for some finite L . Such a basis is easily formed from locally supported functions. We then define the set $\{v_j\} \subset M^h$ so that $v_j = U\psi_j$, where U is the linear mapping constructed above. The validity of A4(ii) is then clear, and the inequalities in A4(iv) and (v) are easily proved using scaling arguments and (4.9) and (4.10). \square

We conclude this section by stating a convergence result for the Lagrange multiplier method (1.4) when $M^h = M_{\beta,k}^h$, $N^h = N_{\beta,k}^h$. We need the following result.

LEMMA 4.2. *Let $\tau_\beta^h(\partial\Omega) = \{S_1, \dots, S_p\}$ and let $r \geq -\frac{1}{2}$. Then if $h \in (0, h_0)$, $h_0 = (0, 1) \in (0, 1)$, and $\psi \in N_{\beta,k}^h$ or $\psi \in M_{\beta,k|\partial\Omega}^h$, $0 \leq \beta_i < 1$, $i = 1, \dots, l$, there exist the positive constants C_1 and C_2 depending on k , ε and $r\beta$ such that*

$$C_1 h^{2r} \int_{\partial\Omega} \varphi_\beta^{2r} \psi^2 \, ds \leq \sum_{j=1}^p d(S_j)^{2r} \int_{S_j} \psi^2 \, ds \leq C_2 h^{2r} \int_{\partial\Omega} \varphi_\beta^{2r} \psi^2 \, ds.$$

Proof. If \bar{S}_j does not contain any of the corners of $\partial\Omega$, then B(i) and B(ii) imply that, if h is small enough and $\psi \in L_2(\partial\Omega)$, then

$$(4.11) \quad C_1 h^{2r} \int_{S_j} \varphi_\beta^{2r} \psi^2 \, ds \leq d(S_j)^{2r} \int_{S_j} \psi^2 \, ds \leq C_2 h^{2r} \int_{S_j} \varphi_\beta^{2r} \psi^2 \, ds,$$

where C_1 and C_2 depend on ε . Assume then that $z_i \in \bar{S}_j$, $1 \leq i < I$. Using the fact that

$$C_1 \int_0^1 t^\alpha p^2 dt \leq \int_0^1 p^2 dt \leq C_2 \int_0^1 t^\alpha p^2 dt$$

for $\alpha > -1$ and for any polynomial p of degree k , with C_1 and C_2 depending on α and k , we conclude by scaling that if $\psi \in N_{\beta,k}^h$ or $\psi \in M_{\beta,k|\partial\Omega}^h$ and h is sufficiently small, then

$$\begin{aligned} C_1 d(S_j)^{-\alpha} \int_{S_j} |x - z_i|^\alpha \psi^2 ds \\ \leq \int_{S_j} \psi^2 ds \leq C_2 d(S_j)^{-\alpha} \int_{S_j} |x - z_i|^\alpha \psi^2 ds, \quad \alpha > -1. \end{aligned}$$

Choosing $\alpha = 2r\beta_i$ and noting that, by B(i) and (4.1),

$$C_1 h^{2r} \leq d(S_j)^{2r(1-\beta_i)} \leq C_2 h^{2r},$$

we again obtain (4.11), with C_1 and C_2 now depending on ε , $r\beta_i$ and k .

Since (4.11) holds for all $S_j \in \tau_\beta^h(\partial\Omega)$, the assertion follows by summing over j . \square

We note that, by Lemma 2.4 and Lemma 2.1, we have the inequality

$$\int_{\partial\Omega} \varphi_\beta^{-1} u^2 ds \leq C \|u\|_{H^1(\Omega)}^2, \quad u \in H^1(\Omega).$$

In view of this, if we define

$$\|(u, \psi)\|_{\beta,h}^2 = \int_\Omega |D^1 u|^2 dx + h^{-1} \int_{\partial\Omega} \varphi_\beta^{-1} u^2 ds + h \int_{\partial\Omega} \varphi_\beta \psi^2 ds,$$

then $\|\cdot\|_{\beta,h}$ is a norm in the space $H^1(\Omega) \times W^{0,\beta/2}(\partial\Omega)$. We denote the normed space $(H^1(\Omega) \times W^{0,\beta/2}(\partial\Omega), \|\cdot\|_{\beta,h})$ by $X_{\beta,h}$.

Let the bilinear form \mathfrak{B} be as in (3.1). We replace from here on the weak formulation (1.2) by the following more convenient variational problem:

$$(4.12) \quad (u, \psi) \in X_{\beta,h}: \mathfrak{B}(u, \psi; v, \phi) = \int_\Omega f v dx + \int_{\partial\Omega} g \phi ds, \quad (v, \phi) \in X_{\beta,h}.$$

In the sequel it is of importance that (4.12) is solvable under weaker assumptions on f and g than (1.2). From (3.1) one concludes immediately that the bilinear form \mathfrak{B} is bounded on $X_{\beta,h} \times X_{\beta,h}$:

$$(4.13) \quad |\mathfrak{B}(u, \psi; v, \phi)| \leq \|(u, \psi)\|_{\beta,h} \|(v, \phi)\|_{\beta,h}.$$

Moreover, by Theorem 3.1 and Lemma 4.2,

$$(4.14) \quad \inf_{(u,\psi) \in M_{\beta,k}^h \times N_{\beta,k}^h} \sup_{(v,\phi) \in M_{\beta,k}^h \times N_{\beta,k}^h} \frac{\mathfrak{B}(u, \psi; v, \phi)}{\|(u, \psi)\|_{\beta,h} \|(v, \phi)\|_{\beta,h}} > C > 0.$$

From (1.4) and (4.12) through (4.14) we conclude, by classical arguments (see [3]), the following result:

THEOREM 4.1. *Assume that problem (4.11) has a solution $(u, \psi) \in H^1(\Omega) \times W^{0,\beta/2}(\partial\Omega)$, $0 < \beta_i < 1$, $i = 1, \dots, I$. Then, if (u_h, ψ_h) is the solution of (1.4) with $M^h = M_{\beta,k}^h$, $N^h = N_{\beta,k}^h$, $h \in (0, h_0)$, $h_0 = h_0(\Omega) \in (0, 1)$, we have the error estimate*

$$\|(u, \psi) - (u_h, \psi_h)\|_{\beta,h} \leq C \min_{(v,\phi) \in M_{\beta,k}^h \times N_{\beta,k}^h} \|(u, \psi) - (v, \phi)\|_{\beta,h},$$

where C depends on β , k and ε .

5. Rate of Convergence. We state first some lemmas which pertain to the approximation properties of polynomials in weighted Sobolev spaces. In what follows, T denotes a triangle with vertices at $(0, 0)$, $(1, 0)$ and $(0, 1)$.

Our first lemma is from [4].

LEMMA 5.1. *Let $\alpha \neq 0$ and let u be defined on T such that $\int_T |x|^\alpha |D^1 u|^2 dx < \infty$. Then there exists a constant q , depending on u and α , and a constant $C > 0$, depending on α , such that*

$$(5.1) \quad \int_T |x|^{\alpha-2} |u - q|^2 dx \leq C \int_T |x|^\alpha |D^1 u|^2 dx.$$

LEMMA 5.2. *Let k be an integer, $k \geq 1$, $\alpha \in R^1$, $\alpha \notin \{0, 2, \dots, 2k - 2\}$, and let u be defined on T such that $\int_T |x|^\alpha |D^k u|^2 dx < \infty$. Then there exists a polynomial p of degree $k - 1$ and a constant C , depending on α , k , such that*

$$(5.2) \quad \sum_{j=0}^{k-1} \int_T |x|^{\alpha+2j-2k} |D^j(u - p)|^2 dx \leq C \int_T |x|^\alpha |D^k u|^2 dx.$$

Proof. We first apply Lemma 5.1 to find constants q_{ij} , $i + j = k - 1$, such that

$$\int_T |x|^{\alpha-2} \left(\frac{\partial^{k-1} u}{\partial x_1^i \partial x_2^j} - q_{ij} \right)^2 dx \leq C \int_T |x|^\alpha |D^k u|^2 dx.$$

Then define

$$p_{k-1}(x) = \sum_{i+j=k-1} \frac{q_{ij}}{i!j!} x_1^i x_2^j.$$

Let $u_{k-1} = u - p_{k-1}$, and apply again Lemma 5.1 to find constants q_{ij} , $i + j = k - 2$, such that

$$\int_T |x|^{\alpha-4} \left(\frac{\partial^{k-2} u_{k-1}}{\partial x_1^i \partial x_2^j} - q_{ij} \right)^2 dx \leq C \int_T |x|^{\alpha-2} |D^{k-1} u_{k-1}|^2 dx.$$

Further, define $u_{k-2} = u_{k-1} - p_{k-2}$, where

$$p_{k-2} = \sum_{i+j=k-2} \frac{q_{ij}}{i!j!} x_1^i x_2^j.$$

Continuing in this way, one finally finds the homogeneous polynomials p_j of degree j , $0 \leq j \leq k - 1$, such that if $u_j = u - \sum_{l=j}^{k-1} p_l$ for $0 \leq j \leq k - 1$, and $u_k = u$, then

$$\begin{aligned} & \int_T |x|^{\alpha+2j-2k} |D^j u_j|^2 dx \\ & \leq C \int_T |x|^{\alpha+2j+2-2k} |D^{j+1} u_{j+1}|^2 dx, \quad j = 0, \dots, k - 1. \end{aligned}$$

Noting that $D^j u_j = D^j u_0$, the assertion follows by taking $p = \sum_{j=0}^{k-1} p_j$. \square

Let Σ_k denote the set of nodal points on \bar{T} associated to a reference Lagrange element of degree k , $k \geq 1$ (cf. [7]). For a continuous function u , defined on T , we let $(u)_k$ denote the k th order Lagrange interpolant to u , i.e., $(u)_k$ is a polynomial of degree k such that $(u)_k(x) = u(x)$ for $x \in \Sigma_k$. We have

LEMMA 5.3. *Let $k \geq 2$, let $2k - 4 < \alpha < 2k - 2$, and let u be defined on T such that $\int_T |x|^\alpha |D^k u|^2 dx < \infty$. Then (5.2) holds for a polynomial $p = (u)_k$.*

Proof. Let p be the polynomial of Lemma 5.2. Then we have

$$(5.3) \quad (u)_k = (u - p)_k + p.$$

From Lemma 2.2 and from (5.2), we have

$$(5.4) \quad \begin{aligned} |(u - p)(x)|^2 &\leq C \int_T |x|^\alpha |D^k u|^2 dx, \quad x \in T, \\ (u - p)(0) &= 0. \end{aligned}$$

Then also $(u - p)_k(0) = 0$, so we have

$$\sum_{l=0}^{k-1} \int_T |x|^{\alpha+2l-2k} |D^l(u - p)_k|^2 dx < \infty, \quad \alpha > 2k - 4.$$

Further, by the definition of the interpolant and by (5.4),

$$(5.5) \quad \begin{aligned} \sum_{l=0}^{k-1} \int_T |x|^{\alpha+2l-2k} |D^l(u - p)_k|^2 dx \\ \leq C \sum_{x \in \Sigma_k} |(u - p)(x)|^2 < C_1 \int_T |x|^\alpha |D^k u|^2 dx. \end{aligned}$$

The assertion now follows by combining (5.3), (5.5), and (5.2) and using the triangle inequality. \square

Using the above results we now prove the first approximation theorem.

THEOREM 5.1. *Let $k \geq 1$, and let $\beta \in R^I$ be such that $0 < \beta_i < 1$, $i = 1, \dots, I$. Further, let u be defined on Ω such that $\int_\Omega \varphi_\beta^{2k} |D^{k+1} u|^2 dx < \infty$. Then, if $h \in (0, h_0)$, $h_0 = h_0(\Omega) \in (0, 1)$, there exists a constant C , depending on k, β, ε , such that*

$$\begin{aligned} \min_{u \in M_{\beta,k}^h} \left\{ \int_T |D^1(u - v)|^2 dx + h^{-1} \int_{\partial\Omega} \varphi_\beta^{-1} |u - v|^2 ds \right\} \\ \leq Ch^{2k} \int_\Omega \varphi_\beta^{2k} |D^{k+1} u|^2 dx. \end{aligned}$$

Proof. We begin by introducing an interpolant of u . Let T be the reference triangle as above. Then, if $T_j \in \tau_\beta^h(\Omega)$, we can write

$$(5.6) \quad T = (G_j \circ A_j)(T_j), \quad T_j \in \tau_\beta^h(\Omega),$$

where A_j is an affine transformation which maps the corners of T_j onto those of T , and G_j is a smooth transformation which straightens the curved sides (if any) of $A_j(T_j)$. If \bar{T}_j contains a corner z_i of $\partial\Omega$, then we assume A_j to be chosen so that $A_j(z_i) = \{0\}$.

With the nodal set Σ_k defined as above, let $\Sigma_k^j = (G_j \circ A_j)^{-1}(\Sigma_k)$, and define u^j as the polynomial of degree k on T_j such that $u^j(x) = u(x)$ for $x \in \Sigma_k^j$. Since the sides of the triangles $T_j \in \tau_\beta^h(\Omega)$ become straight in the limit $h \rightarrow 0$, we see that, if h is sufficiently small, then u^j is uniquely defined for all $T_j \in \tau_\beta^h(\Omega)$. We let $u_{\beta,k}^h \in M_{\beta,k}^h$ denote the interpolant defined as

$$u_{\beta,k}^h(x) = u^j(x), \quad x \in T_j \in \tau_\beta^h(\Omega).$$

We will first prove that, under the assumptions made,

$$(5.7) \quad \int_\Omega |D^1(u - u_{\beta,k}^h)|^2 dx \leq Ch^{2k} \int_\Omega \varphi_\beta^{2k} |D^{k+1} u|^2 dx.$$

Let $T_j \in \tau_\beta^h(\Omega)$ be a triangle which does not touch any of the corners of $\partial\Omega$. Then we have the classical result

$$\int_{T_j} |D^1(u - u^j)|^2 dx \leq Cd(T_j)^{2k} \int_{T_j} |D^{k+1}u|^2 dx,$$

where C depends only on k and ε for h sufficiently small. Using B(ii), this can be further written as

$$(5.8) \quad \int_{T_j} |D^1(u - u^j)|^2 dx \leq Ch^{2k} \int_{T_j} \varphi_\beta^{2k} |D^{k+1}u|^2 dx.$$

We show next that (5.8) also holds when \bar{T}_j contains a corner z_i of $\partial\Omega$. We note first that Lemma 5.1 holds also when T is replaced by $A_j(T_j)$, A_j as in (5.6), provided that h is small enough. Moreover, the constant C in (5.1) does not depend on j . This follows, since $A_j(T_j) = G_j^{-1}(T)$, and the Jacobian of G_j^{-1} tends to unity uniformly in j as $h \rightarrow 0$. Then Lemma 5.2 holds also when T is replaced by $A_j(T_j)$, and so does Lemma 5.3, if Σ_k is replaced by $G_j^{-1}(\Sigma_k)$ in the definition of the interpolant $(u)_k$.

Now choose any $\alpha \in R^1$ so that $2k - 2 < \alpha < 2k$, $\alpha \geq 2k\beta_i$. Then, using the above modifications in Lemma 5.3, we conclude that, if $\tilde{u}(x) = u(A_j^{-1}(x))$, $\tilde{u}^j(x) = u^j(A_j^{-1}(x))$, $x \in A_j(T_j)$, then

$$\int_{A_j(T_j)} |x|^{\alpha-2k} |D^1(\tilde{u} - \tilde{u}^j)|^2 dx \leq C \int_{A_j(T_j)} |x|^\alpha |D^{k+1}\tilde{u}|^2 dx.$$

This inequality is invariant in scaling. Therefore, and taking into account the assumption B(i), the affine transformation A_j^{-1} can only introduce a dependence on ε in the constant C . Thus, we have

$$(5.9) \quad \int_{T_j} |x - z_i|^{\alpha-2k} |D^1(u - u^j)|^2 dx \leq C \int_{T_j} |x - z_i|^\alpha |D^{k+1}u|^2 dx,$$

where C depends on k, α, ε . Using (4.1), we have here the further estimates

$$\begin{aligned} |x - z_i|^{\alpha-2k} &\geq Ch^{(\alpha-2k)/(1-\beta_i)}, \\ |x - z_i|^\alpha &\leq C[\varphi_\beta(x)]^{2k} h^{(\alpha-2k\beta_i)/(1-\beta_i)}, \quad x \in T_j, C > 0. \end{aligned}$$

Together with (5.9), these prove (5.8).

Having verified that (5.8) holds for all $T_j \in \tau_\beta^h(\Omega)$, it suffices to sum over j to prove (5.7).

We prove next the estimate

$$(5.10) \quad h^{-1} \int_{\partial\Omega} \varphi_\beta^{-1} |u - u_{\beta,k}^h|^2 ds \leq Ch^{2k} \int_{\partial\Omega} \varphi_\beta^{2k} |D^{k+1}u|^2 dx.$$

Let $T_j \in \tau_\beta^h(\Omega)$ be a triangle such that T_j has a curved side S_j on $\partial\Omega$, and let \tilde{u} and \tilde{u}^j be defined on $A_j(T_j) = G_j^{-1}(T)$ as above. Assume first that S_j does not touch any of the corners of $\partial\Omega$. Then we start from the inequality

$$\int_{A_j(S_j)} |\tilde{u} - \tilde{u}^j|^2 ds \leq C \|\tilde{u} - \tilde{u}^j\|_{H^1(A_j(T_j))}^2 \leq C_1 \int_{A_j(T_j)} |D^{k+1}\tilde{u}|^2 dx,$$

where C_1 depends only on k and ε for h sufficiently small. Applying to this the affine transformation A_j^{-1} , one gets

$$d(T_j)^{-1} \int_{S_j} |u - u^j|^2 ds \leq Cd(T_j)^{2k} \int_{T_j} |D^{k+1}u|^2 dx,$$

and further, using B(ii),

$$(5.11) \quad h^{-1} \int_{S_j} \varphi_\beta^{-1} |u - u^j|^2 ds \leq Ch^{2k} \int_{T_j} \varphi_\beta^{2k} |D^{k+1}u|^2 dx.$$

If $z_i \in \bar{S}_j$, we take $\alpha \in (2k - 2, 2k)$, $\alpha \geq 2k\beta_i$, and derive first from (2.6) the inequality

$$\begin{aligned} & \int_{A_j(S_j)} |x|^{\alpha-2k-1} |\tilde{u} - \tilde{u}^j|^2 ds \\ & \leq C \int_{A_j(T_j)} \{ |x|^{\alpha-2k-2} |\tilde{u} - \tilde{u}^j|^2 + |x|^{\alpha-2k} |D^1(\tilde{u} - \tilde{u}^j)|^2 \} dx. \end{aligned}$$

Applying on the right side the same reasoning as above and performing the affine transformation A_j^{-1} , we conclude that

$$\int_{S_j} |x - z_i|^{\alpha-2k-1} |u - u^j|^2 ds \leq C \int_{T_j} |x - z_i|^\alpha |D^{k+1}u|^2 dx.$$

Since, by (4.1),

$$\begin{aligned} |x - z_i|^{\alpha-2k-1} & \geq Ch^{(\alpha-2k-1+\beta_i)/(1-\beta_i)} [\varphi_\beta(x)]^{-1}, \\ |x - z_i|^\alpha & \leq Ch^{(\alpha-2k\beta_i)/(1-\beta_i)} [\varphi_\beta(x)]^{2k}, \\ & 2k\beta_i < \alpha < 2k, x \in S_j, C > 0, \end{aligned}$$

we again obtain the estimate (5.11).

Upon summing over j in (5.11), the estimate (5.10) follows. Finally, combining (5.7) and (5.10), the proof of Theorem 5.1 is complete. \square

The second approximation theorem is as follows.

THEOREM 5.2. *Let $k \geq 1$, and let $\beta \in R^I$ be such that $0 < \beta_i < 1$, $i = 1, \dots, I$. Then, if $u \in W^{k+1, k\beta}(\Omega)$ and ψ is defined by (1.3), there exists a constant C , depending on k , β , ε and Ω , such that*

$$\min_{\phi \in N_{\beta, k}^h} \left\{ h \int_{\partial\Omega} \varphi_\beta |\psi - \psi|^2 ds \right\} \leq Ch^{2k} \|u\|_{W^{k+1, k\beta}(\Omega)}^2.$$

Proof. Consider first a function ψ defined on $\partial\Omega$ and sufficiently smooth on each Γ_i , $i = 1, \dots, I$. We use the notation of (4.2) and (4.3) to define an interpolant of ψ in $N_{\beta, k}^h$. First, let $\psi_i(t) = \psi(J_i(t))$, $t \in (0, 1)$, $i = 1, \dots, I$, and let $\psi_{i, k}$ denote the interpolant of ψ_i in the partitioning $\{I_j^{(i)}\}_{j=1}^{m_i}$, defined so that (i) if $2 \leq j \leq m_i - 1$, then $\psi_{i, k|I_j}$ equals the Lagrange interpolant of ψ_i of degree $\max\{1, k - 1\}$ on I_j^i , and (ii) if $j = 1$ or $j = m_i$, then $\psi_{i, k|I_j}$ is defined as a polynomial of degree $k - 1$ such that

$$(5.12) \quad \begin{aligned} & (\psi_i^{(l)} - \psi_{i, k}^{(l)})(t_1^i - 0) = 0, \\ & (\psi_i^{(l)} - \psi_{i, k}^{(l)})(t_{m_i-1}^i + 0) = 0, \quad l = 0, \dots, k - 1. \end{aligned}$$

We let $\psi_{\beta,k}^h$ be a function defined on $\partial\Omega$ such that

$$\psi_{\beta,k}^h(x) = \psi_{i,k}(J_i^{-1}(x)), \quad x \in \Gamma_i, i = 1, \dots, I.$$

It is clear from the definition of $N_{\beta,k}^h$ that $\psi_{\beta,k}^h \in N_{\beta,k}^h$. We will establish first the interpolation error estimates

$$(5.13) \quad h \int_{\partial\Omega} \varphi_{\beta} |\psi - \psi_{\beta,k}^h|^2 ds \leq Ch^{2k+1} \|\psi\|_{W^{k,(k+1/2)\beta}(\partial\Omega)}^2, \\ \psi \in W^{k,(k+1/2)\beta}(\partial\Omega), k = 1, 2, \dots$$

To prove (5.13), let $\nu = \nu(i)$ and $\mu = \mu(i)$ be as in (2.4). Then B(ii) implies the inequalities

$$C_1 t^{\beta_{\nu}} (1-t)^{\beta_{\mu}} \leq d(I_j^i) \leq C_2 t^{\beta_{\nu}} (1-t)^{\beta_{\mu}}, \quad t \in I_j^i, j = 2, \dots, m_i - 1,$$

where C_1 and C_2 depend only on ε . Combining these with the classical estimates of the error of Lagrange interpolation we get

$$(5.14) \quad h \int_{I_j^i} t^{\beta_{\nu}} (1-t)^{\beta_{\mu}} |\psi_i - \psi_{i,k}|^2 dt \\ \leq Ch^{2k+1} \int_{I_j^i} t^{(2k+1)\beta_{\nu}} (1-t)^{(2k+1)\beta_{\mu}} |\psi_i^{(k)}|^2 dt, \quad j = 2, \dots, m_i - 1.$$

Let us now show that (5.14) also holds for $j = 1$ and $j = m_i$. We need the following consequence of Hardy's inequality [8]: Let $\alpha \geq 0$, $k \geq 0$, and let f be defined on $(0, 1)$ so that $\int_0^1 t^{\alpha+2k} |f^{(k)}|^2 dt < \infty$, and so that $f^{(l)}(1) = 0$, $l = 0, \dots, k-1$. Then

$$(5.15) \quad \int_0^1 t^{\alpha} f^2 dt \leq C \int_0^1 t^{\alpha+2k} |f^{(k)}|^2 dt,$$

where C depends only on k .

Since (5.15) is invariant in scaling, we conclude from (5.12) and (5.15) that

$$(5.16) \quad \int_{I_j^i} t^{\beta_{\nu}} (1-t)^{\beta_{\mu}} |\psi_i - \psi_{i,k}|^2 dt \\ \leq C \int_{I_j^i} t^{\beta_{\nu}+2k} (1-t)^{\beta_{\mu}+2k} |\psi_i^{(k)}|^2 dt, \quad j \in \{1, m_i\}.$$

On the other hand, B(iii) implies that

$$t \leq Ch^{1/(1-\beta_{\nu})}, \quad t \in I_1^i,$$

and

$$1-t \leq Ch^{1/(1-\beta_{\mu})}, \quad t \in I_{m_i}^i.$$

Using these estimates in (5.16), it follows that (5.14) holds also for $j = 1$ and $j = m_i$. It then suffices to sum over i and j to prove (5.13).

In view of (5.13), the following estimates hold:

$$\min_{\phi \in N_{\beta,k}^h} \left\{ h \int_{\partial\Omega} \varphi_{\beta} |\psi - \phi|^2 ds \right\} \\ \leq Ch^{2k+1} \|\psi\|_{W^{k,(k+1/2)\beta}(\partial\Omega)}^2, \quad \psi \in W^{k,(k+1/2)\beta}(\partial\Omega) \\ \leq Ch^{2k-1} \|\psi\|_{W^{k-1,(k-1/2)\beta}(\partial\Omega)}^2, \quad \psi \in W^{k-1,(k-1/2)\beta}(\partial\Omega), \\ k = 1, 2, \dots$$

Here the second estimate is trivial if $k = 1$, and for $k > 1$ it follows from (5.13) and from the inclusion $N_{\beta,k-1}^h \subset N_{\beta,k}^h$. Upon interpolating between these estimates and using Lemma 2.3, we have

$$(5.17) \quad \min_{\phi \in N_{\beta,k}^h} \left\{ h \int_{\partial\Omega} \varphi_\beta |\psi - \phi|^2 ds \right\} \\ \leq Ch^{2k} \|\psi\|_{W^{k-1/2,k\beta}(\partial\Omega)}^2, \quad \psi \in W^{k-1/2,k\beta}(\partial\Omega), \quad k = 1, 2, \dots$$

To complete the proof, take ψ in (5.17) to be defined by (1.3), with $u \in W^{k+1,k\beta}(\Omega)$. Since n_i in (1.3) are smooth functions on each arc Γ_i , $i = 1, \dots, I$, and since the operator of multiplication by a smooth function is bounded in the weighted Sobolev spaces considered, we conclude from Lemma 2.4 that ψ satisfies

$$\|\psi\|_{W^{k-1/2,k\beta}(\partial\Omega)} \leq C \|u\|_{W^{k+1,k\beta}(\Omega)}.$$

Upon combining this with (5.17), the proof is complete. \square

Using the above approximation results, we can now estimate the error of the Lagrange multiplier method in terms of weighted norms.

THEOREM 5.3. *Let $\beta \in R^I$ be such that $\beta_i \geq 0$, $1 > \beta_i > 1 - \pi/\omega_i k$, $i = 1, \dots, I$. Further, let u be the solution of problem (1.1) with $f = f_1 + f_2$, $f_1 \in H^{k-1}(\Omega)$, $f_2 \in W^{k-1,k\beta}(\Omega)$ and $g = g_1 + g_2$, $g_1 \in H^{k+1}(\Omega)$, $g_2 \in W^{k+1,k\beta}(\Omega)$, and let ψ be defined by (1.3). Then, if $h \in (0, h_0)$, $h_0 = h_0(\Omega) \in (0, 1)$, and $M^h = M_{\beta,k}^h$, $N^h = N_{\beta,k}^h$, (1.4) has a unique solution (u_h, ψ_h) , and there exists a constant C , depending on k, β, ε and Ω , such that*

$$\int_{\Omega} \varphi_\beta^{-2} |u - u_h|^2 dx + h^2 \int |D^1(u - u_h)|^2 dx \\ + h \int_{\partial\Omega} \varphi_\beta^{-1} |u - u_h|^2 ds + h^3 \int_{\partial\Omega} \varphi_\beta |\psi - \psi_h|^2 ds \\ \leq Ch^{2k+2} \left\{ \|f_1\|_{H^{k-1}(\Omega)}^2 + \|f_2\|_{W^{k-1,k\beta}(\Omega)}^2 + \|g_1\|_{H^{k+1}(\Omega)}^2 + \|g_2\|_{W^{k+1,k\beta}(\Omega)}^2 \right\}.$$

Proof. Given k and β , which satisfy the assumptions of the theorem, we have, by Theorem 2.1,

$$(5.18) \quad u = u_1 + u_2, \quad u_1 \in H^{k+1}(\Omega), \quad u_2 \in W^{k+1,k\beta}(\Omega), \\ \|\mathbf{u}_1\|_{H^{k+1}(\Omega)}^2 + \|\mathbf{u}_2\|_{W^{k+1,k\beta}(\Omega)}^2 \\ \leq C \left\{ \|f_1\|_{H^{k-1}(\Omega)}^2 + \|f_2\|_{W^{k-1,k\beta}(\Omega)}^2 + \|g_1\|_{H^{k+1}(\Omega)}^2 + \|g_2\|_{W^{k+1,k\beta}(\Omega)}^2 \right\}.$$

Also, by (1.3) and Lemma 2.4 and by the trace properties of functions in Sobolev spaces $H^m(\Omega)$, ψ satisfies

$$(5.19) \quad \psi = \psi_1 + \psi_2, \quad \psi_1 \in H^{k-1/2}(\partial\Omega), \quad \psi_2 \in W^{k-1/2,k\beta}(\partial\Omega), \\ \|\psi_1\|_{H^{k-1/2}(\partial\Omega)}^2 + \|\psi_2\|_{W^{k-1/2,k\beta}(\partial\Omega)}^2 \leq C \left\{ \|\mathbf{u}_1\|_{H^{k+1}(\Omega)}^2 + \|\mathbf{u}_2\|_{W^{k+1,k\beta}(\Omega)}^2 \right\}.$$

From (5.19) and (2.5), we conclude that $\psi_2 \in W^{0,k\beta-(k-1/2)\theta}(\partial\Omega)$. Since $\psi_1 \in L_2(\partial\Omega)$ and $0 < \beta_i < 1$, $i = 1, \dots, I$, it follows that $\psi \in W^{0,\beta/2}(\partial\Omega)$. Hence, (u, ψ) is the solution of problem (4.11), and the error bound of Theorem 4.1 applies to the Lagrange multiplier method.

By Theorems 5.1 and 5.2, we have

$$(5.20) \quad \min_{v \in M_{\beta,k}^h} \left\{ \int_{\Omega} |D^1(u - v)|^2 dx + h^{-1} \int_{\partial\Omega} \varphi_{\beta} |u - v|^2 ds \right\} \\ \leq Ch^{2k} \int_{\Omega} \varphi_{\beta}^{2k} |D^{k+1}u|^2 dx,$$

and

$$(5.21) \quad \min_{\phi \in N_{\beta,k}^h} \left\{ h \int_{\partial\Omega} \varphi_{\beta} |\psi_2 - \phi|^2 ds \right\} \leq Ch^{2k} \|u_2\|_{W^{k+1,k,\theta}(\Omega)}^2.$$

Also, using the same type of reasoning as in the proof of Theorem 5.2, one gets

$$(5.22) \quad \min_{\phi \in N_{\beta,k}^h} \left\{ h \int_{\partial\Omega} \varphi_{\beta} |\psi_1 - \phi|^2 ds \right\} \\ \leq C \min_{\phi \in N_{\beta,k}^h} \left\{ h \int_{\partial\Omega} |\psi_1 - \phi|^2 ds \right\} \leq C_1 h^{2k} \|\psi_1\|_{H^{k-1/2}(\partial\Omega)}^2.$$

We finally estimate $\int_{\Omega} \varphi_{\beta}^{-2} |u - u_h|^2 dx$ using a duality argument. Consider the auxiliary problem

$$-\Delta V = \varphi_{\beta}^{-2}(u - u_h), \quad x \in \Omega, \\ V = 0, \quad x \in \partial\Omega,$$

the solution of which satisfies [8]

$$(5.23) \quad V \in W^{2,\beta}(\Omega); \\ \|V\|_{W^{2,\beta}(\Omega)} \leq C \|\varphi_{\beta}^{-2}(u - u_h)\|_{W^{0,\beta}(\Omega)} = C \left\{ \int_{\Omega} \varphi_{\beta}^{-2} |u - u_h|^2 dx \right\}^{1/2}.$$

By Lemma 2.4, if we define $\Xi = -\partial V / \partial n$, $x \in \partial\Omega$, then

$$(5.24) \quad \|\Xi\|_{W^{1/2,\beta}(\partial\Omega)} \leq C \|V\|_{W^{2,\beta}(\Omega)}.$$

By (2.5) and (5.24), we have $\Xi \in W^{1/2,\beta-\theta/2}(\partial\Omega)$, so $\Xi \in W^{0,\beta/2}(\partial\Omega)$. Noting also that, by Lemma 2.1,

$$\int_{\Omega} \varphi_{\beta}^{-2}(u - u_h)v dx \leq C \|u - u_h\|_{H^1(\Omega)} \|v\|_{H^1(\Omega)}, \quad v \in H^1(\Omega),$$

we see that the auxiliary problem admits the variational formulation

$$(5.25) \quad (V, \Xi) \in H^1(\Omega) \times W^{0,\beta/2}(\partial\Omega): \\ \mathfrak{B}(v, \phi; V, \Xi) = \int_{\Omega} \varphi_{\beta}^{-2}(u - u_h)v dx \quad \forall (v, \phi) \in H^1(\Omega) \times W^{0,\beta/2}(\partial\Omega),$$

where the bilinear form \mathfrak{B} is defined by (3.1).

By the same reasoning as above, we get, from (5.24) and (5.25), that

$$(5.26) \quad \min_{(v,\phi) \in M_{\beta,k}^h \times N_{\beta,k}^h} \left\{ \|(V - v, \Xi - \phi)\|_{\beta,h}^2 \right\} \leq Ch^2 \int_{\Omega} \varphi_{\beta}^{-2} |u - u_h|^2 dx.$$

Then, since (1.4), (5.25), and (4.13) imply that

$$(5.27) \quad \int_{\Omega} \varphi_{\beta}^{-2} |u - u_h|^2 dx = \mathfrak{B}(u - u_h, \psi - \psi_h; V - v, \Xi - \phi) \\ \leq \|(u - u_h, \psi - \psi_h)\|_{\beta,h} \|(V - v, \Xi - \phi)\|_{\beta,h} \quad \forall (v, \phi) \in M_{\beta,k}^h \times N_{\beta,k}^h,$$

we obtain by combining (5.26) and (5.27) the estimate

$$(5.28) \quad \int_{\Omega} \varphi_{\beta}^{-2} |u - u_h|^2 dx \leq Ch^2 \|(u - u_h, \psi - \psi_h)\|_{\beta, h}^2.$$

The asserted error bound now follows from the estimates (5.18) through (5.22), (5.28), and from Theorem 4.1. \square

Acknowledgment. The author would like to acknowledge a remark made by the referee, which led to an essential improvement in the proof of Lemma 4.1.

Institute of Mathematics
Helsinki University of Technology
SF-02150 Espoo 15, Finland

1. R. ADAMS, *Sobolev Spaces*, Academic Press, New York, 1975.
2. I. BABUŠKA, "The finite element method with Lagrange multipliers," *Numer. Math.*, v. 20, 1973, pp. 179–192.
3. I. BABUŠKA & A. K. AZIZ, "Survey lectures on the mathematical foundations of the finite element method," *The Mathematical Foundations of the Finite Element Method with Applications to Partial Differential Equations* (A. K. Aziz, Ed.), Academic Press, New York, 1972.
4. I. BABUŠKA, R. B. KELLOGG & J. PITKÄRANTA, "Direct and inverse error estimates for finite elements with mesh refinements," *Numer. Math.*, v. 33, 1979, pp. 447–471.
5. J. BERGH & J. LÖTSTRÖM, *Interpolation Spaces*, Springer-Verlag, Berlin, 1976.
6. F. BREZZI, "On the existence, uniqueness and approximations of saddle point problems arising from Lagrange multipliers," *RAIRO Anal. Numér.*, v. 8, 1974, pp. 129–151.
7. P. G. CIARLET, *The Finite Element Method for Elliptic Problems*, North-Holland, Amsterdam, 1978.
8. V. A. KONDRAT'EV, "Boundary problems for elliptic equations in domains with conical or angular points," *Trans. Moscow Math. Soc.*, v. 16, 1967, pp. 227–313.
9. J. PITKÄRANTA, "Boundary subspaces for the finite element method with Lagrange multipliers," *Numer. Math.*, v. 33, 1979, pp. 273–289.
10. J. PITKÄRANTA, "Local stability conditions for the Babuška method of Lagrange multipliers," *Math. Comp.*, v. 35, 1980, pp. 1113–1129.
11. H. TRIEBEL, "Interpolation theory for function spaces of Besov type defined in domains. II," *Math. Nachr.*, v. 58, 1973, pp. 63–86.