

A Class of Cubic Splines Obtained Through Minimum Conditions

By D. Bini and M. Capovani

Abstract. A class of cubic spline minimizing some special functional is investigated. This class is determined by the solution of a quadratic programming problem in which the minimizing function depends linearly on a parameter $\alpha < 2$. For $\alpha = 1/2$ natural splines are obtained. For $\alpha = -1$ the spline minimizing the mean value of the third derivative is obtained. It is shown that this spline has the best convergence order.

1. Introduction. Suppose that we are given a function $f \in C^4[a, b]$ and a mesh $\Delta_n = \{a = x_0 < x_1 < \dots < x_n = b\}$, such that $x_i = x_0 + ih$, $h = (b - a)/n$. A function $s_n(x)$ is called a cubic spline interpolant to $f(x)$ with respect to Δ_n if:

$$(1.1) \quad \begin{aligned} & s_n \text{ is a cubic polynomial on } [x_{i-1}, x_i], \quad i = 1, 2, \dots, n; \\ & s_n \in C^2[a, b]; \\ & s_n(x_i) = y_i, \quad y_i = f(x_i), \quad i = 0, 1, \dots, n. \end{aligned}$$

The importance of cubic spline interpolation is described in detail in [1]. Conditions (1.1) lead to a system of $n - 1$ equations and $n + 1$ unknowns. Two more equations can be given in various ways. For example, if we know $f'(x_0)$ and $f'(x_n)$ or $f''(x_0)$ and $f''(x_n)$ we can set $s'_n(x_i) = f'(x_i)$ or $s''_n(x_i) = f''(x_i)$, $i = 0, n$, obtaining in this way convergence of $s_n^{(i)}(x)$ to $f^{(i)}(x)$ of order $O(h^{4-i})$, $i = 0, 1, 2, 3$ uniformly over $[a, b]$ [5]. In the absence of such information, usually the following equations are considered

$$(1.2) \quad s''_n(x_0) = s''_n(x_n) = 0.$$

The kind of spline function obtained in this way is known as natural spline, and it is the spline interpolant to $f(x)$ minimizing the functional

$$F_2(s) = \int_a^b (s''(x))^2 dx,$$

which is related to the mean curvature of the graph of the function $s(x)$. For natural spline functions the following result holds [4]

$$\begin{aligned} |s_n^{(i)}(x) - f^{(i)}(x)| &= O(h^{2-i}), \quad a \leq x \leq b, \quad i = 0, 1; \\ |s_n^{(i)}(x) - f^{(i)}(x)| &= O(h^{4-i}), \quad \tilde{a} \leq x \leq \tilde{b}, \quad i = 0, 1, 2; \\ \tilde{a} - a &= b - \tilde{b} = O(h \log h). \end{aligned}$$

Received December 13, 1983; revised February 20, 1984 and February 4, 1985.
 1980 *Mathematics Subject Classification*. Primary 41A15, 65D05, 65D07; Secondary 90C20.

Recently, another variational condition has been analyzed [3]. The spline minimizing the functional

$$F_1(s) = \int_a^b (s'(x))^2 dx,$$

has been shown useful for computational purposes.

In this paper we consider cubic splines which minimize some special functionals. This set of splines is obtained by considering the functionals F_1 and F_2 , two other functionals F_0 and F_3 , involving the functions $s(x)$ and $s'''(x)$ respectively, and convex combinations of F_0, F_1, F_2 and F_3 .

In Section 2, we show that such splines can be obtained by solving a quadratic programming problem in which the minimizing function depends linearly on a parameter $\alpha, -1 \leq \alpha \leq 31/32$. In particular, for $\alpha = 1/2$ we find the natural spline, for $\alpha = 7/8$ the function minimizing F_1 . We widen the class by showing that for any $\alpha \leq 1$ the quadratic programming problem has a solution and that there exists a positive integer n_0 such that the quadratic programming problem has a solution for any $n \geq n_0$ if and only if $\alpha < 2$.

In Section 3, using the properties of a special class of matrices defined in [2], we determine the best spline interpolating f in the class of splines depending on $\alpha, \alpha < 2$. We show that the best spline in this class is the spline which minimizes the functional

$$F_3(s) = \sum_{i=0}^{n-1} \int_{x_i}^{x_{i+1}} (s'''(x))^2 dx,$$

obtained with $\alpha = 1$. For this spline we get the following result

$$\begin{aligned} |s_n^{(i)}(x) - f^{(i)}(x)| &= O(h^{3-i}), & a \leq x \leq b, & i = 0, 1, 2; \\ |s_n^{(i)}(x) - f^{(i)}(x)| &= O(h^{4-i}), & \tilde{a} \leq x \leq \tilde{b}, & i = 0, 1, 2, 3, \\ \tilde{a} - a &= b - \tilde{b} = O(h \log h). \end{aligned}$$

Thus, one gains one order of convergence and the convergence of one more derivative, compared to natural splines.

2. Preliminaries. Following Stoer & Bulirsch [5] we set

$$(2.1) \quad \begin{aligned} s_n(x) &= M_j(x_{j+1} - x)^3/(6h) + M_{j+1}(x - x_j)^3/(6h) + \alpha_j(x - x_j) + \beta_j, \\ x_j &\leq x \leq x_{j+1}, \quad j = 0, 1, \dots, n - 1; \end{aligned}$$

so that, from (1.1), we obtain

$$(2.2) \quad \begin{aligned} \beta_j &= f(x_j) - M_j h^2/6; \\ \alpha_j &= (f(x_{j+1}) - f(x_j))/h - h(M_{j+1} - M_j)/6; \end{aligned}$$

where the $(n + 1)$ -vector $M = (M_j)$ fulfills the relation

$$(2.3) \quad AM = b,$$

Then (2.6) has a solution if $-1 \leq \alpha \leq 31/32$ and this solution defines, through (2.1), (2.2), a cubic spline interpolating $f(x)$ which minimizes the functional

$$\begin{aligned} &F_0 \text{ if } \alpha = 31/32, \quad F_1 \text{ if } \alpha = 7/8, \\ &F_2 \text{ if } \alpha = 1/2, \quad F_3 \text{ if } \alpha = -1, \\ &\sum_{i=0}^3 \theta_i F_i, \text{ if } -1 \leq \alpha \leq 31/32, \text{ for some } \theta_i = \theta_i(\alpha) \geq 0. \end{aligned}$$

A similar result holds also in the case in which the knots x_i are not uniformly spaced.

It is interesting to point out that the same matrix formulation (2.6) allows us to deal with all the different conditions given in (2.5).

Two questions arise at this point:

– Are there other values of α , besides those in the range $[-1, 31/32]$, for which the problem (2.6) has a solution?

– Is there a best value of α in the range of the feasible values, for which the convergence of $s_n^{(i)}$ to $f^{(i)}$ is best possible?

In order to deal with these two questions we introduce a special class of matrices which has been used in [2].

Let τ_m be the linear space spanned by the set $\{I, H, H^2, \dots, H^{m-1}\}$, where $H = (h_{i,j})$ is the $m \times m$ matrix such that $h_{i,i+1} = h_{i+1,i} = 1, i = 1, 2, \dots, m - 1, h_{i,j} = 0$ otherwise. Observe that τ_m is closed under the row-column product, and $AB = BA$ for any $A, B \in \tau_m$, that is, τ_m is a commutative algebra; moreover, $A \in \tau_m$ if and only if

$$(2.7) \quad AH - HA = 0.$$

Writing down condition (2.7), we get

$$\begin{aligned} a_{i,j+1} + a_{i,j-1} &= a_{i+1,j} + a_{i-1,j}, \\ a_{0,j} = a_{i,0} = a_{m+1,j} &= a_{i,m+1} = 0, \quad i, j = 1, 2, \dots, m. \end{aligned}$$

The relation, called cross-sum condition, allows us to build up all the entries of any matrix $A \in \tau_m$ starting with the first row, or the first column, of A . Moreover, we have the following properties [2]:

If $A \in \tau_m$, then:

$$(2.8) \quad \begin{aligned} &A \text{ is symmetric, i.e., } a_{i,j} = a_{j,i}; \\ &A \text{ is persymmetric, i.e., } a_{i,j} = a_{m-i+1, m-j+1}; \\ &\text{If } F = \left(\sqrt{2/(m+1)} \sin(\pi ij/(m+1))\right) \text{ then } F \text{ is} \\ &\text{symmetric and orthogonal and } FAF \text{ is diagonal;} \\ &\text{The eigenvalues of } H \text{ are given by } 2 \cos(\pi i/(m+1)). \end{aligned}$$

Now consider problem (2.6) and partition the matrices A and A_α in the following way:

$$A_\alpha = \left[\begin{array}{c|c|c} 1 & \alpha & \\ \alpha & & \\ \hline & & \tilde{A}_\alpha \\ \hline & & & \alpha \\ & & & & 1 \end{array} \right], \quad A = \left[\begin{array}{c|c} 1 & \\ 0 & \\ \vdots & \\ 0 & \end{array} \right] \tilde{A} \left[\begin{array}{c} 0 \\ \vdots \\ 0 \\ 1 \end{array} \right].$$

Since

$$(2.9) \quad \tilde{A}_\alpha = 2I + \alpha H, \quad \tilde{A} = 4I + H,$$

we have $\tilde{A}_\alpha, \tilde{A} \in \tau_m, m = n - 1$. Expressing the components M_1, M_2, \dots, M_{n-1} as a function of M_0 and M_n , by using the constraints $AM = b$ and substituting in the quadratic function, we get

$$(2.10a) \quad M^T A_\alpha M = (M_0, M_n) \begin{bmatrix} \varphi & \psi \\ \psi & \varphi \end{bmatrix} \begin{bmatrix} M_0 \\ M_n \end{bmatrix} - 2(M_0, M_n) \begin{bmatrix} p_1 \\ p_2 \end{bmatrix};$$

$$\varphi = e_1^T B_\alpha e_1 = e_{n-1} B_\alpha e_{n-1}; \quad \psi = e_1^T B_\alpha e_{n-1} = e_{n-1}^T B_\alpha e_1;$$

$$(2.10b) \quad p_1 = e_1^T (B_\alpha + \alpha \tilde{A}^{-1} - I) b; \quad p_2 = e_{n-1}^T (B_\alpha + \alpha \tilde{A}^{-1} - I) b;$$

$$B_\alpha = I - 2\alpha \tilde{A}^{-1} + \tilde{A}^{-1} \tilde{A}_\alpha \tilde{A}^{-1} \in \tau_{n-1};$$

where e_1, e_{n-1} are the first and the last columns of the $(n - 1) \times (n - 1)$ identity matrix. Relations (2.10) hold in view of the symmetry and persymmetry of the matrix $B_\alpha \in \tau_{n-1}$.

Since the 2×2 matrix in (2.10a) is a principal submatrix of B_α , it is positive definite whenever B_α is positive definite. This fact allows us to prove the following

PROPOSITION 2. *For any $\alpha \leq 1$, the problem (2.6) has a solution.*

Proof. It is sufficient to prove that if $\alpha \leq 1$ the matrix B_α is positive definite. Now the eigenvalues λ_i of B_α can be obtained by (2.10b), (2.9) and (2.8). We have in fact

$$\lambda_i = 1 + (2 - 2\alpha(4 + c_i))/(4 + 2c_i)^2, \quad c_i = \cos(\pi i/n).$$

Therefore, B_α is positive definite if and only if $\alpha < (9 + 2c_i^2 + 8c_i)/(4 + c_i)$. Now, since $(9 + 2c_i^2 + 8c_i)/(4 + c_i) > 1, i = 1, 2, \dots, n - 1$, we get the result.

We can further widen the set of α for which problem (2.6) has a solution if we look at this issue from an asymptotic point of view. First we must compute the values to which φ and ψ converge as n tends to infinity.

LEMMA 1. *We have*

$$\varphi = (2 - \alpha)/\sqrt{3} + \theta_n, \quad |\theta_n| < (2/(3n^2))(1 + |\alpha|);$$

$$\psi = 0 + \theta'_n, \quad |\theta'_n| < (16/(3n^2))(1 + |\alpha|).$$

Proof. Since $B_\alpha \in \tau_{n-1}$, from (2.8), (2.9), (2.10b) we have

$$FB_\alpha F = \text{Diag}(\lambda_1, \lambda_2, \dots, \lambda_{n-1}),$$

$$\lambda_i = 1 + (2 - 2\alpha(4 + c_i))/(4 + 2c_i)^2, \quad c_i = \cos(\pi i/n).$$

Therefore, $\varphi = \varphi_1 + \alpha\varphi_2, \psi = \psi_1 + \alpha\psi_2$, where

$$(2.11a) \quad \varphi_1 = (2/n) \sum_{i=1}^{n-1} (1 + 2/(4 + 2c_i)^2) s_i^2,$$

$$\varphi_2 = (2/n) \sum_{i=1}^{n-1} ((8 + 2c_i)/(4 + 2c_i)^2) s_i^2, \quad s_i = \sin(\pi i/n);$$

$$(2.11b) \quad \psi_1 = (2/n) \sum_{i=1}^{n-1} (1 + 2/(4 + 2c_i)^2) (-1)^{i+1} s_i^2,$$

$$\psi_2 = (2/n) \sum_{i=1}^{n-1} ((8 + 2c_i)/(4 + 2c_i)^2) (-1)^{i+1} s_i^2.$$

Moreover, the quantities (2.11a) can be viewed as the result of applying the trapezoidal quadrature formula to suitable continuous functions with step π/n on the interval $[0, \pi]$. Therefore, we get (see [5, p. 121])

$$\begin{aligned} \varphi_1 &= (2/\pi) \int_0^\pi (1 + 2/(4 + 2 \cos x)^2) \sin^2 x \, dx + k_1/(6n^2); \\ \varphi_2 &= (2/\pi) \int_0^\pi ((8 + 2 \cos x)/(4 + 2 \cos x)^2) \sin^2 x \, dx + k_2/(6n^2); \end{aligned}$$

where $|k_i|$, $i = 1, 2$, can be bounded by the maximum absolute value which the second derivative of the corresponding integrand function takes on over $[0, \pi]$. Therefore, by evaluating the primitives and the second derivatives of these functions, we get

$$(2.12) \quad \begin{aligned} \varphi_1 &= 2/\sqrt{3} + k_1/(6n^2), & |k_1| &\leq 4, \\ \varphi_2 &= -1/\sqrt{3} + k_2/(6n^2), & |k_2| &\leq 3. \end{aligned}$$

Concerning ψ_1 and ψ_2 , observe that each formula in (2.11b) can be viewed as the difference of two quadrature formulas applied to the same function with the same step, but with different knots. Therefore, we have

$$(2.13) \quad \begin{aligned} \psi_1 &= 0 + 4k_3/(3n^2), & |k_3| &\leq 4, \\ \psi_2 &= 0 + 4k_4/(3n^2), & |k_4| &\leq 3. \end{aligned}$$

Now we are ready to prove the following

PROPOSITION 3. *There exists a positive integer n_0 such that, for every $n \geq n_0$, the problem (2.6) has a solution for any $f \in C^4[a, b]$ if and only if $\alpha < 2$.*

Proof. The assertion follows from Lemma 1, since the existence of a solution, for any $f \in C^4[a, b]$, is equivalent to the positive definiteness of the matrix $\begin{bmatrix} \varphi & \psi \\ \psi & \varphi \end{bmatrix}$.

3. Convergence. From relation (2.10a), under the hypothesis $\alpha < 2$, we have that the point in which the function $M^T A_\alpha M$ takes on its minimum value is given by:

$$\begin{bmatrix} M_0 \\ M_n \end{bmatrix} = \begin{bmatrix} \varphi & \psi \\ \psi & \varphi \end{bmatrix}^{-1} \begin{bmatrix} p_1 \\ p_2 \end{bmatrix},$$

that is,

$$(3.1) \quad M_0 = -(\psi p_2 - \varphi p_1)/(\varphi^2 - \psi^2), \quad M_n = -(\psi p_1 - \varphi p_2)/(\varphi^2 - \psi^2).$$

Our aim is to find out how M_0 and M_n depend on α , then determine the value of α for which one obtains best convergence. Since we already know φ and ψ , we have to compute p_1 and p_2 . As a first step of this evaluation, observe that from (2.10b) we have

$$p_1 = \sum_{i=1}^{n-1} u_i b_i, \quad u = (u_i), \quad u = (B_\alpha + \alpha \tilde{A}^{-1} - I) e_1.$$

Now, since $B_\alpha + \alpha \tilde{A}^{-1} - I = 2(1 - 2\alpha) \tilde{A}^{-2}$ we have

$$u = 2(1 - 2\alpha) \tilde{A}^{-2} e_1.$$

Moreover, since $\tilde{A}^{-2} \in \tau_{n-1}$ is symmetric and persymmetric, we have that the i th component of $\tilde{A}^{-2}e_{n-1}$ is equal to the i th component of $\tilde{A}^{-2}e_1$; therefore

$$(3.2) \quad p_1 = 2(1 - 2\alpha) \sum_{i=1}^{n-1} v_i b_i, \quad p_2 = 2(1 - 2\alpha) \sum_{i=1}^{n-1} v_{n-i} b_i, \\ v = (v_i), \quad v = \tilde{A}^{-2}e_1.$$

Now our task has been reduced to the computation of b and v . As far as the vector b is concerned we have the following

LEMMA 2. *If $f \in C^4[a, b]$ and $k = \max_{a \leq x \leq b} |f^{(4)}(x)|$, then*

$$b_i = 6(f''(a) + ihf'''(a) + h^2\gamma_i), \quad b_{n-i} = 6(f''(b) - ihf'''(b) + h^2\delta_i), \\ |\gamma_i|, |\delta_i| < (1/12 + i^2/2)k.$$

Proof. In view of (2.4), applying Taylor's formula to $f(x)$ at the point x_i with step h and $-h$ yields

$$b_i = 6(f''(x_i) + \xi_i h^2), \quad |\xi_i| < k/12.$$

Applying Taylor's formula again to the function $f''(x)$ at the point a and b with step ih and $-ih$, respectively, yields

$$f''(x_i) = f''(a) + ihf'''(a) + ((ih)^2/2)\eta_i, \\ f''(x_{n-i}) = f''(b) - ihf'''(b) + ((ih)^2/2)\nu_i, \quad |\eta_i|, |\nu_i| \leq k,$$

which completes the proof.

Concerning the computation of the vector v , we use the properties of class τ_m to prove the following

LEMMA 3. *Let $d = (2 - \sqrt{3})$. Then the vector $v = \tilde{A}^{-2}e_1$ satisfies the relation*

$$v_i = (-1)^{i+1} id^{i+1}(1 - d^{2n-2i}) / (1 - d^2) + \mu_i, \quad |\mu_i| < n^2 d^{n+1}.$$

Proof. First observe that $\tilde{A} = 4I + H$ and that the matrix $C = 4I - H$ satisfies $C = D\tilde{A}D$, where $D = \text{Diag}(1, -1, 1, -1, \dots, (-1)^n)$. Therefore,

$$C^{-2} = (1/16) \left(\sum_{j=0}^{\infty} H^j / 4^j \right)^2$$

has positive entries and, since $\tilde{A}^{-2} = DC^{-2}D$, we have $v_i = (-1)^{i+1}|v_i|$ where $|v_i|$ are the elements of the first row of C^{-2} .

Now, in order to compute $|v_i|$, observe that the elements c_i of the first row of C^{-1} are given by $c_i = d_{n-i-1}/d_{n-1}$, where d_i satisfies the difference equation

$$d_{i+1} = 4d_i - d_{i-2}, \quad d_0 = 1, \quad d_1 = 4.$$

By solving the above recurrence, we get $d_i = (d^{-i-1} - d^{i+1}) / (2\sqrt{3})$, where $d = (2 - \sqrt{3})$. Hence

$$(3.3) \quad c_i = (d^i - d^{2n-i}) / (1 - d^{2n}) = d^i + \gamma_i, \quad |\gamma_i| < d^{n+1}.$$

Now set $C^{-1} = G + E$, where $G \in \tau_{n-1}$ is defined by its first row $g_{1,j} = d^j$. Since $C \in \tau_{n-1}$, then $E \in \tau_{n-1}$, hence $GE = EG$ and, setting $|v| = (|v_i|)$, we have $|v| = (G + E)^2 e_1 = (G^2 e_1 + w)$, $w = 2EGe_1 + E^2 e_1$. Therefore $\|w\| \leq 2\|E\| \|Ge_1\| + \|E\| \cdot \|Ee_1\|$, where $\|w\| = \max |w_i|$ and $\|E\| = \max_i \sum_{j=1}^{n-1} |e_{i,j}|$. Now, from (3.3) we have $\|Ee_1\| < d^{n+1}$, $\|Ge_1\| \leq d$ and, using the cross-sum condition, $\|E\| \leq d^{n+1}n^2/4$, whence $\|w\| \leq d^{n+2}n^2/2 + d^{2n+2}n^2/4 < n^2 d^{n+1}$.

Now, to complete the proof, we must still show that the elements of $z = G^2e_1$ are given by $z_i = id^{i+1}(1 - d^{2n-2i})/(1 - d^2)$.

We proceed by induction on i . The result holds for $i = 1$ and $i = 2$, by construction of G . Suppose that it holds for r and $r - 1$; by the cross-sum condition we have

$$\begin{aligned} z_{r+1} &= \sum_{i=1}^{n-1} d^i g_{r+1,i} = \sum_{i=1}^{n-1} d^i (g_{r,i-1} + g_{r,i+1} - g_{r-1,i}) \\ &= d(z_r - d^{n-1}g_{r,n-1}) + d^{-1}(z_r - dg_{r,1}) - z_{r-1}. \end{aligned}$$

Since $g_{r,n-1} = d^{n-r}$, we get

$$z_{r+1} = z_r(d + d^{-1}) - z_{r-1} - d^{2n-r} - d^r,$$

and, by the inductive assumptions

$$\begin{aligned} z_{r+1} &= rd^{r+1}(1 - d^{2n-2r})(d + d^{-1})/(1 - d^2) \\ &\quad - (r - 1)d^r(1 - d^{2n-2r+2})/(1 - d^2) - d^{2n-r} - d^r \\ &= (r + 1)d^{r+2}(1 - d^{2n-2r-2})/(1 - d^2), \end{aligned}$$

which completes the proof.

From the above lemma we get the following relation:

$$v_i = \left((-1)^{i+1}id^{i+1}/(1 - d^2) + \theta_i \right), \quad |\theta_i| < 2n^2d^{n+1}.$$

Now we are ready to compute p_1 and p_2 . From the above relation, Lemma 2 and (3.2) we have

$$\begin{aligned} p_1 &= 12(1 - 2\alpha)(\sigma f''(a) + h\rho f'''(a) + \omega_1 h^2) + O(n^3d^n), \\ p_2 &= 12(1 - 2\alpha)(\sigma f''(b) + h\rho f'''(b) + \omega_2 h^2) + O(n^3d^n), \end{aligned}$$

where

$$\begin{aligned} \sigma &= \sum_{i=1}^{n-1} (-1)^{i+1}id^{i+1}/(1 - d^2) + \sum_{i=1}^{n-1} \theta_i, \\ \rho &= \sum_{i=1}^{n-1} (-1)^{i+1}i^2d^{i+1}/(1 - d^2) + \sum_{i=1}^{n-1} i\theta_i, \\ |\omega_1|, |\omega_2| &\leq k \sum_{i=1}^{n-1} (1/12 + i^2/2)id^{i+1}/(1 - d^2). \end{aligned}$$

Now, by using the difference equation technique (see [5, p. 438]) it is easy to prove that

$$\begin{aligned} &\sum_{i=1}^{n-1} (-1)^{i+1}id^{i+1}/(1 - d^2) \\ &= \sqrt{3}/36 + d^{n+1}(n - d/(d + 1))/((1 + d)(1 - d^2)), \\ &\sum_{i=1}^{n-1} (-1)^{i+1}i^2d^{i+1}/(1 - d^2) \\ &= 1/36 + \left(n^2/(1 + d) - 2nd/(1 - d)^2 + d(d - 1)/(1 + d)^3 \right) d^{n+1}/(1 - d^2), \\ &|\omega_1|, |\omega_2| \leq k/12 \sum_{i=1}^{\infty} id^{i+1}/(1 - d^2) + k/2 \sum_{i=1}^{\infty} i^3d^{i+1}/(1 - d^2) \\ &= (k/\sqrt{12})(1/24 + 1) = 25k/(24\sqrt{12}) < k/3 \end{aligned}$$

where, for the sake of simplicity, we have assumed n even. Therefore, we have

$$(3.4) \quad \begin{aligned} p_1 &= (\sqrt{3}/3)(1 - 2\alpha)(f''(a) + (1/\sqrt{3})hf'''(a) + O(h^2)), \\ p_2 &= (\sqrt{3}/3)(1 - 2\alpha)(f''(b) + (1/\sqrt{3})hf'''(b) + O(h^2)). \end{aligned}$$

From relations (3.4), (3.1) and Lemma 1, we finally get the following

PROPOSITION 4. *The second derivatives M_0, M_n at the points a and b , respectively, of the spline function obtained by (2.6), are given by*

$$\begin{aligned} M_0 &= ((2\alpha - 1)/(\alpha - 2))(f''(a) + (1/\sqrt{3})hf'''(a) + O(h^2)), \\ M_n &= ((2\alpha - 1)/(\alpha - 2))(f''(b) + (1/\sqrt{3})hf'''(b) + O(h^2)). \end{aligned}$$

We are now ready to determine the value of α for which one obtains best convergence of $s_n(x)$ to $f(x)$. For this purpose we use the following result [4].

PROPOSITION 5. *If $s_n(x)$ is any cubic spline interpolating $f(x) \in C^4[a, b]$, then*

$$\max_{x_i \leq x \leq x_{i+1}} |s_n^{(j)}(x) - f^{(j)}(x)| \leq (1/2^{3-j})h^{2-j}L_i, \quad j = 0, 1, 2,$$

where $L_i = (1/8)h^2k + \max\{|f''(x_i) - M_i|, |f''(x_{i+1}) - M_{i+1}|\}$ and $|f^{(4)}(x)| \leq k, x \in [a, b]$.

In view of Proposition 5 we have to determine α in such a way that the values $|f''(x_i) - M_i|$ are as small as possible.

Let R be the $(n + 1)$ -vector whose components are $f''(x_i), i = 0, 1, \dots, n$, and consider the vector $R - M$. Using the cubic spline condition (2.3) and the definition of \tilde{A} in (2.9) yields

$$(3.5) \quad \begin{aligned} R - M &= (f''(a) - M_0, w^T \tilde{A}^{-1}, f''(b) - M_n)^T, \\ w &= \tilde{A}\tilde{R} - b + M_0e_1 + M_n e_{n-1}, \\ \tilde{R} &= (f''(x_1), f''(x_2), \dots, f''(x_{n-1}))^T. \end{aligned}$$

Since, by Taylor's formula,

$$\begin{aligned} f''(x_i) &= (f(x_{i-1}) - 2f(x_i) + f(x_{i+1}))/h^2 + (h^2/24)f^{(4)}(\xi_i) + (h^2/24)f^{(4)}(\eta_i), \\ & \quad x_i < \xi_i < x_{i+1}, \quad x_{i-1} < \eta_i < x_i, \end{aligned}$$

we can write

$$\begin{aligned} \tilde{A}\tilde{R} - b &= (-2f''(x_1) + f''(x_2) + (h^2/2)\theta_1, \\ & \quad f''(x_1) - 2f''(x_2) + f''(x_3) + (h^2/2)\theta_2, \dots, \\ & \quad f''(x_{n-3}) - 2f''(x_{n-2}) + f''(x_{n-1}) + (h^2/2)\theta_{n-2}, \\ & \quad f''(x_{n-2}) - 2f''(x_{n-1}) + (h^2/2)\theta_{n-1})^T, \\ |\theta_i| &\leq k = \max_{a \leq x \leq b} |f^{(4)}(x)|. \end{aligned}$$

Again using Taylor's formula, we find

$$\begin{aligned} f''(x_{i+1}) - 2f''(x_i) + f''(x_{i-1}) &= (h^2/2)(f^{(4)}(\tilde{\xi}_i) + f^{(4)}(\tilde{\eta}_i)), \\ x_i &< \tilde{\xi}_i < x_{i+1}, \quad x_{i-1} < \tilde{\eta}_i < x_i, \quad i = 1, 2, \dots, n - 1. \end{aligned}$$

Therefore, we have

$$(3.6) \quad \begin{aligned} w = & (M_0 - 2f''(x_1) + f''(x_2) + (h^2/2)\theta_1, (3/2)h^2\tilde{\theta}_2, \dots, (3/2)h^2\tilde{\theta}_{n-2}, \\ & f''(x_{n-2}) - 2f''(x_{n-1}) + M_n + (h^2/2)\theta_{n-1})^T, \\ & |\tilde{\theta}_i| \leq k. \end{aligned}$$

Now, from (3.5) and (3.6), the value of α for which we have the best uniform convergence on $[a, b]$ is $\alpha = -1$. In fact, in this case, since from Proposition 4

$$\begin{aligned} M_0 &= f''(a) + (h/\sqrt{3})f'''(a) + O(h^2), \\ M_n &= f''(b) + (h/\sqrt{3})f'''(b) + O(h^2), \end{aligned}$$

we have

$$(3.7) \quad \begin{aligned} R - M &= (-(h/\sqrt{3})f'''(a) + O(h^2), w^T\tilde{A}^{-1}, -(h/\sqrt{3})f'''(b) + O(h^2))^T, \\ w &= ((h/\sqrt{3})f'''(a) + O(h^2), (3/2)h^2\tilde{\theta}_2, \dots, \\ & (3/2)h^2\tilde{\theta}_{n-2}, (h/\sqrt{3})f'''(b) + O(h^2))^T. \end{aligned}$$

Using relations (3.7), we can prove the following

PROPOSITION 6. *The spline function $s_n(x)$ obtained by (2.6) with $\alpha = -1$ fulfills the conditions:*

$$|s_n^{(i)}(x) - f^{(i)}(x)| \leq (h/2)^{3-i}(\gamma + O(h)), \quad i = 0, 1, 2, a \leq x \leq b,$$

where $\gamma = \max(|f'''(a)|, |f'''(b)|)$.

Proof. Since

$$\begin{aligned} \|\tilde{A}^{-1}\| &= \left\| (1/4) \sum_{i=0}^{\infty} (-1)^{i+1} H^i / 4^i \right\| \\ &\leq (1/4) \sum_{i=0}^{\infty} (1/4^i) \|H\|^i = (1/4) \sum_{i=0}^{\infty} 1/2^i = 1/2, \end{aligned}$$

we have $\|\tilde{A}^{-1}w\| < (1/2)\|w\|$. Therefore, from (3.7) we get

$$\max_i |f''(x_i) - M_i| \leq (h/\sqrt{3}) \max(|f'''(a)|, |f'''(b)|) + O(h^2),$$

which, in view of Proposition 5, completes the proof.

We are now looking for numbers a', b' , such that $a < a' < b' < b$ and

$$\max_{a' \leq x \leq b'} |s_n^{(i)}(x) - f^{(i)}(x)| = O(h^{4-i}).$$

For this purpose, observe that from (3.7) we have

$$(3.8) \quad \begin{aligned} z = \tilde{A}^{-1}w &= ((h/\sqrt{3})f'''(a) + O(h^2))\tilde{A}^{-1}e_1 \\ &+ ((h/\sqrt{3})f'''(b) + O(h^2))\tilde{A}^{-1}e_{n-1} + (3/2)h^2\tilde{A}^{-1}y, \\ & y = (0, \tilde{\theta}_2, \tilde{\theta}_3, \dots, \tilde{\theta}_{n-1}, 0)^T. \end{aligned}$$

Moreover, from the proof of Lemma 3, we have that the elements of the first column of \tilde{A}^{-1} are given by $(-1)^{i+1}(d^i - d^{2n-i})/(1 - d^{2n})$, $d = 2 - \sqrt{3}$. Therefore, since $\|\tilde{A}^{-1}\| \leq 1/2$, $\|y\| \leq k$ and $\tilde{A} \in \tau_{n-1}$, we have

$$|z_i| \leq (3/4)h^2k + (h/\sqrt{3})(\gamma + O(h))(d^i - d^{2n-i} + d^{n-i} - d^{n+i})/(1 - d^{2n}),$$

where $\gamma = \max\{|f'''(a)|, |f'''(b)|\}$.

Now, if $\log h/\log d \leq i \leq n - \log h/\log d$, then

$$(d^i - d^{2n-i} + d^{n-i} - d^{n+i})/(1 - d^{2n}) \leq d^i + d^{n-i} + O(d^n) \leq 2h + O(d^n);$$

whence

$$(3.9) \quad |z_i| \leq h^2((3/4)k + (2/\sqrt{3})\gamma) + O(h^3),$$

if $\log h/\log d \leq i \leq n - \log h/\log d$.

Now we can apply Proposition 5 to get the following result.

PROPOSITION 7. *The spline function obtained by (2.6) with $\alpha = -1$ fulfills the conditions*

$$|s_n^{(i)}(x) - f^{(i)}(x)| \leq (1/2^{3-i})h^{4-i}((7/8)k + (2/\sqrt{3})\gamma + O(h)),$$

$a' \leq x \leq b', i = 0, 1, 2;$

$$|s_n^{(3)}(x) - f^{(3)}(x)| \leq 2h(k + 2\gamma/\sqrt{3}), \quad a' \leq x \leq b', x \neq x_j, j = 0, 1, \dots, n;$$

where $a' - a = b - b' = h \log h/\log d$ and $k = \max_{a \leq x \leq b} |f^{(4)}(x)|$, $\gamma = \max\{|f'''(a)|, |f'''(b)|\}$.

Proof. In view of (3.9) and (3.5) we have

$$\max_{a' \leq x_j \leq b'} |f''(x_j) - M_j| \leq h^2((3/4)k + (2/\sqrt{3})\gamma) + O(h^3).$$

Therefore, from Proposition 5, we have the result relative to $s_n^{(i)}(x) - f^{(i)}(x)$, $i = 0, 1, 2$. In the case of the third derivative, observe that the function $g(x) = s_n''(x) - f''(x)$ belongs to $C^2[x_i, x_{i+1}]$, therefore, applying Taylor's formula at the point $x \in [x_i, x_{i+1}]$ with increment

$$\tilde{h} = \begin{cases} -h/2 & \text{if } x > (x_{i+1} + x_i)/2, \\ h/2 & \text{if } x \leq (x_{i+1} + x_i)/2, \end{cases}$$

we get

$$s_n'''(x) - f'''(x) = (s_n''(x + \tilde{h}) - f''(x + \tilde{h}) - s_n''(x) + f''(x))/h - (\tilde{h}/2)f^{(4)}(\xi),$$

where ξ belongs to the interval with endpoints x and $x + \tilde{h}$. Therefore,

$$|s_n'''(x) - f'''(x)| \leq \max_{a' \leq x \leq b'} |s_n''(x) - f''(x)|/h + (h/4)k \leq 2h(k + (2/\sqrt{3})\gamma).$$

Final Remarks. We have shown that, among the splines interpolating to $f(x)$ and satisfying (2.6), the best one is the spline $S(x)$ minimizing the functional F_3 . The computation of $S(x)$ can be performed according to the following steps:

- (1) Compute $p_1 = 6 \sum_{i=1}^{n-1} v_i b_i$, $p_2 = 6 \sum_{i=1}^{n-1} v_{n-i} b_i$, (see (3.2));
- (2) Compute M_0 and M_n by means of (3.1);
- (3) Solve the linear system $\tilde{A}M = b - M_0 e_1 - M_n e_{n-1}$.

Compared to natural splines, here we have to compute M_0 and M_n with a computational overhead of $2n + O(1)$ multiplications. In this analysis we have assumed that the numbers φ, ψ and v_i , which do not depend on the data b_i , are given constants which can be precomputed once and for all. Actually, in view of (2.11a), the evaluation of φ and ψ costs a linear time. Moreover, since the matrix vector product Fa , where F is the matrix in (2.8), can be computed by means of fast Fourier transform algorithms, the evaluation of $v = \tilde{A}^{-2}e_1 = FD^{-2}Fe_1$, $D = \text{diag}(4 + 2 \cos(\pi i/(n + 1)))$, can be carried out in $O(n \log n)$ arithmetic operations.

Dipartimento di Matematica
Università di Pisa
56100 Pisa, Italy

Dipartimento di Informatica
Università di Pisa
56100 Pisa, Italy

1. J. H. AHLBERG, E. N. NILSON & J. L. WALSH, *The Theory of Splines and Their Applications*, Academic Press, New York, 1967.

2. D. BINI & M. CAPOVANI, "Spectral and computational properties of band symmetric Toeplitz matrices," *Linear Algebra Appl.*, v. 52/53, 1983, pp. 99–126.

3. A. GHIZZETTI, "Interpolazione con splines verificanti una opportuna condizione," *Calcolo*, v. 20, 1983, pp. 53–65.

4. D. KERSHAW, "A note on the convergence of interpolatory cubic splines," *SIAM J. Numer. Anal.*, v. 8, 1971, pp. 67–74.

5. J. STOER & R. BULIRSCH, *Introduction to Numerical Analysis*, Springer, Berlin, 1980.