

# An Optimal-Order Multigrid Method for P1 Nonconforming Finite Elements

By Susanne C. Brenner

**Abstract.** An optimal-order multigrid method for solving second-order elliptic boundary value problems using P1 nonconforming finite elements is developed.

**1. Introduction.** Let  $\Omega$  be a convex polygon in  $\mathbf{R}^2$ . Let  $f \in L^2(\Omega)$ ,  $\alpha \in C^1(\bar{\Omega})$  and  $\beta \in C^0(\bar{\Omega})$ . We assume there exist constants  $\alpha_0, \beta_0$  such that  $\alpha \geq \alpha_0 > 0$  and  $\beta \geq \beta_0$ , where  $\beta_0$  depends on the boundary condition. In this paper we develop an optimal-order multigrid method for solving the Dirichlet problem ( $\beta_0 = 0$ )

$$(1.1) \quad \begin{aligned} -\nabla \cdot (\alpha \nabla u) + \beta u &= f && \text{in } \Omega, \\ u &= 0 && \text{on } \partial\Omega \end{aligned}$$

and the Neumann problem ( $\beta_0 > 0$ )

$$(1.2) \quad \begin{aligned} -\nabla \cdot (\alpha \nabla u) + \beta u &= f && \text{in } \Omega, \\ \frac{\partial u}{\partial n} &= 0 && \text{on } \partial\Omega, \end{aligned}$$

using P1 nonconforming finite elements (cf. [3], [4], [10]). We refer the reader to [1], [7] and [9] for conforming multigrid methods. As in [1], our multigrid method will be described in a coordinate-free fashion.

Our method consists of smoothing on the current-grid and coarser-grid correction, as in the conforming multigrid method. The important difference in the nonconforming case is that  $V_{k-1} \not\subseteq V_k$ , where  $V_k$ 's are the finite element spaces on mesh level  $k$ . Hence we can no longer simply use the natural injection for the intergrid transfer of grid functions. The key idea is to define an operator  $I_{k-1}^k: V_{k-1} \rightarrow V_k$  that reduces to natural injection on continuous piecewise linear functions. By doing so, we can use the well-known analysis of the conforming multigrid method. We will show that the approximate solution satisfies the same type of error estimates as the discretization error and that it can be obtained in  $\mathcal{O}(n)$  steps, where  $n$  is the dimension of the discretized finite element space. Since our intergrid transfer operator does not preserve either the energy or the  $L^2$ -norm, the standard proof of convergence (cf. [2]) for the  $\mathcal{V}$ -cycle does not carry over directly. We will therefore only discuss a  $\mathcal{W}$ -cycle method, even though the  $\mathcal{V}$ -cycle method may be convergent.

The paper is organized as follows. We begin with a discussion of the notation and fundamental estimates from the theory of finite elements. The intergrid transfer operator is discussed in Section 3. Section 4 contains the results on the contracting

---

Received July 2, 1987; revised February 2, 1988.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 65N30, 65F10.

©1989 American Mathematical Society  
0025-5718/89 \$1.00 + \$.25 per page

property of the  $k$ th level iteration followed by the convergence theorems for the nested iteration in Section 5. The singular Neumann problem ( $\beta_0 = 0$  in (1.2)) will be treated in Section 6. A piecewise quadratic nonconforming finite element is discussed in the last section.

**2. Preliminaries and Notation.** Let  $V = W_2^1(\Omega)$  for the Neumann problem (1.2) and  $V = \{v \in W_2^1(\Omega) : v|_{\partial\Omega} = 0\}$  for the Dirichlet problem (1.1). Here,  $W_2^1(\Omega)$  denotes the usual Sobolev space (cf. [3]). The variational formulation for (1.1) and (1.2) is to find  $u \in V$  such that

$$(2.1) \quad a(u, v) = F(v) \quad \forall v \in V,$$

where

$$(2.2) \quad a(u, v) = \int_{\Omega} \alpha \nabla u \cdot \nabla v + \beta uv \quad \text{and} \quad F(v) = \int_{\Omega} f v.$$

Let  $\{\mathcal{T}^k\}$ ,  $k \geq 1$ , be a family of triangulations of  $\Omega$ , where  $\mathcal{T}^{k+1}$  is obtained by connecting the midpoints of the edges of the triangles in  $\mathcal{T}^k$ . Let  $h_k := \max_{T \in \mathcal{T}^k} \text{diam } T$  (therefore  $h_k = 2h_{k+1}$ ). Then there exist positive constants  $C_1, C_2$ , independent of  $k$ , such that

$$(2.3) \quad C_2 h_k^2 \leq |T| \leq C_1 h_k^2 \quad \forall T \in \mathcal{T}^k,$$

where  $|T|$  denotes the area of the triangle  $T$ . These constants depend only on the angles that appear in  $\mathcal{T}^1$ . Throughout this paper we let  $C$  and  $C_i$  denote generic constants independent of  $k$ .

For the Dirichlet problem (1.1), define the finite element space

$$(2.4) \quad V_k := \{v : v|_T \text{ is linear for all } T \in \mathcal{T}^k, v \text{ is continuous at the midpoints of the edges and } v = 0 \text{ at the midpoints on } \partial\Omega\}.$$

For the Neumann problem (1.2), we define  $V_k$  similarly but without any restrictions on  $v$  along the boundary of  $\Omega$ . Note that functions in  $V_k$  are not continuous. In other words,  $V_k$  is a nonconforming finite element space.

We also use a conforming finite element space for our analysis. Define

$$(2.5) \quad W_k := \{w : w|_T \text{ is linear for all } T \in \mathcal{T}^k, w \text{ is continuous on } \Omega \text{ and } w|_{\partial\Omega} = 0\}$$

for the Dirichlet problem (1.1). For the Neumann problem (1.2), we make no assumptions on  $w$  along the boundary of  $\Omega$ . The space  $W_k$  will only be used to obtain our theoretical estimates. We emphasize that it will *not* play any role in the actual multigrid algorithm. Observe that  $W_k = V_k \cap V = V_k \cap V_{k+1}$ .

Let  $\{\phi_1^k, \dots, \phi_{n_k}^k\}$  be the basis of  $V_k$  such that each  $\phi_j^k$  equals 1 at exactly one midpoint and equals 0 at all other midpoints. For any linear functions  $\psi, \phi$  on a triangle  $K$ ,

$$(2.6) \quad \int_K \psi \phi = \frac{1}{3} |K| \left( \sum_{i=1}^3 \psi(m_i) \phi(m_i) \right),$$

where the  $m_i$ 's are the midpoints of the sides of  $K$  (cf. [3, p. 183]). It follows that the  $\phi_i$ 's are orthogonal with respect to the  $L^2$ -inner product.

Similarity is an equivalence relation on triangles. For each equivalence class  $\mathcal{R}$ , there exist constants  $C_{\mathcal{R}} > 0$  and  $C'_{\mathcal{R}} > 0$  such that for any triangle  $T \in \mathcal{R}$  and  $v \in \mathcal{P}_1(T)$ , the space of first-degree polynomials on  $T$ , we have

$$C_{\mathcal{R}}\Theta(v) \leq \int_T |\nabla v|^2 \leq C'_{\mathcal{R}}\Theta(v).$$

Here,  $\Theta(v) = [v(m_1) - v(m_2)]^2 + [v(m_2) - v(m_3)]^2 + [v(m_3) - v(m_1)]^2$  and  $m_1, m_2, m_3$  are the midpoints of the sides of  $T$ . Since any triangle in  $\mathcal{T}^k$  ( $k = 1, 2, \dots$ ) is similar to a triangle in  $\mathcal{T}^1$ , there exist  $C_3, C_4 > 0$  such that

$$(2.7) \quad C_3\Theta(v) \leq \int_T |\nabla v|^2 \leq C_4\Theta(v)$$

for any  $v \in \mathcal{P}_1(T)$ ,  $T \in \mathcal{T}^k$ ,  $k = 1, 2, \dots$ . Moreover, as a consequence of (2.3), (2.6) and (2.7), there exists  $C > 0$  such that

$$(2.8) \quad \int_T |\nabla v|^2 \leq Ch_k^{-2} \|v\|_{L^2}^2$$

for  $v \in \mathcal{P}_1(T)$  and  $T \in \mathcal{T}^k$ .

For each  $k$ , define (on  $V_k + W_2^1(\Omega)$ )

$$(2.9) \quad a_k(u, v) := \sum_{T \in \mathcal{T}^k} \int_T (\alpha \nabla u \cdot \nabla v + \beta uv)$$

and

$$(2.10) \quad \|u\|_k := \sqrt{a_k(u, u)}.$$

The bilinear form  $a_k(\cdot, \cdot)$  is obviously symmetric and positive definite on  $V_k$ . The stiffness matrix representing  $a_k(\cdot, \cdot)$  with respect to the basis  $\{\phi_1^k, \dots, \phi_{n_k}^k\}$  has at most five entries per row. As a consequence of (2.8), we have

$$(2.11) \quad \|u\|_k \leq Ch_k^{-1} \|u\|_{L^2} \quad \forall u \in V_k.$$

We also note that if  $u, v \in W_2^1(\Omega)$ , then  $a_k(u, v) = a(u, v)$ .

We now recall some fundamental estimates from the theory of finite elements.

Let  $\Pi_k$  and  $\tilde{\Pi}_k$  be the interpolation operators associated with  $V_k$  and  $W_k$ , respectively. If  $u \in W_2^2(\Omega)$ , we have the following estimates for the interpolation error:

$$(2.12) \quad \|u - \Pi_k u\|_{L^2} + h_k \|u - \Pi_k u\|_k \leq Ch_k^2 \|u\|_{W_2^2}$$

and

$$(2.13) \quad \|u - \tilde{\Pi}_k u\|_{L^2} + h_k \|u - \tilde{\Pi}_k u\|_k \leq Ch_k^2 \|u\|_{W_2^2}$$

(cf. [3]).

Since  $f \in L^2(\Omega)$ , elliptic regularity implies that  $u \in W_2^2(\Omega)$  (cf. [6]). For the same  $f$ , let  $u_k \in V_k$  satisfy

$$a_k(u_k, v) = \int_{\Omega} f v \quad \forall v \in V_k$$

and let  $\tilde{u}_k \in W_k$  satisfy

$$a_k(\tilde{u}_k, v) = \int_{\Omega} f v \quad \forall v \in W_k.$$

Since  $V_k$  satisfies the patch test (cf. [8], [10]), we have the following estimate for the discretization error:

$$(2.14) \quad \|u - u_k\|_{L^2} + h_k \|u - u_k\|_k \leq Ch_k^2 \|u\|_{W_2^2}$$

(cf. [4], [10]). The estimate for the conforming discretization error is, of course, well known (cf. [3]):

$$(2.15) \quad \|u - \tilde{u}_k\|_{L^2} + h_k \|u - \tilde{u}_k\|_k \leq Ch_k^2 \|u\|_{W_2^2}.$$

In [1], it was shown that  $\tilde{u}_k$  could be calculated by an iterative procedure to within an accuracy comparable to the error estimated by (2.15) using an amount of work that is proportional to the number of unknowns, namely the dimension of  $W_k$ . Our main goal in this paper is to prove a corresponding result for the computation of  $u_k$ .

From the spectral theorem, there exist eigenvalues  $0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_{n_k}$  and eigenfunctions  $\psi_1, \psi_2, \dots, \psi_{n_k} \in V_k$ ,  $(\psi_i, \psi_j)_{L^2} = \delta_{ij}$  (= the Kronecker delta), such that  $a_k(\psi_i, v) = \lambda_i (\psi_i, v)_{L^2}$  for all  $v \in V_k$ . From (2.11), there exists  $C_5 > 0$  such that

$$(2.16) \quad \lambda_i \leq C_5 h_k^{-2}.$$

If  $v \in V_k$ , we can write  $v = \sum_{i=1}^{n_k} \nu_i \psi_i$ . The norm  $\|v\|_{s,k}$  is defined (cf. [1]) as follows:

$$(2.17) \quad \|v\|_{s,k} := \left( \sum_{i=1}^{n_k} \nu_i^2 \lambda_i^s \right)^{1/2}.$$

Note that  $\|v\|_{0,k} = \|v\|_{L^2}$  and  $\|v\|_{1,k} = \|v\|_k$ .

Finally, it follows from the Cauchy-Schwarz inequality that

$$(2.18) \quad |a_k(v, w)| \leq \|v\|_{1+t,k} \|w\|_{1-t,k}$$

for any  $t \in \mathbf{R}$  and  $v, w \in V_k$ .

**3. The Intergrid Transfer Operator.** For  $v \in V_{k-1}$  the intergrid transfer operator  $I_{k-1}^k: V_{k-1} \rightarrow V_k$  is defined as follows. Let  $p$  be a midpoint of a side of a triangle in  $\mathcal{T}^k$ . If  $p$  lies in the interior of a triangle in  $\mathcal{T}^{k-1}$ , then we define

$$(I_{k-1}^k v)(p) := v(p).$$

Otherwise, if  $p$  lies on the common edge of two adjacent triangles  $T_1$  and  $T_2$  in  $\mathcal{T}^{k-1}$ , then we define

$$(I_{k-1}^k v)(p) := \frac{1}{2} [v|_{T_1}(p) + v|_{T_2}(p)].$$

Note that the matrix for  $I_{k-1}^k$  with respect to the bases  $\{\phi_1^{k-1}, \dots, \phi_{n_{k-1}}^{k-1}\}$  and  $\{\phi_1^k, \dots, \phi_{n_k}^k\}$  has at most five entries per row.

From the definition of  $I_{k-1}^k$ , it is clear that

$$(3.1) \quad I_{k-1}^k v = v \quad \forall v \in W_{k-1} = V_k \cap V_{k-1} \subseteq V.$$

In other words,  $I_{k-1}^k|_{W_{k-1}}$  is just the natural injection.

LEMMA 1. *There exists  $C > 0$  such that*

$$(3.2) \quad \|I_{k-1}^k v\|_k \leq C \|v\|_{k-1}$$

and

$$(3.3) \quad \|I_{k-1}^k v\|_{L^2} \leq C \|v\|_{L^2}$$

for all  $v \in V_{k-1}$ .

*Proof.* The inequality (3.3) follows immediately from (2.6), the definition of  $I_{k-1}^k$  and the quasi-uniformity of the triangulations.

Inequality (3.2) can be deduced from (3.3) as follows. Given  $v \in V_{k-1}$ , define  $g \in V_{k-1}$  by

$$(3.4) \quad \int_{\Omega} g \phi = a_{k-1}(v, \phi) \quad \forall \phi \in V_{k-1},$$

$w \in W_{k-1}$  by

$$(3.5) \quad a_k(w, \phi) = \int_{\Omega} g \phi \quad \forall \phi \in W_{k-1},$$

and  $z \in V$  by

$$(3.6) \quad a(z, \phi) = \int_{\Omega} g \phi \quad \forall \phi \in V.$$

Then (3.3), (2.11), (2.14), (2.15) and elliptic regularity imply that

$$(3.7) \quad \begin{aligned} \|I_{k-1}^k v\|_k &\leq \|I_{k-1}^k(v-w)\|_k + \|w\|_k \\ &\leq Ch_k^{-1} \|I_{k-1}^k(v-w)\|_{L^2} + \|w-v\|_{k-1} + \|v\|_{k-1} \\ &\leq Ch_k^{-1} \|v-w\|_{L^2} + \|v\|_{k-1} \\ &\leq Ch_k^{-1} [\|v-z\|_{L^2} + \|w-z\|_{L^2}] + \|v\|_{k-1} \\ &\leq Ch_k \|z\|_{W_2^2} + \|v\|_{k-1} \leq Ch_k \|g\|_{L^2} + \|v\|_{k-1}. \end{aligned}$$

But

$$\|g\|_{L^2}^2 = a_{k-1}(v, g) \leq \|v\|_{k-1} \|g\|_{k-1} \leq Ch_{k-1}^{-1} \|v\|_{k-1} \|g\|_{L^2}.$$

Therefore,

$$(3.8) \quad \|g\|_{L^2} \leq Ch_{k-1}^{-1} \|v\|_{k-1}.$$

Combining (3.7) and (3.8), we obtain (3.2).  $\square$

**4. Contracting Properties of the  $k$ th Level Iteration.** The  $k$ th level iteration with initial guess  $z_0$  yields  $MG(k, z_0, G)$  as an approximate solution to the following problem.

Find  $z \in V_k$  such that  $a_k(z, v) = G(v) \quad \forall v \in V_k$ , where  $G \in V_k'$ .

For  $k = 1$ ,  $MG(1, z_0, G)$  is the solution obtained from a direct method. For  $k > 1$ ,  $MG(k, z_0, G) = z_m + I_{k-1}^k q_p$ , where the approximation  $z_m \in V_k$  is constructed recursively from the initial guess  $z_0$  and the equations

$$(z_i - z_{i-1}, v)_{L^2} = (\Lambda_k)^{-1} (G(v) - a_k(z_{i-1}, v)), \quad \forall v \in V_k, \quad 1 \leq i \leq m.$$

Here,  $\Lambda_k = C_5 h_k^{-2}$  (cf. (2.16)), which is greater than or equal to  $\max_{1 \leq i \leq n_k} \lambda_i$ , and  $m$  is an integer to be determined later. With respect to  $\{\phi_1^k, \dots, \phi_{n_k}^k\}$ ,  $z_m$

can be obtained from  $z_0$  by iterating a sparse band matrix because the  $\phi_i^k$ 's are  $L^2$ -orthogonal. The coarser-grid correction  $q_p \in V_{k-1}$  is obtained by applying the  $(k-1)$ -level iteration  $p$  times ( $p = 2, 3$ ). More precisely,

$$\begin{aligned} q_0 &= 0, \\ q_i &= MG(k-1, q_{i-1}, \overline{G}), \quad 1 \leq i \leq p, \end{aligned}$$

where  $\overline{G} \in V'_{k-1}$  is defined by  $\overline{G}(v) := G(I_{k-1}^k v) - a_k(z_m, I_{k-1}^k v) \forall v \in V_{k-1}$ .

The main result in this section is the following theorem.

**THEOREM 1.** *If the number of smoothing steps is large enough, then the  $k$ th level iteration is a contraction for both the energy norm and the  $L^2$ -norm. Moreover, the contraction number is independent of  $k$ .*

Theorem 1 is a trivial consequence of the following lemmas. In order to simplify the notation, we define the statements

$(S_k)$ : When the  $k$ th level iteration is applied to the problem of finding  $z \in V_k$  such that  $a_k(z, v) = G(v) \forall v \in V_k$ , we have  $\|z - MG(k, z_0, G)\|_k \leq \gamma \|z - z_0\|_k$ ,

and

$(\overline{S}_k)$ : When the  $k$ th level iteration is applied to the problem of finding  $z \in V_k$  such that  $a_k(z, v) = G(v) \forall v \in V_k$ , we have  $\|z - MG(k, z_0, G)\|_{L^2} \leq \overline{\gamma} \|z - z_0\|_{L^2}$ .

**LEMMA 2.** *There exists  $\gamma \in (0, 1)$  and an integer  $m \geq 1$ , both independent of  $k$ , such that*

$$(S_{k-1}) \text{ implies } (S_k).$$

**LEMMA 3.** *There exists  $\overline{\gamma} \in (0, 1)$  and an integer  $m \geq 1$ , both independent of  $k$ , such that*

$$(\overline{S}_{k-1}) \text{ implies } (\overline{S}_k).$$

Our analysis is based on estimates of the following errors. Let  $e_0 := z - z_0$ ,  $e_m := z - z_m$  and  $e_f := z - MG(k, z_0, G)$ . Also let  $e \in V_{k-1}$  satisfy

$$(4.1) \quad a_{k-1}(e, v) = \overline{G}(v) = a_k(e_m, I_{k-1}^k v) \quad \forall v \in V_{k-1},$$

and let  $\tilde{e} \in W_{k-1}$  satisfy

$$(4.2) \quad a_{k-1}(\tilde{e}, v) = \overline{G}(v) = a_k(e_m, I_{k-1}^k v) = a_k(e_m, v) \quad \forall v \in W_{k-1}.$$

As in the conforming case (cf. [1]), we have the following effects of the smoothing steps.

**LEMMA 4.** *There exists  $C > 0$  such that*

$$(4.3) \quad \|e_m\|_{L^2} \leq \|e_0\|_{L^2},$$

$$(4.4) \quad \|e_m\|_k \leq \|e_0\|_k$$

and

$$(4.5) \quad \| \|e_m\| \|_{2,k} \leq Ch_k^{-1} m^{-1/2} \| \|e_0\| \|_{1,k} = Ch_k^{-1} m^{-1/2} \|e_0\|_k.$$

From (4.1) and (3.2) we have

$$\|e\|_{k-1}^2 = a_k(e_m, I_{k-1}^k e) \leq \|e_m\|_k \|I_{k-1}^k e\|_k \leq C \|e_m\|_k \|e\|_{k-1}.$$

Therefore, there exists  $C > 0$  such that

$$(4.6) \quad \|e\|_{k-1} \leq C \|e_m\|_k \leq C \|e_0\|_k.$$

Since  $e$  and  $\tilde{e}$  are approximate solutions in different spaces to the same problem (cf. (4.1) and (4.2)), they are close to each other.

LEMMA 5. *There exists  $C > 0$  such that*

$$(4.7) \quad \|e - \tilde{e}\|_{L^2} \leq Ch_k m^{-1/2} \|e_0\|_{L^2}$$

and

$$(4.8) \quad \|e - \tilde{e}\|_{k-1} \leq Cm^{-1/2} \|e_0\|_k.$$

*Proof.* Let  $f_0 \in V_{k-1}$  satisfy

$$(f_0, v)_{L^2} = a_k(e_m, I_{k-1}^k v) \quad \forall v \in V_{k-1}.$$

It follows from (2.18) and (3.2) that

$$\begin{aligned} \|f_0\|_{L^2}^2 &= a_k(e_m, I_{k-1}^k f_0) \leq \|e_m\|_{2,k} \|I_{k-1}^k f_0\|_{0,k} \\ &\leq C \|e_m\|_{2,k} \|f_0\|_{L^2}. \end{aligned}$$

Therefore,

$$(4.9) \quad \|f_0\|_{L^2} \leq C \|e_m\|_{2,k}.$$

Let  $v_0 \in W_2^2(\Omega)$  satisfy

$$-\nabla \cdot (\alpha \nabla v_0) + \beta v_0 = f_0 \quad \text{in } \Omega, \quad v_0 = 0 \quad \text{on } \partial\Omega \quad \left( \frac{\partial v_0}{\partial n} = 0 \quad \text{on } \partial\Omega \right)$$

in the Dirichlet (Neumann) case. Note that  $e$  and  $\tilde{e}$  are the finite element (Galerkin) approximations to  $v_0$  in  $V_{k-1}$  and  $W_{k-1}$ , respectively. By (2.14) and (2.15), we have  $\|v_0 - e\|_{L^2} + h_{k-1} \|v_0 - e\|_{k-1} \leq Ch_{k-1}^2 \|v_0\|_{W_2^2}$  and  $\|v_0 - \tilde{e}\|_{L^2} + h_{k-1} \|v_0 - \tilde{e}\|_{k-1} \leq Ch_{k-1}^2 \|v_0\|_{W_2^2}$ .

Since  $h_{k-1} = 2h_k$ , it follows from the triangle inequality that

$$\|e - \tilde{e}\|_{L^2} + h_k \|e - \tilde{e}\|_{k-1} \leq Ch_k^2 \|v_0\|_{W_2^2}.$$

By elliptic regularity,  $\|v_0\|_{W_2^2} \leq C \|f_0\|_{L^2}$ . Therefore, from (4.9) we obtain

$$(4.10) \quad \|e - \tilde{e}\|_{L^2} + h_k \|e - \tilde{e}\|_{k-1} \leq Ch_k^2 \|e_m\|_{2,k}.$$

The inequalities (4.7) and (4.8) now follow from (4.10) and (4.5).  $\square$

Next, observe that from (4.2) and the fact that  $W_{k-1} \subseteq V$  we have an orthogonality relation,

$$(4.11) \quad a_k(e_m - \tilde{e}, v) = 0 \quad \forall v \in W_{k-1}.$$

The analysis of  $e_m - \tilde{e}$  is similar to the one used in conforming multigrid methods.

LEMMA 6. *There exists  $C > 0$  such that*

$$(4.12) \quad \|e_m - \tilde{e}\|_{L^2} \leq Cm^{-1/2} \|e_0\|_{L^2}$$

and

$$(4.13) \quad \|e_m - \tilde{e}\|_k \leq Cm^{-1/2} \|e_0\|_k.$$

*Proof.* By (4.11), (2.18) and (4.5) we have

$$(4.14) \quad \begin{aligned} \|e_m - \tilde{e}\|_k^2 &= a_k(e_m - \tilde{e}, e_m - \tilde{e}) = a_k(e_m - \tilde{e}, e_m) \\ &\leq \| |e_m - \tilde{e}| \|_{0,k} \| |e_m| \|_{2,k} \\ &\leq Ch_k^{-1} m^{-1/2} \| |e_m - \tilde{e}| \|_{0,k} \|e_0\|_k. \end{aligned}$$

We will use a duality argument to estimate  $\| |e_m - \tilde{e}| \|_{0,k} = \|e_m - \tilde{e}\|_{L^2}$ . Let  $w \in W_2^2(\Omega)$  satisfy

$$-\nabla \cdot (\alpha \nabla w) + \beta w = e_m - \tilde{e} \quad \text{in } \Omega, \quad w = 0 \quad \text{on } \partial\Omega \quad \left( \frac{\partial w}{\partial n} = 0 \text{ on } \partial\Omega \right)$$

in the Dirichlet (Neumann) case. Let  $w_k \in V_k$  satisfy

$$a_k(w_k, v) = \int_{\Omega} (e_m - \tilde{e})v \quad \forall v \in V_k.$$

Then

$$\begin{aligned} \|e_m - \tilde{e}\|_{L^2}^2 &= \int_{\Omega} (e_m - \tilde{e})(e_m - \tilde{e}) = a_k(w_k, e_m - \tilde{e}) \\ &= a_k(w - v, e_m - \tilde{e}) + a_k(w_k - w, e_m - \tilde{e}) \end{aligned}$$

for any  $v \in W_{k-1}$  by the orthogonality relation (4.11). Therefore, by the approximation property (2.13), the discretization error estimate (2.14) and elliptic regularity we have

$$\begin{aligned} \|e_m - \tilde{e}\|_{L^2}^2 &\leq \inf_{v \in W_{k-1}} \|w - v\|_k \|e_m - \tilde{e}\|_k + \|w_k - w\|_k \|e_m - \tilde{e}\|_k \\ &\leq Ch_k \|w\|_{W_2^2} \|e_m - \tilde{e}\|_k \leq Ch_k \|e_m - \tilde{e}\|_{L^2} \|e_m - \tilde{e}\|_k. \end{aligned}$$

Thus,

$$(4.15) \quad \|e_m - \tilde{e}\|_{L^2} \leq Ch_k \|e_m - \tilde{e}\|_k.$$

The inequality (4.13) now follows by combining (4.14) and (4.15).

From (4.15), (4.13) and (2.11) it follows that

$$\begin{aligned} \|e_m - \tilde{e}\|_{L^2} &\leq Ch_k \|e_m - \tilde{e}\|_k \\ &\leq Cm^{-1/2} h_k \|e_0\|_k \leq Cm^{-1/2} \|e_0\|_{L^2}. \quad \square \end{aligned}$$

*Proof of Lemma 2.* Recall that  $e_f = z - MG(k, z_0, G)$  and  $e_0 = z - z_0$ . We have by (3.1)

$$\begin{aligned} \|e_f\|_k &= \|e_m - I_{k-1}^k q_p\|_k \\ &\leq \|e_m - \tilde{e}\|_k + \|I_{k-1}^k(\tilde{e} - e)\|_k + \|I_{k-1}^k(e - q_p)\|_k. \end{aligned}$$

From (4.13), (4.8), (3.2) and  $(S_{k-1})$ , it follows that

$$\|e_f\|_k \leq Cm^{-1/2} \|e_0\|_k + C\gamma^p \|e\|_{k-1}.$$

But (4.6) yields

$$\|e_f\|_k \leq (Cm^{-1/2} + C\gamma^p) \|e_0\|_k.$$



If  $\gamma \in (0, 1)$  is small enough, then  $C\gamma^p < \gamma/2$  (since  $p > 1$ ). If  $m$  is large enough, then  $Cm^{-1/2} < \gamma/2$ . For such choices, we have

$$\|e_f\|_k \leq \gamma \|e_0\|_k. \quad \square$$

*Proof of Lemma 3.* We have by (3.1)

$$\|e_f\|_{L^2} \leq \|e_m - \tilde{e}\|_{L^2} + \|I_{k-1}^k(\tilde{e} - e)\|_{L^2} + \|I_{k-1}^k(e - q_p)\|_{L^2}.$$

By (4.12), (4.7), (3.3) and  $(\bar{S}_{k-1})$  it follows that

$$(4.16) \quad \|e_f\|_{L^2} \leq Cm^{-1/2} \|e_0\|_{L^2} + C\bar{\gamma}^p \|e\|_{L^2}.$$

But

$$\begin{aligned} \|e\|_{L^2} &\leq \|e - \tilde{e}\|_{L^2} + \|\tilde{e} - e_m\|_{L^2} + \|e_m\|_{L^2} \\ &\leq Cm^{-1/2} \|e_0\|_{L^2} + \|e_0\|_{L^2} \end{aligned}$$

by (4.7), (4.12) and (4.3). Therefore,  $\|e\|_{L^2} \leq C\|e_0\|_{L^2}$ . Hence from (4.16),

$$\|e_f\|_{L^2} \leq (Cm^{-1/2} + C\bar{\gamma}^p) \|e_0\|_{L^2}.$$

If  $\bar{\gamma} \in (0, 1)$  is small enough, then  $C\bar{\gamma}^p < \bar{\gamma}/2$ . If  $m$  is large enough, then  $Cm^{-1/2} < \bar{\gamma}/2$ . For such choices we have

$$\|e_f\|_{L^2} \leq \bar{\gamma} \|e_0\|_{L^2}. \quad \square$$

**5. Nested Iteration.** We have a sequence of discretizations for the problem (1.1) or (1.2). For each  $k$ , we want to find an approximate solution  $\hat{u}_k$  to the problem of finding  $u_k \in V_k$  such that

$$a_k(u_k, v) = \int_{\Omega} f v \quad \forall v \in V_k.$$

In the overall multigrid strategy,  $\hat{u}_1$  is obtained by a direct method. The approximations  $\hat{u}_k$  ( $k \geq 2$ ) are obtained recursively by

$$\begin{aligned} u_0^j &= I_{j-1}^j \hat{u}_{j-1}, \\ u_l^j &= MG(j, u_{l-1}^j, F), \quad 1 \leq l \leq r, \quad F(v) = \int_{\Omega} f v, \\ \hat{u}_j &= u_r^j. \end{aligned}$$

Here  $r$  is an integer to be determined.

Define  $\hat{e}_k := u_k - \hat{u}_k$ . In particular,  $\hat{e}_1 = 0$ . Lemma 2, (2.14), (2.13), (3.1) and (3.2) imply that

$$\begin{aligned} \|\hat{e}_k\|_k &\leq \gamma^r \|u_k - I_{k-1}^k \hat{u}_{k-1}\|_k \\ &\leq \gamma^r \{ \|u_k - u\|_k + \|u - \tilde{\Pi}_{k-1} u\|_k + \|I_{k-1}^k(\tilde{\Pi}_{k-1} u - \hat{u}_{k-1})\|_k \} \\ &\leq C\gamma^r \{ h_k \|u\|_{W_2^2} + \|\tilde{\Pi}_{k-1} u - \hat{u}_{k-1}\|_{k-1} \} \\ &\leq C\gamma^r \{ h_k \|u\|_{W_2^2} + \|\tilde{\Pi}_{k-1} u - u\|_{k-1} + \|u - u_{k-1}\|_{k-1} + \|u_{k-1} - \hat{u}_{k-1}\|_{k-1} \} \\ &\leq C\gamma^r \{ h_k \|u\|_{W_2^2} + h_{k-1} \|u\|_{W_2^2} + \|\hat{e}_{k-1}\|_{k-1} \}. \end{aligned}$$

Since  $h_{k-1} = 2h_k$ ,

$$(5.1) \quad \|\hat{e}_k\|_k \leq Ch_k \gamma^r \|u\|_{W_2^2} + C\gamma^r \|\hat{e}_{k-1}\|_{k-1}.$$

By iterating (5.1) we have

$$\begin{aligned} \|\hat{e}_k\|_k &\leq Ch_k\gamma^r \|u\|_{W_2^2} + C^2 h_{k-1}\gamma^{2r} \|u\|_{W_2^2} + \cdots + C^k h_1\gamma^{kr} \|u\|_{W_2^2} \\ &\leq \|u\|_{W_2^2} \frac{h_k C \gamma^r}{1 - 2C\gamma^r} \end{aligned}$$

if  $2C\gamma^r < 1$ . Therefore,

$$(5.2) \quad \|\hat{e}_k\|_k \leq Ch_k \|u\|_{W_2^2}.$$

In summary, we have proved the following theorem.

**THEOREM 2.** *If  $r$  is large enough, then there exists a constant  $C > 0$  such that*

$$\|u - \hat{u}_k\|_k \leq Ch_k \|u\|_{W_2^2}.$$

Similarly, we can prove the following theorem for the  $L^2$ -error.

**THEOREM 3.** *If  $r$  is large enough, then there exists a constant  $C > 0$  such that*

$$\|u - \hat{u}_k\|_k \leq Ch_k^2 \|u\|_{W_2^2}.$$

**THEOREM 4.** *The cost for obtaining  $\hat{u}_k$  is  $\mathcal{O}(n_k)$ , where  $n_k$  is the dimension of  $V_k$ .*

*Proof.* This is a consequence of the fact that  $p = 2, 3$  and that the number of nonzero entries in the stiffness matrices, the smoothing iteration matrices and the intergrid transfer matrices are all proportional to  $n_k$ . The proof is the same as the one in [1].  $\square$

**6. The Singular Neumann Problem.** When  $\beta \equiv 0$  in (1.2), the necessary and sufficient condition for the existence of a solution is

$$(6.1) \quad \int_{\Omega} f = 0.$$

If (6.1) is satisfied, there exists a unique solution  $u$  in  $\hat{V} = \{v \in W_2^1(\Omega) : \int_{\Omega} v = 0\}$ . The multigrid method developed in earlier sections can be modified to yield approximate solutions of  $u$ . Let  $\hat{V}_k = \{v \in V_k : \int_{\Omega} v = 0\}$  and  $\hat{W}_k = \{w \in W_k : \int_{\Omega} w = 0\}$ . Let  $u_k \in \hat{V}_k$  satisfy

$$(6.2) \quad a_k(u_k, v) = \int_{\Omega} f v \quad \forall v \in \hat{V}_k$$

and  $\tilde{u}_k \in \hat{W}_k$  satisfy

$$(6.3) \quad a_k(\tilde{u}_k, v) = \int_{\Omega} f v \quad \forall v \in \hat{W}_k.$$

Then

$$\begin{aligned} \|u - \tilde{u}_k\|_k &= \inf_{v \in \hat{W}_k} \|u - v\|_k \\ (6.4) \quad &\leq \left\| u - \left( \tilde{\Pi}_k u - \frac{1}{|\Omega|} \int_{\Omega} \tilde{\Pi}_k u \right) \right\|_k \\ &= \|u - \tilde{\Pi}_k u\|_k \leq Ch_k \|u\|_{W_2^2}. \end{aligned}$$

A duality argument shows that

$$(6.5) \quad \|u - \tilde{u}_k\|_{L^2} \leq Ch_k^2 \|u\|_{W_2^2}.$$

The analog of (2.15) therefore holds for the space  $\hat{W}_k$ .

To estimate  $\|u - u_k\|_k$ , we start with the formula (cf. [8])

$$(6.6) \quad \|u - u_k\|_k \leq \inf_{v \in \hat{V}_k} \|u - v\|_k + \sup_{v \in \hat{V}_k} \frac{|a_k(u - u_k, v)|}{\|v\|_k}.$$

By arguments similar to those that led to (6.4), we see that the first term is bounded by  $Ch_k\|u\|_{W_2^2}$ . The second term is also bounded by  $Ch_k\|u\|_{W_2^2}$ ; the analysis is similar to that in [4]. Therefore,

$$(6.7) \quad \|u - u_k\|_k \leq Ch_k\|u\|_{W_2^2}.$$

Again, a duality argument (cf. the proof of (4.15)) shows that

$$(6.8) \quad \|u - u_k\|_{L^2} \leq Ch_k^2\|u\|_{W_2^2}.$$

Thus the analog of (2.14) holds for  $\hat{V}_k$ .

The operator  $I_{k-1}^k$  defined in Section 3 must be modified so that it maps  $\hat{V}_{k-1}$  into  $\hat{V}_k$ . We define  $\hat{I}_{k-1}^k: \hat{V}_{k-1} \rightarrow \hat{V}_k$  by

$$(6.9) \quad \hat{I}_{k-1}^k v := I_{k-1}^k v - \frac{1}{|\Omega|} \int_{\Omega} I_{k-1}^k v.$$

The computation of the integral involves only  $\mathcal{O}(n_k)$  operations (using the quadrature formula (2.6)). Note that  $\hat{I}_{k-1}^k w = w$  for all  $w \in \hat{W}_{k-1}$ .

We have, by (3.2),

$$(6.10) \quad \|\hat{I}_{k-1}^k v\|_k = \|I_{k-1}^k v\|_k \leq C\|v\|_{k-1}$$

and, by the Cauchy-Schwarz inequality and (3.3),

$$(6.11) \quad \begin{aligned} \|\hat{I}_{k-1}^k v\|_{L^2} &= \left\| I_{k-1}^k v - \frac{1}{|\Omega|} \int_{\Omega} I_{k-1}^k v \right\|_{L^2} \\ &\leq \|I_{k-1}^k v\|_{L^2} + \left\| \frac{1}{|\Omega|} \int_{\Omega} I_{k-1}^k v \right\|_{L^2} \\ &\leq 2\|I_{k-1}^k v\|_{L^2} \leq C\|v\|_{L^2}. \end{aligned}$$

Because of (6.4), (6.5), (6.7), (6.8), (6.10) and (6.11), the theory developed in earlier sections carries over to this case if we replace  $V, V_k, W_k$  and  $I_{k-1}^k$  by  $\hat{V}, \hat{V}_k, \hat{W}_k$  and  $\hat{I}_{k-1}^k$ , respectively.

In practice we can use the same scheme with  $I_{k-1}^k$  replaced by  $\hat{I}_{k-1}^k$  and  $V_1$  replaced by  $\hat{V}_1$ . The solution obtained is in  $\hat{V}_k$  since the zero mean value is preserved by the intergrid transfer operator, the smoothing steps and the coarser-grid correction.

**7. Extension to a Quadratic Nonconforming Finite Element.** The principles which led to our optimal-order multigrid method for P1 nonconforming elements can also be applied to higher-order nonconforming finite elements. In this section we will indicate how this can be done for a quadratic nonconforming finite element (cf. [5]). For simplicity, we restrict our discussion to the Dirichlet problem for the Laplace equation

$$(7.1) \quad \begin{aligned} -\Delta u &= f \quad \text{in } \Omega, \\ u &= 0 \quad \text{on } \partial\Omega, \end{aligned}$$

where  $\Omega$  is a convex polygon.

Let  $V = \{v \in W_2^1(\Omega) : v|_{\partial\Omega} = 0\}$  and  $a(u, v) = \int_{\Omega} \nabla u \cdot \nabla v$  for  $u, v \in V$ . Then the variational formulation for (7.1) is to find  $u \in V$  such that

$$(7.2) \quad a(u, v) = \int f v \quad \forall v \in V.$$

We assume that  $u \in W_2^3(\Omega) \cap V$  in order to fully exploit the properties of the quadratic element.




FIGURE 1

Let  $K$  be a triangle. The barycentric coordinates of the six Gauss-Legendre points  $p_1, p_2, p_3, p_4, p_5, p_6$  along the sides of  $K$  (cf. Figure 1) are obtained by permuting

$$\left( \frac{1}{2} \left( 1 + \frac{\sqrt{3}}{3} \right), \frac{1}{2} \left( 1 - \frac{\sqrt{3}}{3} \right), 0 \right).$$

If  $g$  is a quadratic polynomial on  $K$ , then

$$(7.3) \quad g(p_6) - g(p_5) + g(p_4) - g(p_3) + g(p_2) - g(p_1) = 0$$

(cf. [5]).

Let  $\beta, \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6 \in \mathbf{R}$  such that  $\sum_{i=1}^6 (-1)^i \alpha_i = 0$ . Then there is exactly one quadratic polynomial  $g$  such that  $g(p_i) = \alpha_i$  and  $g(\text{centroid}) = \beta$ .

Let  $\mathcal{T}^k$  be the family of triangulations described in Section 2. Define

$$(7.4) \quad V_k := \{v : v|_T \text{ is a quadratic polynomial for each } T \in \mathcal{T}^k, v \text{ is continuous at the Gauss-Legendre points of interelement boundaries and } v \text{ vanishes at the Gauss-Legendre points on } \partial\Omega\}$$

and

$$(7.5) \quad W_k := \{w : w|_T \text{ is a quadratic polynomial for each } T \in \mathcal{T}^k, w \text{ is continuous on } \Omega \text{ and } w \equiv 0 \text{ on } \partial\Omega\}.$$

Again, note that  $W_k = V_k \cap V = V_k \cap V_{k+1}$ .

For each  $k$ , define

$$(7.6) \quad a_k(v_1, v_2) = \sum_{T \in \mathcal{T}^k} \int_T \nabla v_1 \cdot \nabla v_2 \quad \text{for } v_1, v_2 \in V + V_k$$

and

$$(7.7) \quad \|v\|_k = a_k(v, v)^{1/2}.$$




FIGURE 2

Let  $u_k \in V_k$  satisfy

$$a_k(u_k, v) = \int f v \quad \forall v \in V_k,$$

and  $\tilde{u}_k \in W_k$  satisfy

$$a_k(\tilde{u}_k, v) = \int f v \quad \forall v \in W_k.$$

Then we have the estimates

$$(7.8) \quad \|u - u_k\|_{L^2} + h_k \|u - u_k\|_k \leq Ch_k^3 \|u\|_{W_2^3}$$

and

$$(7.9) \quad \|u - \tilde{u}_k\|_{L^2} + h_k \|u - \tilde{u}_k\|_k \leq Ch_k^3 \|u\|_{W_2^3}.$$

Inequality (7.9) is well known. Inequality (7.8) holds because the nonconforming element satisfies the patch test (cf. [4], [5]). The theory for P1 nonconforming finite elements can be extended to this case in a straightforward manner if we can find an operator  $I_{k-1}^k: V_{k-1} \rightarrow V_k$  such that (i) it is a bounded linear operator with respect to both the  $L^2$  norm and the energy norm and (ii) it reduces to the natural injection on  $W_{k-1}$ .

For  $v \in V_{k-1}$ , the operator  $I_{k-1}^k: V_{k-1} \rightarrow V_k$  is defined as follows.

Let  $p$  be a Gauss-Legendre point of a side of a triangle in  $\mathcal{T}^k$ . If  $p$  lies on the common edge of two adjacent triangles  $\Delta_1$  and  $\Delta_2$  in  $\mathcal{T}^{k-1}$ , then

$$(I_{k-1}^k v)(p) := \frac{1}{2} [v|_{\Delta_1}(p) + v|_{\Delta_2}(p)].$$

If  $p$  lies inside a triangle  $T$  in  $\mathcal{T}^{k-1}$ , then there are two cases to consider.  $I_{k-1}^k v$  assumes the same values as  $v$  at the points  $b_2, b_4$  and  $b_6$ ; the value of  $I_{k-1}^k v$  at the points  $b_1, b_3$  and  $b_5$  is determined by condition (7.3) applied to the three outer triangles  $T_1, T_2, T_3$  (cf. Figure 2). To complete the definition of the intergrid transfer operator we must verify that the values of  $I_{k-1}^k v$  at the six Gauss-Legendre points of  $T_4$  satisfy (7.3). Then we let  $I_{k-1}^k v$  take the same value as  $v$  at the centroids of  $T_1, T_2, T_3$  and  $T_4$ .

LEMMA 7. Let  $p_1, \dots, p_{12}$  be the Gauss-Legendre points along the boundary of  $T \in \mathcal{T}^k$  (cf. Figure 2). Then

$$(7.10) \quad \sum_{i=1}^{12} (-1)^i (I_{k-1}^k v)(p_i) = \sum_{i=1}^{12} (-1)^i v|_T(p_i).$$

*Proof.* Let  $e$  denote the edge that contains  $p_1, p_2, p_3$  and  $p_4$ . We have

$$(7.11) \quad \sum_{i=1}^4 (-1)^i (I_{k-1}^k v)(p_i) = \sum_{i=1}^4 (-1)^i v|_T(p_i) + \sum_{i=1}^4 \frac{(-1)^i}{2} [v|_{T'}(p_i) - v|_T(p_i)],$$

where  $T'$  is the neighboring triangle in  $\mathcal{T}^{k-1}$  that also contains  $e$ . On  $e$ ,  $q := v|_{T'} - v|_T$  is a quadratic polynomial that vanishes at the two Gauss-Legendre points of the original triangle  $T$ , which are symmetric with respect to the midpoint. Therefore  $q$  is symmetric with respect to the midpoint of  $e$ . Thus  $q(p_4) = q(p_1)$  and  $q(p_3) = q(p_2)$ . Hence the second sum on the right-hand side of (7.11) adds up to zero.

Since a similar equality holds for each edge, Lemma 7 is now proved.  $\square$

Because  $v$  satisfies (7.3) on  $T_1, T_2$  and  $T_3$ ,

$$(7.12) \quad \sum_{i=1}^{12} (-1)^i v(p_i) + \sum_{j=1}^6 (-1)^j v(b_j) = 0.$$

But  $v$  also satisfies (7.3) on  $T_4$ , hence

$$(7.13) \quad \sum_{j=1}^6 (-1)^j v(b_j) = 0.$$

Equalities (7.12), (7.13) and Lemma 7 then imply that

$$\sum_{i=1}^{12} (-1)^i (I_{k-1}^k v)(p_i) = 0.$$

But by construction,

$$\sum_{i=1}^{12} (-1)^i (I_{k-1}^k v)(p_i) + \sum_{j=1}^6 (-1)^j (I_{k-1}^k v)(b_j) = 0.$$

Therefore,

$$\sum_{j=1}^6 (-1)^j (I_{k-1}^k v)(b_j) = 0,$$

and the compatibility condition on  $T_4$  is satisfied. Hence  $I_{k-1}^k v$  is well defined.

By our definition of  $I_{k-1}^k v$ , it is obvious that  $I_{k-1}^k w = w$  for  $w \in W_{k-1}$ .

The proof of Lemma 1 is still applicable for this intergrid transfer operator. Therefore,  $I_{k-1}^k$  is bounded with respect to both the  $L^2$  and energy norms.

As indicated earlier, once  $I_{k-1}^k$  has these properties we can obtain a multigrid method for finding approximate solutions  $\hat{u}_k$  of  $u_k$  such that

$$\|u - \hat{u}_k\|_{L^2} + h_k \|u - \hat{u}_k\|_k \leq Ch_k^3 \|u\|_{W_2^3}.$$

**Acknowledgments.** Special thanks go to the author's advisor, Professor Ridgway Scott, for his advice and encouragement. I would also like to thank the referee for helpful suggestions and for the simple proof of Lemma 1.

Department of Mathematics  
University of Michigan  
Ann Arbor, Michigan 48109

1. R. E. BANK & T. DUPONT "An optimal order process for solving finite element equations," *Math. Comp.*, v. 36, 1981, pp. 35-51.
2. D. BRAESS & W. HACKBUSCH, "A new convergence proof for the multigrid method including the V-cycle," *SIAM J. Numer. Anal.*, v. 20, 1983, pp. 967-975.
3. P. CIARLET, *The Finite Element Method for Elliptic Problems*, North-Holland, Amsterdam, 1978.
4. M. CROUZEIX & P.-A. RAVIART, "Conforming and nonconforming finite element methods for solving the stationary Stokes equations I," *RAIRO Anal. Numér. Sér. Rouge*, v. 7, no. R-3, 1973, pp. 33-75.
5. M. FORTIN & M. SOULIE, "A non-conforming piecewise quadratic element on triangles," *Internat. J. Numer. Methods Engrg.*, v. 19, 1983, pp. 505-520.
6. P. GRISVARD, "Behavior of solutions of an elliptic boundary value problem in polygonal or polyhedral domains," *Numerical Solution of Partial Differential Equations - III* (Synspade 1975) (B. Hubbard, ed.), Academic Press, New York, 1976, pp. 207-274.
7. W. HACKBUSCH, *Multi-Grid Methods and Applications*, Springer-Verlag, Berlin and New York, 1985.
8. G. STRANG & G. FIX, *An Analysis of the Finite Element Method*, Prentice-Hall, Englewood Cliffs, N.J., 1973.
9. K. STÜBEN & U. TROTTEMBERG, "Multigrid methods: fundamental algorithms, model problem analysis and applications," *Multigrid Methods* (Cologne, 1981), Lecture Notes in Math., vol. 960 (W. Hackbusch & U. Trottenberg, eds.), Springer-Verlag, Berlin and New York, 1982, pp. 1-176.
10. F. THOMASSET, *Implementation of Finite Element Methods for Navier-Stokes Equations*, Springer-Verlag, New York, 1981.