

## FINITE ELEMENT APPROXIMATION OF THE $p$ -LAPLACIAN

JOHN W. BARRETT AND W. B. LIU

**ABSTRACT.** In this paper we consider the continuous piecewise linear finite element approximation of the following problem: Given  $p \in (1, \infty)$ ,  $f$ , and  $g$ , find  $u$  such that

$$-\nabla \cdot (|\nabla u|^{p-2} \nabla u) = f \quad \text{in } \Omega \subset \mathbb{R}^2, \quad u = g \quad \text{on } \partial\Omega.$$

The finite element approximation is defined over  $\Omega^h$ , a union of regular triangles, yielding a polygonal approximation to  $\Omega$ . For sufficiently regular solutions  $u$ , achievable for a subclass of data  $f$ ,  $g$ , and  $\Omega$ , we prove optimal error bounds for this approximation in the norm  $W^{1,q}(\Omega^h)$ ,  $q = p$  for  $p < 2$  and  $q \in [1, 2]$  for  $p > 2$ , under the additional assumption that  $\Omega^h \subseteq \Omega$ . Numerical results demonstrating these bounds are also presented.

### 1. INTRODUCTION

Let  $\Omega$  be a bounded open set in  $\mathbb{R}^2$  with a Lipschitz boundary  $\partial\Omega$ . Given  $p \in (1, \infty)$ ,  $f \in L^2(\Omega)$ , and  $g \in W^{1-1/p,p}(\partial\Omega)$ , we consider the following problem:

( $\mathcal{P}$ ) Find  $u \in W_g^{1,p}(\Omega) \equiv \{v \in W^{1,p}(\Omega) : v = g \text{ on } \partial\Omega\}$  such that

$$\int_{\Omega} |\nabla u|^{p-2} (\nabla u, \nabla v)_{\mathbb{R}^2} d\Omega = \int_{\Omega} f v d\Omega \quad \forall v \in W_0^{1,p}(\Omega),$$

where  $|v|^2 = (v, v)_{\mathbb{R}^2}$ . Throughout we adopt the standard notation  $W^{m,q}(D)$  for Sobolev spaces on  $D$  with norm  $\|\cdot\|_{W^{m,q}(D)}$  and seminorm  $|\cdot|_{W^{m,q}(D)}$ . We note that the seminorm  $|\cdot|_{W^{1,q}(D)}$  and the norm  $\|\cdot\|_{W^{1,q}(D)}$  are equivalent on  $W_0^{1,q}(D)$ .

Problem ( $\mathcal{P}$ ) above is the weak formulation of the Dirichlet problem for the  $p$ -Laplacian

$$(1.1) \quad -\nabla \cdot (|\nabla u|^{p-2} \nabla u) = f \quad \text{in } \Omega, \quad u = g \quad \text{on } \partial\Omega.$$

The well-posedness of ( $\mathcal{P}$ ) is well established, and one can refer to, for example, Glowinski and Marrocco [5] or the account in Ciarlet [4]. Of course, one can study more general boundary conditions and the presence of lower-order terms in the differential operator. However, for ease of exposition, we just consider ( $\mathcal{P}$ ), although most of our results can be adapted to more general

---

Received by the editor May 28, 1991 and, in revised form, March 25, 1992 and September 10, 1992.

1991 *Mathematics Subject Classification.* Primary 65N30.  
 Research supported by SERC research grant GR/F81255.

problems. From Glowinski and Marrocco [5], or Ciarlet [4],  $(\mathcal{P})$  is equivalent to the following minimization problem:

( $\mathcal{Q}$ ) Find  $u \in W_g^{1,p}(\Omega)$  such that

$$(1.2a) \quad J_\Omega(u) \leq J_\Omega(v) \quad \forall v \in W_g^{1,p}(\Omega),$$

where

$$(1.2b) \quad J_\Omega(v) \equiv \frac{1}{p} \int_\Omega |\nabla v|^p d\Omega - \int_\Omega f v d\Omega.$$

It is easily established that  $J_\Omega(\cdot)$  is strictly convex and continuous on  $W_g^{1,p}(\Omega)$ . Further,  $J_\Omega(\cdot)$  is Gateaux differentiable with

$$(1.3) \quad J'_\Omega(u)(v) \equiv \int_\Omega |\nabla u|^{p-2} (\nabla u, \nabla v)_{\mathbb{R}^2} d\Omega - \int_\Omega f v d\Omega \quad \forall v \in W_0^{1,p}(\Omega).$$

Hence, there exists a unique solution to ( $\mathcal{Q}$ ), and ( $\mathcal{Q}$ ) is equivalent to  $(\mathcal{P})$ , its Euler equation. In addition, we have that

$$(1.4) \quad \|u\|_{W^{1,p}(\Omega)} \leq C[\|f\|_{L^2(\Omega)}^{1/(p-1)} + \|g\|_{W^{1-1/p,p}(\partial\Omega)}].$$

The problem  $(\mathcal{P})$  occurs in many mathematical models of physical processes: nonlinear diffusion and filtration, see Philip [8]; power-law materials, see Atkinson and Champion [1]; and quasi-Newtonian flows, see Atkinson and Jones [2], for example.

It is the purpose of this paper to analyze the finite element approximation of  $(\mathcal{P})$ . Let  $\Omega^h$  be a polygonal approximation to  $\Omega$  defined by  $\overline{\Omega^h} \equiv \bigcup_{\tau \in T^h} \overline{\tau}$ , where  $T^h$  is a partitioning of  $\Omega^h$  into a finite number of disjoint open regular triangles  $\tau$ , each of maximum diameter bounded above by  $h$ . In addition, for any two distinct triangles, their closures are either disjoint, or have a common vertex, or a common side. Let  $\{P_j\}_{j=1}^J$  be the vertices associated with the triangulation  $T^h$ , where  $P_j$  has coordinates  $(x_j, y_j)$ . Throughout we assume that  $P_j \in \partial\Omega^h$  implies  $P_j \in \partial\Omega$ , and that  $\Omega^h \subseteq \Omega$ . We note that, owing to the elliptic degeneracy of the  $p$ -Laplacian and the limited regularity of the solution  $u$ , see below, it is not a simple matter to extend the results in this paper to the case  $\Omega^h \not\subseteq \Omega$ . Associated with  $T^h$  is the finite-dimensional space

$$(1.5) \quad S^h \equiv \{\chi \in C(\overline{\Omega^h}) : \chi|_\tau \text{ is linear } \forall \tau \in T^h\} \subset W^{1,p}(\Omega^h).$$

Let  $\pi_h : C(\overline{\Omega^h}) \rightarrow S^h$  denote the interpolation operator such that for any  $v \in C(\overline{\Omega^h})$ , the interpolant  $\pi_h v \in S^h$  satisfies  $\pi_h v(P_j) = v(P_j)$ ,  $j = 1, \dots, J$ . We recall the following standard approximation results. For  $m = 0$  or  $1$ , and for all  $\tau \in T^h$ , we have (a) for  $q \in [1, \infty]$ ,  $s \in [1, \infty]$ , provided  $W^{2,s}(\tau) \hookrightarrow W^{m,q}(\tau)$ ,

$$(1.6a) \quad |v - \pi_h v|_{W^{m,q}(\tau)} \leq Ch^{2(1/q-1/s)} h^{2-m} |v|_{W^{2,s}(\tau)} \quad \forall v \in W^{2,s}(\tau);$$

and (b) for  $q > 2$ ,

$$(1.6b) \quad |v - \pi_h v|_{W^{m,q}(\tau)} \leq Ch^{1-m} |v|_{W^{1,q}(\tau)} \quad \forall v \in W^{1,q}(\tau).$$

In (1.6a) we have noted the imbedding  $W^{2,1}(\tau) \hookrightarrow C(\overline{\tau})$ ; see, for example, p. 300 in Kufner et al. [6].

The finite element approximation of  $(\mathcal{P})$  that we wish to consider is:

( $\mathcal{P}^h$ ) Find  $u^h \in S_g^h$  such that

$$(1.7a) \quad \int_{\Omega^h} |\nabla u^h|^{p-2} (\nabla u^h, \nabla v^h)_{\mathbb{R}^2} d\Omega^h = \int_{\Omega^h} f v^h d\Omega^h \quad \forall v^h \in S_0^h,$$

where

$$(1.7b) \quad S_0^h \equiv \{\chi \in S^h : \chi = 0 \text{ on } \partial\Omega^h\}$$

and

$$(1.7c) \quad S_g^h \equiv \{\chi \in S^h : \chi = g^h \text{ on } \partial\Omega^h\},$$

where  $g^h \in S^h$  is chosen to approximate the Dirichlet boundary data. If  $p > 2$ , then  $u \in W^{1,p}(\Omega)$  implies  $u \in C(\bar{\Omega})$ , and so we set  $g^h \equiv \pi_h u$ . For the explicit error bounds derived in §3 for  $p < 2$ , we assume that  $u \in C(\bar{\Omega})$ , and so once again set  $g^h \equiv \pi_h u$ . However, for the abstract analysis of this and the next section,  $g^h$  can be arbitrary. The corresponding minimization problem is:

( $\mathcal{E}^h$ ) Find  $u^h \in S_g^h$  such that

$$(1.8) \quad J_{\Omega^h}(u^h) \leq J_{\Omega^h}(v^h) \quad \forall v^h \in S_g^h.$$

The well-posedness of ( $\mathcal{P}^h$ )  $\equiv$  ( $\mathcal{E}^h$ ) follows in an analogous way to that of ( $\mathcal{P}$ ) and ( $\mathcal{E}$ ), see Glowinski and Marrocco [5] or Ciarlet [4], and

$$(1.9) \quad \|u^h\|_{W^{1,p}(\Omega^h)} \leq C[\|f\|_{L^2(\Omega^h)}^{1/(p-1)} + \|g^h\|_{W^{1,p}(\Omega^h)}].$$

We note that for  $p = 2$ , problem ( $\mathcal{P}$ ) reduces to the weak formulation of the linear Laplacian, and hence the regularity of  $u$  and the finite element error analysis are well established in this case. For  $p \neq 2$ , the regularity of  $u$  is less well established, as (1.1) is then a degenerate quasi-linear elliptic problem. It is well known, see Example 3.1 in §3, that  $u$  has limited regularity for infinitely smooth data  $f$ ,  $g$ , and  $\Omega$ . Therefore, there is no benefit in considering higher-order finite element approximations, and hence our restriction to continuous piecewise linears from the outset. Lieberman [7] has proved that if  $\partial\Omega \in C^{1,\beta}$ , then  $g$  is the trace of a function  $\in C^{1,\gamma}(\bar{\Omega})$  for  $\beta, \gamma \in (0, 1)$ , and if  $f \in L^\infty(\Omega)$ , then  $u \in C^{1,\alpha}(\bar{\Omega})$  for some  $\alpha \in (0, 1)$ . However, for explicit finite element error bounds one requires global regularity results on the second, or maybe higher, derivatives of  $u$ . Unfortunately, such results are not available at present in the literature, but it is an active area of research worldwide.

The following error bounds were proved in Glowinski and Marrocco [5] for the case  $\Omega^h \equiv \Omega$  and  $g \equiv 0$ :

If  $u \in W_0^{1,p}(\Omega) \cap W^{2,p}(\Omega)$ , then

$$(1.10) \quad \|u - u^h\|_{W^{1,p}(\Omega)} \leq \begin{cases} Ch^{1/(3-p)} & \text{if } p \leq 2, \\ Ch^{1/(p-1)} & \text{if } p \geq 2, \end{cases}$$

where throughout this paper  $C$  denotes a generic positive constant independent of  $h$ . Chow [3], employing an approach of Tyukhtin [9], improved these error bounds. He proved that

$$(1.11a) \quad \|u - u^h\|_{W^{1,p}(\Omega)} \leq C \|u - v^h\|_{W^{1,p}(\Omega)}^{p/2} \quad \forall v^h \in S_0^h \quad \text{if } p \leq 2,$$

and

$$(1.11b) \quad \|u - u^h\|_{W^{1,p}(\Omega)} \leq C (\|\pi_h u\|_{W^{1,p}(\Omega)}) \|u - \pi_h u\|_{W^{1,p}(\Omega)}^{2/p} \quad \text{if } p > 2,$$

and hence, if  $u \in W_0^{1,p}(\Omega) \cap W^{2,p}(\Omega)$ , it follows from (1.6a) that

$$(1.12) \quad \|u - u^h\|_{W^{1,p}(\Omega)} \leq \begin{cases} Ch^{p/2} & \text{if } p \leq 2, \\ Ch^{2/p} & \text{if } p \geq 2. \end{cases}$$

It is the purpose of this paper to prove optimal error bounds. The layout is as follows. In the next section we prove an abstract error bound for the approximation  $(\mathcal{P}^h)$  of  $(\mathcal{P})$ . In §3 we study the case  $p \in (1, 2)$  and prove an optimal  $W^{1,p}$  error bound, that is,  $O(h)$ , provided that  $u \in W^{3,1}(\Omega) \cap C^{2,(2-p)/p}(\overline{\Omega})$ . Thus, this optimal error bound requires a stronger regularity assumption on  $u$  than that for the bound (1.12) in the case  $p < 2$ . In §4 we study the case  $p > 2$  and first show that the bound (1.12) for  $p > 2$  can be achieved under the weaker regularity requirement  $u \in W^{1,\infty}(\Omega) \cap W^{2,2}(\Omega)$ . Second, under the additional assumption  $|f| \geq \rho > 0$  a.e. in  $\Omega$ , we prove an optimal  $W^{1,4/3}$  error bound. We note that the above regularity requirements on  $u$  for these optimal error bounds are achievable for a subclass of data  $f$ ,  $g$ , and  $\Omega$ . In §5 we show that the error bounds derived in the previous sections hold for the fully practical scheme of employing numerical integration on the right-hand side of (1.7) if  $f$  is sufficiently smooth. Finally, we report on some numerical examples, which confirm these optimal error bounds.

## 2. AN ABSTRACT ERROR BOUND

We first prove a lemma, which is a generalization of Lemmas 5.1, 5.2, 5.3 and 5.4 in Glowinski and Marrocco [5].

**Lemma 2.1.** *For all  $p > 1$  and  $\delta \geq 0$  there exist positive constants  $C_1$  and  $C_2$  such that for all  $\xi, \eta \in \mathbb{R}^2$ ,  $\xi \neq \eta$ ,*

$$(2.1a) \quad \left| |\xi|^{p-2}\xi - |\eta|^{p-2}\eta \right| \leq C_1 |\xi - \eta|^{1-\delta} (|\xi| + |\eta|)^{p-2+\delta}$$

and

$$(2.1b) \quad (|\xi|^{p-2}\xi - |\eta|^{p-2}\eta, \xi - \eta)_{\mathbb{R}^2} \geq C_2 |\xi - \eta|^{2+\delta} (|\xi| + |\eta|)^{p-2-\delta}.$$

*Proof.* The approach is similar to that in Glowinski and Marrocco [5]. For all  $\xi, \eta \in \mathbb{R}^2$ ,  $\xi \neq \eta$ , let

$$(2.2) \quad G_1(\xi, \eta) \equiv \left| |\xi|^{p-2}\xi - |\eta|^{p-2}\eta \right| / [|\xi - \eta|^{1-\delta} (|\xi| + |\eta|)^{p-2+\delta}].$$

We wish to prove that  $G_1$  is bounded above. For any  $\varepsilon > 0$ ,  $G_1$  is continuous on

$$D_\varepsilon \equiv \{(\xi, \eta) : |\xi - \eta| \geq \varepsilon \text{ and } (|\xi| + |\eta|) \leq 1/\varepsilon\}.$$

In addition, we note that for all  $\xi, \eta \in \mathbb{R}^2$ ,  $\xi \neq \eta$ ,

$$(2.3) \quad \begin{aligned} G_1(\xi, \eta) &\equiv G_1(\eta, \xi), & G_1(\lambda\xi, \lambda\eta) &\equiv G_1(\xi, \eta) \quad \text{for all } \lambda \in \mathbb{R}^+, \\ G_1(0, \eta) &= 1 \quad \text{and} \quad G_1(A\xi, A\eta) &\equiv G_1(\xi, \eta) \quad \text{if } A^T A = I, \end{aligned}$$

i.e.,  $A$  is a rotation matrix. Therefore, without loss of generality we can take  $\xi = e_1 \equiv (1, 0)$ . Since

$$(2.4) \quad G_1(e_1, \eta) \rightarrow 1 \quad \text{as } |\eta| \rightarrow \infty,$$

it remains to show that  $\limsup G_1(e_1, \eta) < \infty$  as  $|e_1 - \eta| \rightarrow 0$ .

Let  $\eta = (1 + \rho \cos \theta, \rho \sin \theta)$ . Then a simple calculation yields that

$$(2.5) \quad \lim_{\rho \rightarrow 0} G_1(e_1, \eta) = \begin{cases} 0 & \text{if } \delta > 0, \\ 2^{2-p} \{1 + p(p-2) \cos^2 \theta\}^{1/2} & \text{if } \delta = 0. \end{cases}$$

Hence the desired result (2.1a).

Similarly, we prove (2.1b). Let

$$(2.6) \quad G_2(\xi, \eta) \equiv |\xi - \eta|^{2+\delta} (|\xi| + |\eta|)^{p-2-\delta} / (|\xi|^{p-2}\xi - |\eta|^{p-2}\eta, \xi - \eta)_{\mathbb{R}^2}.$$

From Glowinski and Marrocco [5] we have that

$$(2.7) \quad (|\xi|^{p-2}\xi - |\eta|^{p-2}\eta, \xi - \eta)_{\mathbb{R}^2} > 0 \quad \text{if } \xi \neq \eta.$$

Therefore, we only need to prove that  $G_2$  is bounded above. In addition, the results (2.3) and (2.4) hold for  $G_2$ .

Setting  $\eta = (1 + \rho \cos \theta, \rho \sin \theta)$ , a simple calculation yields that

$$(2.8) \quad \lim_{\rho \rightarrow 0} G_2(e_1, \eta) = \begin{cases} 0 & \text{if } \delta > 0, \\ 2^{p-2} \{1 + (p-2) \cos^2 \theta\}^{-1} & \text{if } \delta = 0. \end{cases}$$

Hence the desired result (2.1b).  $\square$

The inequality (2.1a) was proved in Glowinski and Marrocco [5] for  $p \in (1, 2]$  with  $\delta = 2 - p$ , and for  $p \geq 2$  with  $\delta = 0$ ; similarly, (2.1b) was proved for  $p \in (1, 2]$  with  $\delta = 0$ , and for  $p \geq 2$  with  $\delta = p - 2$ .

For  $p \in (1, \infty)$  and  $\sigma \geq 0$  we define for any  $v \in W^{1,p}(\Omega^h)$

$$(2.9) \quad |v|_{(p,\sigma)} \equiv \int_{\Omega^h} (|\nabla u| + |\nabla v|)^{p-\sigma} |\nabla v|^\sigma d\Omega^h,$$

where  $u$  is the solution of  $(\mathcal{P})$ . We prove the following results for later use.

**Lemma 2.2.** For  $p \in (1, \sigma]$  we have

$$(2.10a) \quad |v|_{(p,\sigma)}^{\sigma/p} \leq |v|_{W^{1,p}(\Omega^h)}^\sigma \leq C[|u|_{W^{1,p}(\Omega^h)} + |v|_{W^{1,p}(\Omega^h)}]^{p-\sigma} |v|_{(p,\sigma)},$$

and for  $p \in [\sigma, \infty)$ ,

$$(2.10b) \quad |v|_{W^{1,p}(\Omega^h)}^p \leq |v|_{(p,\sigma)} \leq C[|u|_{W^{1,p}(\Omega^h)} + |v|_{W^{1,p}(\Omega^h)}]^{p-\sigma} |v|_{W^{1,p}(\Omega^h)}^\sigma.$$

Hence, (2.9) is well defined for  $v \in W^{1,p}(\Omega^h)$ .

*Proof.* Setting  $w \equiv (|\nabla u| + |\nabla v|)^{p-\sigma}$ , we first consider the case  $p \in (1, \sigma]$ . The left inequality in (2.10a) follows immediately from noting that  $w \leq |\nabla v|^{p-\sigma}$ . Applying Hölder's inequality, we have

$$\begin{aligned} |v|_{W^{1,p}(\Omega^h)}^\sigma &\equiv \left\{ \int_{\Omega^h} w^{-p/\sigma} [w^{1/\sigma} |\nabla v|]^p d\Omega^h \right\}^{\sigma/p} \\ &\leq \left\{ \int_{\Omega^h} w^{-p/(\sigma-p)} d\Omega^h \right\}^{(\sigma-p)/p} |v|_{(p,\sigma)} \\ &= \left\{ \int_{\Omega^h} (|\nabla u| + |\nabla v|)^p d\Omega^h \right\}^{(\sigma-p)/p} |v|_{(p,\sigma)}. \end{aligned}$$

The right inequality in (2.10a) follows by noting that for all  $\gamma \in [0, \infty)$  there exists  $C_\gamma > 0$  such that  $|a + b|^\gamma \leq C_\gamma(|a|^\gamma + |b|^\gamma)$  for all  $a, b \in \mathbb{R}$ .

The inequalities (2.10b) can be proved in a similar manner.  $\square$

The next theorem is the natural generalization of the result in §7 of Chow [3]. We use the minimization property of  $u^h$  and Lemma 2.1, whereas Chow uses the Glowinski and Marrocco version of Lemma 2.1.

**Theorem 2.1.** *Let  $u$  and  $u^h$  be the unique solutions of  $(\mathcal{P}) \equiv (\mathcal{Q})$  and  $(\mathcal{P}^h) \equiv (\mathcal{Q}^h)$ , respectively. Then for any  $\delta_1 \in [0, 2)$  and  $\delta_2 \geq 0$ , and any  $v^h \in S_g^h$ , it follows that*

$$(2.11) \quad |u - u^h|_{(p, 2+\delta_2)} \leq C |u - v^h|_{(p, 2-\delta_1)}.$$

*Proof.* We have for any  $v^h \in S_g^h$  that

$$(2.12a) \quad \begin{aligned} J_{\Omega^h}(v^h) - J_{\Omega^h}(u) &= \int_0^1 J'_{\Omega^h}(u + s(v^h - u))(v^h - u) ds \\ &= \int_0^1 [J'_{\Omega^h}(u + s(v^h - u))([u + s(v^h - u)] - u) \\ &\quad - J'_{\Omega^h}(u)([u + s(v^h - u)] - u)] \frac{ds}{s} \\ &\quad + J'_{\Omega^h}(u)(v^h - u) \\ &\equiv A(v^h) + J'_{\Omega^h}(u)(v^h - u), \end{aligned}$$

where from (1.3)

$$(2.12b) \quad \begin{aligned} A(v^h) &\equiv \int_0^1 \left[ \int_{\Omega^h} \{[|\nabla(u + s(v^h - u))|]^{p-2} \right. \\ &\quad \left. \times \nabla(u + s(v^h - u)) - |\nabla u|^{p-2} \nabla u\} \nabla(v^h - u) \right] d\Omega^h ds. \end{aligned}$$

From (2.12b) and (2.1a) we have that

$$(2.13) \quad \begin{aligned} |A(v^h)| &\leq C_1 \int_0^1 s^{1-\delta_1} \int_{\Omega^h} (|\nabla[u + s(v^h - u)]| + |\nabla u|)^{p-2+\delta_1} \\ &\quad \times |\nabla(v^h - u)|^{2-\delta_1} d\Omega^h ds \\ &\leq C |u - v^h|_{(p, 2-\delta_1)}, \end{aligned}$$

where we have noted that for all  $v_1, v_2$ , and  $s \in [0, 1]$

$$(2.14) \quad \frac{1}{2}s(|\nabla v_1| + |\nabla v_2|) \leq |\nabla[v_1 + sv_2]| + |\nabla v_1| \leq 2(|\nabla v_1| + |\nabla v_2|).$$

From (2.12b), (2.1b), and (2.14) we have that

$$(2.15) \quad \begin{aligned} |A(v^h)| &\geq C_2 \int_0^1 s^{1+\delta_2} \int_{\Omega^h} (|\nabla[u + s(v^h - u)]| + |\nabla u|)^{p-2+\delta_2} \\ &\quad \times |\nabla(v^h - u)|^{2+\delta_2} d\Omega^h ds \\ &\geq C |u - v^h|_{(p, 2+\delta_2)}. \end{aligned}$$

From (1.8) and (2.12) we have that for all  $v^h \in S_g^h$

$$(2.16) \quad \begin{aligned} A(u^h) + J'_{\Omega^h}(u)(u^h - u) &\equiv J_{\Omega^h}(u^h) - J_{\Omega^h}(u) \leq J_{\Omega^h}(v^h) - J_{\Omega^h}(u) \\ &\equiv A(v^h) + J'_{\Omega^h}(u)(v^h - u). \end{aligned}$$

Therefore, it follows from (2.16), (2.13), and (2.15) that

$$(2.17) \quad |u - u^h|_{(p, 2+\delta_2)} \leq C |u - v^h|_{(p, 2-\delta_1)} + J'_{\Omega^h}(u)(v^h - u^h).$$

As  $\Omega^h$  is Lipschitz,  $\Omega^h \subseteq \Omega$  and  $\chi \equiv v^h - u^h \in S_0^h$ , we can extend  $\chi$  to be zero on  $\Omega \setminus \Omega^h$ . Denoting this extension by  $\hat{\chi}$ , we have that  $\hat{\chi} \in W_0^{1,p}(\Omega)$  and hence from (P) that  $J'_{\Omega^h}(u)(\chi) \equiv J'_\Omega(u)(\hat{\chi}) = 0$ . Therefore, the desired result (2.11) follows from (2.17).  $\square$

### 3. ERROR BOUNDS FOR $p \in (1, 2)$

Assuming that  $u \in W^{2,1}(\Omega)$ , which implies that  $u \in C(\bar{\Omega})$ , we can set  $g^h \equiv \pi_h u$  in (1.7c). Choosing  $\delta_2 = 0$  in (2.11) and noting (2.10a), (1.4), (1.9), and (1.6a), we have for all  $\delta_1 \in [0, 2)$  and for all  $v^h \in S_g^h$

$$(3.1) \quad \|u - u^h\|_{W^{1,p}(\Omega^h)}^2 \leq C \|u - u^h\|_{(p,2)} \leq C \|u - v^h\|_{(p,2-\delta_1)}.$$

Choosing  $\delta_1 = 2 - p$  and noting (2.10a) yield that for all  $v^h \in S_g^h$

$$(3.2) \quad \|u - u^h\|_{W^{1,p}(\Omega^h)}^2 \leq C \|u - v^h\|_{(p,p)} \leq C \|u - v^h\|_{W^{1,p}(\Omega^h)}^p.$$

From a Poincaré inequality we have for all  $q \in [1, \infty)$ , for all  $v \in W^{1,q}(\Omega)$ , and for all  $v^h, w^h \in S_g^h$  that

$$(3.3) \quad \|v - w^h\|_{W^{1,q}(\Omega^h)} \leq C \|v - v^h\|_{W^{1,q}(\Omega^h)} + C \|v - w^h\|_{W^{1,q}(\Omega^h)}.$$

Hence, from (3.2) with  $v^h \equiv \pi_h u$ , (3.3), and (1.6a) we have that

$$(3.4a) \quad \|u - u^h\|_{W^{1,p}(\Omega^h)} \leq C \|u - \pi_h u\|_{W^{1,p}(\Omega^h)} + C \|u - \pi_h u\|_{W^{1,p}(\Omega^h)}^{p/2}$$

$$(3.4b) \quad \leq Ch^{p/2} \quad \text{if } u \in W^{2,p}(\Omega),$$

the generalization of the results (1.11a) and (1.12), for  $p < 2$ , of Chow [3] to the case of nonhomogeneous boundary data  $g$  and  $\Omega^h \subseteq \Omega$ . Below we prove an optimal  $W^{1,p}$  error bound for sufficiently regular  $u$ , based on choosing  $\delta_1 = 0$  in (3.1).

**Lemma 3.1.** *Let  $\alpha \in (-1, 0)$ . If  $v \in W^{2,1}(\Omega)$ , then*

$$(3.5) \quad \int_{\Omega} |v|^\alpha |\nabla v|^2 d\Omega < \infty.$$

*Proof.* We have that

$$\begin{aligned} \int_{\Omega} |v|^\alpha |v_x|^2 dx dy &\equiv \frac{1}{\alpha + 1} \int_{\Omega} (\text{sign}(v) |v|^{\alpha+1})_x v_x dx dy \\ &\equiv \frac{1}{\alpha + 1} \left\{ \int_{\partial\Omega} \text{sign}(v) |v|^{\alpha+1} v_x dy - \int_{\Omega} \text{sign}(v) |v|^{\alpha+1} v_{xx} dx dy \right\} \end{aligned}$$

and a similar identity with  $v_x$  replaced by  $v_y$ . The desired result (3.5) then follows from the imbedding  $W^{2,1}(\Omega) \hookrightarrow C(\bar{\Omega})$  and the trace inequality  $\|\cdot\|_{L^1(\partial\Omega)} \leq C \|\cdot\|_{W^{1,1}(\Omega)}$ .  $\square$

**Theorem 3.1.** *If  $u \in W^{3,1}(\Omega) \cap C^{2,\alpha}(\bar{\Omega})$ , with  $\alpha > 0$ , then it follows that*

$$(3.6a) \quad \|u - u^h\|_{W^{1,p}(\Omega^h)}^2 \leq C[h^2 + h^{p(1+\alpha)}],$$

and hence, if  $u \in W^{3,1}(\Omega) \cap C^{2,(2-p)/p}(\bar{\Omega})$ , then

$$(3.6b) \quad \|u - u^h\|_{W^{1,p}(\Omega^h)} \leq Ch.$$

*Proof.* As  $u \in C^{2,\alpha}(\overline{\Omega})$ , we have from (1.6a) that for all  $\tau \in T^h$  and for all  $(x, y) \in \bar{\tau}$

$$(3.7) \quad |\nabla(u - \pi_h u)(x, y)| \leq Ch|H[u]|_{L^\infty(\tau)} \leq ChH[u](x, y) + Ch^{1+\alpha},$$

where  $H[u] \equiv |u_{xx}| + |u_{xy}| + |u_{yy}|$ .

It is easy to check that the function  $q(t) \equiv (a + t)^{p-2}t^2$  with  $a \geq 0$  is increasing on  $\mathbb{R}^+$  and hence that  $q(|t_1 + t_2|) \leq 2[q(|t_1|) + q(|t_2|)]$  for all  $t_1, t_2 \in \mathbb{R}$ . Therefore, we have from (3.1) with  $\delta_1 = 0$  and  $v^h \equiv \pi_h u$ , (3.7), and the above that

$$(3.8) \quad \begin{aligned} |u - u^h|_{W^{1,p}(\Omega^h)}^2 &\leq C \int_{\Omega^h} (|\nabla u| + |\nabla(u - \pi_h u)|)^{p-2} |\nabla(u - \pi_h u)|^2 d\Omega^h \\ &\leq Ch^2 \int_{\Omega^h} (|\nabla u| + ChH[u])^{p-2} (H[u])^2 d\Omega^h \\ &\quad + Ch^{2(1+\alpha)} \int_{\Omega^h} (|\nabla u| + Ch^{1+\alpha})^{p-2} d\Omega^h \\ &\leq Ch^{p(1+\alpha)} + Ch^2 \int_{\Omega^h} (|\nabla u|)^{p-2} (H[u])^2 d\Omega^h. \end{aligned}$$

Setting  $v_1 \equiv u_x$  and  $v_2 \equiv u_y$ , we have from (3.5), as  $v_1, v_2 \in W^{2,1}(\Omega)$ , that

$$(3.9) \quad \begin{aligned} \int_{\Omega^h} (|\nabla u|)^{p-2} (H[u])^2 d\Omega^h &\leq C \int_{\Omega^h} (v_1^2 + v_2^2)^{(p-2)/2} (|\nabla v_1|^2 + |\nabla v_2|^2) d\Omega^h \\ &\leq C \int_{\Omega^h} [|v_1|^{p-2} |\nabla v_1|^2 + |v_2|^{p-2} |\nabla v_2|^2] d\Omega^h < \infty. \end{aligned}$$

Combining (3.8) and (3.9) yields the result (3.6a) and hence (3.6b) with  $\|\cdot\|_{W^{1,p}(\Omega^h)}$  replaced by  $|\cdot|_{W^{1,p}(\Omega^h)}$ . The results (3.6) then follow by noting (3.3), (1.6a), and that  $u \in W^{3,1}(\Omega)$  implies  $u \in W^{2,p}(\Omega)$ .  $\square$

We note that one can prove (3.6b) under alternative regularity requirements on  $u$ , e.g.,  $u \in W^{3,p}(\Omega)$ . However, we will not exploit this here. We now show that the regularity requirements on  $u$  in Theorem 3.1 hold for a model problem.

**Example 3.1.** We consider a radially symmetric version of problem  $(\mathcal{P})$ . Let  $\Omega \equiv \{r: r < 1\}$ ,  $f(x, y) \equiv F(r)$ ,  $f \in L^q(\Omega)$  for  $q > 2$ , and  $g$  be constant, where  $r \equiv (x^2 + y^2)^{1/2}$ . Then

$$(3.10a) \quad u(x, y) \equiv U(r) \equiv - \int_r^1 \text{sign}(Z(t)) |Z(t)|^{1/(p-1)} dt + g,$$

$$(3.10b) \quad U'' \equiv (|Z|^{(2-p)/(p-1)} Z') / (p-1)$$

and

$$(3.10c) \quad U''' \equiv C_1 \text{sign}(Z) |Z|^{(3-2p)/(p-1)} (Z')^2 + C_2 |Z|^{(2-p)/(p-1)} Z''$$

for some constants  $C_i$ , where

$$(3.10d) \quad z(x, y) \equiv Z(r) \equiv (|U'|^{p-2} U')(r) = -\frac{1}{r} \int_0^r t F(t) dt.$$



It is a simple matter to deduce from (3.10d) that

$$(3.11a) \quad f \in C^{0,q}(\overline{\Omega}) \Rightarrow z \in C^{1,q}(\overline{\Omega}) \quad \forall q \in [0, 1]$$

and

$$(3.11b) \quad f \in W^{1,q}(\Omega) \quad \text{for } q > 1 \Rightarrow z \in W^{2,1}(\Omega) \Rightarrow z \in C(\overline{\Omega}).$$

It follows from (3.10b) and (3.11a) that

$$(3.12a) \quad p \in (1, \frac{3}{2}] \text{ and } f \in C^{0,\beta}(\overline{\Omega}) \quad \text{for } \beta \in [0, 1] \Rightarrow u \in C^{2,\beta}(\overline{\Omega}),$$

$$(3.12b) \quad p \in [\frac{3}{2}, 2) \text{ and } f \in C^{0,\beta}(\overline{\Omega}) \text{ for } \beta \in [0, (2-p)/(p-1)] \\ \Rightarrow u \in C^{2,\beta}(\overline{\Omega}),$$

and from (3.10c), (3.11b), and Lemma 3.1 that

$$(3.12c) \quad p \in (1, 2) \text{ and } f \in W^{1,q}(\Omega) \text{ for } q > 1 \Rightarrow u \in W^{3,1}(\Omega).$$

#### 4. ERROR BOUNDS FOR $p > 2$

Let  $g^h \equiv \pi_h u$  in (1.7c). From (2.11) with  $\delta_1 = 0$  and  $\delta_2 = p - 2$ , (2.10b), (1.4), (1.9), and (1.6b) it follows that

$$(4.1) \quad |u - u^h|_{W^{1,p}(\Omega^h)}^p \leq |u - u^h|_{(p,p)} \leq C|u - \pi_h u|_{(p,2)} \leq C|u - \pi_h u|_{W^{1,p}(\Omega^h)}^2,$$

and hence it follows from (3.3) and (1.6a) that

$$(4.2a) \quad \|u - u^h\|_{W^{1,p}(\Omega^h)} \leq C\|u - \pi_h u\|_{W^{1,p}(\Omega^h)} + C|u - \pi_h u|_{W^{1,p}(\Omega^h)}^{2/p}$$

$$(4.2b) \quad \leq Ch^{2/p} \quad \text{if } u \in W^{2,p}(\Omega),$$

the generalization of the results (1.11b) and (1.12), for  $p > 2$ , of Chow [3] to the case of nonhomogeneous boundary data  $g$  and  $\Omega^h \subseteq \Omega$ . Alternatively, assuming  $u \in W^{1,\infty}(\Omega)$ , we have from (2.11) with  $\delta_1 = 2 - s$ ,  $s \in [1, 2]$ , and  $\delta_2 = p - 2$  and (1.6b) that

$$(4.3) \quad |u - u^h|_{W^{1,p}(\Omega^h)}^p \leq |u - u^h|_{(p,p)} \leq C|u - \pi_h u|_{(p,s)} \leq C|u - \pi_h u|_{W^{1,s}(\Omega^h)}^s.$$

In addition, we note from (1.6b) that for  $u \in W^{1,\infty}(\Omega)$

$$(4.4) \quad \|u - \pi_h u\|_{W^{1,q}(\Omega^h)} \leq C\|u - \pi_h u\|_{W^{1,s}(\Omega^h)}^{s/q} \quad \text{if } q > s.$$

Hence, it follows from (4.3), (3.3), (4.4), and (1.6a) that if  $u \in W^{1,\infty}(\Omega) \cap W^{2,s}(\Omega)$ ,  $s \in [1, 2]$ , then

$$(4.5a) \quad \|u - u^h\|_{W^{1,p}(\Omega^h)} \leq C\|u - \pi_h u\|_{W^{1,p}(\Omega^h)} + C|u - \pi_h u|_{W^{1,s}(\Omega^h)}^{s/p}$$

$$(4.5b) \quad \leq C\|u - \pi_h u\|_{W^{1,s}(\Omega^h)}^{s/p} \leq Ch^{s/p}.$$

Choosing  $f \equiv 1$  and  $g \equiv 0$  in Example 3.1 yields that  $u(x, y) \equiv C(1 - r^{p/(p-1)})$ , and so  $u \in W^{2,s}(\Omega)$  only if  $s < 2(p-1)/(p-2)$ . Therefore, in general  $u$  rarely belongs to  $W^{2,p}(\Omega)$  in order for (4.2b) to guarantee that the error converges at least at the rate of  $h^{2/p}$  in  $W^{1,p}$ . However, from (4.5b) we see that this rate is ensured under the far weaker regularity requirement of  $u \in W^{1,\infty}(\Omega) \cap W^{2,2}(\Omega)$ , and this is satisfied by the example above.

Below we prove error bounds in weaker norms,  $\|\cdot\|_{W^{1,q}(\Omega^h)}$  with  $q \in [1, p)$ .

**Lemma 4.1.** For all  $t \in [2, p]$  and  $q \in [1, t]$  for which

$$(4.6a) \quad \int_{\Omega} |\nabla u|^{-(p-t)q/(t-q)} d\Omega < \infty \quad \text{if } q \in [1, t)$$

and

$$(4.6b) \quad |\nabla u|^{-(p-t)} \in L^\infty(\Omega) \quad \text{if } q = t,$$

we have for  $u \in W^{1,\infty}(\Omega) \cap W^{2,s}(\Omega)$  with  $s \in [1, 2]$  that

$$(4.7) \quad \|u - u^h\|_{W^{1,q}(\Omega^h)} \leq Ch^{s/t}.$$

*Proof.* Choosing  $\delta_1 = 2 - s$  and  $\delta_2 = t - 2$  in (2.11), noting (4.3) and (4.6), and applying a Hölder inequality, we obtain that

$$(4.8) \quad |u - u^h|_{W^{1,q}(\Omega^h)}^t \leq C|u - u^h|_{(p,t)} \leq C|u - \pi_h u|_{(p,s)} \leq C|u - \pi_h u|_{W^{1,s}(\Omega^h)}^s.$$

The desired result (4.7) then follows from (4.8), (1.6a), (3.3), and (4.4).  $\square$

To improve on the  $h^{s/p}$  convergence rate for the error in (4.5b), we wish to take  $t \in [2, p)$ , which gives rise to the restrictions (4.6) on  $u$ ; that is, we require  $\{(x, y) \in \Omega: |\nabla u(x, y)| = 0\}$  to have zero measure and a growth condition on  $|\nabla u|^{-1}$ . From inspection we see that the weakest growth restriction on  $|\nabla u|^{-1}$  for a fixed  $t$  is needed when  $q = 1$ . We now look for sufficient conditions on  $u$  and the data  $f$  in order for these restrictions to hold.

**Lemma 4.2.** If  $u \in W^{1,\infty}(\Omega) \cap W^{2,s}(\Omega)$ ,  $s \in [1, \infty]$ , then there exists an  $M \in L^s(\Omega)$  such that

$$(4.9) \quad |f| \leq M|\nabla u|^{p-2} \quad \text{a.e. in } \Omega.$$

*Proof.* Let  $\nabla u \equiv (v_1, v_2) \in [W^{1,s}(\Omega)]^2$  and  $v \equiv (v_1^2 + v_2^2)^{1/2} \equiv |\nabla u| \in L^\infty(\Omega)$ . As  $|v_1/v| + |v_2/v|$  is bounded and  $\nabla v \equiv (v_1 \nabla v_1 + v_2 \nabla v_2)/v$ , it follows that  $v \in W^{1,s}(\Omega)$ . In addition, we have that

$$(4.10) \quad \begin{aligned} f &= -\operatorname{div}(v^{p-2}v_1, v^{p-2}v_2) \\ &= -v^{p-2}\{[(v_1)_x + (v_2)_y] + (p-2)[v_1v_x + v_2v_y]/v\}. \end{aligned}$$

Hence the desired result (4.9).  $\square$

Under the assumption that  $\{(x, y) \in \Omega: f(x, y) = 0\}$  has zero measure, the inequality (4.9), for example, yields for  $t \geq 2$  and  $1 \leq q < t < p$  that

$$(4.11) \quad \int_{\Omega} |\nabla u|^{-(p-t)q/(t-q)} d\Omega \leq \int_{\Omega} [M/|f|]^{(p-t)q/[(p-2)(t-q)]} d\Omega.$$

Therefore, with  $M \in L^s(\Omega)$ , for a given  $s \in [1, \infty]$ , and imposing a growth condition on  $|f|^{-1}$ , one can choose appropriate  $t$  and  $q$  so that (4.6a) and hence (4.7) hold. Below we give an example of such a result.

**Theorem 4.1.** Let  $u \in W^{1,\infty}(\Omega) \cap W^{2,s}(\Omega)$ ,  $s \in [1, \infty]$ . If  $|f|^{-\gamma} \in L^1(\Omega)$  for some  $\gamma \in (0, \infty)$ , or if  $|f|^{-1} \in L^\infty(\Omega)$  we set  $\gamma = \infty$ , then we have for  $q \in [1, p)$  that

$$(4.12a) \quad \|u - u^h\|_{W^{1,q}(\Omega^h)} \leq \begin{cases} Ch^{2/t} & \text{if } s \geq 2, \\ Ch^{s/t} & \text{if } s \in [1, 2), \end{cases}$$

where

$$(4.12b) \quad t = \max\{2, q[(s + \gamma)p + (p - 2)\gamma s]/[(s + \gamma)q + (p - 2)\gamma s]\}.$$

*Proof.* First a simple calculation yields that  $t$  satisfying (4.12b) is such that  $t \in [2, p)$  and  $t > q$ . Setting

$$\eta \equiv q(p - t)/[(p - 2)(t - q)],$$

we conclude that  $\eta \leq \gamma s/(s + \gamma)$  and hence  $s\eta \leq \gamma(s - \eta)$ , and if  $\gamma$  is finite then  $\eta < s$ . Therefore, from (4.11), the assumptions on  $f$  and Hölder's inequality we have

$$(4.13) \quad \begin{aligned} \int_{\Omega} |\nabla u|^{-(p-t)q/(t-q)} d\Omega &\leq \int_{\Omega} (M|f|^{-1})^{\eta} d\Omega \\ &\leq \left( \int_{\Omega} M^s d\Omega \right)^{\eta/s} \left( \int_{\Omega} |f|^{-\eta s/(s-\eta)} d\Omega \right)^{(s-\eta)/s} \leq C. \end{aligned}$$

Similarly, (4.13) holds if  $\gamma$  is infinite, as  $\eta \leq s$ . The desired result (4.12a) then follows from (4.6a) and (4.7).  $\square$

We note that for fixed  $q, \gamma$ , and  $s$  the right-hand side of (4.12b) tends to  $\max\{2, q[(s + \gamma) + \gamma s]/\gamma s\}$  as  $p \rightarrow \infty$ . Therefore, the error bound (4.12a) does not degenerate as  $p \rightarrow \infty$ , unlike (4.2b) and (4.5b).

**Corollary 4.1.** *Let  $u \in W^{1,\infty}(\Omega) \cap W^{2,s}(\Omega)$ ,  $s \in [1, \infty]$ . Suppose that there exists a constant  $\rho > 0$  such that  $|f| \geq \rho$  a.e. in  $\Omega$ ; then for  $q \in [1, p)$  we have that*

$$(4.14a) \quad \|u - u^h\|_{W^{1,q}(\Omega^h)} \leq \begin{cases} Ch^{2/t} & \text{if } s \geq 2, \\ Ch^{s/t} & \text{if } s \in [1, 2), \end{cases}$$

where

$$(4.14b) \quad t = \max\{2, q[p + (p - 2)s]/[q + (p - 2)s]\}.$$

Hence, we have that for  $q = 2s/(1 + s)$

$$(4.14c) \quad \|u - u^h\|_{W^{1,q}(\Omega^h)} \leq \begin{cases} Ch & \text{if } s \geq 2, \\ Ch^{s/2} & \text{if } s \in [1, 2). \end{cases}$$

*Proof.* The result (4.14a,b) follows directly from setting  $\gamma = \infty$  in (4.12). The result (4.14c) then follows from (4.14a,b) by noting that  $t = 2$  if  $q = 2s/(1 + s)$ .  $\square$

## 5. NUMERICAL EXAMPLES

The standard Galerkin method analyzed in the previous sections requires the term  $\int_{\Omega^h} f v^h d\Omega^h$  for all  $v^h \in S_0^h$  to be integrated exactly. This is difficult in practice, and it is computationally more convenient to consider a scheme where numerical integration is applied to this term. With  $\bar{\Omega}^h \equiv \bigcup_{\tau \in \mathcal{T}^h} \bar{\tau}$  and  $\{a_i\}_{i=1}^3$  being the vertices of a triangle  $\tau$ , we define the quadrature rule

$$(5.1) \quad Q_{\tau}(v) \equiv \frac{1}{3} \text{meas}(\tau) \sum_{i=1}^3 v(a_i) \equiv \int_{\tau} \pi_h v d\tau$$

approximating  $\int_{\tau} v d\tau$  for  $v \in C(\bar{\tau})$ . Then, for  $v, w \in C(\bar{\Omega}^h)$ , we set

$$(5.2) \quad (v, w)^h \equiv \sum_{\tau \in T^h} Q_{\tau}(vw) \equiv \int_{\Omega^h} \pi_h(vw) d\Omega^h$$

as an approximation to  $\int_{\Omega^h} vw d\Omega^h$ .

The fully practical finite element approximation of  $(\mathcal{P})$  that we wish to consider is:

$(\hat{\mathcal{P}}^h)$  Find  $\hat{u}^h \in S_g^h$  such that

$$(5.3) \quad \int_{\Omega^h} |\nabla \hat{u}^h|^{p-2} (\nabla \hat{u}^h, \nabla v^h)_{\mathbb{R}^2} d\Omega^h = (f, v^h)^h \quad \forall v^h \in S_0^h.$$

The corresponding minimization problem is:

$(\hat{\mathcal{E}}^h)$  Find  $\hat{u}^h \in S_g^h$  such that

$$(5.4a) \quad \hat{J}_{\Omega^h}(\hat{u}^h) \leq \hat{J}_{\Omega^h}(v^h) \quad \forall v^h \in S_g^h,$$

where

$$(5.4b) \quad \hat{J}_{\Omega^h}(v^h) \equiv \frac{1}{p} \int_{\Omega^h} |\nabla v^h|^p d\Omega^h - (f, v^h)^h.$$

The well-posedness of  $(\hat{\mathcal{P}}^h) \equiv (\hat{\mathcal{E}}^h)$  follows in an analogous way to that of  $(\mathcal{P})$  and  $(\mathcal{E})$ , and

$$(5.5) \quad \|\hat{u}^h\|_{W^{1,p}(\Omega^h)} \leq C[\|f\|_{L^{\infty}(\Omega^h)}^{1/(p-1)} + \|g^h\|_{W^{1,p}(\Omega^h)}].$$

We now bound the error  $\hat{u}^h - u$ . First we have the analogue of Theorem 2.1.

**Theorem 5.1.** *Let  $u$  and  $\hat{u}^h$  be the unique solutions of  $(\mathcal{P}) \equiv (\mathcal{E})$  and  $(\hat{\mathcal{P}}^h) \equiv (\hat{\mathcal{E}}^h)$ , respectively. Let  $f \in W^{1,\infty}(\Omega) \cap W^{2,2}(\Omega)$ . Then for any  $\delta_1 \in [0, 2)$  and  $\delta_2 \geq 0$ , and any  $v^h \in S_g^h$ , it follows that*

$$(5.6) \quad |u - \hat{u}^h|_{(p, 2+\delta_2)} \leq C|u - v^h|_{(p, 2-\delta_1)} + Ch^2 \|v^h - \hat{u}^h\|_{W^{1,1}(\Omega^h)}.$$

*Proof.* The proof follows exactly that of Theorem 2.1 with  $\hat{u}^h$  and  $\hat{J}_{\Omega^h}$  instead of  $u^h$  and  $J_{\Omega^h}$  in (2.16). However, whereas  $J'_{\Omega^h}(u)(v^h - u^h) = 0$ , we now have for all  $v^h \in S_g^h$

$$(5.7a) \quad \begin{aligned} \hat{J}'_{\Omega^h}(u)(v^h - \hat{u}^h) &\equiv J'_{\Omega^h}(u)(v^h - \hat{u}^h) \\ &\quad - \int_{\Omega^h} f(v^h - \hat{u}^h) d\Omega^h + (f, v^h - \hat{u}^h)^h \\ &\equiv - \int_{\Omega^h} f(v^h - \hat{u}^h) d\Omega^h + (f, v^h - \hat{u}^h)^h \end{aligned}$$

and

$$(5.7b) \quad \begin{aligned} \left| \int_{\Omega^h} f(v^h - \hat{u}^h) d\Omega^h - (f, v^h - \hat{u}^h)^h \right| &\leq C\|(I - \pi_h)[f(v^h - \hat{u}^h)]\|_{L^1(\Omega^h)} \\ &\leq Ch^2 |f(v^h - \hat{u}^h)|_{W^{2,1}(\Omega^h)} \leq Ch^2 \|v^h - \hat{u}^h\|_{W^{1,1}(\Omega^h)}, \end{aligned}$$

provided  $f \in W^{1,\infty}(\Omega) \cap W^{2,2}(\Omega)$ . Hence, we obtain the desired result (5.6).  $\square$

In particular, assuming  $u \in W^{2,1}(\Omega)$  if  $p < 2$ , we have for  $v^h \equiv g^h \equiv \pi_h u$  that for any  $\delta_1 \in [0, 2)$  and  $\delta_2 \geq 0$

$$(5.8) \quad |u - \hat{u}^h|_{(p, 2+\delta_2)} \leq C|u - \pi_h u|_{(p, 2-\delta_1)} + Ch^2.$$

Hence, it is a simple matter to check that the results of the previous sections hold for  $\hat{u}^h$  as well as  $u^h$  if  $f \in W^{1,\infty}(\Omega) \cap W^{2,2}(\Omega)$ . We note that this constraint on  $f$  can be weakened and is imposed here for ease of exposition only.

We now report on some numerical results with the fully practical approximation (5.3). For computational ease we took  $\Omega$  to be the square  $[0, 1] \times [0, 1]$ . This was partitioned into uniform right-angled triangles by dividing it first into equal squares of sides of length  $1/N$  and then into triangles by inserting the SW-NE diagonals. We imposed homogeneous Neumann data on the sides  $x = 0$  and  $y = 0$  and Dirichlet data on the sides  $x = 1$  and  $y = 1$ . Therefore, the problem can be viewed as a Dirichlet problem over  $[-1, 1] \times [-1, 1]$ , and so our error analysis applies directly.

We computed our approximation (5.3) by solving the equivalent minimization problem (5.4). We used a Polak-Ribière conjugate gradient method, which worked reasonably well for the values of  $p$  reported here. We did not experiment with the augmented Lagrangian approach advocated by Glowinski and Marrocco [5], but this conjugate gradient approach was far superior to the gradient method suggested by Wei [10].

For our test problems we consider solutions of the radially symmetric problem, Example 3.1, extended to the unit square. In the first three examples we took for various values of  $p$  and  $\gamma$

$$(5.9) \quad \begin{aligned} f &\equiv F(r) \equiv r^\sigma \quad \text{and} \\ u &\equiv U(r) \equiv (p-1)[1/(\sigma+2)]^{1/(p-1)}[1-r^{(\sigma+p)/(p-1)}]/(\sigma+p). \end{aligned}$$

In all the examples,  $f$  is sufficiently smooth, so that the error bounds for  $u^h$  in the previous sections hold for  $\hat{u}^h$  as well.

**Example 5.1.** This is (5.9) with  $\sigma = 0$  and  $p = 1.5$ . It follows from (3.12) that  $u \in W^{3,1}(\Omega) \cap C^{2,1}(\bar{\Omega})$ , and so from Theorem 3.1 we expect  $O(h)$  convergence in  $W^{1,1.5}(\Omega)$ . This is certainly achieved by inspecting Table 5.1, where we adopt the notation  $0.8233(-3) \equiv 0.8233 \times 10^{-3}$ . In fact,  $\hat{u}^h$  is converging to  $u$  at the rate  $O(h^2)$  in  $L^\infty(\Omega)$ , and there is a superconvergence for  $\pi_h u - \hat{u}^h$  in  $W^{1,1}(\Omega)$  and  $W^{1,p}(\Omega)$ .  $\square$

TABLE 5.1

$N$	$\ \pi_h u - \hat{u}^h\ _{W^{1,1}(\Omega)}$	$\ \pi_h u - \hat{u}^h\ _{W^{1,p}(\Omega)}$	$\ \pi_h u - \hat{u}^h\ _{L^\infty(\Omega)}$
10	0.8233(-3)	0.4823(-3)	0.8150(-3)
20	0.2061(-3)	0.1207(-3)	0.2034(-3)
40	0.5196(-4)	0.3043(-4)	0.5109(-4)
80	0.1235(-4)	0.723(-5)	0.1263(-4)

**Example 5.2.** This is (5.9) with  $\sigma = 0$  and  $p = 4$ . It follows from Example 3.1 and §4 that  $u \in W^{2,s}(\Omega)$ , with  $s < 3$ , and from (4.14c) we expect  $O(h)$  convergence in  $W^{1,1}(\Omega)$ . From Table 5.2 we see this is achieved. In fact,  $\pi_h u - \hat{u}^h$  exhibits superconvergence in  $W^{1,1}(\Omega)$ .  $\square$

TABLE 5.2

$N$	$\ \pi_h u - \hat{u}^h\ _{W^{1,1}(\Omega)}$	$\ \pi_h u - \hat{u}^h\ _{W^{1,p}(\Omega)}$	$\ \pi_h u - \hat{u}^h\ _{L^\infty(\Omega)}$
10	0.1789(-2)	0.4486(-2)	0.3790(-2)
20	0.5049(-3)	0.2519(-2)	0.1585(-2)
40	0.1376(-3)	0.1414(-2)	0.6493(-3)
80	0.3659(-4)	0.7936(-3)	0.2625(-3)

**Example 5.3.** Here we take (5.9) with  $\sigma = 7$  and  $p = 4$ . It follows from Example 3.1 that  $u \in W^{2,\infty}(\Omega)$ . From (4.12), as  $s = \infty$  and  $\gamma < 2/7$ , we have with  $q = 1$  that  $t > 32/11$ . Therefore, for all  $\varepsilon > 0$  we have that

$$(5.10) \quad \|u - \hat{u}^h\|_{W^{1,1}(\Omega)} \leq Ch^{(11-\varepsilon)/16}.$$

We note that a sharper bound,  $h^{(14-\varepsilon)/19}$ , can be obtained by noting that for this model problem  $|U'(r)| \geq Cr^{8/3}$  and applying (4.6) and (4.7) directly. From Table 5.3 we see that the above bounds are pessimistic. In fact, we have  $O(h^2)$  convergence in  $L^\infty(\Omega)$  and  $\pi_h u - \hat{u}^h$  exhibits superconvergence in  $W^{1,1}(\Omega)$ .  $\square$

TABLE 5.3

$N$	$\ \pi_h u - \hat{u}^h\ _{W^{1,1}(\Omega)}$	$\ \pi_h u - \hat{u}^h\ _{W^{1,p}(\Omega)}$	$\ \pi_h u - \hat{u}^h\ _{L^\infty(\Omega)}$
10	0.8153(-3)	0.5988(-2)	0.5014(-2)
20	0.2164(-2)	0.1931(-2)	0.1235(-2)
40	0.5918(-3)	0.8893(-3)	0.2989(-3)
80	0.1429(-3)	0.1952(-3)	0.6449(-4)

Finally we consider an example for  $p > 2$ , where  $\{(x, y) \in \Omega: f(x, y) = 0\}$  does not have zero measure.

**Example 5.4.** We take

$$(5.11a) \quad F(r) \equiv \begin{cases} 0 & \text{for } r \leq a, \\ 4^{p-1}(r-a)^{(3p-4)}[2 + (a/r) - 3p] & \text{for } r \geq a \end{cases}$$

and

$$(5.11b) \quad U(r) \equiv \begin{cases} 0 & \text{for } r \leq a, \\ (r-a)^4 & \text{for } r \geq a \end{cases}$$

with  $a = 0.3$  and  $p = 4$ . At present the only global error estimate we have for this case is the result (4.5b). Clearly, this is pessimistic from inspecting Table 5.4, where once again we see  $O(h^2)$  convergence in  $L^\infty(\Omega)$  and  $\pi_h u - \hat{u}^h$  is superconvergent in  $W^{1,1}(\Omega)$ . We note that the maximum error did not occur in the disc  $\{r: r \leq 0.3\}$ .  $\square$

TABLE 5.4

$N$	$\ \pi_h u - \hat{u}^h\ _{W^{1,1}(\Omega)}$	$\ \pi_h u - \hat{u}^h\ _{W^{1,p}(\Omega)}$	$\ \pi_h u - \hat{u}^h\ _{L^\infty(\Omega)}$
10	0.5879(-1)	0.4653(-1)	0.3080(-1)
20	0.1553(-1)	0.1182(-1)	0.7930(-2)
40	0.4332(-2)	0.3118(-2)	0.1992(-2)
80	0.1139(-2)	0.1706(-2)	0.4923(-3)

## BIBLIOGRAPHY

1. C. Atkinson and C. R. Champion, *Some boundary-value problems for the equation  $\nabla \cdot (|\nabla \phi|^N \nabla \phi) = 0$* , Quart. J. Mech. Appl. Math. **37** (1984), 401–419.
2. C. Atkinson and C. W. Jones, *Similarity solutions in some non-linear diffusion problems and in boundary-layer flow of a pseudo plastic fluid*, Quart. J. Mech. Appl. Math. **27** (1974), 193–211.
3. S.-S. Chow, *Finite element error estimates for non-linear elliptic equations of monotone type*, Numer. Math. **54** (1989), 373–393.
4. P. G. Ciarlet, *The finite element method for elliptic problems*, North-Holland, Amsterdam, 1978.
5. R. Glowinski and A. Marrocco, *Sur l'approximation par éléments finis d'ordre un, et la résolution, par pénalisation-dualité, d'une classe de problèmes de Dirichlet non lineaires*, RAIRO Anal. Numér. **2** (1975), 41–76.
6. A. Kufner, O. John, and S. Fučík, *Function spaces*, Noordhoff, Leyden, 1977.
7. G. M. Lieberman, *Boundary regularity for solutions of degenerate elliptic equations*, Non-linear Anal. **12** (1988), 1203–1219.
8. J. R. Philip, *N-diffusion*, Austral. J. Phys. **14** (1961), 1–13.
9. V. B. Tyukhtin, *The rate of convergence of approximation methods of solution of one-sided variational problems. I*, Vestnik Leningrad. Univ. Mat. Mekh. Astronom. **13** (1982), 111–113.
10. D. Wei, *Finite element approximations of solutions to  $p$ -harmonic equations with Dirichlet data*, Numer. Funct. Anal. Optim. **10** (1989), 1235–1251.

DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE, LONDON SW7 2BZ, ENGLAND

E-mail address: j.barrett@ic.ac.uk