

THE p AND hp VERSIONS OF THE FINITE ELEMENT METHOD FOR PROBLEMS WITH BOUNDARY LAYERS

CHRISTOPH SCHWAB AND MANIL SURI

Dedicated to Professor Ivo Babuška on the occasion of his seventieth birthday

ABSTRACT. We study the uniform approximation of boundary layer functions $\exp(-x/d)$ for $x \in (0, 1)$, $d \in (0, 1]$, by the p and hp versions of the finite element method. For the p version (with fixed mesh), we prove super-exponential convergence in the range $p + 1/2 > e/(2d)$. We also establish, for this version, an overall convergence rate of $\mathcal{O}(p^{-1}\sqrt{\ln p})$ in the energy norm error which is *uniform* in d , and show that this rate is sharp (up to the $\sqrt{\ln p}$ term) when *robust* estimates uniform in $d \in (0, 1]$ are considered. For the p version with variable mesh (i.e., the hp version), we show that exponential convergence, uniform in $d \in (0, 1]$, is achieved by taking the first element at the boundary layer to be of size $\mathcal{O}(pd)$.

Numerical experiments for a model elliptic singular perturbation problem show good agreement with our convergence estimates, even when few degrees of freedom are used and when d is as small as, e.g., 10^{-8} . They also illustrate the superiority of the hp approach over other methods, including a low-order h version with optimal “exponential” mesh refinement.

The estimates established in this paper are also applicable in the context of corresponding spectral element methods.

1. INTRODUCTION

Our goal in this paper is to develop the approximation theory for *boundary layer functions*

$$(1.1) \quad u(x) = \exp(-ax/d), \quad 0 < x < L,$$

where $d \in (0, 1]$ is a small parameter that can approach zero, $a > 0$ is a constant and $L \geq 1$ is a typical length scale of the problem under consideration. We are interested in obtaining convergence estimates that are *robust*, i.e., *uniform* in d , when (1.1) is approximated by piecewise polynomials via p and hp type numerical schemes.

Boundary layers (1.1) arise as solution components in singularly perturbed elliptic boundary value problems, a model example of which is

$$(1.2) \quad L_d u_d := -d^2 u_d''(x) + a^2 u_d(x) = f(x), \quad x \in I = (-1, 1),$$

Received by the editor March 7, 1995.

1991 *Mathematics Subject Classification.* Primary 65N30, 35B30, 65N15.

Key words and phrases. Boundary layer, singularly perturbed problem, p version, hp version, spectral element method.

This work was supported in part by the Air Force Office of Scientific Research, Air Force Systems Command, USAF under Grant F49620-92-J-0100.

$$(1.3) \quad u_d(\pm 1) = \alpha^\pm.$$

A large body of literature has been devoted to their effective resolution. Most available references analyze the convergence of finite difference or finite element schemes of *fixed* (usually low) polynomial degree in conjunction with various mesh refinements (the *h* version); see, e.g., [4, 6, 13, 18, 19], and the references therein.

If the mesh refinement is quasi-uniform (or, more generally, independent of d), either on the whole domain or locally near the boundary, then the optimal (algebraic) decrease in the global error is observed *provided* a condition of the form $h \leq Cd$ is met (h being the mesh spacing parameter). Such methods are *nonrobust*, in a sense made precise in §3. In practical terms, the amount of discretization required with such schemes for satisfactory resolution of the boundary layers may be infeasible when d is very small. On the other hand, strongly graded d -dependent mesh refinement, like the one from [20] presented in §6, does lead to robust convergence, at an optimal rate that is *algebraic* (see [4, 16, 18, 19, 20], where this and other graded meshes are discussed).

An alternative approach is to increase the polynomial degree and keep the mesh fixed, i.e., use a p version or spectral element method. In [5], various such schemes (Galerkin, Tau and Collocation) have been considered for the special case of (1.2)–(1.3) where $f \equiv 0$, $\alpha^+ = 1$, $\alpha^- = 0$, using a Chebyshev-weighted spectral approximation. In this paper, we consider the *unweighted* Galerkin p version/spectral element approximation. We provide a detailed study of the approximation theory for this method, showing that an *asymptotic superexponential convergence rate* for the error in the energy norm is achieved for $\tilde{p} := p + 1/2 > e/(2d)$. We also provide estimates for this error in the preasymptotic phase when d is small, showing that (A) for $(3/(4d))^{1/2} \leq \tilde{p} \leq 2/d$, the error is bounded by $C \exp(-\tilde{p}^2 d/3)$ and (B) for $\tilde{p} \leq Kd^{-1/2}$, the error is bounded by Cp^{-1} (numerical experiments in §6 are in agreement with these rates). The results we prove for a single element also hold when a *fixed* mesh with several elements is used. Using our various estimates, we establish that for the pure p version on fixed meshes, the overall robust rate, uniform in d , is $\mathcal{O}(p^{-1}\sqrt{\ln p})$ and, up to the $\sqrt{\ln p}$ term, this is the best possible. Note that this rate is essentially *double* the uniform rate of $\mathcal{O}(h^{1/2})$ achievable (for the global error) by the h version with quasi-uniform meshes (Theorem A.1(ii) of [13]). It is also double the uniform rate for p version/spectral element methods that can be established from the results in [5]. (Since the methods in [5] involve a weighted projection, the estimates there are in (stronger) weighted norms.)

The p -type results in [5] can be considerably improved by using special “mapped” polynomials in the spectral element method. This is shown in [8, 7], where singular mappings of appropriately high order are used to establish algebraic rates of convergence that deteriorate relatively slowly as $d \rightarrow 0$. However, these estimates are still not uniform in d , and therefore, not robust in our sense.

Our main result in this paper shows that excellent robust rates for the uniform approximation of functions (1.1) can be achieved by using, instead of the pure p /spectral version, a *variable* mesh with only one more element. More precisely, a *robust exponential rate* can be obtained by using the p version on two elements, where the first one is of size $\mathcal{O}(pd)$. (For problems like (1.2)–(1.3), three elements are needed, owing to boundary layers at either end — see §6.) We call this an *hp* version since the size (though not the number) of elements changes, as does p . (More appropriately, it is an “*rp*” method.) Note that an exponential rate is not

possible with either the h version or the pure p /spectral version — the estimates obtained in the papers above are all *algebraic*. Finite element computations for (1.2)–(1.3) presented in §6 confirm the theoretical convergence estimates obtained here and clearly show the dramatic superiority of this robust hp FEM over other methods, especially for small d .

Although we concentrate here only on the approximation theory for the one-dimensional function (1.1) applied to one-dimensional problems like (1.2), the scope of our results is wider. This is due to the fact that solutions to singularly perturbed problems over *two-dimensional* domains ω , arising, e.g., in beam, plate and shell theory, as well as in reaction-diffusion and certain fluid dynamics problems, also exhibit boundary layers, which are of the form

$$(1.4) \quad u_{BL}(s, x) = C(s) \exp(-a\rho(x)/d), \quad 0 < s < T, \quad 0 < \rho < \rho_0.$$

Here s, ρ denote, respectively, the arc length and normal distance to the boundary, of a point x in a neighborhood of $\partial\omega$, and the function $C(s)$ is smooth. For several problems of practical interest, decompositions of the solution into a regular part and such boundary layers $u_{BL}(s, x)$ have been obtained in the literature; cf. [1] for the Reissner-Mindlin plate, [17] for beam theory, [11, 12] for shells. Similar decompositions arise also in three-dimensional problems (then, however, $s = (s_1, s_2)$ are coordinates in the boundary manifold). The key observation from (1.4) is that since $C(s)$ is smooth, the boundary layer phenomenon is essentially one-dimensional, namely, in the direction normal to $\partial\omega$. Hence, the crucial aspect of the FE approximation of such functions is how the FE spaces are designed in the ρ direction, i.e., how the function (1.1) is approximated in one dimension. Using our results, therefore, we can construct two- and three-dimensional FE spaces (with robust exponential convergence) for the functions (1.4), e.g., using tensor product spaces in the (s, ρ) coordinates. See [15, 16]. Note that “brute force” mesh refinement will be even less competitive in two dimensions and practically impossible in three dimensions.

The outline of this paper is as follows. In §2 we present an asymptotic expansion for the solution of the model problem (1.2)–(1.3) which includes the boundary layers. The proof uses standard techniques and is provided for completeness in the Appendix. In §3, we describe the finite element methods and error measures to be analyzed. We also define the concept of robustness, using a definition from [2]. Section 4 is devoted to the convergence analysis of the p version FEM. In §5, we consider an hp version for which we prove a robust exponential convergence rate in various norms. Finally, in §6 we present numerical experiments comparing, in particular, the p and hp version FEMs analyzed here with an h version from [20], based on asymptotically optimal meshes. We show that the hp version consistently outperforms the other versions and that high accuracy can be achieved with few degrees of freedom for arbitrarily small d (we take values of d as small as 10^{-8}).

Throughout, $H^k(I)$ will denote the Sobolev space of order $k \in \mathbb{N}_0$ on an interval $I \subset \mathbb{R}$, with $H^0(I) = L_2(I)$ and $\|\cdot\|_{k,I}$, $|\cdot|_{k,I}$ denoting the norm and seminorm as usual. Whenever there is no confusion about the domain, we omit the subscript I . For $u, v \in L_2(I)$, we denote by (u, v) the L_2 inner product. Also, $H_0^1(I) = \{u \in H^1(I) : u(\pm 1) = 0\}$, $H_D^1(I) = \{u \in H^1(I) : u(\pm 1) = \alpha^\pm\}$ and $H^{-1}(I) = (H_0^1(I))^*$, the dual space. Throughout the paper, C, K will denote generic constants, while C_i, \tilde{C}_i will denote constants that are explicitly given or can be easily estimated from the exposition.

2. REGULARITY OF THE MODEL PROBLEM

The variational formulation of the model problem (1.2)–(1.3) reads: Find $u_d \in H_D^1(I)$ such that

$$(2.1) \quad B_d(u_d, v) = (f, v) \quad \forall v \in H_0^1(I).$$

Here, $f \in H^{-1}(I)$ and

$$(2.2) \quad B_d(u, v) := \int_I \{d^2 u' v' + a^2 uv\} dx.$$

For every $f \in H^{-1}(I)$ the problem (2.1) admits a unique solution $u_d \in H_D^1(I)$, and if $f \in H^k(I)$, then $u_d \in H^{k+2}(I) \cap H_D^1(I)$. This regularity, however, is nonuniform in d since in the a priori “shift” estimate

$$(2.3) \quad \|u_d\|_{k+2} \leq C(k, d) \|f\|_k, \quad k = 0, 1, 2, \dots,$$

the constant C strongly depends on d . The following theorem, the proof of which can be found in the Appendix, presents a decomposition of u_d into a smooth part $u_d^M(x)$ and boundary layers

$$(2.4) \quad u_{a,d}(x) = \exp(-a(1+x)/d), \quad \bar{u}_{a,d}(x) = \exp(-a(1-x)/d).$$

Theorem 2.1. *Let $f \in H^{4M+2}(I)$ for some $M \in \mathbb{N}$. Then*

$$(2.5) \quad u_d(x) = u_d^M(x) + A_d^M u_{a,d}(x) + B_d^M \bar{u}_{a,d}(x),$$

where $u_d^M(x)$ satisfies the following regularity estimate uniformly in d for $\ell = 0, 1, \dots, 2M$:

$$(2.6) \quad |u_d^M|_\ell \leq a^{-1} (d/a)^{2M+2-\ell} |f|_{(2M+2)} + 2a^{-2} \sum_{k=0}^{M+1} (d/a)^{2k} |f|_{2k+\ell}.$$

Further,

$$(2.7) \quad |A_d^M| + |B_d^M| \leq C(a) \left\{ |\alpha^+| + |\alpha^-| + \sum_{k=0}^M (d/a)^{2k} \left(|f^{(2k)}(-1)| + |f^{(2k)}(+1)| \right) \right\},$$

where $C(a)$ is independent of M and d .

For any interval \tilde{I} let $\Pi_n(\tilde{I})$ denote the set of polynomials on \tilde{I} of degree $\leq n$. The following result follows by Remark 6.1 in the Appendix.

Corollary 2.1. *Let $f \in \Pi_{2M+1}(I)$; then $u_d^M \in \Pi_{2M+1}(I)$ in (2.5).*

Remark 2.1. Analogous results hold when the Dirichlet end conditions (1.3) are replaced by Neumann or mixed boundary conditions.

For f smooth enough (i.e., M large enough), we see from Theorem 2.1 that the regularity of u_d (in terms of d) will be determined by the boundary layer terms. We have, in fact, by (2.5)–(2.7),

$$(2.8) \quad |u_d|_\ell \leq |u_d^M|_\ell + |A_d^M| |u_{a,d}|_\ell + |B_d^M| |\bar{u}_{a,d}|_\ell \leq C(1 + |u_{a,d}|_\ell + |\bar{u}_{a,d}|_\ell),$$

where the constant C depends upon a , f and α^\pm but is independent of d . For the function $u_{a,d}$, we have for $\ell = 0, 1, 2, \dots$

$$(2.9) \quad |u_{a,d}|_\ell = \left(\frac{d}{a}\right)^{\frac{1}{2}-\ell} \left[\frac{1 - e^{-4a/d}}{2} \right]^{1/2} \approx C d^{\frac{1}{2}-\ell},$$

so that (2.9) and its analog for $|\bar{u}_{\alpha,d}|_\ell$, substituted in (2.8), gives an upper bound for $|u_d|_\ell$. Since, except for special cases, the coefficients A_d^M, B_d^N are nonzero, we see that the following *equivalence* generally holds:

$$(2.10) \quad |u_d|_\ell \approx C(1 + d^{\frac{1}{2}-\ell}), \quad \ell = 0, 1, \dots, 2M.$$

To conclude this section, we define the following *solution spaces*, which will be used later:

$$H_{d,M}^B = \{u_d | u_d \text{ is a solution of (1.2)-(1.3) with } f \in H^{4M+2}(I), \\ \|f\|_{4M+2} \leq B, |\alpha^\pm| \leq B\},$$

$$H_{d,\Pi_n}^B = \{u_d | u_d \text{ is a solution of (1.2)-(1.3) with } f \in \Pi_n(I) \text{ such that} \\ \text{all coefficients in } f \text{ are absolutely bounded by } B, |\alpha^\pm| \leq B\}.$$

3. THE FINITE ELEMENT METHOD

For any finite-dimensional subspace S of $H^1(I)$, denote $S_D = S \cap H_D^1(I)$, $S_0 = S \cap H_0^1(I)$. Then a finite element approximation u_d^S of u_d is obtained by restricting both sides of the weak formulation (2.1) to finite-dimensional subspaces: Find $u_d^S \in S_D$ such that

$$(3.1) \quad B_d(u_d^S, v) = (f, v) \quad \forall v \in S_0.$$

For every $d \in (0, 1]$ there exists a unique solution $u_d^S \in S_D$ of (3.1).

We will be interested in spaces S of piecewise polynomials on I characterized by the mesh-degree combination $\Sigma = (\Delta, \vec{p})$, defined as follows. Let the $m + 1 \geq 2$ nodal points

$$(3.2) \quad -1 =: x_0 < x_1 < x_2 < \dots < x_{m-1} < x_m := 1$$

be given; then the mesh Δ is defined by

$$(3.3) \quad \Delta = \{I_i\}_{i=1}^m, \quad I_i = (x_{i-1}, x_i), \quad h_i = |I_i| = x_i - x_{i-1}$$

(we will also write $\Delta = \{x_0, x_1, \dots, x_m\}$ where convenient).

The degree vector \vec{p} is defined by

$$(3.4) \quad \vec{p} = (p_1, p_2, \dots, p_m).$$

Then

$$(3.5) \quad S(\Sigma) = \{u : u|_{I_i} \in \Pi_{p_i}(I_i), \quad I_i \in \Delta\} \cap C^0[-1, 1].$$

Obviously, $S(\Sigma) \subset H^1(I)$ and

$$(3.6) \quad \dim S(\Sigma) = 1 + \sum_{i=1}^m p_i, \quad N = \dim S_D(\Sigma) = \dim S(\Sigma) - 2.$$

By (2.1), (3.1),

$$(3.7) \quad B_d(u_d - u_d^S, v) = 0 \quad \forall v \in S_0(\Sigma),$$

so that $e_d^S = u_d - u_d^S$ satisfies

$$(3.8) \quad \|e_d^S\|_d = \inf_{\chi \in S_D(\Sigma)} \|u_d - \chi\|_d.$$

Here the energy norm $\|\cdot\|_d$, $0 < d \leq 1$, is defined by

$$(3.9) \quad \|v\|_d = (B_d(v, v))^{1/2} = (d^2|v|_1^2 + \|v\|_0^2)^{1/2} \approx d|v|_1 + \|v\|_0.$$

The question we wish to explore here is the design of the spaces $S(\Sigma)$ such that $\|e_d^S\|_d$ has a good convergence rate $g(N) \rightarrow 0$ as $N \rightarrow \infty$ independent of d . To do this, we recall the definition of *robustness* from [2].

Definition 3.1. The FEM for problem (3.1) using spaces $S_D(\Sigma)$ is *robust* with uniform order $g(N)$ for $0 < d \leq 1$ with respect to solution sets $\mathcal{H}_d = H_{d,M}^B$ (or H_{d,Π_n}^B) and error measures $E_d = \|\cdot\|_d$ if and only if

$$\lim_{N \rightarrow \infty} \left(\left(\sup_{d \in (0,1]} \sup_{u_d \in \mathcal{H}_d} E_d(u_d - u_d^N) \right) \frac{1}{g(N)} \right) = C < \infty.$$

Although we concentrate here primarily on the energy norm, other error measures could be considered as well: the L_2 norm obviously follows as a corollary, while the maximum norm is considered in Corollary 5.1. Note that by (2.10), the unscaled H^1 norm of u_d is not bounded uniformly for $d \in (0, 1]$, so that we cannot expect robustness with uniform order in this norm (see, e.g., estimates (4.3) in [13]).

Let $u_d \in H_{d,M}^B$. Using Theorem 2.1 and (3.8), we see immediately that for the energy norm,

$$\begin{aligned} (3.10) \quad E_d(u_d - u_d^N) &= \|e_d^S\|_d \leq \inf_{\chi \in S_D} \|(u_d^M + A_d^M u_{a,d} + B_d^M \bar{u}_{a,d}) - \chi\|_d \\ &\leq \inf_{\substack{\chi_1 \in S \\ \chi_1(\pm 1) = u_d^M(\pm 1)}} \|u_d^M - \chi_1\|_d \\ &\quad + |A_d^M| \inf_{\substack{\chi_2 \in S \\ \chi_2(\pm 1) = u_{a,d}(\pm 1)}} \|u_{a,d} - \chi_2\|_d + |B_d^M| \inf_{\substack{\chi_3 \in S \\ \chi_3(\pm 1) = \bar{u}_{a,d}}} \|\bar{u}_{a,d} - \chi_3\|_d. \end{aligned}$$

Assume the space $S(\Sigma)$ has the following approximation property:

$$(3.11) \quad \inf_{\substack{\chi \in S \\ \chi(\pm 1) = u(\pm 1)}} \|u - \chi\|_1 \leq F(N, k) \|u\|_k, \quad k = 1, 2, \dots,$$

where $F(N, k)$ is some (optimal) approximation order (i.e., $F(N, k) \rightarrow 0$ as $N \rightarrow \infty$). Then for $u_d \in H_{d,M}^B$, by Theorem 2.1, the first infimum in (3.10) will tend to zero at the rate $KF(N, 2M)$ as $N \rightarrow \infty$, where K is a constant independent of d (K only depends upon B and M). Also, we may assume by Theorem 2.1 that $|A_d^M|, |B_d^M| \leq K$, so that the second infimum in (3.10) will decrease at the rate $K\sqrt{a} \Phi(\frac{d}{a}, S)$, where

$$\begin{aligned} (3.12) \quad \Phi(d, S) &= \inf_{\substack{\chi \in S \\ \chi(\pm 1) = u_{1,d}(\pm 1)}} \|u_{1,d} - \chi\|_d \\ &= \inf_{\substack{\chi \in S \\ \chi(\pm 1) = u_{1,d}(\pm 1)}} \{d^2 |u_{1,d} - \chi|_1^2 + \|u_{1,d} - \chi\|_0^2\}^{1/2}. \end{aligned}$$

By symmetry about $x = 0$, the last term in (3.10) will also have the same bound. Then it may be shown that our FEM will be robust in the sense of Definition 3.1 if and only if $\Phi(d, S)$ in (3.12) can be bounded independently of d , i.e.,

$$(3.13) \quad \sup_{d \in (0,1]} \Phi(d, S) \leq G(N).$$

In that case, by (3.10)–(3.13) and Definition 3.1, our FEM will be robust with uniform order

$$(3.14) \quad g(N) = C \max\{F(N, 2M), G(N)\}.$$

We will use the following related definition.

Definition 3.2. The spaces $S(\Sigma)$ will be said to approximate boundary layers $u_{1,d}$ robustly at the rate $G(N)$ in the energy norm if and only if (3.13) holds.

Remark 3.1. Our main concern in (3.14) is the rate $G(N)$, i.e., finding spaces $S(\Sigma)$ such that (3.13) holds with $G(N) \rightarrow 0$ uniformly at a sufficiently fast rate. This is because in general, $G(N)$ will be the dominant term in (3.14), the idea being that M is large enough so that $F(N, 2M)$ is sufficiently small. For the hp spaces in §5, however, $G(N) \rightarrow 0$ exponentially, so that the algebraic rate $F(N, 2M)$ achieved by assuming regularity in terms of finite M will dominate as N becomes sufficiently large. This technical problem could be overcome by restricting the set of solutions \mathcal{H}_d in Definition 3.1 to those for which the first infimum in (3.10) decays exponentially (or sufficiently fast). In particular, choosing $\mathcal{H}_d = H_{d,\Pi_n}^B$ will make this infimum vanish for suitable $S(\Sigma)$ (see Theorem 5.2 ahead).

Remark 3.2. The FE spaces satisfying (3.13) constructed in this paper and the estimates $G(N)$ established for them are also applicable to various other problems where the solution can be decomposed into boundary layers and smooth terms.

4. APPROXIMATION RESULTS FOR THE p VERSION

In this section, we will prove asymptotic error estimates for $\Phi(d, S)$ given by (3.12) as $p \rightarrow \infty$, in the case that a single element $I = (-1, 1)$ is used, i.e., $S(\Sigma) = \Pi_p(I)$. Our first estimate (4.1) will be valid uniformly in d for the range $\tilde{p} > e/2d$. (For any integer k , we write $\tilde{k} = k + \frac{1}{2}$.) We will also provide separate estimates (again uniform in d) for the preasymptotic ranges $\sqrt{3/4d} \leq \tilde{p} \leq 2/d$ and $1 < \tilde{p} < Kd^{-\frac{1}{2}}$. Our final theorem will establish a uniform robustness rate of $Cp^{-1}\sqrt{\ln p}$ for the p version over a fixed mesh, which will be shown to be *optimal* (up to the factor $\sqrt{\ln p}$).

In order to estimate $\Phi(d, S)$, we will use the following lemma from [3, Chapter 3], that will give a *concurrent* approximation of $u_{1,d}(x)$ in the $L_2(I)$ norm and $H^1(I)$ seminorm.

Lemma 4.1. Let $u, u' \in L_2(I)$ and denote by

$$(4.1) \quad a_n = \tilde{n} \int_{-1}^1 u'(x) P_n(x) dx$$

the Legendre coefficients of $u'(x)$. Then there exists $\chi \in \Pi_p(I)$ such that

$$(4.2) \quad \chi(\pm 1) = u(\pm 1),$$

$$(4.3) \quad \|u' - \chi'\|_{0,I}^2 = \sum_{n=p}^{\infty} \frac{|a_n|^2}{\tilde{n}},$$

$$(4.4) \quad \|u - \chi\|_{0,I}^2 \leq \sum_{n=p}^{\infty} \frac{|a_n|^2}{n(n+1)\tilde{n}},$$

$$(4.5) \quad \|u' - \chi'\|_{0,I} \leq \|u' - \xi'\|_{0,I}$$

for any $\xi \in \Pi_p(I)$ satisfying $\xi(\pm 1) = u(\pm 1)$.

For a proof, we refer to [3, Theorem 3.3.4].

Remark 4.1. The polynomial χ above is obtained as an antiderivative of the truncated Legendre expansion of u' , of degree $p - 1$. While by (4.5), this is optimal in the $H^1(I)$ seminorm, it is nonoptimal in the $\|\cdot\|_d$ norm. Nevertheless, Lemma 4.1 will be sufficient for our purposes here.

The estimates in (4.3), (4.4) obviously depend on the size of the Legendre coefficients a_n in dependence on d and n . The following lemma gives precise bounds for these coefficients for our function $u \equiv u_{1,d}$.

Lemma 4.2. *Let $u \equiv u_{1,d}$ and a_n be defined by (4.1). Then with $\tilde{n} = n + \frac{1}{2}$,*

$$(4.6) \quad \left(1 - \frac{2\nu_0}{\tilde{n}}\right) \leq \frac{a_n(d)}{\phi(n,d)} \leq \left(1 - \frac{2\nu_0}{\tilde{n}}\right)^{-1} \text{ for } n = 1, 2, \dots,$$

where

$$(4.7) \quad \phi(n,d) = (-1)^{n+1} \frac{d^{-\frac{1}{2}} \tilde{n}^{\frac{1}{2}}}{(1+z^2)^{\frac{1}{4}}} e^{-\tilde{n}(z-\xi(z))}, \quad z = (\tilde{n}d)^{-1},$$

$$(4.8) \quad \xi(z) = (1+z^2)^{\frac{1}{2}} + \ln\left(\frac{z}{1+(1+z^2)^{1/2}}\right),$$

$$\nu_0 = \frac{1}{6\sqrt{5}} + \frac{1}{12} \approx 0.158.$$

Proof. Using (4.1) and the fact that $u'_{1,d} = -d^{-1}u_{1,d}$, we have

$$\begin{aligned} a_n &= -\tilde{n} d^{-1} e^{-1/d} \int_{-1}^1 e^{-x/d} \frac{1}{2^n n!} \frac{d^n}{dx^n} ((x^2 - 1)^n) dx \\ &= (-1)^{n+1} d^{-n-1} e^{-1/d} \frac{\tilde{n}}{2^n n!} \int_{-1}^1 (1-x^2)^n e^{-x/d} dx. \end{aligned}$$

Hence, by formula 3.387 of [10],

$$(4.9) \quad \begin{aligned} a_n(d) &= (-1)^{n+1} d^{-n-1} e^{-1/d} \frac{\tilde{n}}{2^n} \sqrt{\pi} (2d)^{\tilde{n}} I_{\tilde{n}}(d^{-1}) \\ &= (-1)^{n+1} d^{-1/2} \sqrt{2\pi} \tilde{n} e^{-1/d} I_{\tilde{n}}(d^{-1}), \end{aligned}$$

where $I_{\tilde{n}}(d^{-1})$ is the modified Bessel function ([10, 8.406]). Thus, to obtain the asymptotic behavior of $a_n(d)$, we must investigate $I_{\tilde{n}}(d^{-1})$. To this end, we use asymptotic expansions of $I_\nu(\nu z)$ that are *uniform* for $z > 0$. Such uniform expansions have been obtained by F.W.J. Olver (see [9] and the references therein).

Let $\nu = \tilde{n} = n + \frac{1}{2}$ and $z = (\nu d)^{-1}$; then

$$(4.10) \quad e^{-1/d} I_{\tilde{n}}(d^{-1}) = e^{-\nu z} I_\nu(\nu z).$$

It is shown in [9] that

$$(4.11) \quad e^{-\nu z} I_\nu(\nu z) = \left(\frac{t}{2\pi\nu}\right)^{\frac{1}{2}} e^{-\nu(z-\xi(z))} \frac{\sum_{s=0}^m \frac{\mathcal{U}_s(t)}{\nu^s} + \epsilon_m(\nu, t)}{1 + \epsilon_m(\nu, 0)},$$

where $t = (1+z^2)^{-1/2}$, $m \geq 0$ is an integer, and the $\mathcal{U}_s(t)$ are certain polynomials of degree $3s$ in t (see [9]), the first two of which are given by

$$(4.12) \quad \mathcal{U}_0(t) = 1, \quad \mathcal{U}_1(t) = (3t - 5t^3)/24.$$

The ϵ_m in (4.10) are estimated by ([9])

$$(4.13) \quad |\epsilon_m(\nu, t)| \leq \frac{\nu}{(\nu - \nu_0)} \frac{\mathcal{V}_t^1(\mathcal{U}_{m+1})}{\nu^{m+1}},$$

where

$$\mathcal{V}_a^b(\mathcal{U}) = \int_a^b |\mathcal{U}'(t)| dt \text{ and } \nu_0 = \mathcal{V}_0^1(\mathcal{U}_1) = \frac{1}{6\sqrt{5}} + \frac{1}{12}.$$

Simplifying (4.9) and using (4.10)–(4.13) with $m = 0$ yields, with $\phi(n, d)$ as in (4.7), that

$$a_n = \phi(n, d) \frac{1 + \epsilon_0(\tilde{n}, t)}{1 + \epsilon_0(\tilde{n}, 0)}.$$

The assertion then follows since

$$0 < \frac{1 + \epsilon_0(\tilde{n}, t)}{1 + \epsilon_0(\tilde{n}, 0)} \leq \left(1 - \frac{2\nu_0}{\tilde{n}}\right)^{-1}. \quad \square$$

Remark 4.2. The bounds (4.6) are quite sharp, since, for example, for $n \geq 1$ we obtain that

$$0.7895 \leq 1 - \frac{2\nu_0}{\tilde{n}}, \quad \left(1 - \frac{2\nu_0}{\tilde{n}}\right)^{-1} \leq 1.2667.$$

Lemma 4.2 reduces the description of the asymptotic behavior of $a_n(d)$ to a discussion of the function $\phi(n, d)$. We then obtain the following bounds on the approximation errors (4.3), (4.4).

Lemma 4.3. *We have*

$$(4.14) \quad \|u' - \chi'\|_0^2 \leq \sum_{n=p}^{\infty} \theta^+(n, d) e^{-2\tilde{n}(z-\xi(z))},$$

$$(4.15) \quad \|u - \chi\|_0^2 \leq \sum_{n=p}^{\infty} \frac{1}{n(n+1)} \theta^+(n, d) e^{-2\tilde{n}(z-\xi(z))},$$

$$(4.16) \quad \|u' - \chi'\|_0^2 \geq \sum_{n=p}^{\infty} \theta^-(n, d) e^{-2\tilde{n}(z-\xi(z))},$$

where $z = (\tilde{n}d)^{-1}$, $\xi(z)$ is as in (4.8) and

$$(4.17) \quad \theta^\pm(n, d) := \left(1 - \frac{2\nu_0}{\tilde{n}}\right)^{\mp 2} (d^2 + \tilde{n}^{-2})^{-\frac{1}{2}}.$$

As is readily apparent from the expression for $\phi(n, d)$ in (4.7), we can expect exponential decay of a_n as $n \rightarrow \infty$ provided the function $z - \xi(z)$ is positive and of reasonable size. The following lemma provides bounds for $z - \xi(z)$ in terms of the asymptotes shown in Figure 1. The proof follows by elementary arguments.

Lemma 4.4. *For any $z > 0$, we have $z - \xi(z) \geq 0$. Moreover, the following bounds hold:*

$$(4.18) \quad -(1 + \ln(z/2)) \leq z - \xi(z) \leq z - (1 + \ln(z/2)),$$

$$(4.19) \quad \frac{1}{2z} - \frac{1}{24z^3} \leq z - \xi(z) \leq \frac{1}{2z}.$$




FIGURE 1. The function $z - \xi(z)$ (---) and its asymptotes $1/(2z) - 1/(24z^3)$ (—) and $-(1 + \ln(z/2))$ (···)

We prove now an error bound for sufficiently large p ($\tilde{p} > e/2d$).

Theorem 4.1. *Let $r := e/(2\tilde{p}d) < 1$. Then for $u \equiv u_{1,d}$ there exists a polynomial $\chi \in \Pi_p(I)$ such that $\chi(\pm 1) = u(\pm 1)$ and*

$$(4.20) \quad \|u' - \chi'\|_0 \leq C_1 d^{-1/2} r^{\tilde{p}} (1 - r^2)^{-1/2},$$

$$(4.21) \quad \|u - \chi\|_0 \leq C_0 d^{1/2} r^{\tilde{p}} (1 - r^2)^{-1/2}.$$

Here, C_i are independent of p and d (numerical values can be read off the proof).

Proof. By (4.14), (4.17), we must estimate the sum

$$(4.22) \quad \begin{aligned} S^+ &= \sum_{n=p}^{\infty} \left(1 - \frac{2\nu_0}{\tilde{n}}\right)^{-2} (d^2 + \tilde{n}^{-2})^{-\frac{1}{2}} e^{-2\tilde{n}(z-\xi(z))} \\ &\leq \left(1 - \frac{4\nu_0}{3}\right)^{-2} \sum_{n=p}^{\infty} d^{-1} e^{-2\tilde{n}(z-\xi(z))}. \end{aligned}$$

Using the lower bound in (4.18), we get

$$(4.23) \quad \begin{aligned} S^+ &\leq (1 - 4\nu_0/3)^{-2} d^{-1} \sum_{n=p}^{\infty} \left(\frac{e}{2\tilde{n}d}\right)^{2\tilde{n}} \\ &= C_1^2 d^{-1} \sum_{n=p}^{\infty} r^{2\tilde{n}} = C_1^2 d^{-1} r^{2\tilde{p}} (1 - r^2)^{-1}. \end{aligned}$$

This is (4.20). To prove (4.21), we observe that $r < 1$ implies that

$$\frac{1}{p(p+1)} \leq \frac{9}{8\tilde{p}^2} \leq \frac{9}{8} \left(\frac{4d^2}{e^2} \right) = \frac{9d^2}{2e^2}.$$

Hence,

$$\frac{1}{p(p+1)} S^+ \leq C_0^2 d r^{2\tilde{p}} (1-r^2)^{-1}, \quad C_0^2 = \frac{9}{2e^2} C_1^2,$$

and (4.21) follows. \square

Corollary 4.1. *Let $\Phi(d, S)$ be as in (3.12). Then for $r = e/(2\tilde{p}d) < 1$,*

$$(4.24) \quad \Phi(d, S) \leq C_2 d^{1/2} r^{\tilde{p}} (1-r^2)^{-1/2}, \quad C_2 = (C_0^2 + C_1^2)^{1/2}.$$

Remark 4.3. The asymptotic rate of convergence with respect to p in (4.20) and hence (4.24) is optimal up to a constant depending on d , since by (4.16),

$$\|u' - \chi'\|_0^2 \geq \sum_{n=p}^{\infty} \theta^-(n, d) e^{-2\tilde{n}(z-\xi(z))} =: S^-.$$

Using the upper bound in (4.18) yields

$$\begin{aligned} S^- &\geq \left(1 - \frac{2\nu_0}{\tilde{p}}\right)^2 \sum_{n=p}^{\infty} (d^2 + \tilde{n}^{-2})^{-\frac{1}{2}} e^{-2/d} r^{2\tilde{n}} \\ &\geq \left(1 - \frac{4\nu_0}{3}\right)^2 e^{-2/d} d^{-1} \sum_{n=p}^{\infty} (1+z^2)^{-1/2} r^{2\tilde{n}} \\ &\geq \left(1 - \frac{4\nu_0}{3}\right)^2 (1+4/e^2)^{-\frac{1}{2}} e^{-2/d} d^{-1} r^{2\tilde{p}} (1-r^2)^{-1}, \end{aligned}$$

since $z = (\tilde{n}d)^{-1} \leq (\tilde{p}d)^{-1} < 2/e$. Hence,

$$\|u' - \chi'\|_0 \geq C_3 e^{-1/d} d^{-1/2} r^{\tilde{p}} (1-r^2)^{-1/2}.$$

Now χ is the same polynomial as in Lemma 4.1, so that (4.5) holds. Comparing the above estimate with (4.20), we see that (4.20), (4.24) are optimal in p for any fixed $d > 0$ as $p \rightarrow \infty$.

The estimates in Theorem 4.1 are useful for the case that \tilde{p} is large compared to $e/(2d)$. Such a situation arises in the next section, where this theorem will be applied. In actual practice, if d is small, then it can be difficult to ever be in this asymptotic range of \tilde{p} . The computational results in §6 show that convergence is observed in the preasymptotic range $\tilde{p} \leq e/(2d)$ as well. Therefore, we now obtain estimates for the rate of decrease of the error in the range $\sqrt{3/(4d)} \leq \tilde{p} \leq e/(2d)$.

In Theorem 4.1, we used the bounds (4.18) as well as (4.19), the latter being sharper for the range $\tilde{p} \leq e/(2d)$, i.e., $z \geq 2/e$ (see Figure 1). It is seen that the two lower bounds for $z - \xi(z)$ in Lemma 4.4 intersect at the root of

$$\frac{1}{2z} - \frac{1}{24z^3} + 1 + \ln\left(\frac{z}{2}\right) = 0,$$

i.e., at $z^* = 0.51388\dots$, which is close to 0.5. Our estimate will therefore be valid for $z \geq 0.5$, i.e., in the extended range $\sqrt{3/(4d)} \leq \tilde{p} \leq 2/d$.

Theorem 4.2. *Assume that $\sqrt{3/(4d)} \leq \tilde{p} \leq 2/d$. Then, for $u \equiv u_{1,d}$, there exists a polynomial $\chi \in \Pi_p(I)$ such that $\chi(\pm 1) = u(\pm 1)$ and*

$$(4.25) \quad \|u' - \chi'\|_0^2 \leq C_1^2 \left(\tilde{p} + \frac{3}{4d} \right) \exp(-2\tilde{p}^2 d/3) + C_3 d^{-1} (e/4)^{4/d},$$

$$(4.26) \quad \|u - \chi\|_0^2 \leq \frac{9}{8} C_1^2 \tilde{p}^{-2} \left(\tilde{p} + \frac{3}{4d} \right) \exp(-2\tilde{p}^2 d/3) + \frac{4}{15} C_3 d (e/4)^{4/d}.$$

Here the constants C_i are independent of p and d (and are given in the proof below).

Proof. Let us define the index sets

$$I_1(d) = \{n \in \mathbb{N} : \tilde{p} \leq \tilde{n} < 2/d\}, \quad I_2(d) = \{n \in \mathbb{N} : \tilde{n} \geq 2/d\}.$$

Then, taking χ to be the polynomial in Lemma 4.1, we have by (4.14)

$$\|u' - \chi'\|_0^2 \leq S_1 + S_2, \quad S_i = \sum_{n \in I_i} \theta^+(n, d) e^{-2\tilde{n}(z-\xi(z))}.$$

The quantities S_1 and S_2 will be estimated using the lower bounds in (4.19) and (4.18), respectively. First, by (4.19),

$$(4.27) \quad S_1 \leq \sum_{n \in I_1(d)} \theta^+(n, d) \exp\left(-2\tilde{n} \left(\frac{1}{2z} - \frac{1}{24z^3}\right)\right),$$

where, since $n \in I_1(d)$, we have $0 < z^{-1} = \tilde{n}d < 2$, so that $0 < z^{-3} < 4z^{-1}$. Hence,

$$\frac{1}{2z} - \frac{1}{24z^3} \geq \frac{1}{2z} - \frac{1}{6z} = \frac{1}{3z},$$

and (4.27), (4.17) give (with $C_1 = (1 - 4\nu_0/3)^{-1}$ as in (4.23))

$$S_1 \leq C_1 \sum_{n \in I_1(d)} \tilde{n}(1 + \tilde{n}^2 d^2)^{-1/2} e^{-\frac{2}{3}\tilde{n}^2 d}.$$

Now the function $x e^{-\frac{2}{3}x^2 d}$ attains its global maximum at $x = \sqrt{3/(4d)}$ and is decreasing for $x > \sqrt{3/(4d)}$. Hence,

$$\begin{aligned} S_1 &\leq C_1 \sum_{n \in I_1(d)} \tilde{n} e^{-\frac{2}{3}\tilde{n}^2 d} \leq C_1 \left(\tilde{p} e^{-\frac{2}{3}\tilde{p}^2 d} + \int_{\tilde{p}}^{2/d} x e^{-\frac{2}{3}x^2 d} dx \right) \\ &\leq C_1 \left(\tilde{p} + \frac{3}{4d} \right) e^{-\frac{2}{3}\tilde{p}^2 d}. \end{aligned}$$

For the term S_2 , we use (4.18). Noting that $\tilde{n}^{-1} \leq d/2$, we have

$$S_2 \leq \sum_{\tilde{n} \geq 2/d} \theta^+(n, d) \left(\frac{e}{2\tilde{n}d} \right)^{2\tilde{n}} \leq \tilde{C}_3 d^{-1} \sum_{\tilde{n} \geq 2/d} \left(\frac{e}{4} \right)^{2\tilde{n}},$$

where $\tilde{C}_3 = (1 - \nu_0)^{-2} \left(\frac{4}{5}\right)^{1/2}$. Summing the geometric series leads to the last term in (4.25), where $C_3 = \tilde{C}_3(1 - e^2/16)^{-1}$.

For the L_2 estimates, (4.15) gives

$$\|u - \chi\|_0^2 \leq \tilde{S}_1 + \tilde{S}_2, \quad \tilde{S}_i = \sum_{n \in I_i(d)} \frac{1}{n(n+1)} \theta^+(n, d) e^{-2\tilde{n}(z-\xi(z))}, \quad i = 1, 2.$$

It is easy to see that

$$\tilde{S}_1 \leq \frac{1}{p(p+1)} S_1 \leq \frac{9}{8\tilde{p}^2} S_1.$$

Also, for $n \in I_2(d)$, we have $n(n+1) \geq \frac{16-d^2}{4d^2} \geq \frac{15}{4d^2}$ for $d \leq 1$, so that

$$\tilde{S}_2 \leq \frac{4d^2}{15} S_2.$$

The theorem follows. □

Theorem 4.2 leads to the following corollary.

Corollary 4.2. For $\sqrt{3/(4d)} \leq \tilde{p} \leq 2/d$,

$$(4.28) \quad \Phi(d, S) \leq C_4 e^{-\frac{\tilde{p}^2 d}{3}} + C_5 \left(\frac{e}{4}\right)^{\frac{2}{d}},$$

where the constants C_i are independent of p, d .

Proof. Using the definition of $\Phi(d, S)$, we obtain from Theorem 4.2,

$$(\Phi(d, S))^2 \leq C_1^2 \left(\tilde{p} + \frac{3}{4d}\right) \left(d^2 + \frac{9}{8}\tilde{p}^{-2}\right) e^{-\frac{2}{3}\tilde{p}^2 d} + \frac{19}{15} C_3 \left(\frac{e}{4}\right)^{\frac{4}{d}}.$$

Hence, (4.28) holds with $C_5^2 = \frac{19}{15} C_3$ and

$$C_1^2 \left(\tilde{p}d^2 + \frac{3}{4}d + \frac{9}{8}\tilde{p}^{-1} + \frac{27}{32}\frac{\tilde{p}^{-2}}{d}\right) \leq C_1^2 \left(4\tilde{p}^{-1} + \frac{3}{2}\tilde{p}^{-1} + \frac{9}{8}\tilde{p}^{-1} + \frac{9}{8}\right) \leq C_4^2.$$

□

We see from Corollary 4.2 that for small d , since the term $C_5(e/4)^{2/d}$ is negligible, the first term $C_4 \exp(-(\tilde{p}^2 d)/3)$ in (4.28) will dominate. Hence, the error will decrease at an exponential rate in this range when $\tilde{p}^2 d/3$ is large enough. For $\tilde{p} > e/(2d)$, a better estimate may be provided by Theorem 4.1. When $\tilde{p}^2 d/3$ is small (i.e., $\tilde{p} = Kd^{-\frac{1}{2}}$), the estimate (4.28) deteriorates. We will therefore establish another bound, which is valid in the range $1 < \tilde{p} \leq Kd^{-\frac{1}{2}}$. First, we prove the following lemma.

Lemma 4.5. *There exists a unique polynomial $\chi \in \Pi_p(I)$ that minimizes $\|\chi\|_0$ subject to the constraints $\chi(\pm 1) = \alpha^\pm$. This polynomial χ satisfies*

$$(4.29) \quad \frac{1}{C} \frac{\max(|\alpha^+|, |\alpha^-|)}{p} \leq \|\chi\|_0 \leq C \frac{\max(|\alpha^+|, |\alpha^-|)}{p},$$

$$(4.30) \quad \|\chi\|_1 \leq C \max(|\alpha^+|, |\alpha^-|) p,$$

with $C > 1$ a constant independent of α^\pm, p .

Proof. We may write χ in the Legendre series expansion satisfying the end constraints,

$$\chi(x) = \sum_{k=0}^p a_k P_k(x), \quad \sum_{k=0}^p a_k (\pm 1)^k = \alpha^\pm.$$

Introducing Lagrange multipliers for the constraints, we get the minimization problem

$$\min_{\vec{a}, \lambda_+, \lambda_-} F(\vec{a}, \lambda_+, \lambda_-) = \sum_{k=0}^p \omega_k a_k^2 + \lambda_+ \left(\sum_{k=0}^p a_k - \alpha^+ \right) + \lambda_- \left(\sum_{k=0}^p a_k (-1)^k - \alpha^- \right),$$

where $\omega_k = 2/(2k + 1)$. Let $A = p(p + 1)^2(p + 2)/4$. Then it may be shown that the unique minimizer for the above is given by

$$\lambda_{\pm} = \frac{2}{A} \left(\alpha^{\mp} (-1)^p \frac{(p + 1)}{2} - \alpha^{\pm} \frac{(p + 1)^2}{2} \right),$$

$$a_k = (\alpha^+ + (-1)^k \alpha^-) \frac{((p + 1)^2 - (-1)^{p+k} (p + 1))}{2A\omega_k},$$

from which the bounds for $\|\chi\|_0$ in (4.29) follow easily. The bound for $\|\chi\|_1$ follows by the inverse inequality for polynomials,

$$|\chi|_1 \leq Cp^{-2} \|\chi\|_0. \quad \square$$

Theorem 4.3. *Assume that $1 < \tilde{p} \leq Kd^{-\frac{1}{2}}$ for some K (which may depend upon \tilde{p}, d). Then for such p, d ,*

$$(4.31) \quad \Phi(d, S) \leq CKp^{-1},$$

where the constant C is independent of K, p and d .

Proof. We note that for any $\chi \in \Pi_p(I)$ satisfying $\chi(\pm 1) = u_{1,d}(\pm 1)$,

$$\Phi(d, S) \leq C(\|u_{1,d}\|_d + d|\chi|_1 + \|\chi\|_0).$$

We use (2.9) to bound $\|u_{1,d}\|_d$ and choose χ as in Lemma 4.5, with $\alpha^{\pm} = u_{1,d}(\pm 1)$. Then we obtain by (4.29), (4.30),

$$(4.32) \quad \Phi(d, S) \leq C(d^{\frac{1}{2}} + dp + p^{-1}).$$

Now since $\tilde{p} \leq Kd^{-\frac{1}{2}}$, we have $d^{\frac{1}{2}} \leq K\tilde{p}^{-1}$. Substituting this in (4.32) gives (4.31). \square

Let us now put together the results of Theorems 4.1–4.3. The following theorem shows that the spaces $S(\Sigma) = \Pi_p(I)$ approximate boundary layers $u_{1,d}$ robustly at the rate $G(p) = Cp^{-1}\sqrt{\ln p}$ in the energy norm (in the sense of Definition 3.2). Moreover, the best robust rate possible is Cp^{-1} , so that the result established is optimal up to a factor $\sqrt{\ln p}$.

Theorem 4.4. *Let $S(\Sigma) = \Pi_p(I)$. Then*

$$Cp^{-1} \leq G(p) = \sup_{d \in (0,1]} \Phi(d, S) \leq Cp^{-1}\sqrt{\ln p},$$

where C is a constant independent of p .

Proof. Let $d \in (0, 1]$ be arbitrary. Suppose first that

$$1 \leq \frac{\tilde{p}}{2\sqrt{\ln p}} \leq \sqrt{\frac{3}{4d}}, \quad \text{i.e.,} \quad 1 \leq \tilde{p} \leq \sqrt{\frac{3 \ln p}{d}}.$$

Then by Theorem 4.3, with $K = \sqrt{3 \ln p}$, we have in this range

$$\Phi(d, S) \leq CKp^{-1} \leq Cp^{-1}\sqrt{\ln p}.$$

Next, for $\sqrt{3/(4d)} \leq \tilde{p}/(2\sqrt{\ln p}) \leq 2/(2d\sqrt{\ln p})$ we may use Corollary 4.2, by which

$$\Phi(d, S) \leq C_4 e^{-\tilde{p}^2 d/3} + C_5 \left(\frac{e}{4}\right)^{\frac{2}{d}}.$$

Since $\tilde{p}/(2\sqrt{\ln p}) \geq \sqrt{3/(4d)}$, we have $\tilde{p}^2 d/3 \geq \ln p$, so that $\exp(-\tilde{p}^2 d/3) \leq 1/p$. Also, since $2/d \geq \tilde{p}$, it follows that $(e/4)^{2/d} \leq (e/4)^{\tilde{p}} \leq p^{-1}$. Hence,

$$(4.33) \quad \Phi(d, S) \leq Cp^{-1}$$

in this range. Finally, it is easy to see that the estimate for $\Phi(d, S)$ for the range $\tilde{p} > 2/d$, given by Theorem 4.1, also satisfies (4.33).

To establish the lower bound, we note that by the triangle inequality, for any $\chi \in \Phi_p(I)$ with $\chi(\pm 1) = u(\pm 1)$,

$$\begin{aligned} \Phi(d, S) &\geq \|\chi\|_d - \|u\|_d \\ &\geq \|\chi\|_0 - \|u\|_d. \end{aligned}$$

If $d \rightarrow 0$, then $\|u\|_d \rightarrow 0$. But, by Lemma 4.5, $\|\chi\|_0 \geq Cp^{-1}$, giving the result. \square

Suppose the p version is used with a *fixed* mesh for problems (3.1). Then the rate $F(N, k)$ in (3.11) satisfies

$$F(N, k) \leq Cp^{-(k-1)},$$

so that the first infimum on the right side of (3.10) will certainly be less than Cp^{-1} , uniformly in d whenever $k \geq 2$. Using Theorem 4.4 on the whole interval $[-1, 1]$, we can bound the infima involving boundary layers uniformly by $Cp^{-1}\sqrt{\ln p}$ (the fact that we have more than one interval can only enhance this rate). Hence, the p version is robust with uniform order $Cp^{-1}\sqrt{\ln p}$. Moreover, this robust rate is optimal up to $\sqrt{\ln p}$, since for $d \rightarrow 0$, the approximation of the boundary layer terms in the end intervals cannot be better than Cp^{-1} by Theorem 4.4. We therefore have the following result.

Theorem 4.5. *The p version with fixed mesh for problems (3.1), $0 < d \leq 1$, is robust with uniform order $g(p)$ satisfying*

$$Cp^{-1} \leq g(p) \leq Cp^{-1}\sqrt{\ln p}$$

with respect to solution sets $H_{d,M}^B$ (or H_{d,Π_n}^B) and error measure the energy norm.

Remark 4.4. In terms of the number of degrees of freedom N , we see that $g(N) \approx N^{-1}\sqrt{\ln N}$. This is essentially *twice* the best uniform rate of $N^{-1/2}$ that can be attained using the h version with a quasiuniform mesh [13]. Hence, the “doubling” phenomenon for the rate of convergence for the p version, which is well known for the case that $(x + 1)^\alpha$ type singularities are present at $x = -1$ (see, e.g., [3]), also occurs when the solution contains boundary layer components of the type $\exp(-(x + 1)/d)$ at $x = -1$.

5. APPROXIMATION RESULTS FOR AN hp VERSION

In the previous section, we showed that the p version over a single element yields a super-exponential rate of convergence for $\tilde{p} > e/2d$. Also, the error decreases at the (exponential) rate $\exp(-\tilde{p}^2 d/3)$ in the preasymptotic range $\sqrt{3/(4d)} \leq \tilde{p} \leq 2/d$ for small d . Unfortunately, in practice both these ranges may be difficult to achieve if d is small and p is restricted ($p \leq 8$ is typical in programs such as MSC/PROBE

and STRESSCHECK), so that all that may be observed is the uniform rate of $\mathcal{O}(p^{-1}\sqrt{\ln p})$ predicted by Theorems 4.3, 4.4. In this section, we show that if only one extra element of size $\mathcal{O}(pd)$ is inserted in the boundary layer, then *robust exponential convergence* is achieved uniformly for $0 < d \leq 1$ as p increases. Since the mesh is changed at each step when p is increased, we call this an *hp* version FEM (more appropriately, it is an *rp* version FEM). Naturally, if the polynomial degree p is sufficiently large, we have to allow a transition to the single-element mesh analyzed in Theorem 4.1.

A more general question that could be considered is, given N degrees of freedom (N as in (3.6)), for what mesh-degree combination Σ (i.e., choice of S_D) is the error minimized? We do not consider this theoretical question here, since the simple two-element mesh below already gives exponential convergence, uniformly in d . This mesh is easier to implement than a general *hp* version, and moreover, in computational experiments performed using meshes with more elements, we were unable to achieve better convergence rates (see §6 and [20]). Note that the mesh-degree combination we propose is similar to the *optimal* mesh-degree combination obtained for a related problem by Scherer in [14] (see Remark 5.2).

The following theorem is our main result in this section.

Theorem 5.1. *Let $I = (-1, 1)$ and $u(x) = u_{1,d} = \exp(-(x + 1)/d)$. Let further $\Sigma = (\Delta, \tilde{p})$ be such that for $p \geq 1$*

$$\begin{aligned} \tilde{p} &= \{p, 1\}, \quad \Delta = \{-1, -1 + \kappa\tilde{p}d, 1\} \quad \text{if } \kappa\tilde{p}d < 2, \\ \tilde{p} &= \{p\}, \quad \Delta = \{-1, 1\} \quad \text{if } \kappa\tilde{p}d \geq 2, \end{aligned}$$

where $0 < \kappa_0 \leq \kappa < 4/e$ is a constant independent of p and d . Then there exists $u_p \in S(\Sigma)$ such that $u_p(\pm 1) = u(\pm 1)$ and

$$(5.1) \quad \|u - u_p\|_d \leq d^{1/2}C_6 \alpha^{\tilde{p}}, \quad \|u - u_p\|_0 \leq d^{1/2}C_7 \alpha^{\tilde{p}}, \quad \|u' - u'_p\|_0 \leq d^{-1/2}C_8 \alpha^{\tilde{p}}.$$

Here the constants are independent of p and d but depend on κ_0 and

$$(5.2) \quad \alpha := \left\{ \begin{array}{ll} e/(2\tilde{p}d) & \text{if } \kappa\tilde{p}d \geq 2, \\ \max\{\kappa e/4, e^{-(\kappa-\epsilon)}\} & \text{if } \kappa\tilde{p}d < 2 \end{array} \right\} < 1,$$

with $\epsilon > \ln p/(2p)$ arbitrary.

Proof. If $\kappa\tilde{p}d \geq 2$, we have that $r = e/(2\tilde{p}d) < 1$, owing to our assumption that $\kappa < 4/e$. Therefore, Theorem 4.1 is applicable, and a p -increase in the single-element mesh $\Delta = \{-1, 1\}$ yields exponential convergence with the number r decreasing with p .

Consider now the case $\kappa\tilde{p}d < 2$, i.e., the two-element mesh $\Delta = \{-1, -1 + \kappa\tilde{p}d, 1\}$. We assume first that $\tilde{p} \geq 2/\kappa_0$ and construct the function $u_p(x) \in S(\Sigma)$ elementwise. Denote $I_1 = (-1, a)$, where $a = -1 + \kappa\tilde{p}d$, $\kappa_0 \leq \kappa < 4/e$, and let $s_1 \in \Pi_p(I_1)$. Transforming I_1 to $I = (-1, 1)$, we see that for $t = 0, 1$,

$$\int_{-1}^a \left(\frac{d^t}{dx^t} (u - s_1) \right)^2 dx = \left(\frac{2}{\kappa\tilde{p}d} \right)^{2t-1} \int_{-1}^1 \left(\frac{d^t}{dy^t} (\tilde{u} - \tilde{s}_1) \right)^2 dy.$$

Here, $\tilde{f}(y)$ denotes the image on I of any function $f(x)$ defined on I_1 . Consequently, we obtain that $\tilde{u}(y) = \exp(-(y + 1)\kappa\tilde{p}/2) = u_{1,\tilde{d}}(y)$, where $\tilde{d} = 2/\kappa\tilde{p}$.

Since $\kappa < 4/e$, we have $r := e/(2\tilde{p}\tilde{d}) = \kappa e/4 < 1$. Now Theorem 4.1 and Corollary 4.1 apply uniformly to functions $u \equiv u_{1,d}$ for all $d \in (0, 1]$. Since $\tilde{p} \geq 2/\kappa_0 \geq 2/\kappa$ we have that $\tilde{d} < 1$, and hence Theorem 4.1 and Corollary 4.1 will apply when d is chosen to be \tilde{d} . Then, since $r < 1$, we obtain a polynomial $s_p \in \Pi_p(I_1)$ satisfying

$$(5.3) \quad s_p(-1) = u(-1), \quad s_p(a) = u(a),$$

and

$$(5.4) \quad \left\| \frac{d^t}{dx^t} (u - s_p) \right\|_{0,I_1}^2 \leq C_t^2 \left(\frac{2}{\kappa\tilde{p}\tilde{d}} \right)^{2t-1} \tilde{d}^{1-2t} \frac{r^{2\tilde{p}}}{(1-r^2)}, \quad t = 0, 1,$$

$$(5.5) \quad (d^2 \|u' - s'_p\|_{0,I_1}^2 + \|u - s_p\|_{0,I_1}^2)^{1/2} \leq C_2 d^{1/2} \frac{r^{\tilde{p}}}{(1-r^2)^{1/2}}.$$

This gives the asserted bound on I_1 in the case $\tilde{p} > 2/\kappa_0$. Since this excludes only finitely many values of p , these estimates hold for all p after possibly adjusting the constants C_t , $t = 0, 1, 2$ in (5.4), (5.5).

As noted in Remark 4.1, the approximation $s_p(x)$ constructed via Lemma 4.1 is optimal in the $|\cdot|_1$ seminorm but not in the $\|\cdot\|_d$ norm. For fixed $d > 0$, s_p yields the optimal-order error as $p \rightarrow \infty$, but is suboptimal as $d \rightarrow 0$, owing to the enforcement of the interpolation condition (5.3). Therefore, we modify s_p as follows: let $u_p = s_p - s_1 + \tilde{s}_1$, where s_1 is the linear interpolant of $u_{1,d}(x)$ at $x = -1$ and $x = a$, and \tilde{s}_1 is a linear function such that $\tilde{s}_1(-1) = u(-1)$ and $\tilde{s}_1(a) = \max\{d^{1/2}u(a), u(1)\}$. Then

$$(5.6) \quad \begin{aligned} \|u - u_p\|_{d,I_1} &= \|u - (s_p - s_1 + \tilde{s}_1)\|_{d,I_1} \\ &\leq \|u - s_p\|_{d,I_1} + \|s_1 - \tilde{s}_1\|_{d,I_1}. \end{aligned}$$

The first term was estimated in (5.5), so we estimate the second term. We have

$$\begin{aligned} \int_{-1}^a (s_1 - \tilde{s}_1)^2 dx &\leq \max_{-1 \leq x \leq a} |(s_1 - \tilde{s}_1)(x)|^2 (1+a) \\ &\leq |(1 - \sqrt{d})u(a)|^2 (1+a). \end{aligned}$$

Since $1 + a = \kappa\tilde{p}d$ and $u(a) = \exp(-(a + 1)/d) = \exp(-\kappa\tilde{p})$, we get

$$\|s_1 - \tilde{s}_1\|_{0,I_1}^2 \leq e^{-2\kappa\tilde{p}} \kappa\tilde{p}d.$$

Also,

$$\int_{-1}^a (s'_1 - \tilde{s}'_1)^2 dx \leq (1 - \sqrt{d})^2 |u(a)|^2 (1+a)^{-1} \leq \frac{e^{-2\kappa\tilde{p}}}{\kappa\tilde{p}d}.$$

Hence,

$$d \|s'_1 - \tilde{s}'_1\|_{0,I_1} \leq \frac{d^{1/2}}{\sqrt{\kappa\tilde{p}}} e^{-\kappa\tilde{p}}$$

and altogether

$$(5.7) \quad \begin{aligned} \|s_1 - \tilde{s}_1\|_d &\leq d^{\frac{1}{2}} \left(\kappa\tilde{p} + \frac{1}{\kappa\tilde{p}} \right)^{\frac{1}{2}} e^{-\kappa\tilde{p}} \\ &\leq d^{\frac{1}{2}} \left(\kappa + \frac{1}{\kappa} \right)^{\frac{1}{2}} e^{-(\kappa-\epsilon)\tilde{p}} \end{aligned}$$

for any $\epsilon > \frac{\ln \tilde{p}}{2\tilde{p}}$ ($\epsilon = \frac{1}{2e}$ works for all p). Then, from (5.5)–(5.7),

$$(5.8) \quad \|u - u_p\|_{d,I_1}^2 \leq \tilde{C}_0^2 d \left\{ \left(\frac{\kappa e}{4}\right)^{2\tilde{p}} + e^{-2(\kappa-\epsilon)\tilde{p}} \right\}.$$

Next we consider I_2 . Here we select $u_p \in \Pi_1(I_2)$ to be the linear interpolant between $\max\{d^{1/2}u(a), u(1)\}$ at $x = a$ and $u(1)$ at $x = 1$. One verifies that

$$(5.9) \quad u_p(x) = (u(1) - \max\{u(1), \sqrt{d} u(a)\}) \frac{(x-a)}{1-a} + \max\{u(1), \sqrt{d} u(a)\}.$$

Now let $\frac{2}{\kappa d} \geq \tilde{p} \geq \frac{2}{\kappa d} - \frac{|\ln d|}{2\kappa}$. Then, since $\frac{|\ln d|}{2\kappa} \leq \frac{e^{-1}}{2\kappa d}$, we have $\tilde{p} \geq \frac{(4 - e^{-1})}{2\kappa d}$ in this range. Also,

$$u(1) \geq \sqrt{d} u(a) \text{ and } 1 - a \leq \frac{|\ln d|d}{2} \leq \ln \left(\frac{2\tilde{p}\kappa}{(4 - e^{-1})} \right) \frac{d}{2}.$$

Hence,

$$(5.10) \quad \int_a^1 u_p^2 dx \leq u^2(1) \frac{|\ln d|d}{2} \leq \frac{d}{2} \ln \left(\frac{2\tilde{p}\kappa}{(4 - e^{-1})} \right) e^{-2\kappa\tilde{p}} \leq \tilde{C}_1 d e^{-2(\kappa-\epsilon)\tilde{p}},$$

$$\int_a^1 (u'_p)^2 = 0.$$

Next, for $\tilde{p} < \frac{2}{\kappa d} - \frac{|\ln d|}{2\kappa}$,

$$u(1) < \sqrt{d} u(a) \quad \text{and} \quad \frac{1}{1-a} \leq \frac{2}{d|\ln d|}.$$

Hence,

$$(5.11) \quad \int_a^1 u_p^2 dx \leq (\sqrt{d} u(a))^2 (1-a) \leq 2d e^{-2\kappa\tilde{p}},$$

$$\int_a^1 (u'_p)^2 dx \leq \frac{2d u^2(a)}{|\ln d| d} \leq \frac{2e^{-2\kappa\tilde{p}}}{|\ln d|}.$$

For $d \leq e^{-1}$, this gives

$$(5.12) \quad d^2 \int_a^1 (u'_p)^2 dx \leq 2d^2 e^{-2\kappa\tilde{p}}.$$

For $e^{-1} < d \leq 1$, we have by (5.9) that

$$\int_a^1 (u'_p)^2 dx \leq 2 \left(\frac{(u(1) - u(a))^2}{1-a} + \frac{(1 - \sqrt{d})^2 u^2(a)}{1-a} \right).$$

By the Mean Value Theorem, there exists $\xi \in [a, 1]$ such that $u(1) - u(a) = u'(\xi)(1-a)$, so that

$$(5.13) \quad d^2 \int_a^1 (u'_p)^2 dx \leq 2e^{-2(\frac{\xi+1}{d})} (1-a) + 4d \frac{(1 - \sqrt{d})^2 u^2(a)}{|\ln d|}$$

$$\leq \tilde{C}_2 d e^{-2\kappa\tilde{p}}$$

uniformly as $d \rightarrow 1$, where \tilde{C}_2 may be explicitly evaluated. Hence, we conclude by (5.10)–(5.13) that

$$(5.14) \quad \|u_p\|_{d,I_2}^2 \leq \tilde{C}_3 d e^{-2(\kappa-\epsilon)\bar{p}}.$$

Also, it is easy to verify that

$$(5.15) \quad \|u\|_{0,I_2}^2 \leq \frac{d}{2} e^{-2\kappa\bar{p}}, \quad \|u'\|_{0,I_2}^2 \leq \frac{2}{d} e^{-2\kappa\bar{p}},$$

so that by (5.14), (5.15) and the triangle inequality,

$$(5.16) \quad \|u - u_p\|_{d,I_2}^2 \leq \tilde{C}_4 d e^{-2(\kappa-\epsilon)\bar{p}}.$$

Then the first inequality in (5.1) follows from Theorem 4.1, (5.8), (5.16). The other two inequalities also follow from the estimates above. \square

Remark 5.1. The constant κ in Theorem 5.1 could be selected such that $\kappa^* e = 4e^{-\kappa^*}$, which yields $\kappa^* \approx 0.71$. This gives $\alpha \approx e^{-\kappa^*}$ in (5.2) when two elements are being used. This value for κ^* is, however, not optimal since it is obtained by optimizing some upper bounds. The optimal choice of κ is numerically addressed in §6 ahead. Use of the above value of κ^* , however, simplifies the bounds above.

Remark 5.2. The choice of $\Sigma = (\Delta, \bar{p})$ used in Theorem 5.1 is similar to that obtained by Scherer in [14]. He considered the best mesh-degree combination (for a fixed number of degrees of freedom N) that would minimize the L^∞ error of best approximation (by discontinuous piecewise polynomials) of the function e^{-x} on the interval $[0, \infty)$. He was able to solve this problem explicitly — the asymptotically optimal Σ was given by $\Delta = \{0, q_0(p + 1), \infty\}$, $\bar{p} = \{p, 1\}$, where $p = N - 2$ and $q_0 = 0.89548641\dots$. For this Σ , Scherer showed that the asymptotic L^∞ convergence rate was $e^{-q_0 N} = e^{-q_0(p+2)}$, which (up to an algebraic factor in N) was, asymptotically, the best possible for any mesh-degree combination.

We can also deduce pointwise error bounds.

Corollary 5.1. *Under the assumptions of Theorem 5.1 we have*

$$(5.17) \quad \|u - u_p\|_{L^\infty(I)} \leq C_9 \alpha^{\bar{p}}$$

with α as in Theorem 5.1.

Proof. This follows from (5.1) and the interpolation inequality

$$\|v\|_{L^\infty(I)} \leq 2\|v\|_{L_2(I)}^{1/2} \|v'\|_{L_2(I)}^{1/2}. \quad \square$$

Remark 5.3. The estimates in Theorem 5.1, Corollary 5.1 are obtained using polynomials of degree 1 in I_2 . They evidently remain valid if I_2 is subdivided and/or the degree p is greater than 1 in I_2 .

Theorem 5.1 says that it is sufficient to use two intervals of the type described to resolve boundary layers with a robust exponential convergence rate. As discussed in §2, the solution will typically have other (smoother) components as well. For the approximation of these components, the mesh-degree combination of Theorem 5.1 will typically not be sufficient and will have to be enhanced (e.g., by subdivision or p -increase in element 2). This enhancement will ensure that the rate $F(N, k)$ in (3.11), which measures the approximation of these smoother components, is sufficiently rapid. For solutions in $H_{d,M}^B$, the robust rate of convergence $g(N)$ of the hp version will then be given by (3.14), where $G(N)$ represents the exponential

rate (5.1). As noted in Remark 3.1, the overall rate will be exponential only if the smooth components are also approximated exponentially. One such case occurs when f is a polynomial, as noted in the theorem below. (Note that we have a boundary layer at each end-point now.)

Theorem 5.2. *Consider the hp version for problem (3.1), $0 < d \leq 1$, with $\Sigma = (\Delta, \bar{p})$ given by*

$$(5.18) \quad \begin{aligned} \bar{p} &= \{p, p, p\}, \quad \Delta = \{-1, -1 + \kappa\tilde{p}d, 1 - \kappa\tilde{p}d, 1\} \text{ if } \kappa\tilde{p}d < 1, \\ \bar{p} &= \{p, p\}, \quad \Delta = \{-1, 0, 1\} \text{ if } \kappa\tilde{p}d \geq 1, \end{aligned}$$

where κ is as in Theorem 5.1. Let $\alpha < 1$ be defined as in (5.2) (with the condition $\kappa\tilde{p}d < 2$ replaced by $\kappa\tilde{p}d < 1$). Then there exists a constant $C > 0$ independent of p and d such that with respect to solution sets H_{d, Π_n}^B and error measure the energy norm, this version is robust with uniform order $g(p) = Cd^{1/2}\alpha^{\bar{p}}$ for $p \geq n$.

Proof. The theorem follows easily by (3.10), Corollary 2.1 and Theorem 5.1. □

6. NUMERICAL RESULTS

In this section, we present the results of numerical computations for the model problem (1.2)–(1.3), where:

$$(6.1) \quad f(x) \equiv 1, \quad \alpha^+ = \alpha^- = 0, \quad a = 1.$$

The exact solution is then given by

$$(6.2) \quad u_d(x) = 1 - \frac{\cosh(x/d)}{\cosh(1/d)},$$

so that

$$(6.3) \quad \|u_d\|_d^2 = B_d(u_d, u_d) = (1, u_d) = 2 - 2d \tanh(1/d) = \mathcal{O}(1).$$

Note that since $f(x)$ is a polynomial of degree 0, Corollary 2.1 applies. Noting (6.3), we conclude that the relative error in the energy norm,

$$E_R(d) = \|u_d - u_d^S\|_d / \|u_d\|_d,$$

should behave like $\Phi(d, S)$ given by (3.12). All graphs shown in this section will depict $E_R(d)$ versus the number of degrees of freedom in the finite element method. The value of d (and, where applicable, of κ) will be stated with the figures. All computations were done in double precision on an SGI indigo2 workstation using MATLAB 4.1.

We first consider the p version over a single element. Figure 2 shows $E_R(d)$ plotted versus the number of degrees of freedom $N = p - 1$, for various values of d , in a semilog scale. By Corollary 4.1, for $\tilde{p} > e/2d$, the error will be in the asymptotic (superexponential) range. This is only reached, however, when $p \approx 13$ for $d = 0.1$, $p \approx 136$ for $d = 0.01$, $p \approx 1359$ for $d = 0.001$ and $p \approx 13,591$ for $d = 0.0001$. We see that, except for the first value, none of the rest can be considered as within a practical range of p . For $d = 0.1$, however, the graph in the semilog scale of Figure 2 is close to a straight line for p in this range, showing agreement with the theory.

Turning to the case $d = 0.01$, we note that Corollary 4.2 predicts for d small and $\sqrt{3/(4d)} \leq \tilde{p} \leq 2/d$, i.e., $5 \leq p \leq 200$, that $\log(E_R)$ should behave like $-\gamma \tilde{p}^2 d$, where $\gamma > 0$ is independent of d (the value of γ in Corollary 4.2 is $1/3$, but this may




FIGURE 2. The p version with one element

not be optimal). Hence, we should observe a parabolic curve for $d = 0.01$ when p is large enough. Again, the graph in Figure 2 is consistent with this bound.

As d becomes even smaller, the error in Figure 2 is seen to deteriorate further. For $1 \leq \tilde{p} \leq Kd^{-\frac{1}{2}}$, Theorem 4.3 predicts a convergence rate of only CKp^{-1} . This is precisely what is observed in Figure 6 ($d = 10^{-3}$) and Figure 7 ($d = 10^{-6}$) ahead. The graphs are now in a log – log scale, and we observe straight lines with slope -1 . The “doubling” over the rate of convergence with the uniform h version is also clearly apparent from these figures.

Let us now consider the hp version, i.e., the p version on a variable mesh. Since our model solution (6.1) has a boundary layer at each endpoint of the domain, the minimum number of elements as in Theorem 5.2 will now be 3, with Σ given by (5.18). (Since f is a polynomial of degree 0, we can actually take the minimal degree vector to be $\vec{p} = \{p, 1, p\}$.) From Theorem 5.2, we have the error estimate

$$(6.4) \quad E_R \leq C(\kappa)d^{1/2}\alpha^{N/2}, \quad N = \dim(S_0(\Sigma)) = 2p + 1,$$

with α given by (5.2). The experiments in Figure 3, obtained with $\kappa = \kappa^* = 0.71$, clearly show the uniform exponential convergence as well as the factor $d^{1/2}$, since $\log(E_R(d))$ plotted against N is a straight line, which translates downwards as d decreases. By Remark 5.1, for $\kappa = 0.71$, we have $\alpha \approx e^{-0.71}$ for p large — this is the same value that emerges by measuring the slopes in Figure 3. Note that the striking accuracy obtained for small d is not possible with a comparable number of





FIGURE 3. The hp version for three elements with $\kappa = 0.71$

degrees of freedom and methods based on a single element (see, e.g., the results in [7, 5]).

In Figure 4, we investigate the convergence of the three-element hp version for different values of κ , when $d = 10^{-6}$ (other values of d show similar results). We observe that $\kappa = \kappa^* = 0.71$ is not quite optimal, since $\kappa = 1$ gives better results. Careful examination shows that the graph for $\kappa = 1$ consists of two linear pieces with different slopes. This is due to the fact that initially, the error in the central interval is dominant, so that the value of α in (5.2), (6.3) is close to $e^{-\kappa}$. As p increases, the size of this interval decreases and the error in the other two intervals eventually dominates, with α behaving like $\kappa e/4$. (Recall that we obtained κ^* by setting $e^{-\kappa}$ equal to $\kappa e/4$, so that only one straight line is observed in this case.)

Finally, in Figures 5–7, we show a performance comparison between the various methods for $d = 10^{-2}$, 10^{-3} and 10^{-6} , respectively (smaller values of d up to 10^{-8} were tested, for which the behavior was similar to $d = 10^{-6}$). In these figures, we have shown the results with four methods: (a) the p version with one element, (b) the h version with $p = 1$, (c) the hp version with 3 elements taking $\vec{p} = (p, p, p)$ and $\kappa = 1$ and (d) the h version (taking $p = 1$) with the exponential mesh $\Delta = \{-1, x_1, \dots, x_{m-1}, 1\}$, where, for m even,

$$(6.5) \quad x_{\frac{m}{2} \pm i} = \mp d\tilde{p} \ln \left(1 - c \frac{2i}{m} \right), \quad i = 0, \dots, \frac{m}{2},$$

FIGURE 4. The dependence on parameter κ

with $c = 1 - \exp(-1/(d\tilde{p}))$. The mesh (6.5) is derived in [16, 20]. We observe the following.

- (i) The uniform h version converges with order $\mathcal{O}(N^{-1/2})$, the p version on a single element with order $\mathcal{O}(N^{-1})$, and the h version with exponential mesh at the optimal algebraic rate of $\mathcal{O}(N^{-1})$.
- (ii) Both the h version with exponential mesh and the hp version have an error which behaves like $\mathcal{O}(d^{1/2})$ in dependence on d . The other two versions do not display this translation as $d \rightarrow 0$.
- (iii) For $d = 10^{-2}$, the p version rapidly reaches a superexponential rate, and eventually becomes the method with the fastest convergence. Asymptotically, i.e., for $\kappa\tilde{p}d > 2$ and fixed d , the p version with a single element will always have the best convergence rate according to Theorems 4.1 and 5.1. Accordingly, Theorem 5.1 indicates that at about $\kappa\tilde{p}d = 2$ one must switch from the hp version to a single-element p version. For $d = 10^{-2}$, this is apparent in Figure 5, where the one-element p version becomes superior at some point. However, as is clearly visible in Figures 6 and 7, this point may occur so late that the only feasible method (in the practical range of p) is the three-element hp version.




FIGURE 5. Comparison of various methods, $d = 10^{-2}$

APPENDIX

We prove Theorem 2.1. For $M \in \mathbb{N}$, we define

$$(6.6) \quad \omega_d^M(x) = \sum_{k=0}^M d^{2k} a^{-2k-2} f^{(2k)}(x).$$

Then we see, using (1.2), that $R_d^M = u_d - \omega_d^M$ satisfies

$$(6.7) \quad L_d R_d^M = f - L_d \omega_d^M = (d/a)^{2M+2} f^{(2M+2)}(x) =: g(x).$$

For M large, we see that ω_d^M will satisfy (1.2) up to the correction $g(x)$. However, in general, the boundary conditions (1.3) will not be satisfied. We therefore introduce appropriate boundary layer terms to enforce (1.3). For this purpose, we define $u_k^{BL}(x)$ to be the unique solution of

$$(6.8) \quad L_d u_k^{BL}(x) = 0 \quad \text{on } I,$$

$$(6.9) \quad u_k^{BL}(\pm 1) = C_k^\pm := \delta_{0,k} \alpha^\pm - a^{-2k-2} f^{(2k)}(\pm 1).$$

Then with $u_{a,d}$ and $\bar{u}_{a,d}$ as defined in (2.4), we may verify that

$$(6.10) \quad u_k^{BL}(x) = A_k u_{a,d}(x) + B_K \bar{u}_{a,d}(x),$$




FIGURE 6. Comparison of various methods, $d = 10^{-3}$

where

$$(6.11) \quad A_k = \frac{C_k^- - C_k^+ e^{-2a/d}}{1 - e^{-4a/d}}, \quad B_k = \frac{C_k^+ - C_k^- e^{-2a/d}}{1 - e^{-4a/d}}.$$

Then we write

$$U_M^{BL}(x) = \sum_{k=0}^M d^{2k} u_k^{BL}(x) = A_d^M u_{a,d}(x) + B_d^M \bar{u}_{a,d}(x),$$

where $A_d^M = \sum_{k=0}^M d^{2k} A_k$ and $B_d^M = \sum_{k=0}^M d^{2k} B_k$. We see that

$$(6.12) \quad |A_d^M| \leq \sum_{k=0}^M d^{2k} |A_k| \leq (1 - e^{-4a})^{-1} \sum_{k=0}^M d^{2k} (|C_k^-| + |C_k^+|),$$

with a similar bound holding for $|B_d^M|$. Equation (2.7) follows from (6.12) and (6.9).

We now define

$$r_d^M = u_d - \omega_d^M - A_d^M u_{a,d} - B_d^M \bar{u}_{a,d} = R_d^M - U_M^{BL}.$$

Then $r_d^M \in H_0^1(I)$ and r_d^M satisfies (6.7). Hence,

$$\|r_d^M\|_E^2 := B_d(r_d^M, r_d^M) = (g, r_d^M) \leq \|g\|_0 \|r_d^M\|_E.$$




FIGURE 7. Comparison of various methods, $d = 10^{-6}$

From this we deduce (using (6.7)) that

$$(6.13) \quad \|r_d^M\|_0 \leq a^{-1}(d/a)^{2M+2}\|f^{(2M+2)}\|_0, \quad |r_d^M|_1 \leq a^{-1}(d/a)^{2M+1}\|f^{(2M+2)}\|_0.$$

Since r_d^M satisfies (6.7), we may differentiate (6.7) successively to obtain, using (6.13),

$$(6.14) \quad |r_d^M|_\ell \leq a^{-1}(d/a)^{2M+2-\ell}\|f^{(2M+2)}\|_0 + a^{-2} \sum_{k=0}^{\ell-2} (d/a)^{2M+2-\ell+k}\|f^{(2M+2+k)}\|_0,$$

where $\ell = 0, 1, \dots, 2M$. Moreover, from (6.6), we see that for $\ell = 0, 1, \dots, 2M$

$$(6.15) \quad |\omega_d^M|_\ell \leq a^{-2} \sum_{k=0}^M (d/a)^{2k}\|f^{(2k+\ell)}\|_0.$$

Define $u_d^M = \omega_d^M + r_d^M$. Then (2.5) holds. Also, using (6.14)–(6.15), we may establish (2.6).

Remark 6.1. Suppose $f \in \Pi_{2M+1}(I)$. Then in (6.7), we have $g(x) \equiv 0$, so that, since $r_d^M \in H_0^1(I)$ satisfies (6.7), we must have $r_d^M = 0$. Hence, $u_d^M = \omega_d^M$, and it is seen by (6.6) that $u_d^M \in \Pi_{2M+1}(I)$.

ACKNOWLEDGEMENTS

The authors would like to thank Christos Xenophontos for the computations performed in Section 6, and Professor F. W. J. Olver for helpful discussions regarding the reference [9].

REFERENCES

1. D. N. Arnold and R. S. Falk. Asymptotic analysis of the boundary layer for the Reissner-Mindlin plate model. *SIAM J. Math. Anal.* 27: 486–514, 1996.
2. I. Babuška and M. Suri. On locking and robustness in the finite element method. *SIAM J. Numer. Anal.*, 29:1261–1293, 1992. MR **94c**:65128
3. I. Babuška and B. A. Szabo. Lecture notes on finite element analysis, (to appear).
4. I. A. Blatov and V. V. Strygin. On estimates best possible in order in the Galerkin finite element method for singularly perturbed boundary value problems. *Russian Acad. Sci. Dokl. Math.*, 47:93–96, 1993.
5. C. Canuto. Spectral methods and a maximum principle. *Math. Comp.*, 51:615–629, 1988. MR **89d**:65099
6. E. C. Gartland. Uniform high-order difference schemes for a singularly perturbed two-point boundary value problem. *Math. Comp.*, 48:551–564, 1987. MR **89a**:65116
7. W. B. Liu and J. Shen. A new efficient spectral Galerkin method for singular perturbation problems, Preprint, Department of Mathematics, Penn State University, State College Pa (1994).
8. W. B. Liu and T. Tang. Boundary layer resolving methods for singularly perturbed problems, submitted to *I.M.A. J. Numer. Anal.*
9. F. W. J. Olver. Error bounds for the Liouville-Green (or WKB) approximation. *Proc. Cambridge Philos. Soc.*, 57:790–810, 1961. MR **24**:A313
10. I. S. Gradshteyn and I. M. Ryzhik. *Table of Series, Integrals and Products, enlarged edition*. Wiley, New York, 1980. MR **81g**:33001 /
11. H. Hakula, Y. Leino, and J. Pitkäranta. Scale resolution, layers and high-order numerical modeling of shells, to appear in *Comp. Meth. Appl. Mech. Eng.*, 1996.
12. H. Kraus. Thin elastic shells: an introduction to the theoretical foundations and the analysis of their static and dynamic behavior. New York, Wiley 1967.
13. A. H. Schatz and L. B. Wahlbin. On the finite element method for singularly perturbed reaction-diffusion problems in two and one dimensions. *Math. Comp.*, 40:47–89, 1983. MR **84c**:65137
14. K. Scherer. On optimal global error bounds obtained by scaled local error estimates. *Numer. Math.*, 36:151–176, 1981. MR **82e**:65053
15. C. Schwab and M. Suri. Locking and boundary layer effects in the finite element approximation of the Reissner-Mindlin plate model. *Proc. Symp. Appl. Math.*, 48:367–371, 1994. MR **95m**:65195
16. C. Schwab, M. Suri, and C. Xenophontos. The hp finite element method for problems in mechanics with boundary layers (to appear).
17. C. Schwab and S. Wright. Boundary layers in hierarchical beam and plate models, report. *Journal of Elasticity* 38 (1995), 1–40. MR **96d**:73044
18. G.I. Shishkin. Grid approximation of singularly perturbed parabolic equations with internal layers. *Soviet J. Numer. Anal. Math. Modelling*, 3:393–407, 1988. MR **89k**:65109
19. R. Vulanović, D. Herceg, and N. Petrović. On the extrapolation for a singularly perturbed boundary value problem. *Computing*, 36:69–79, 1986. MR **88a**:65008
20. C. A. Xenophontos. The hp version of the finite element method for singularly perturbed problems in unsmooth domains. *Ph.D. Dissertation, UMBC*, 1996.

SEMINAR FOR APPLIED MATHEMATICS, ETH ZÜRICH, RÄMISTRASSE 101, CH-8092, ZÜRICH, SWITZERLAND

E-mail address: schwab@sam.math.ethz.ch

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MARYLAND BALTIMORE COUNTY, 5401 WILKENS AVENUE, BALTIMORE, MARYLAND 21228

E-mail address: suri@math.umbc.edu