

## ALMOST PERIODIC FACTORIZATION OF CERTAIN BLOCK TRIANGULAR MATRIX FUNCTIONS

ILYA M. SPITKOVSKY AND DARRYL YONG

ABSTRACT. Let

$$G(x) = \begin{bmatrix} e^{i\lambda x} I_m & 0 \\ c_{-1} e^{-i\nu x} + c_0 + c_1 e^{i\alpha x} & e^{-i\lambda x} I_m \end{bmatrix},$$

where  $c_j \in \mathbb{C}^{m \times m}$ ,  $\alpha, \nu > 0$  and  $\alpha + \nu = \lambda$ . For rational  $\alpha/\nu$  such matrices  $G$  are periodic, and their Wiener-Hopf factorization with respect to the real line  $\mathbb{R}$  always exists and can be constructed explicitly. For irrational  $\alpha/\nu$ , a certain modification (called an almost periodic factorization) can be considered instead. The case of invertible  $c_0$  and commuting  $c_1 c_0^{-1}$ ,  $c_{-1} c_0^{-1}$  was disposed of earlier—it was discovered that an almost periodic factorization of such matrices  $G$  does not always exist, and a necessary and sufficient condition for its existence was found.

This paper is devoted mostly to the situation when  $c_0$  is not invertible but the  $c_j$  commute pairwise ( $j = 0, \pm 1$ ). The complete description is obtained when  $m \leq 3$ ; for an arbitrary  $m$ , certain conditions are imposed on the Jordan structure of  $c_j$ . Difficulties arising for  $m = 4$  are explained, and a classification of both solved and unsolved cases is given.

The main result of the paper (existence criterion) is theoretical; however, a significant part of its proof is a constructive factorization of  $G$  in numerous particular cases. These factorizations were obtained using Maple; the code is available from the authors upon request.

### 1. INTRODUCTION

Let  $AP$  be the Bohr algebra of almost periodic functions, that is, the smallest  $C^*$ -algebra of  $L^\infty(\mathbb{R})$  containing all the functions  $e_\lambda(x) = e^{i\lambda x}$ ,  $\lambda \in \mathbb{R}$ . It is well known (the standard references for these and other properties of  $AP$  are [3, 11, 12]) that for every  $f \in AP$ ,

1. there exists the *Bohr mean value*

$$\mathbf{M}(f) = \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T f(x) dx,$$

and

2. the *Fourier coefficients*  $\widehat{f}(\lambda) \stackrel{\text{def}}{=} \mathbf{M}(f e_{-\lambda})$  are different from zero for at most countably many values of  $\lambda \in \mathbb{R}$ .

---

Received by the editor March 12, 1997 and, in revised form, September 18, 1998.

1991 *Mathematics Subject Classification*. Primary 47A68, 47-04, 42A75.

*Key words and phrases*. Almost periodic matrix functions, factorization, explicit computation.

The first author's research was partially supported by NSF Grant DMS-9800704.

The second author's research was started during a Research Experience for Undergraduates sponsored by the NSF at the College of William and Mary during the summer of 1995.

The set  $\Omega(f) = \{\mu: \widehat{f}(\mu) \neq 0\}$  is called the *Fourier spectrum* of  $f$ , and

$$(1.1) \quad \sum_{\mu \in \Omega(f)} \widehat{f}(\mu)e_{\mu}$$

is its (formal) *Fourier series*.

We say that  $f \in AP_W$  if the Fourier series (1.1) converges absolutely:

$$\sum_{\mu \in \Omega(f)} |\widehat{f}(\mu)| < \infty.$$

Finally, let

$$AP^{\pm} = \{f \in AP: \Omega(f) \subset \mathbb{R}_{\pm}\} \quad \text{and} \quad AP_W^{\pm} = AP^{\pm} \cap AP_W.$$

Here, as usual,  $\mathbb{R}_{\pm} = \{x \in \mathbb{R}: \pm x \geq 0\}$ .

For matrix functions  $f$ , conditions  $f \in AP, AP^{\pm}, AP_W$ , etc. are understood entrywise, and  $\mathbf{M}(f), \widehat{f}(\mu), \Omega(f)$  are defined by exactly the same formulas as for scalar functions.

Following [6], we introduce an *AP factorization* of an  $n \times n$  matrix function  $G$  as its representation in the form

$$(1.2) \quad G = G_+ \Lambda G_-,$$

where  $\Lambda(x) = \text{diag}[e_{\lambda_1}, \dots, e_{\lambda_n}]$ ,

$$(1.3) \quad G_+^{\pm 1} \in AP^+, \quad G_-^{\pm 1} \in AP^-,$$

and  $\lambda_1, \dots, \lambda_n \in \mathbb{R}$ . We say that (1.2) is an  $AP_W$  factorization of  $G$  if conditions (1.3) are replaced by the (more restrictive) conditions  $G_+^{\pm 1} \in AP_W^+, G_-^{\pm 1} \in AP_W^-$ .

If  $G$  is *AP factorable*, the numbers  $\lambda_1, \dots, \lambda_n$  are defined uniquely; they are called the *partial AP indices* (of  $G$ ). Of course, for an *AP* ( $AP_W$ ) factorization (1.2) to exist it is necessary that  $G^{\pm 1} \in AP$  (respectively,  $AP_W$ ). However, this necessary condition is not sufficient and, except for the case of periodic matrix functions  $G$  (in which an *AP* factorization by a simple change of variable reduces to the usual Wiener-Hopf factorization), the theory of *AP* factorization is “under construction”. Its connections with integral equations, completion problems, and signal processing are discussed in [6, 7], [17, 15, 1], and [14] respectively. Explicit formulas for the factors in (1.2) for certain special types of  $G$  are obtained in [6, 9, 8]. Most of them refer to matrices  $G$  of the following block triangular form:

$$(1.4) \quad G_f = \begin{bmatrix} e_{\lambda} I_m & 0 \\ f & e_{-\lambda} I_m \end{bmatrix}$$

(so that  $n = 2m$ ) arising in the treatment of convolution type equations on finite intervals of length  $\lambda$ .

In particular, the following two statements were established in [6].

**Lemma 1.1.** *Let  $f$  be an  $AP_W$  matrix function, and let*

$$f_0 = \sum_{\mu \in \Omega(f) \cap (-\lambda, \lambda)} \widehat{f}(\mu)e_{\mu}.$$

*Then the matrices  $G_f$  and  $G_{f_0}$  are  $AP$  ( $AP_W$ ) factorable only simultaneously, and their partial *AP* indices coincide.*

Due to Lemma 1.1, for any  $f \in AP_W$  in (1.4) we may suppose without loss of generality that  $\Omega(f) \subset (-\lambda, \lambda)$ .

**Theorem 1.2.** *Let  $\Omega(f) \cap (-\lambda, \lambda)$  consist of at most two points, say  $\mu$  and  $\sigma$ . Then  $G_f$  is  $AP_W$  factorable. Its partial  $AP$  indices all equal zero if and only if  $\mu\sigma = 0$ ,  $\widehat{f}(0)$  is invertible, or  $\mu\sigma < 0$ ,  $\frac{\lambda}{\mu-\sigma} \in \mathbb{Z}$  and both  $\widehat{f}(\mu)$ ,  $\widehat{f}(\sigma)$  are invertible.*

The next logical step is to consider a trinomial  $f$  with  $\Omega(f) \subset (-\lambda, \lambda)$ . However, with no additional restrictions on the location of  $\Omega(f)$  this remains an open problem. In this paper, we concentrate on the case  $\Omega(f) = \{-\nu, 0, \alpha\}$ , that is,

$$(1.5) \quad f = c_{-1}e_{-\nu} + c_0 + c_1e_{\alpha},$$

where  $\alpha, \nu > 0$  and  $\alpha + \nu = \lambda$ .

If  $\beta = \frac{\nu}{\alpha}$  is rational, then the matrix  $G_f$  is periodic, and its  $AP_W$  factorization exists and can be easily constructed. Thus, we suppose in what follows that  $\beta$  is irrational. The next result applies to the case when the matrices  $c_j$  in (1.5) commute with each other. In this case there exists a similarity  $T$  such that

$$(1.6) \quad T^{-1}c_jT = \text{diag}[c_{j1}, \dots, c_{jr}], \quad c_{jk} \in \mathbb{C}^{l_k \times l_k}, \quad k = 1, \dots, r; \quad j = 0, \pm 1,$$

and each diagonal block  $c_{jk}$  has a singleton spectrum (see [13, Section 4.4]):

$$\sigma(c_{jk}) = \{\xi_{jk}\} \quad (j = 0, \pm 1; k = 1, \dots, r).$$

As in [2], we call  $\{\xi_{jk}\}_{j=-1}^1$  the *bonded eigenvalue triples* of  $c_j$ .

**Theorem 1.3.** *Let  $G_f$  be of the form (1.4) with  $f$  given by (1.5) and commuting coefficients  $c_j$ . Then  $G_f$  is  $AP$  factorable with zero partial  $AP$  indices if and only if, for all bonded triples  $\{\xi_{-1,k}, \xi_{0,k}, \xi_{1,k}\}$ ,*

$$(1.7) \quad |\xi_{1,k}^{\nu} \xi_{-1,k}^{\alpha}| \neq |\xi_{0,k}|^{\lambda} \quad (k = 1, \dots, r).$$

In this form, Theorem 1.3 was established in [2, Theorem 7.2]; the case of invertible  $c_j$  was disposed of earlier in [9]. In fact, the result of [9] contains an additional important piece of information: if all  $c_j$  are invertible and (1.7) fails for at least one value of  $k$ , then  $G_f$  does not admit any  $AP$  factorization, even if non-zero partial  $AP$  indices are allowed. Also, it was shown in [16] that an  $AP$  factorization with zero partial  $AP$  indices of an  $AP_W$  matrix function is automatically its  $AP_W$  factorization. Hence, the following result holds.

**Corollary 1.4.** *Let  $G_f$  be as in Theorem 1.3 and, in addition, let  $c_0$  be invertible. Then  $G_f$  is  $AP_W$  factorable with zero partial  $AP$  indices if condition (1.7) holds, and is not  $AP$  factorable otherwise.*

Of course, it would now be natural to consider an  $AP$  factorization of  $G_f$  with trinomial  $f$ , pairwise commuting  $c_j$ , and no restrictions imposed on the invertibility of  $c_0$  and the values of partial  $AP$  indices. We will see, however, that this problem embraces a general setting of a trinomial  $f$  with arbitrary (not necessarily commuting) coefficients  $c_j$  and is therefore too difficult to handle at the present stage of the development. Our paper is a report on several partial results on the  $AP$  factorability of matrices (1.4), (1.5) with non-invertible  $c_0$ .

The paper is structured as follows. Section 2 contains an auxiliary result on the factorization of block diagonal matrices. It also describes a procedure which allows us to replace a matrix of the form (1.4), (1.5) with invertible  $c_{-1}$  (and no commutativity conditions on  $c_j$ ) by another matrix of the same type without changing its factorability properties. This procedure is, in fact, a variation of the one introduced in [2] for matrices (1.4) with a finite number (not limited to three) of

points  $\mu_j$  in  $\Omega(f) \cap (-\lambda, \lambda)$  but pairwise commuting  $\widehat{f}(\mu_j)$ . As a direct application of this procedure,  $AP_W$  factorability is established for matrices (1.4), (1.5) with  $m = 2$ , invertible  $c_{-1}$  (or  $c_1$ ) and nilpotent  $c_0 c_{-1}^{-1}$  (respectively,  $c_0 c_1^{-1}$ ).

Section 3 contains necessary and sufficient factorability conditions for matrices (1.4), (1.5) with commuting  $c_j$  under certain additional restrictions on their Jordan structure. This covers, in particular, all matrices of size  $m \leq 3$ , invertible  $c_1$  or  $c_{-1}$  of size  $m \leq 4$ , and matrices of arbitrary size, provided that each eigenvalue of at least one of the  $c_j$  corresponds to one Jordan cell. An application to difference equations is given.

In Section 4, we concentrate on  $4 \times 4$  matrices  $c_j$ . An example is given explaining why this case cannot be covered in general before the  $AP$  factorability of matrices (1.4), (1.5) with arbitrary invertible non-commuting  $c_j$  is understood. All possible cases are classified, and those for which the  $AP$  factorability remains unknown are singled out.

Proofs of the results in Sections 3 and 4 are partially theoretical and partially consist in exhausting a large number of cases in which an  $AP_W$  factorization can be constructed explicitly. These cases are relegated to Section 5 the supplement at the end of this volume, where final formulas are listed. Of course, they can be checked by straightforward calculations. We emphasize, however, that a symbolic manipulation Maple program was used to obtain these formulas, and without it this paper could hardly have been completed.

## 2. AUXILIARY RESULTS

Suppose  $G$  is a block diagonal  $AP$  matrix:  $G = \text{diag}[G_1, G_2]$ . If its diagonal blocks  $G_1, G_2$  are  $AP$  factorable, then  $G$  itself is  $AP$  factorable. Moreover, an  $AP$  factorization of  $G$  can be obtained by “pasting together”  $AP$  factorizations of  $G_1$  and  $G_2$ :  $G_1 = G_+^{(1)} \Lambda^{(1)} G_-^{(1)}$ ,  $G_2 = G_+^{(2)} \Lambda^{(2)} G_-^{(2)}$  imply

$$G = \text{diag}[G_+^{(1)}, G_+^{(2)}] \text{diag}[\Lambda^{(1)}, \Lambda^{(2)}] \text{diag}[G_-^{(1)}, G_-^{(2)}].$$

It is natural to ask whether the converse is true. The answer is positive provided that  $G \in AP_W$  and partial  $AP$  indices of  $G$  equal zero. Indeed, a matrix  $F \in AP_W$  admits an  $AP$  factorization with zero partial  $AP$  indices if and only if the corresponding Toeplitz operator  $T_F$  is invertible on  $L^2$  [5] (see also [7]). Since  $T_G$  is a direct sum of  $T_{G_1}$  with  $T_{G_2}$ , the invertibility of  $T_G$  is equivalent to simultaneous invertibility of  $T_{G_1}$  and  $T_{G_2}$ .

We are not aware of any equivalent of  $AP$  factorability (with non-zero partial  $AP$  indices) in operator terms. Probably, the answer to the question is still positive, but we restrict our consideration to a somewhat weaker version.

**Lemma 2.1.** *Let  $G = \text{diag}[G_1, G_2]$ . If  $G$  and one of its diagonal blocks  $G_1, G_2$  are  $AP$  factorable, then the other diagonal block is also  $AP$  factorable.*

*Proof.* Consider first the case when  $G_1 = 1$ . Then an  $AP$  factorization of  $G$  can be rewritten as

$$(2.1) \quad F_+ \begin{bmatrix} 1 & 0 \\ 0 & G_2 \end{bmatrix} = \Lambda G_-,$$

where  $F_+ = G_+^{-1} \in AP^+$ . Denote  $F_+ = (f_{ij})_{i,j=1}^n$ . From (2.1),  $e_{-\lambda_j} f_{j1} \in AP^-$ , so that

$$(2.2) \quad \Omega(f_{j1}) \subset [0, \lambda_j].$$

In particular,  $f_{j1} = 0$  for all  $j$  (if there are any) such that  $\lambda_j < 0$ . Rewriting (2.1) as

$$\begin{bmatrix} 1 & 0 \\ 0 & G_2^{-1} \end{bmatrix} G_+ \Lambda = G_-^{-1},$$

we find similarly that

$$(2.3) \quad \Omega(g_{1j}) \subset [0, -\lambda_j],$$

where  $G_+ = (g_{ij})_{i,j=1}^n$ . Therefore,  $g_{1j} = 0$  for all  $j$  (if there are any) such that  $\lambda_j > 0$ . Observe also that  $G_+ F_+ = I$  implies that  $\sum_{j=1}^n g_{1j} f_{j1} = 1$ . Since for non-zero  $\lambda_j$  at least one of the entries  $g_{1j}, f_{j1}$  is equal to zero, the latter equality proves the existence of zero partial  $AP$  indices  $\lambda_j$ . Due to (2.2), (2.3), the corresponding functions  $g_{1j}, f_{j1}$  are constant, and for at least one value of  $j$ ,  $g_{1j} f_{j1} \neq 0$ .

Applying an appropriate permutation of columns of  $G_+$  and rows of  $G_-$ , we may suppose without loss of generality that  $\lambda_1 = 0, g_{11} = c \neq 0, f_{11} = d \neq 0$ . Partitioning  $G_+, F_+$  as

$$G_+ = \begin{bmatrix} c & g_1^+ \\ g_2^+ & G_2^+ \end{bmatrix}, F_+ = \begin{bmatrix} d & f_1^+ \\ f_2^+ & F_2^+ \end{bmatrix},$$

we conclude that  $c = \det F_2^+ / \det F_+ = \det F_2^+ \det G_+$ . Since  $c \neq 0$ , the matrix  $F_2^+$  is invertible in  $AP^+$  simultaneously with  $G_+$ . From (2.1) and (2.2) it follows that the left-upper entry of  $G_-$  and  $H_- = G_-^{-1}$  equals  $d$  and  $c$ , respectively. Thus,

$$G_- = \begin{bmatrix} d & g_1^- \\ g_2^- & G_2^- \end{bmatrix}, H_- = \begin{bmatrix} c & h_1^- \\ h_2^- & H_2^- \end{bmatrix},$$

and  $c = \det G_2^- / \det G_-$ . Since  $c \neq 0$ , the matrix  $G_2^-$  is invertible in  $AP^-$  together with  $G_-$ . Now partition  $\Lambda = \begin{bmatrix} 1 & 0 \\ 0 & \Lambda_2 \end{bmatrix}$ . Then (2.1) yields  $F_2^+ G_2 = \Lambda_2 G_2^-$ , or  $G_2 = (F_2^+)^{-1} \Lambda_2 G_2^-$ . Since  $(F_2^+)^{\pm 1} \in AP^+$  and  $(G_2^-)^{\pm 1} \in AP^-$ , the latter formula delivers an  $AP$  factorization of  $G_2$ . This proves the desired statement in the case  $G_1 = 1$ .

If  $G_1 = e_\lambda$ , then the matrix  $e_{-\lambda} G = \text{diag}[1, e_{-\lambda} G_2]$  is  $AP$  factorable together with  $G$ . According to the already proven particular case,  $e_{-\lambda} G_2$  is  $AP$  factorable. But then  $G_2$  is  $AP$  factorable as well.

An induction argument allows us to consider  $G_1$  of the form  $\text{diag}[e_{\lambda_1}, \dots, e_{\lambda_k}] = \Lambda_1$ . Finally, for an arbitrary  $AP$  factorable  $G_1 = G_+^{(1)} \Lambda_1 G_-^{(1)}$  we can write

$$G = \text{diag}[G_+^{(1)}, I] \text{diag}[\Lambda_1, G_2] \text{diag}[G_-^{(1)}, I]$$

and consider an ( $AP$  factorable) matrix  $\text{diag}[\Lambda_1, G_2]$  instead of the original matrix  $G$ . □

Another technical tool we need applies to matrix functions  $G_f$  with a trinomial  $f$  containing an invertible  $c_{-1}$  coefficient.

**Lemma 2.2.** *Let  $G$  be of the form (1.4) with  $f$  given by (1.5). If  $c_{-1}$  is invertible, then  $G$  is AP ( $AP_W$ ) factorable only simultaneously with (and has the same partial AP indices as) the matrix function*

$$(2.4) \quad G_1 = \begin{bmatrix} e_{\lambda_1} I_m & 0 \\ f_1 & e_{-\lambda_1} I_m \end{bmatrix},$$

where

$$(2.5) \quad f_1 = c_{-1}^{(1)} e_{-\nu_1} + c_0^{(1)} + c_1^{(1)} e_{\alpha_1},$$

$$c_{-1}^{(1)} = (-1)^s (c_{-1}^{-1} c_0)^{s+1}, \quad c_0^{(1)} = c_{-1}^{-1} c_1, \quad c_1^{(1)} = (-1)^{s+1} (c_{-1}^{-1} c_0)^{s+2},$$

$$\lambda_1 = \nu, \quad \nu_1 = \alpha - s\nu, \quad \alpha_1 = (s + 1)\nu - \alpha,$$

and finally,  $s$  is the integral part of  $\frac{\alpha}{\nu}$ :  $s \in \mathbb{Z}$  and  $s < \frac{\alpha}{\nu} < s + 1$ .

*Proof.* It suffices to construct matrix functions  $X_+$  and  $X_-$  such that  $X_{\pm}^{\pm 1} \in AP_W^{\pm}$ ,  $X_{\pm}^{\pm 1} \in AP_{\bar{W}}$  and

$$(2.6) \quad X_+ G X_- = G_1.$$

To this end, let

$$(2.7) \quad \begin{aligned} X_+ &= \begin{bmatrix} c_{-1}^{-1} f e_{\nu} & -e_{\lambda+\nu} I \\ e_{-\lambda-\nu} I + (g - e_{-\lambda} I) c_{-1}^{-1} f & I - g e_{\lambda} \end{bmatrix} \begin{bmatrix} I & 0 \\ 0 & c_{-1}^{-1} \end{bmatrix}, \\ X_- &= \begin{bmatrix} I & 0 \\ 0 & c_{-1} \end{bmatrix} \begin{bmatrix} e_{-\alpha} I + \sum_{j=1}^s (-1)^j (c_{-1}^{-1} c_0)^j e_{j\nu-\alpha} & I \\ -I & 0 \end{bmatrix}, \end{aligned}$$

where  $g = c_{-1}^{-1} c_1 - \sum_{j=1}^{s+2} (-1)^j (c_{-1}^{-1} c_0)^j e_{(j-1)\nu-\alpha}$ . Directly from the definition of  $s$  it follows that  $X_- \in AP^-$ . Since  $\det X_- = \det c_{-1}$  is a non-zero constant,  $X_-^{-1}$  belongs to  $AP^-$  together with  $X_-$ .

A straightforward computation shows that

$$\begin{aligned} X_+ G X_- &= \begin{bmatrix} c_{-1}^{-1} f e_{\nu} & -e_{\lambda+\nu} I \\ e_{-\lambda-\nu} I + (g - e_{-\lambda} I) c_{-1}^{-1} f & I - g e_{\lambda} \end{bmatrix} \\ &\quad \times \begin{bmatrix} e_{\lambda} I & 0 \\ c_{-1}^{-1} f & e_{-\lambda} I \end{bmatrix} \begin{bmatrix} e_{-\alpha} I + \sum_{j=1}^s (-1)^j (c_{-1}^{-1} c_0)^j e_{j\nu-\alpha} & I \\ -I & 0 \end{bmatrix} \\ &= \begin{bmatrix} c_{-1}^{-1} f e_{\nu} & -e_{\lambda+\nu} I \\ e_{-\lambda-\nu} I + (g - e_{-\lambda} I) c_{-1}^{-1} f & I - g e_{\lambda} \end{bmatrix} \\ &\quad \times \begin{bmatrix} e_{\nu} I + \sum_{j=1}^s (-1)^j (c_{-1}^{-1} c_0)^j e_{(j+1)\nu} & e_{\lambda} I \\ c_{-1}^{-1} e_{-\alpha} f + c_{-1}^{-1} f \sum_{j=1}^s (-1)^j (c_{-1}^{-1} c_0)^j e_{j\nu-\alpha} - e_{-\lambda} I & c_{-1}^{-1} f \end{bmatrix} \\ &= (y_{ij})_{i,j=1}^2, \end{aligned}$$

where

$$\begin{aligned}
 y_{11} &= c_{-1}^{-1} f e_{2\nu} + c_{-1}^{-1} f \sum_{j=1}^s (-1)^j (c_{-1}^{-1} c_0)^j e_{(j+2)\nu} - c_{-1}^{-1} f e_{2\nu} \\
 &\quad - c_{-1}^{-1} f \sum_{j=1}^s (-1)^j (c_{-1}^{-1} c_0)^j e_{(j+2)\nu} + e_\nu I = e_\nu I, \\
 y_{12} &= c_{-1}^{-1} f e_{\lambda+\nu} - c_{-1}^{-1} f e_{\lambda+\nu} = 0, \\
 y_{22} &= e_{-\nu} I + (g e_\lambda - I) c_{-1}^{-1} f + (I - g e_\lambda) c_{-1}^{-1} f = e_{-\nu} I,
 \end{aligned}$$

and finally,

$$\begin{aligned}
 y_{21} &= (e_{-\lambda-\nu} I + (g - e_{-\lambda} I) c_{-1}^{-1} f) \left( e_\nu I + \sum_{j=1}^s (-1)^j (c_{-1}^{-1} c_0)^j e_{(j+1)\nu} \right) \\
 &\quad + (I - g e_\lambda) \left( c_{-1}^{-1} e_{-\alpha} f + c_{-1}^{-1} f \sum_{j=1}^s (-1)^j (c_{-1}^{-1} c_0)^j e_{j\nu-\alpha} - e_{-\lambda} I \right) \\
 &= e_{-\lambda} I + \sum_{j=1}^s (-1)^j (c_{-1}^{-1} c_0)^j e_{j\nu-\lambda} + (g - e_{-\lambda} I) \\
 &\quad \times \left( c_{-1}^{-1} f e_\nu + c_{-1}^{-1} f \sum_{j=1}^s (-1)^j (c_{-1}^{-1} c_0)^j e_{(j+1)\nu} \right. \\
 &\quad \left. - c_{-1}^{-1} f e_\nu - c_{-1}^{-1} f \sum_{j=1}^s (-1)^j (c_{-1}^{-1} c_0)^j e_{(j+1)\nu} + I \right) \\
 &= e_{-\lambda} I + \sum_{j=1}^s (-1)^j (c_{-1}^{-1} c_0)^j e_{j\nu-\lambda} + g - e_{-\lambda} I \\
 &= \sum_{j=1}^s (-1)^j (c_{-1}^{-1} c_0)^j e_{j\nu-\lambda} + c_{-1}^{-1} c_1 - \sum_{j=1}^{s+2} (-1)^j (c_{-1}^{-1} c_0)^j e_{(j-1)\nu-\alpha} \\
 &= (-1)^s (c_{-1}^{-1} c_0)^{s+1} e_{s\nu-\alpha} + c_{-1}^{-1} c_1 + (-1)^{s+1} (c_{-1}^{-1} c_0)^{s+2} e_{(s+1)\nu-\alpha} = f_1.
 \end{aligned}$$

This implies (2.6). Since  $\det G = \det G_1 = 1$ , from (2.6) it follows, in particular, that  $\det X_+ = (\det X_-)^{-1}$  is a non-zero constant. It remains to show that  $X_+ \in AP^+$ , because then  $X_+^{-1} \in AP^+$  as well. Three blocks of  $X_+$  are obviously in  $AP^+$ . The remaining (left-lower) block can be rewritten as

$$\begin{aligned}
 &e_{-\lambda-\nu} I + (g - e_{-\lambda} I) c_{-1}^{-1} f \\
 &= e_{-\lambda-\nu} I + c_{-1}^{-1} c_1 e_{-\nu} + c_{-1}^{-1} c_1 c_{-1}^{-1} c_0 + (c_{-1}^{-1} c_1)^2 e_\alpha \\
 &\quad - \sum_{j=1}^{s+2} (-1)^j (c_{-1}^{-1} c_0)^j e_{(j-2)\nu-\alpha} - \sum_{j=1}^{s+2} (-1)^j (c_{-1}^{-1} c_0)^{j+1} e_{(j-1)\nu-\alpha} \\
 &\quad - \sum_{j=1}^{s+2} (-1)^j (c_{-1}^{-1} c_0)^j e_{(j-1)\nu} (c_{-1}^{-1} c_1) - e_{-\lambda-\nu} I - c_{-1}^{-1} c_0 e_{-\lambda} - c_{-1}^{-1} c_1 e_{-\nu} \\
 &= c_{-1}^{-1} c_1 c_{-1}^{-1} c_0 + (c_{-1}^{-1} c_1)^2 e_\alpha - \sum_{j=1}^{s+2} (-1)^j (c_{-1}^{-1} c_0)^j e_{(j-1)\nu} (c_{-1}^{-1} c_1) \\
 &\quad + (c_{-1}^{-1} c_0) e_{-\lambda} - (-1)^s (c_{-1}^{-1} c_0)^{s+3} e_{(s+1)\nu-\alpha} - (c_{-1}^{-1} c_0) e_{-\lambda}.
 \end{aligned}$$

Cancelling out the terms  $\pm(c_{-1}^{-1}c_0)e_{-\lambda}$  in the last expression, we see that this block belongs to  $AP^+$  as well.  $\square$

Formula (2.6) is a particular case of the transformation introduced in [2] for an arbitrary  $AP$  polynomial (not necessarily a trinomial)  $f$  with invertible Fourier coefficient corresponding to the leftmost point in  $\Omega(f) \cap (-\lambda, \lambda)$ . However, in [2] only the case of commuting coefficients was considered. Also, formulas (2.7) for a trinomial case are more explicit than the general formulas of [2].

The resulting matrix  $G_1$  in general has the same structure as the original matrix  $G$ :  $\Omega(f_1) \subset \{-\nu_1, 0, \alpha_1\}$ , where  $\alpha_1, \nu_1 > 0$ ,  $\alpha_1 + \nu_1 = \lambda_1$  and  $\beta_1 = \nu_1/\alpha_1$  is irrational together with  $\beta$ . In some instances, however,  $G_1$  may be easier to deal with. One such situation is discussed in the next theorem; other applications of Lemma 2.2 can be found in subsequent sections.

**Theorem 2.3.** *Let the matrix  $G$  be given by (1.4), (1.5) with  $c_{-1}$  invertible,  $c_0c_{-1}^{-1}$  nilpotent and having all Jordan cells of the size at most  $\lfloor \frac{\alpha}{\nu} \rfloor + 2$ . Then 1)  $G$  is  $AP_W$  factorable, and 2) its partial  $AP$  indices equal zero if and only if  $c_1$  is invertible.*

*Proof.* Due to Lemma 2.2, we may consider the matrix (2.4) instead of  $G$ . The conditions imposed on the Jordan structure of  $c_0c_{-1}^{-1}$  imply that  $(c_{-1}^{-1}c_0)^{s+2} = 0$ . Thus,  $f_1$  in (2.4) is in fact a binomial with  $\Omega(f_1) \subset \{-\nu_1, 0\}$ . According to Theorem 1.2, the matrix  $G_1$  is  $AP_W$  factorable, and its partial  $AP$  indices equal zero if and only if the constant term  $c_{-1}^{-1}c_1$  of  $f_1$  is invertible. The latter condition is equivalent to the invertibility of  $c_1$ .  $\square$

Recall now the duality between an  $AP$  factorization (1.2) of  $G_f$  and that of  $G_{f^*}$ :

$$(2.8) \quad G_{f^*} = (JG_-^*)\Lambda^*(G_+^*J),$$

where  $J = \begin{bmatrix} 0 & I \\ I & 0 \end{bmatrix}$ . From (2.8) and Theorem 2.3 follows

**Corollary 2.4.** *Let the matrix  $G$  be given by (1.4), (1.5) with  $c_1$  invertible,  $c_0c_1^{-1}$  nilpotent and having all Jordan cells of the size at most  $\lfloor \frac{\nu}{\alpha} \rfloor + 2$ . Then  $G$  is  $AP_W$  factorable, and its partial  $AP$  indices equal zero if and only if  $c_{-1}$  is invertible.*

Observe that the condition on the size of Jordan cells is satisfied automatically if  $m = 2$ . Hence, the following statement holds.

**Corollary 2.5.** *Let the matrix  $G$  be given by (1.4), (1.5) with  $m = 2$ , let one of the coefficients  $c_{\pm 1}$  be non-singular, and let the corresponding product  $c_0c_{\pm 1}^{-1}$  be nilpotent. Then 1)  $G$  is  $AP_W$  factorable, and 2) its partial  $AP$  indices equal zero if and only if the second of the coefficients  $c_{\pm 1}$  is invertible as well.*

### 3. MAIN RESULT

We now turn to matrices (1.4) with the off-diagonal block (1.5) having pairwise commuting coefficients  $c_{\pm 1}, c_0$ . The representation (1.6) is not unique, and we choose one with the maximal possible number  $r$  of diagonal blocks. Each triple  $\{c_{-1,k}, c_{0k}, c_{1k}\}$  is then *irreducible*, that is, does not allow a further reduction to a block diagonal form with the help of a common similarity. Of course, the commutativity property of  $\{c_{-1}, c_0, c_1\}$  is inherited by the triples  $\{c_{-1,k}, c_{0k}, c_{1k}\}$ .

The ambiguity of  $T$  also allows us, for each  $k = 1, \dots, r$ , to put one of the matrices  $c_{jk}$  (with our choice of  $j = 0, \pm 1$ ) in its Jordan canonical form. If, for a



given  $k$ , at least one of the matrices  $c_{jk}$  is *unicellular* (that is, its canonical Jordan form consists of only one cell), then for such a  $T$  all the matrices  $c_{jk}$  with the same  $k$  automatically become upper triangular and, in addition, have a Toeplitz structure. The latter means that  $(p, q)$ -entry of each of the matrices  $c_{-1,k}, c_{0,k}, c_{1,k}$  is the same as its  $(p + 1, q + 1)$ -entry ( $p, q = 1, \dots, l_k - 1$ ). For  $l_k > 1$ , the common value of the entries right above the main diagonal in  $c_{jk}$  for such  $k$  will be denoted by  $\eta_{jk}$  (of course, the common value of the diagonal elements of the  $c_{jk}$  in this case is  $\xi_{jk}$ ).

With this notation at hand, we are ready to formulate our main result.

**Theorem 3.1.** *Let  $G$  be given by (1.4), (1.5) with pairwise commuting coefficients  $c_{\pm 1}, c_0$ . Suppose that in (1.6) for each  $k = 1, \dots, r$  at least one of the following conditions holds: 1)  $\xi_{0k} \neq 0$ , 2)  $\xi_{1,k}\xi_{-1,k} \neq 0$ , 3) one of the blocks  $c_{\pm 1,k}, c_{0k}$  is unicellular, 4)  $l_k \leq 3$ , 5)  $\xi_{1,k}$  or  $\xi_{-1,k}$  differs from zero and  $l_k \leq 4$ . Then  $G$  is not AP factorable if, for at least one value of  $k$ ,*

(3.1)

$$|\xi_{1,k}^\nu \xi_{-1,k}^\alpha| = |\xi_{0k}|^\lambda \neq 0, \text{ or } \xi_{-1,k} = \xi_{0k} = \xi_{1,k} = 0 \text{ and } |\eta_{1,k}^\nu \eta_{-1,k}^\alpha| = |\eta_{0k}|^\lambda \neq 0,$$

and is  $AP_W$  factorable otherwise.

*Proof.* Using (1.6), introduce a matrix

$$\begin{bmatrix} T^{-1} & 0 \\ 0 & T^{-1} \end{bmatrix} G \begin{bmatrix} T & 0 \\ 0 & T \end{bmatrix} = \begin{bmatrix} e_\lambda I_m & 0 \\ \text{diag}[c_{-1,k}e_{-\nu} + c_{0k} + c_{1,k}e_\alpha] & e_{-\lambda} I_m \end{bmatrix},$$

having the same factorization properties as  $G$ . By an appropriate permutation of its rows and columns, this matrix can be further rewritten as a direct sum of the blocks

$$G_k = \begin{bmatrix} e_\lambda I_{l_k} & 0 \\ c_{-1,k}e_{-\nu} + c_{0k} + c_{1,k}e_\alpha & e_{-\lambda} I_{l_k} \end{bmatrix},$$

$k = 1, \dots, r$ . Let  $R = \{1, \dots, r\}$  and denote by  $R_0$  the subset of those  $r \in R$  such that  $\xi_{1,k} = \xi_{-1,k} = \xi_{0k} = 0, l_k > 1$  and (at least) one of the blocks  $c_{\pm 1,k}, c_{0k}$  is unicellular. We now partition  $R$  into a disjoint union  $\bigcup_{j=1}^4 R_j$ , where

$$\begin{aligned} R_1 &= \{k : |\xi_{1,k}^\nu \xi_{-1,k}^\alpha| = |\xi_{0k}|^\lambda \neq 0\}, \\ R_2 &= \{k \in R_0 : |\eta_{1,k}^\nu \eta_{-1,k}^\alpha| = |\eta_{0k}|^\lambda \neq 0\}, \\ R_3 &= R_0 \setminus R_2, \\ R_4 &= R \setminus (R_1 \cup R_0). \end{aligned}$$

For every  $k \in R_0$ , yet another permutation of rows and columns allows us to represent  $G_k$  as a direct sum of  $\begin{bmatrix} e_\lambda & 0 \\ 0 & e_{-\lambda} \end{bmatrix}$  with

$$G'_k = \begin{bmatrix} e_\lambda I_{l_k-1} & 0 \\ c'_{-1,k}e_{-\nu} + c'_{0k} + c'_{1,k}e_\alpha & e_{-\lambda} I_{l_k-1} \end{bmatrix}.$$

Here  $c'_{jk}$  are obtained from  $c_{jk}$  by deleting its first column and last row. The Toeplitz structure of  $c_{jk}$  is inherited by  $c'_{jk}$ . In particular, the  $c'_{jk}$  pairwise commute and  $\sigma(c'_{jk}) = \{\eta_{jk}\}$  ( $j = 0, \pm 1; k \in R_0$ ).

Denote by  $G^{(1)}$  the direct sum of all the blocks  $G_k, k \in R_1$ , and  $G'_k, k \in R_2$ . Let  $G^{(2)}$  be a direct sum of all  $G_k$  ( $k \in R_4$ ),  $G'_k$  ( $k \in R_3$ ), and  $|R_2|$  copies of

the diagonal blocks  $\begin{bmatrix} e_\lambda & 0 \\ 0 & e_{-\lambda} \end{bmatrix}$ . Then  $G$  can be put in the form  $G^{(1)} \oplus G^{(2)}$  by an appropriate permutation of its rows and columns. In turn,  $G^{(1)}$  will become a permutation of a matrix of the type (1.4) with  $f = b_{-1}e_{-\nu} + b_0 + b_1e_\alpha$  and  $b_j = \left(\bigoplus_{k \in R_1} c_{jk}\right) \oplus \left(\bigoplus_{k \in R_2} c'_{jk}\right)$ .

In terms of the sets  $R_j$ , this theorem claims that  $G$  is  $AP_W$  factorable if  $R_1 \cup R_2 = \emptyset$ , and is not  $AP$  factorable otherwise. This follows from Lemma 2.1, provided that  $G^{(2)}$  is  $AP_W$  factorable and, for  $R_1 \cup R_2 \neq \emptyset$ ,  $G^{(1)}$  is not  $AP$  factorable. The latter statement holds due to Corollary 1.4. It remains to prove the former. We will do this by showing that each direct summand of  $G^{(2)}$  is  $AP_W$  factorable. There are five types of these summands:

(i) diagonal blocks  $\begin{bmatrix} e_\lambda & 0 \\ 0 & e_{-\lambda} \end{bmatrix}$ ,

and matrices (1.4) with  $f$  given by (1.5), pairwise commuting  $c_{\pm 1}, c_0$  (slightly abusing the notation, we again denote their size by  $m$ ), singleton spectra  $\sigma(c_j) = \{\xi_j\}$  ( $j = \pm 1, 0$ ) for which

- (ii)  $|\xi_1^\nu \xi_{-1}^\alpha| \neq |\xi_0|^\lambda$ ,
- (iii)  $\xi_0 = 0$ , exactly one of  $\xi_{\pm 1}$  differs from zero and (at least) one of the blocks  $c_{\pm 1}, c_0$  is unicellular,
- (iv)  $\xi_0 = 0$ , exactly one of  $\xi_{\pm 1}$  differs from zero, and  $m \leq 4$ ,
- (v)  $\xi_0 = \xi_1 = \xi_{-1} = 0$  and  $m \leq 3$ .

Indeed, the blocks  $G_k$  with  $k \in R_1$  have no impact on  $G^{(2)}$ ,  $k \in R_2$  generate only summands of type (i),  $k \in R_3$  yield summands of type (i) and (ii) or (iii), and  $k \in R_4$  produce summands of types (ii)-(v).

The summands of type (i) are trivially  $AP_W$  factorable (with partial  $AP$  indices  $\pm\lambda$ ). The summands of type (ii) are  $AP_W$  factorable (with zero partial  $AP$  indices) according to Theorem 1.3. It remains to consider matrices (1.4) of types (iii)-(v).

In cases (iii) and (iv) we may without loss of generality suppose that  $\xi_1 = 0$ ,  $\xi_{-1} \neq 0$ ; otherwise,  $G_{f^*}$  can be considered instead of  $G_f$ . If in addition,  $c_0 = 0$  or  $c_1 = 0$ , then  $f$  is a binomial and the corresponding matrix (1.4) is  $AP_W$  factorable due to Theorem 1.2. This happens, in particular, if  $m = 1$ .

If all three coefficients of  $f$  differ from zero, we consider the matrix (2.4). It can happen that  $c_0^{s+2} = 0$ , in which case the resulting block (2.5) is a binomial. Applying Theorem 1.2 and Lemma 2.2, we conclude that (2.4), and therefore (1.4), are  $AP_W$  factorable. If  $c_0^{s+2} \neq 0$ , we consider cases (iii) and (iv) separately.

(iii) The matrices  $c_j$  have an upper triangular Toeplitz structure which is inherited by the coefficients  $c_j^{(1)}$  of (2.5). Hence,

$$m > \text{rank } c_0^{(1)} = \text{rank } c_1$$

and

$$m > \text{rank } c_{-1}^{(1)} = \text{rank } c_0^{s+1} > \text{rank } c_1^{(1)} = \text{rank } c_0^{s+2} > 0.$$

Let  $q = \max\{\text{rank } c_0^{(1)}, \text{rank } c_{-1}^{(1)}\}$ ,  $p = m - q$ . Then both  $p$  and  $q$  are strictly positive. By a permutation of its rows and columns, the matrix  $G_1$  can be reduced to the form

$$(3.2) \quad \begin{bmatrix} e_\nu I_p & 0 \\ 0 & e_{-\nu} I_p \end{bmatrix} \oplus \begin{bmatrix} e_\nu I_q & 0 \\ f_2 & e_{-\nu} I_q \end{bmatrix},$$

where

$$(3.3) \quad f_2 = c_{-1}^{(2)}e_{-\nu_1} + c_0^{(2)} + c_1^{(2)}e_{\alpha_1}$$

and the matrices  $c_j^{(2)}$  are obtained from  $c_j^{(1)}$  by deleting their first  $p$  columns and last  $p$  rows. It suffices to prove now that the second direct summand in (3.2) is  $AP_W$  factorable.

If  $\text{rank } c_0^{(1)} \geq \text{rank } c_{-1}^{(1)}$ , this summand falls into type (ii). In the opposite case, this is again a matrix of type (iii), but its size is strictly smaller than that of the original matrix:  $q < m$ . By induction we now conclude that all matrices of type (iii) are  $AP_W$  factorable.

(iv) The case of unicellular  $c_0$  is covered by (iii). Since  $m \leq 4$  and  $c_0^{s+2} \neq 0$ , the only remaining case is  $s = 0$ ,  $m = 4$  and  $c_0$  consisting of one  $3 \times 3$  and one  $1 \times 1$  Jordan cell. The same Jordan structure is possessed by the matrix  $c_{-1}^{-1}c_0$ . Without loss of generality we may suppose that in (2.5)

$$(3.4) \quad c_{-1}^{(1)} = c_{-1}^{-1}c_0 = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Then

$$c_1^{(1)} = -(c_{-1}^{-1}c_0)^2 = \begin{bmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

The matrix  $c_0^{(1)} = c_{-1}^{-1}c_1$  is nilpotent and commutes with (3.4). Thus,

$$c_0^{(1)} = \begin{bmatrix} 0 & z & u & b \\ 0 & 0 & z & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & a & 0 \end{bmatrix}.$$

If  $a = b = 0$ , then the matrix  $G_1$  can be split into a direct sum of  $\begin{bmatrix} e_\nu I_2 & 0 \\ 0 & e_{-\nu} I_2 \end{bmatrix}$

and  $G_2 = \begin{bmatrix} e_\nu I_2 & 0 \\ f_2 & e_{-\nu} I_2 \end{bmatrix}$ , where  $f_2$  is given by (3.3) with

$$c_{-1}^{(2)} = I_2, \quad c_0^{(2)} = \begin{bmatrix} z & u \\ 0 & z \end{bmatrix}, \quad c_1^{(2)} = \begin{bmatrix} 0 & -1 \\ 0 & 0 \end{bmatrix}.$$

The matrix  $G_2$  is of type (ii) or (iii) (depending on whether or not  $z$  is zero), and therefore  $AP_W$  factorable. Of course,  $G_1$  is  $AP_W$  factorable together with  $G_2$ .

If  $a$  or  $b$  differs from zero, represent  $G_1$  as a direct sum of  $\text{diag}[e_\nu, e_{-\nu}]$  with  $G_3 = \begin{bmatrix} e_\nu I_3 & 0 \\ f_3 & e_{-\nu} I_3 \end{bmatrix}$ , where  $f_3 = c_{-1}^{(3)}e_{-\nu_1} + c_0^{(3)} + c_1^{(3)}e_{\alpha_1}$  and

$$c_{-1}^{(3)} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad c_0^{(3)} = \begin{bmatrix} z & u & b \\ 0 & z & 0 \\ 0 & a & 0 \end{bmatrix}, \quad c_1^{(3)} = \begin{bmatrix} 0 & -1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

The explicit  $AP_W$  factorization of  $G_3$  is shown in Appendix A of the supplement. Hence, all matrices of type (iv) are  $AP_W$  factorable.

Finally, consider the remaining type (v). If  $m \leq 2$ , then each matrix  $c_j$  either is unicellular or equals zero. In both cases, an  $AP_W$  factorization exists. Therefore, we may suppose that  $m = 3$ . Excluding another trivial case  $c_0 = 0$  (in which  $f$  is a binomial), we are left with the only possible Jordan structure of  $c_0$ : one  $2 \times 2$  and one  $1 \times 1$  block. Then, without loss of generality,

$$c_0 = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

The matrices  $c_{\pm 1}$  commute with  $c_0$  and are nilpotent. Therefore,

$$c_{\pm 1} = \begin{bmatrix} 0 & y_{\pm} & x_{\pm} \\ 0 & 0 & 0 \\ 0 & z_{\pm} & 0 \end{bmatrix}$$

The matrix  $G$  splits into a direct sum of  $\text{diag}[e_{\lambda}, e_{-\lambda}]$  and  $G_1 = \begin{bmatrix} e_{\lambda}I_2 & 0 \\ f_1 & e_{-\lambda}I_2 \end{bmatrix}$ , where  $f_1 = c_{-1}^{(1)}e_{-\nu} + c_0^{(1)} + c_1^{(1)}e_{\alpha}$ ,

$$c_0^{(1)} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad c_{\pm 1}^{(1)} = \begin{bmatrix} x_{\pm} & y_{\pm} \\ 0 & z_{\pm} \end{bmatrix}$$

From commutativity of  $c_1$  with  $c_{-1}$  it follows that  $x_+z_- = x_-z_+$ ; however, later on we will encounter a factorization problem for matrices  $G_1$  with  $c_{\pm 1}^{(1)}$  not satisfying this requirement. Therefore, we do not impose the condition  $x_+z_- = x_-z_+$  in our consideration.

The case  $x_+ = x_- = z_+ = z_- = 0$  is excluded because otherwise the triple  $\{c_{-1}, c_0, c_1\}$  would be reducible. The cases  $x_+z_+ \neq 0$  and  $x_-z_- \neq 0$  are covered by Corollary 2.5. In all the remaining cases an  $AP_W$  factorization of  $G_1$  also exists; it is constructed explicitly in Appendix B of the supplement. Hence, matrices  $G$  of type (v) are also  $AP_W$  factorable.  $\square$

As an application of Theorem 3.1, consider a difference equation

$$(3.5) \quad c_{-1}y(t - \nu) + c_0y(t) + c_1y(t + \alpha) = g(t) \text{ a.e. on } (0, \lambda),$$

where  $g$  is a given vector function in  $L^p(0, \lambda)$ ,  $y$  is an unknown vector function in  $L^p(\mathbb{R})$  with  $\text{supp } y \subset [0, \lambda]$ .

According to standard terminology, we say that (3.5) is normally solvable (in  $L^p$ ) if the set of vector functions  $g$  for which (3.5) has a solution is closed.

**Theorem 3.2.** *In (3.5) let  $\alpha + \nu = \lambda$ , let  $\frac{\alpha}{\nu} (> 0)$  be irrational, and let the coefficients  $c_j \in \mathbb{C}^{m \times m}$  satisfy the conditions of Theorem 3.1. Then the equation (3.5) is normally solvable if and only if, in the notation of Theorem 3.1, condition (3.1) fails for every  $k$ .*

This result does not depend on  $p \in (1, \infty)$ .

*Proof.* As follows from [7, Section 4.1], equation (3.5) is normally solvable if and only if the Wiener-Hopf operator  $W_G$ , the symbol  $G$  of which is given by (1.4), (1.5), has closed range in  $L^p(0, \infty)$ .

If condition (3.1) fails for all  $k$ , then the matrix function  $G$  is  $AP_W$  factorable due to Theorem 3.1. Hence,  $W_G$  has a generalized inverse, and therefore its range is closed.

To prove the converse statement, consider first a particular case when in (1.5) each matrix  $c_j$  has a singleton spectrum  $\{\xi_j\}$ , and

$$|\xi_1^\nu \xi_{-1}^\alpha| = |\xi_0|^\lambda \neq 0.$$

According to Theorem 3.1, the matrix function  $G$  in this case is not  $AP$  factorable.

If  $m = 1$ , the homogeneous equation (3.5) takes the form

$$y(t) = \begin{cases} -\frac{\xi_{-1}}{\xi_0} y(t - \nu) & \text{if } \nu < t < \lambda, \\ -\frac{\xi_1}{\xi_0} y(t + \alpha) & \text{if } 0 < t < \nu, \end{cases}$$

and has at most one linearly independent solution (see, for example, [4]).

For  $m > 1$ , a similarity can be used to put the  $c_j$  simultaneously in a triangular form, with  $\xi_j$  on the diagonal. Therefore, the number of linearly independent solutions of the respective homogeneous equation (3.5) is at most  $m$ . Suppose that this equation is normally solvable. Then the corresponding Wiener-Hopf operator  $W_G$  has a closed range and a finite dimensional kernel; in other words, it is  $n$ -normal. This property, as well as the *index*  $\text{ind } W_G$  of the operator  $W_G$  (the difference between the dimension of its kernel and the codimension of its range), is preserved under small perturbations. Consider such a small perturbation  $W_{G_{f'}}$  with  $f' = c_{-1}e_{-\nu} + (c_0 + \epsilon I) + c_1e_\alpha$ , and  $0 \neq |\xi_0 + \epsilon| \neq |\xi_0|$ . Then  $G' = G_{f'}$  admits an  $AP_W$  factorization with zero partial  $AP$  indices (Corollary 1.4), so that  $W_{G'}$  is invertible. Hence,  $\text{ind } W_G = \text{ind } W_{G'} = 0$ . From here it follows that  $\text{codim Im } W_G$  is finite together with  $\dim \text{Ker } W_G$ ; that is, the operator  $W_G$  is Fredholm. Since  $G \in AP_W$ , Theorem 2.5 of [7] implies that  $G$  is  $AP_W$  factorable. This contradiction shows that in fact the range  $\text{Im } W_G$  of the operator  $W_G$  is not closed.

Finally, consider the general case when (3.1) holds for some  $k$ . Then, as was shown in the proof of Theorem 3.1, the corresponding matrix  $G$  can be split into a direct sum of summands, a non-zero number of which are of the type just considered. Hence,  $W_G$  also splits into a direct sum of operators, some of which have a non-closed range. Therefore,  $\text{Im } W_G$  is not closed.  $\square$

*Remark.* The above reasoning shows that for matrix functions  $G$  satisfying the conditions of Theorem 3.1 the operator  $W_G$  has a closed range if and only if  $G$  is  $AP$  factorable. This is not true in general; examples of not  $AP$  factorable  $2 \times 2$  triangular matrix functions  $G \in AP_W$  for which  $\text{Im } W_G$  is closed can be found in [10].

#### 4. REMARKS ON $4 \times 4$ CASES

Theorem 3.1 covers all matrices (1.4), (1.5) with commuting  $c_j$  of size  $m \leq 3$ . Hence, the case of reducible  $4 \times 4$  triples is also covered. For irreducible  $\{c_{-1}, c_0, c_1\}$ , each  $c_j$  has a singleton spectrum, say  $\sigma(c_j) = \{\xi_j\}$ . The cases when at least one of the  $\xi_j$  differs from zero or  $c_j$  is unicellular also fall into the setting of Theorem 3.1.

This leaves us with the situation of an irreducible triple of  $4 \times 4$  nilpotent matrices  $c_j$  ( $j = 0, \pm 1$ ), none of which is unicellular. We may suppose in addition that none of them is diagonalizable (that is, has only  $1 \times 1$  Jordan cells). Indeed, a diagonalizable nilpotent matrix equals zero, and the corresponding  $G$  is then  $AP_W$  factorable due to Theorem 1.2. There remain three possible Jordan structures: two  $2 \times 2$  cells, one  $2 \times 2$  and two  $1 \times 1$  cells, and one  $3 \times 3$  and one  $1 \times 1$  cells.

The following example demonstrates why the case of two  $2 \times 2$  Jordan cells is hard to handle.

**Example.** Let  $c_j = \begin{bmatrix} 0 & c_j^{(0)} \\ 0 & 0 \end{bmatrix}$ , where the  $c_j^{(0)}$  are arbitrary (not necessarily commuting) non-singular  $2 \times 2$  matrices,  $j = \pm 1, 0$ . Then  $G$  can be split into a direct sum of  $\begin{bmatrix} e_\lambda I_2 & 0 \\ 0 & e_{-\lambda} I_2 \end{bmatrix}$  and  $G_0 = \begin{bmatrix} e_\lambda I_2 & 0 \\ c_{-1}^{(0)} e_{-\nu} + c_0^{(0)} + c_1^{(0)} e_\alpha & e_{-\lambda} I_2 \end{bmatrix}$ . According to Lemma 2.1, the matrices  $G$  and  $G_0$  are  $AP$  factorable only simultaneously. Hence, the  $AP$  factorization problem for  $G$  is reduced to the corresponding problem for matrices of the form (1.4) with non-commuting coefficients of  $f$ . Since the latter problem is still open, it is not surprising that a complete description of the  $AP$  factorability for matrices (1.4), (1.5) with commuting  $4 \times 4$  coefficients  $c_j$  is also missing.

We will now discuss the two remaining possibilities for the Jordan structure of  $c_0$ . First, let  $c_0$  consist of one  $2 \times 2$  and two  $1 \times 1$  Jordan cells. Without loss of generality,  $c_0$  itself is in a Jordan form:

$$(4.1) \quad c_0 = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

From the commutativity of  $c_{\pm 1}$  with  $c_0$  and their nilpotency it follows that

$$c_{\pm 1} = \begin{bmatrix} 0 & a_\pm & b_\pm & d_\pm \\ 0 & 0 & 0 & 0 \\ 0 & f_\pm & h_\pm & l_\pm \\ 0 & g_\pm & j_\pm & k_\pm \end{bmatrix},$$

where  $A_\pm = \begin{bmatrix} h_\pm & l_\pm \\ j_\pm & k_\pm \end{bmatrix}$  are themselves nilpotent.

We may also use a similarity to reduce  $A_+$  to its Jordan canonical form without disturbing  $c_0$  and the structure of  $A_-$ . Thus,  $h_+ = k_+ = j_+ = 0$  and  $l_+ = 0$  or 1.

If  $l_+ = 1$ , then commutativity of  $c_1$  with  $c_{-1}$  implies that  $h_- = k_- = j_- = 0$ . If  $l_+ = 0$  (that is,  $A_+ = 0$ ), then we can use a similarity to reduce  $A_-$  to its Jordan canonical form without changing  $c_0$  and  $A_+$ . Hence, in any case it may be supposed that  $h_\pm = k_\pm = j_\pm = 0$ , that is,

$$(4.2) \quad c_{\pm 1} = \begin{bmatrix} 0 & a_\pm & b_\pm & d_\pm \\ 0 & 0 & 0 & 0 \\ 0 & f_\pm & 0 & l_\pm \\ 0 & g_\pm & 0 & 0 \end{bmatrix}$$

Also, from commutativity of  $c_1$  with  $c_{-1}$  (which is preserved under the similarities applied above),

$$(4.3) \quad l_+ g_- = l_- g_+, \quad l_+ b_- = l_- b_+, \quad b_+ f_- + d_+ g_- = b_- f_+ + d_- g_+.$$

**Theorem 4.1.** *Let  $G$  be given by (1.4), (1.5) with  $c_0, c_{\pm 1}$  as in (4.1) and (4.2), respectively, satisfying (4.3) and forming an irreducible triple  $\{c_{-1}, c_0, c_1\}$ . Then  $G$  is not  $AP$  factorable if*

$$b_+ = b_- = g_+ = g_- = 0, \quad |D_-^\alpha D_+^\nu| = |l_+^\nu l_-^\alpha| \neq 0,$$

where

$$D_{\pm} = \det \begin{bmatrix} a_{\pm} & d_{\pm} \\ f_{\pm} & l_{\pm} \end{bmatrix} = a_{\pm}l_{\pm} - d_{\pm}f_{\pm},$$

and is  $AP_W$  factorable otherwise.

*Proof.* We need to show that  $G$  is  $AP_W$  factorable if

- i) at least one of the numbers  $b_{\pm}, d_{\pm}$  differs from zero, or
- ii)  $b_+ = b_- = g_+ = g_- = l_+l_-D_+D_- = 0$

and that in the case

$$\text{iii) } b_+ = b_- = g_+ = g_- = 0, l_{\pm}D_{\pm} \neq 0$$

it is  $AP$  ( $AP_W$ ) factorable if and only if

$$(4.4) \quad |D_-^{\alpha}D_+^{\nu}| \neq |l_+^{\nu}l_-^{\alpha}|.$$

In case i), rewrite  $G$  as a direct sum of  $\text{diag}[e_{\lambda}, e_{-\lambda}]$  and another matrix of the form (1.4), with  $m = 3$  and

$$c_{\pm 1} = \begin{bmatrix} a_{\pm} & b_{\pm} & d_{\pm} \\ f_{\pm} & 0 & l_{\pm} \\ g_{\pm} & 0 & 0 \end{bmatrix}, \quad c_0 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

If  $c_{-1}$  is invertible, that is,  $b_-g_-l_- \neq 0$ , then Lemma 2.2 can be used. A direct computation shows that

$$c_{-1}^{-1}c_0 = \begin{bmatrix} 0 & 0 & 0 \\ \frac{1}{b_-} & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

and therefore  $(c_{-1}^{-1}c_0)^2 = 0$ . Hence,  $f_1$  in (2.4) is at most a binomial, and the matrix  $G_1$  is  $AP_W$  factorable due to Theorem 1.2. The original matrix  $G$  is then also  $AP_W$  factorable.

Using (2.8) and appropriate transpositions of rows and columns, we can cover the case of invertible  $c_1$ , that is,  $b_+g_+l_+ \neq 0$ . It remains to construct an  $AP_W$  factorization in the cases when, in addition to (4.3),

$$(4.5) \quad b_+g_+l_+ = b_-g_-l_- = 0.$$

This is done in Appendix C.

In cases ii) and iii), we represent  $G$  as a direct sum of  $\begin{bmatrix} e_{\lambda}I_2 & 0 \\ 0 & e_{-\lambda}I_2 \end{bmatrix}$  and another matrix  $G_1$  of the form (1.4), (1.5) with  $m = 2$  and

$$c_{\pm 1}^{(1)} = \begin{bmatrix} a_{\pm} & d_{\pm} \\ f_{\pm} & l_{\pm} \end{bmatrix}, \quad c_0^{(1)} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

If  $l_+ = 0$  and  $d_+f_+ \neq 0$ , then the matrix  $G_1$  is  $AP_W$  factorable due to Corollary 2.5. The same reasoning applies if  $l_- = 0, d_-f_- \neq 0$ . The cases  $l_+ = l_- = d_+f_+ = d_-f_- = 0$  when not all of the four entries  $d_{\pm}, f_{\pm}$  equal zero are covered by Appendix B in the supplement. Observe that the case  $d_{\pm} = f_{\pm} = 0$  is excluded due to the irreducibility of the original triple  $\{c_{-1}, c_0, c_1\}$  given by (4.1), (4.2). Hence, the situation when  $l_+ = l_- = 0$  is covered completely.

In all other cases (when at least one of  $l_+, l_-$  differs from zero) we may use the symmetry (2.8) to suppose without loss of generality that, say,  $l_- \neq 0$ . An obvious

similarity performed on the original  $4 \times 4$  matrices  $c_{\pm 1}$  (and not changing  $c_0$ ) allows us to suppose in addition that  $d_- = f_- = 0$ . This similarity may, of course, change the values of  $a_{\pm}$  and  $d_+, f_+$ ; however,  $\det c_{\pm 1}^{(1)}$  remain the same, so that the new value of  $a_-$  is  $D_-/l_-$ . To simplify the notation, we redenote  $D_+$  by  $D$ .

If  $l_+ = 0$ , then  $d_+, f_+$  do not change under the above mentioned similarity. The only situation left uncovered by previous considerations is the case in which exactly one of  $d_+, f_+$  differs from zero.

In case ii), we are left with only two possibilities: 1)  $l_- \neq 0, l_+ = d_- = f_- = 0$ , exactly one of the entries  $d_+, f_+$  differs from zero, and 2)  $l_+l_- \neq 0, d_- = f_- = 0, a_-D = 0$ . Appendix D in the supplement shows that the corresponding matrix  $G_1$  (and therefore  $G$ ) is  $AP_W$  factorable.

In case iii), the additional condition  $d_- = f_- = 0$  means that  $a_- (= D_-/l_-) \neq 0$ , and (4.4) can be rewritten as

$$(4.6) \quad |a_-^\alpha D^\nu| \neq |l_+^\nu|.$$

A straightforward calculation shows that  $G_1 = X_+ G' X_-$ , where

$$X_+ = \begin{bmatrix} 1 & d_+l_-e_\lambda & 0 & 0 \\ -\frac{f_+}{l_+} & a_-l_+e_\lambda - l_-(e_\nu + a_-) & -e_\nu & \frac{f_+e_\nu}{l_+} \\ 0 & d_+(a_-l_+ + a_+l_-)e_\alpha & -d_+ & a_+ \\ 0 & (a_-l_+^2 + d_+f_+l_-)e_\alpha - l_-l_+ & -l_+ & f_+ \end{bmatrix}$$

is invertible in  $AP_W^+$ ,

$$X_- = \begin{bmatrix} 1 & -\frac{d_+l_-}{a_-l_+} & 0 & -\frac{d_+e_{-\alpha}}{a_-l_+} \\ 0 & \frac{1}{a_-l_+} & 0 & \frac{e_{-\alpha}}{a_-l_+l_-} \\ \frac{f_+(1+a_-e_{-\nu})}{D} & -\frac{a_+l_-(1+a_-e_{-\nu})}{a_-D} & \frac{f_+e_{-\lambda}}{D} & \frac{1}{l_-} - \frac{a_+(a_-e_{-\lambda}+e_{-\alpha})}{a_-D} \\ 0 & 0 & \frac{l_+}{D} & 0 \end{bmatrix}$$

is invertible in  $AP_W^-$ , and

$$G' = \begin{bmatrix} e_\lambda & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ a_-l_+e_{-\nu} + l_+ + De_\alpha & 0 & 0 & e_{-\lambda} \end{bmatrix}$$

can be split into a direct sum of  $I_2$  with

$$G_2 = \begin{bmatrix} e_\lambda & 0 \\ a_-l_+e_{-\nu} + l_+ + De_\alpha & e_{-\lambda} \end{bmatrix}$$

Of course,  $G_1$  is  $AP$  ( $AP_W$ ) factorable only simultaneously with  $G'$ , and in turn,  $G'$  has the same factorability properties as  $G_2$ . The latter matrix satisfies the conditions of Corollary 1.4 with  $m = 1$ . In the notation of this statement,  $\xi_{1,k} = D, \xi_{0k} = l_+$  and  $\xi_{-1,k} = a_-l_+$  with the only value of  $k$  ( $=1$ ), so that condition (1.7), necessary and sufficient for an  $AP$  ( $AP_W$ ) factorization to exist, is equivalent to (4.6).  $\square$

Finally, let  $c_0$  consist of one  $3 \times 3$  and one  $1 \times 1$  Jordan cells

$$c_0 = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$



Then the only possible form of  $c_{\pm 1}$  is

$$c_{\pm 1} = \begin{bmatrix} 0 & d_{\pm} & f_{\pm} & b_{\pm} \\ 0 & 0 & d_{\pm} & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & a_{\pm} & 0 \end{bmatrix},$$

where

$$(4.7) \quad a_+ b_- = a_- b_+.$$

The case  $a_+ = a_- = b_+ = b_- = 0$  is excluded if the triple  $\{c_{-1}, c_0, c_1\}$  is irreducible. Splitting  $G$  into a direct sum of  $\text{diag}[e_{\lambda}, e_{-\lambda}]$  and another matrix of the form (1.4), we may suppose that  $m = 3$  and

$$c_{\pm 1} = \begin{bmatrix} d_{\pm} & f_{\pm} & b_{\pm} \\ 0 & d_{\pm} & 0 \\ 0 & a_{\pm} & 0 \end{bmatrix}, \quad c_0 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

In the case when all four of the coefficients  $a_{\pm}$ ,  $b_{\pm}$  are different from zero, an  $AP_W$  factorization exists and can be explicitly constructed (see Appendix E in the supplement). Due to the commutativity condition (4.7), the number of non-zero entries among  $a_{\pm}$ ,  $b_{\pm}$  cannot equal one. However, there remain cases of exactly two or three non-zero numbers  $a_{\pm}$ ,  $b_{\pm}$ , and in these cases the  $AP$  factorability of the corresponding matrices  $G$  is still unknown.

#### REFERENCES

- [1] M. Bakonyi, L. Rodman, I. Spitkovsky, and H. Woerdeman, *Positive extensions of matrix functions of two variables with support in an infinite band*, C. R. Acad. Sci. Paris Sér. I Math. **323** (1996), no. 8, 859–863. MR **97i**:47023
- [2] M. A. Bastos, Yu. I. Karlovich, I. M. Spitkovsky, and P. M. Tishin, *On a new algorithm for almost periodic factorization*, Operator Theory: Advances and Applications **103** (1998), 53–74. CMP **98**:16
- [3] C. Corduneanu, *Almost periodic functions*, J. Wiley & Sons, 1968. MR **58**:2006
- [4] N. K. Karapetjanc and S. G. Samko, *The functional equation  $\psi(x + \alpha) - b(x)\psi(x) = g(x)$* , Izv. Akad. Nauk Armjan. SSR. Ser. Mat. **5** (1970), no. 5, 441–448. MR **44**:2101
- [5] Yu. I. Karlovich, *On the Haseman problem*, Demonstratio Math. **26** (1993), 581–595. MR **95a**:47048
- [6] Yu. I. Karlovich and I. M. Spitkovsky, *Factorization of almost periodic matrix-valued functions and the Noether theory for certain classes of equations of convolution type*, Mathematics of the USSR, Izvestiya **34** (1990), 281–316. MR **90f**:47034
- [7] ———, *(Semi)-Fredholmness of convolution operators on the spaces of Bessel potentials*, Operator Theory: Advances and Applications **71** (1994), 122–152. MR **95h**:47034
- [8] ———, *Almost periodic factorization: An analogue of Chebotarev's algorithm*, Contemporary Math. **189** (1995), 327–352. MR **96h**:47024
- [9] ———, *Factorization of almost periodic matrix functions*, J. Math. Anal. Appl. **193** (1995), 209–232. MR **96m**:47047
- [10] ———, *Semi-Fredholm properties of certain singular integral operators*, Operator Theory: Advances and Applications **90** (1996), 264–287. MR **97k**:47046
- [11] B. M. Levitan, *Almost periodic functions*, GITTL, Moscow, 1953 (in Russian). MR **15**:700a
- [12] B. M. Levitan and V. V. Zhikov, *Almost periodic functions and differential equations*, Cambridge University Press, 1982. MR **84g**:34004
- [13] G. S. Litvinchuk and I. M. Spitkovsky, *Factorization of measurable matrix functions*, Birkhäuser Verlag, Basel and Boston, 1987. MR **90g**:47030
- [14] Yu. Lyubarskii and I. Spitkovsky, *Sampling and interpolating for a lacunary spectrum*, Royal Society of Edinburgh, Proceedings **126A** (1996), 77–87. MR **97b**:41004

- [15] L. Rodman, I. M. Spitkovsky, and H. J. Woerdeman, *Carathéodory-Toeplitz and Nehari problems for matrix valued almost periodic functions*, Trans. Amer. Math. Soc. **350** (1998), 2185–2227. MR **98h**:47023
- [16] I. M. Spitkovsky, *On the factorization of almost periodic matrix functions*, Math. Notes **45** (1989), no. 5–6, 482–488. MR **90k**:47033
- [17] I. M. Spitkovsky and H. J. Woerdeman, *The Carathéodory-Toeplitz problem for almost periodic functions*, J. Functional Analysis **115** (1993), no. 2, 281–293. MR **94f**:47020

DEPARTMENT OF MATHEMATICS, THE COLLEGE OF WILLIAM AND MARY, WILLIAMSBURG, VA 23187-8795

*E-mail address:* `ilya@math.wm.edu`

DEPARTMENT OF APPLIED MATHEMATICS, UNIVERSITY OF WASHINGTON, SEATTLE, WA 98195

*E-mail address:* `dyong@u.washington.edu`