

**ERROR ESTIMATES
FOR THE FINITE ELEMENT APPROXIMATION
OF LINEAR ELASTIC EQUATIONS
IN AN UNBOUNDED DOMAIN**

HOUDE HAN AND WEIZHU BAO

ABSTRACT. In this paper we present error estimates for the finite element approximation of linear elastic equations in an unbounded domain. The finite element approximation is formulated on a bounded computational domain using a nonlocal approximate artificial boundary condition or a local one. In fact there are a family of nonlocal approximate boundary conditions with increasing accuracy (and computational cost) and a family of local ones for a given artificial boundary. Our error estimates show how the errors of the finite element approximations depend on the mesh size, the terms used in the approximate artificial boundary condition, and the location of the artificial boundary. A numerical example for Navier equations outside a circle in the plane is presented. Numerical results demonstrate the performance of our error estimates.

1. INTRODUCTION

Let Γ_i be a bounded simple closed curve in \mathbb{R}^2 and Ω be the unbounded domain with the boundary Γ_i (see Figure 1). We consider the following linear elastic problem:

$$(1.1) \quad -\mu \Delta u - (\lambda + \mu) \operatorname{grad} \operatorname{div} u = f \quad \text{in } \Omega,$$

$$(1.2) \quad u = 0 \quad \text{on } \Gamma_i,$$

$$(1.3) \quad u \text{ is bounded when } r = \sqrt{x_1^2 + x_2^2} \rightarrow +\infty,$$

where $x = (x_1, x_2)$ is the Cartesian coordinate system and the corresponding polar coordinate system is (r, θ) , $u = (u_1, u_2)^T$ is the displacement, $\lambda, \mu > 0$ are the Lamé constants and $f = (f_1, f_2)^T$ is the density of applied body force whose support is compact.

Since the problem (1.1)-(1.3) is defined in the unbounded domain Ω , in finding the numerical solution of this problem, it is often difficult to use the classical finite element method or finite difference method directly. In the last two decades, several methods were proposed to solve boundary value problems in unbounded domains

Received by the editor September 3, 1998 and, in revised form, September 8, 1999.

2000 *Mathematics Subject Classification*. Primary 65N30.

Key words and phrases. Unbounded domain, finite element approximation, artificial boundary, artificial boundary condition, linear elastic equations.

This work was supported partly by the Climbing Program of National Key Project of Foundation and the National Natural Science Foundation of China. Computation was supported by the State Key Laboratory of Scientific and Engineering Computing in China.

[6]. One of the most popular methods is to introduce an artificial boundary and set up artificial boundary conditions on it. Then the original problem is reduced to a boundary value problem in a bounded computational domain. Thus a numerical approximation of the original problem can be obtained by solving the reduced problem. In recent years many authors have worked on this subject for various problems by different techniques, see Engquist and Majda [3], Goldstein [11], Feng [4], Han and Wu [20, 21], Hagstrom and Keller [12, 13], Halpern and Schatzman [14], Han et al. [18, 19], Han and Bao [15, 16, 17], Givoli et al. [7, 8, 9], and the references therein.

In the above works, several authors also gave error estimates for the numerical solution, see [20, 21, 10]. But their error estimates only depend on the mesh size and the approximate artificial boundary condition. How the error depends on the location of the artificial boundary is unknown. But this is a very interesting problem for engineers. In this paper, we will discuss high-order local artificial boundary conditions and provide error estimates for the finite element approximation of the exterior problem (1.1)-(1.3). Our error estimates depend on not only the mesh size and the approximate artificial boundary condition but also the location of the artificial boundary.

The layout of this paper is as follows. In Section 2 we derive high-order local artificial boundary conditions at a given artificial boundary for the problem (1.1)-(1.3). In Section 3 we introduce the finite element formulation of the problem (1.1)-(1.3) in a bounded computational domain using an approximate nonlocal artificial boundary condition and prove an error estimate for the finite element approximation. In Section 4 we propose the finite element formulation of the problem (1.1)-(1.3) in a bounded computational domain using a high-order local artificial boundary condition and establish an error estimate for the finite element approximation. Finally in Section 5 we report on some numerical experiments.

2. HIGH-ORDER LOCAL ARTIFICIAL BOUNDARY CONDITIONS

In order to derive high-order local artificial boundary conditions, we recall here the derivation of the exact boundary condition at a given artificial boundary for the linear elastic problem (1.1)-(1.3) as described in [7, 21].

Introducing a circle Γ_e with radius R such that $\text{supp } f \subset B_R(0) := \{x \in \mathbb{R}^2 : |x| < R\}$, then Ω is divided into two parts: the unbounded part $\Omega_e := \Omega \setminus B_R(0)$




FIGURE 1

and the bounded part $\Omega_i := \Omega \setminus \bar{\Omega}_e$ (see Figure 1). The restriction of a solution u of the problem (1.1)-(1.3) to the unbounded domain Ω_e is then a solution of the following problem:

$$(2.1) \quad -\mu \Delta u - (\lambda + \mu) \operatorname{grad} \operatorname{div} u = 0 \quad \text{in } \Omega_e,$$

$$(2.2) \quad u|_{\Gamma_e} = u(R, \theta),$$

$$(2.3) \quad u \text{ is bounded when } r = \sqrt{x_1^2 + x_2^2} \rightarrow +\infty.$$

We know that the general solution of (2.1)-(2.3) is (see [21] for detail)

$$(2.4) \quad u_i(r, \theta) = (r^2 - R^2)W_i(r, \theta) + G_i(r, \theta) \quad R \leq r < +\infty \quad 0 \leq \theta \leq 2\pi \quad i = 1, 2,$$

where G_1, G_2, W_1 and W_2 are harmonic functions which satisfy

$$(2.5) \quad G_i(r, \theta) = \frac{a_0^i}{2} + \sum_{n=1}^{\infty} (a_n^i \cos n\theta + b_n^i \sin n\theta) \frac{R^n}{r^n} \\ R \leq r < +\infty \quad 0 \leq \theta \leq 2\pi \quad i = 1, 2,$$

$$(2.6) \quad W_i(r, \theta) = \sum_{n=3}^{\infty} (p_n^i \cos n\theta + q_n^i \sin n\theta) \frac{R^{n-2}}{r^n} \\ R \leq r < +\infty \quad 0 \leq \theta \leq 2\pi \quad i = 1, 2;$$

with $\kappa = \frac{\mu}{\lambda + \mu}$ and

$$(2.7) \quad a_n^i = \frac{1}{\pi} \int_0^{2\pi} G_i(R, \theta) \cos n\theta \, d\theta = \frac{1}{\pi} \int_0^{2\pi} u_i(R, \theta) \cos n\theta \, d\theta \quad i = 1, 2 \quad n \geq 0,$$

$$(2.8) \quad b_n^i = \frac{1}{\pi} \int_0^{2\pi} G_i(R, \theta) \sin n\theta \, d\theta = \frac{1}{\pi} \int_0^{2\pi} u_i(R, \theta) \sin n\theta \, d\theta \quad i = 1, 2 \quad n \geq 1;$$

$$(2.9) \quad p_n^1 = q_n^2 = \frac{n-2}{2+4\kappa} (a_{n-2}^1 - b_{n-2}^2), \quad q_n^1 = -p_n^2 = \frac{n-2}{2+4\kappa} (b_{n-2}^1 + a_{n-2}^2) \quad n \geq 3.$$

Let $\varepsilon(u) = (\varepsilon_{ij}(u))_{2 \times 2}$ and $\sigma(u) = (\sigma_{ij}(u))_{2 \times 2}$ be the strain and stress tensors, respectively, which satisfy

$$(2.10) \quad \varepsilon_{ij}(u) = \frac{1}{2} \left(\frac{\partial u_i}{\partial x_j} + \frac{\partial u_j}{\partial x_i} \right), \quad \sigma_{ij}(u) = \lambda \operatorname{div} u \, \delta_{ij} + 2\mu \varepsilon_{ij}(u) \quad 1 \leq i, j \leq 2.$$

Furthermore let $\sigma_n(u) = (\sigma_{n_1}(u), \sigma_{n_2}(u))^T$ be the normal stress corresponding to the displacement u at the artificial boundary Γ_e , say

$$(2.11) \quad \sigma_{n_i}(u) = \sigma_{i1}(u) \cos \theta + \sigma_{i2}(u) \sin \theta|_{\Gamma_e} \quad i = 1, 2.$$

Combining (2.4) with $r = R$, (2.10) and (2.11), a computation shows (see details in [21])

$$(2.12) \quad \sigma_n(u) = \begin{pmatrix} \sigma_{n_1}(u) \\ \sigma_{n_2}(u) \end{pmatrix} = \begin{pmatrix} \frac{\mu(2+2\kappa)}{1+2\kappa} \frac{\partial G_1}{\partial r} \Big|_{r=R} - \frac{2\mu\kappa}{(1+2\kappa)R} \frac{\partial G_2}{\partial \theta} \Big|_{r=R} \\ \frac{\mu(2+2\kappa)}{1+2\kappa} \frac{\partial G_2}{\partial r} \Big|_{r=R} + \frac{2\mu\kappa}{(1+2\kappa)R} \frac{\partial G_1}{\partial \theta} \Big|_{r=R} \end{pmatrix}.$$

We differentiate (2.5) with respect to r on noting (2.4) and (2.12) and set $r = R$ to obtain

$$(2.13) \quad \begin{aligned} \sigma_{n_1}(u) &= -\frac{2+2\kappa}{1+2\kappa} \frac{\mu}{\pi R} \sum_{n=1}^{\infty} n \int_0^{2\pi} u_1(R, \phi) \cos n(\theta - \phi) d\phi - \frac{2\kappa}{1+2\kappa} \frac{\mu}{R} \frac{\partial u_2(R, \theta)}{\partial \theta} \\ &= \frac{-1}{1+2\kappa} \frac{\mu}{\pi R} \sum_{n=1}^{\infty} n \int_0^{2\pi} [(2+2\kappa)u_1(R, \phi) \cos n(\theta - \phi) \\ &\quad - 2\kappa u_2(R, \theta) \sin n(\theta - \phi)] d\phi \\ &\equiv T_1(u), \end{aligned}$$

$$(2.14) \quad \begin{aligned} \sigma_{n_2}(u) &= \frac{2\kappa}{1+2\kappa} \frac{\mu}{R} \frac{\partial u_1(R, \theta)}{\partial \theta} - \frac{2+2\kappa}{1+2\kappa} \frac{\mu}{\pi R} \sum_{n=1}^{\infty} n \int_0^{2\pi} u_2(R, \phi) \cos n(\theta - \phi) d\phi \\ &= \frac{-1}{1+2\kappa} \frac{\mu}{\pi R} \sum_{n=1}^N n \int_0^{2\pi} [2\kappa u_1(R, \theta) \sin n(\theta - \phi) \\ &\quad + (2+2\kappa)u_2(R, \phi) \cos n(\theta - \phi)] d\phi \\ &\equiv T_2(u). \end{aligned}$$

This is the desired exact boundary condition at Γ_e for the problem (1.1)-(1.3). Thus the restriction of the solution u of the problem (1.1)-(1.3) to the bounded domain Ω_i is a solution of the following problem.

(P) Find u such that

$$(2.15) \quad -\mu \Delta u - (\lambda + \mu) \operatorname{grad} \operatorname{div} u = f \quad \text{in } \Omega_i,$$

$$(2.16) \quad u = 0 \quad \text{on } \Gamma_i,$$

$$(2.17) \quad \sigma_n(u) = (T_1(u), T_2(u))^T \quad \text{on } \Gamma_e.$$

Let

$$(2.18) \quad \begin{aligned} T_1^N(u) &= \frac{-1}{1+2\kappa} \frac{\mu}{\pi R} \sum_{n=1}^N n \int_0^{2\pi} [(2+2\kappa)u_1(R, \phi) \cos n(\theta - \phi) \\ &\quad - 2\kappa u_2(R, \theta) \sin n(\theta - \phi)] d\phi, \end{aligned}$$

$$(2.19) \quad \begin{aligned} T_2^N(u) &= \frac{-1}{1+2\kappa} \frac{\mu}{\pi R} \sum_{n=1}^N n \int_0^{2\pi} [2\kappa u_1(R, \theta) \sin n(\theta - \phi) \\ &\quad + (2+2\kappa)u_2(R, \phi) \cos n(\theta - \phi)] d\phi. \end{aligned}$$

Then we obtain a series of approximate artificial boundary conditions at Γ_e

(2.20)

$$\sigma_n(u) = T^N(u) \equiv (T_1^N(u), T_2^N(u))^T \quad \text{on } \Gamma_e \quad N = 0, 1, 2, \dots,$$

where $T^0(u) = (0, 0)^T$ is the stress free boundary condition which is often used in engineering literature. Then the original problem (1.1)-(1.3) can be reduced to the following problem defined on the bounded domain Ω_i approximately for $N = 0, 1, 2, \dots$

(P_N) Find u_N such that

(2.21)
$$-\mu \Delta u_N - (\lambda + \mu) \text{grad div } u_N = f \quad \text{in } \Omega_i,$$

(2.22)
$$u_N = 0 \quad \text{on } \Gamma_i,$$

(2.23)
$$\sigma_n(u_N) = T^N(u_N) \quad \text{on } \Gamma_e.$$

Now we discuss high-order local artificial boundary conditions at Γ_e for the problem (1.1)-(1.3). We consider a solution u of the problem (1.1)-(1.3), which consists of the first N harmonics at Γ_e . Thus we assume

(2.24)
$$u_i(R, \theta) = \frac{a_0^i}{2} + \sum_{n=1}^N (a_n^i \cos n\theta + b_n^i \sin n\theta) \quad i = 1, 2,$$

where the a_n^1, b_n^1, a_n^2 and b_n^2 are constants (Fourier coefficients, see (2.7) and (2.8)). Substituting (2.24) into (2.13) and (2.14), we get

(2.25)

$$\left\{ \begin{aligned} \sigma_{n_1}(u) &= -\frac{2 + 2\kappa \mu}{1 + 2\kappa R} \sum_{n=1}^N n (a_n^1 \cos n\theta + b_n^1 \sin n\theta) - \frac{2\kappa \mu}{1 + 2\kappa R} \frac{\partial u_2(R, \theta)}{\partial \theta}, \\ \sigma_{n_2}(u) &= \frac{2\kappa \mu}{1 + 2\kappa R} \frac{\partial u_1(R, \theta)}{\partial \theta} - \frac{2 + 2\kappa \mu}{1 + 2\kappa R} \sum_{n=1}^N n (a_n^2 \cos n\theta + b_n^2 \sin n\theta). \end{aligned} \right.$$

It is desired to find a linear differential operator L_N which does not depend on n , such that

(2.26)

$$L_N[1] = 0 \quad L_N[\cos n\theta] = n \cos n\theta \quad L_N[\sin n\theta] = n \sin n\theta \quad n = 1, 2, \dots, N.$$

With such an operator at hand, noting (2.24), then (2.25) can be written

(2.27)

$$\left\{ \begin{aligned} \sigma_{n_1}(u) &= -\frac{2 + 2\kappa \mu}{1 + 2\kappa R} L_N[u_1(R, \theta)] - \frac{2\kappa \mu}{1 + 2\kappa R} \frac{\partial u_2(R, \theta)}{\partial \theta}, \\ \sigma_{n_2}(u) &= \frac{2\kappa \mu}{1 + 2\kappa R} \frac{\partial u_1(R, \theta)}{\partial \theta} - \frac{2 + 2\kappa \mu}{1 + 2\kappa R} L_N[u_2(R, \theta)]. \end{aligned} \right.$$

The equality (2.27) is a local boundary condition at Γ_e which is exact for all solutions consisting of at most the first N harmonics at Γ_e . Noting the fact that

(2.28)

$$\frac{d^{2m}}{d\theta^{2m}} \cos n\theta = (-1)^m n^{2m} \cos n\theta, \quad \frac{d^{2m}}{d\theta^{2m}} \sin n\theta = (-1)^m n^{2m} \sin n\theta$$

$$m \geq 0 \quad n \geq 0,$$

TABLE 1. The coefficients $\alpha_m^{(N)}$ in the first five local artificial boundary conditions

	$\alpha_1^{(N)}$	$\alpha_2^{(N)}$	$\alpha_3^{(N)}$	$\alpha_4^{(N)}$	$\alpha_5^{(N)}$
$N = 1$	1				
$N = 2$	7/6	-1/6			
$N = 3$	74/60	-15/60	1/60		
$N = 4$	533/420	-43/144	11/360	-1/1008	
$N = 5$	3881/3780	-214/643	71/1728	-13/6048	1/25920

we can assume the operator L_N has the following form:

$$(2.29) \quad L_N[u(R, \theta)] = \sum_{m=1}^N (-1)^m \alpha_m^{(N)} \frac{\partial^{2m}}{\partial \theta^{2m}} u(R, \theta).$$

Inserting (2.29) into (2.27), noting (2.25) and (2.24), we obtain

$$(2.30) \quad \sum_{m=1}^N n^{2m} \alpha_m^{(N)} = n \quad n = 1, 2, \dots, N.$$

It is straightforward to check that the linear system (2.30) has a unique solution for any $n \in \mathbb{N}$. Table 1 shows the coefficients $\alpha_m^{(N)}$ in the first five local artificial boundary conditions. In the paper [25] by A. Sidi, a similar linear system was discussed. Combining (2.27) and (2.29), we get high-order local artificial boundary conditions at Γ_e for the problem (1.1)-(1.3):

$$(2.31) \quad \sigma_n(u) = \tilde{T}^N(u) \equiv \left(\tilde{T}_1^N(u), \tilde{T}_2^N(u) \right)^T \quad N = 1, 2, \dots,$$

where

$$(2.32) \quad \left\{ \begin{aligned} \tilde{T}_1^N(u) &= -\frac{2 + 2\kappa \mu}{1 + 2\kappa R} \sum_{m=1}^N (-1)^m \alpha_m^{(N)} \frac{\partial^{2m} u_1(R, \theta)}{\partial \theta^{2m}} - \frac{2\kappa \mu}{1 + 2\kappa R} \frac{\partial u_2(R, \theta)}{\partial \theta}, \\ \tilde{T}_2^N(u) &= \frac{2\kappa \mu}{1 + 2\kappa R} \frac{\partial u_1(R, \theta)}{\partial \theta} - \frac{2 + 2\kappa \mu}{1 + 2\kappa R} \sum_{m=1}^N (-1)^m \alpha_m^{(N)} \frac{\partial^{2m} u_2(R, \theta)}{\partial \theta^{2m}}. \end{aligned} \right.$$

Then the original problem (1.1)-(1.3) can be reduced to the following problem defined on the bounded domain Ω_i approximately for $N = 1, 2, \dots$.
 (\tilde{P}_N) Find \tilde{u}_N such that

$$(2.33) \quad -\mu \Delta \tilde{u}_N - (\lambda + \mu) \text{grad div } \tilde{u}_N = f \quad \text{in } \Omega_i,$$

$$(2.34) \quad \tilde{u}_N = 0 \quad \text{on } \Gamma_i,$$

$$(2.35) \quad \sigma_n(\tilde{u}_N) = \tilde{T}^N(\tilde{u}_N) \quad \text{on } \Gamma_e.$$

3. THE ERROR ESTIMATES FOR THE CASE OF USING NONLOCAL ARTIFICIAL BOUNDARY CONDITIONS

In the work [21], the authors have already given error estimates for the finite element approximation of the problem (P_N) . But from their estimates, we don't know how the errors depend on the location of the artificial boundary. In this section, we will present new error estimates for the finite element approximation of problems (P_N) . These error estimates depend not only the mesh size and the approximate artificial boundary condition but also the location of the artificial boundary. This kind of error estimate is very useful in engineering applications.

Let $H^m(\Omega_i)$ and $H^s(\Gamma_e)$ be the usual Sobolev spaces on the domain Ω_i and the boundary Γ_e with integer m and real number s . Suppose

$$V = \{v = (v_1, v_2)^T \in H^1(\Omega_i) \times H^1(\Omega_i) \mid v|_{\Gamma_i} = 0\}.$$

Then the boundary value problem (P) is equivalent to the following variational problem.

(VP) Find $u \in V$ such that

$$(3.1) \quad a(u, v) + b(u, v) = f(v) \quad \forall v \in V,$$

where

$$(3.2)$$

$$a(u, v) = \int_{\Omega_i} \left[\lambda \operatorname{div} u \operatorname{div} v + 2\mu \sum_{i,j=1}^2 \varepsilon_{ij}(u) \varepsilon_{ij}(v) \right] dx$$

$$\equiv \int_{\Omega_i} [\lambda \operatorname{div} u \operatorname{div} v + 2\mu \varepsilon(u) : \varepsilon(v)] dx \quad \forall u, v \in V,$$

$$(3.3)$$

$$b(u, v) = \frac{2 + 2\kappa}{1 + 2\kappa} \frac{\mu}{\pi} \sum_{n=1}^{\infty} n \int_0^{2\pi} \int_0^{2\pi} [u_1(R, \phi)v_1(R, \theta) + u_2(R, \phi)v_2(R, \theta)] \cos n(\theta - \phi) d\theta d\phi$$

$$+ \frac{2\kappa}{1 + 2\kappa} \frac{\mu}{\pi} \sum_{n=1}^{\infty} n \int_0^{2\pi} \int_0^{2\pi} [u_1(R, \phi)v_2(R, \theta) - u_2(R, \phi)v_1(R, \theta)] \sin n(\theta - \phi) d\theta d\phi$$

$$= \frac{2 + 2\kappa}{1 + 2\kappa} \frac{\mu}{\pi} \sum_{n=1}^{\infty} \int_0^{2\pi} \int_0^{2\pi} \left[\frac{\partial u_1(R, \phi)}{\partial \phi} \frac{\partial v_1(R, \theta)}{\partial \theta} + \frac{\partial u_2(R, \phi)}{\partial \phi} \frac{\partial v_2(R, \theta)}{\partial \theta} \right] \cdot \frac{\cos n(\theta - \phi)}{n} d\theta d\phi$$

$$+ \frac{2\kappa}{1 + 2\kappa} \frac{\mu}{\pi} \sum_{n=1}^{\infty} \int_0^{2\pi} \int_0^{2\pi} \left[\frac{\partial u_1(R, \phi)}{\partial \phi} \frac{\partial v_2(R, \theta)}{\partial \theta} - \frac{\partial u_2(R, \phi)}{\partial \phi} \frac{\partial v_1(R, \theta)}{\partial \theta} \right] \cdot \frac{\sin n(\theta - \phi)}{n} d\theta d\phi \quad \forall u, v \in V,$$

$$(3.4)$$

$$f(v) = \int_{\Omega_i} f \cdot v dx \quad \forall v \in V.$$

Furthermore let

(3.5)

$$\begin{aligned}
 b_N(u, v) = & \frac{2 + 2\kappa}{1 + 2\kappa} \frac{\mu}{\pi} \sum_{n=1}^N n \int_0^{2\pi} \int_0^{2\pi} [u_1(R, \phi)v_1(R, \theta) \\
 & + u_2(R, \phi)v_2(R, \theta)] \cos n(\theta - \phi) d\theta d\phi \\
 & + \frac{2\kappa}{1 + 2\kappa} \frac{\mu}{\pi} \sum_{n=1}^N n \int_0^{2\pi} \int_0^{2\pi} [u_1(R, \phi)v_2(R, \theta) \\
 & - u_2(R, \phi)v_1(R, \theta)] \sin n(\theta - \phi) d\theta d\phi \\
 & \forall u, v \in V.
 \end{aligned}$$

Then the boundary value problem (P_N) is equivalent to the following variational problem.

(VP_N) Find $u_N \in V$ such that

(3.6)
$$a(u_N, v) + b_N(u_N, v) = f(v) \quad \forall v \in V.$$

If we replace V by its finite element subspace V^h in which h represents the mesh size [11], then the finite element approximation of the problem (VP_N) is as follows.

(VP_N^h) Find $u_N^h \in V^h$ such that

(3.7)
$$a(u_N^h, v^h) + b_N(u_N^h, v^h) = f(v^h) \quad \forall v^h \in V^h.$$

We note that the symmetric bilinear form $a(\cdot, \cdot)$ is bounded and coercive on $V \times V$ from the K orn inequality [23] and Poincar e inequality [1], i.e., there exist positive constants M_1, M_2 such that

(3.8)
$$|a(u, v)| \leq M_1 \|u\|_V \cdot \|v\|_V \quad \forall u, v \in V,$$

(3.9)
$$M_2 \|v\|_V^2 \leq a(v, v) \quad \forall v \in V.$$

Thus we can define an equivalent norm on the space V :

(3.10)
$$\|v\|_* = [a(v, v)]^{1/2} \quad \forall v \in V.$$

Therefore we have that

(3.11)
$$|a(u, v)| \leq \|u\|_* \cdot \|v\|_* \quad \forall u, v \in V,$$

(3.12)
$$\|v\|_*^2 \leq a(v, v) \quad \forall v \in V.$$

For the symmetric bilinear forms $b(\cdot, \cdot)$ and $b_N(\cdot, \cdot)$, we have that

Lemma 3.1. *The following inequality holds:*

(3.13)
$$0 \leq b_N(v, v) \leq b(v, v) \leq a(v, v) \equiv \|v\|_*^2 \quad \forall v \in V \quad N \geq 0,$$

(3.14)
$$|b(u, v)| \leq \|u\|_* \cdot \|v\|_*, \quad |b_N(u, v)| \leq \|u\|_* \cdot \|v\|_* \quad \forall u, v \in V \quad N \geq 0.$$

where $b(u, v)$ and $b_N(u, v)$ are defined in (3.3) and (3.5), respectively.

Proof. For any given $u, v \in V$, we formally expand $u|_{\Gamma_e} = u(R, \theta)$ and $v|_{\Gamma_e} = v(R, \theta)$ in Fourier series, i.e.,

(3.15)
$$u_i(R, \theta) = \frac{a_i^i}{2} + \sum_{n=1}^{\infty} (a_n^i \cos n\theta + b_n^i \sin n\theta) \quad i = 1, 2,$$

(3.16)
$$v_i(R, \theta) = \frac{c_i^i}{2} + \sum_{n=1}^{\infty} (c_n^i \cos n\theta + d_n^i \sin n\theta) \quad i = 1, 2,$$

where a_n^i, b_n^i are defined in (2.7) and (2.8), and

$$(3.17) \quad c_n^i = \frac{1}{\pi} \int_0^{2\pi} v_i(R, \theta) \cos n\theta \, d\theta, \quad d_n^i = \frac{1}{\pi} \int_0^{2\pi} v_i(R, \theta) \sin n\theta \, d\theta, \quad i=1, 2 \quad n \geq 0.$$

Inserting (3.15) and (3.16) into (3.3) and (3.5), we get

$$(3.18) \quad b(u, v) = \frac{2\mu\pi}{1 + 2\kappa} \sum_{n=1}^{\infty} n \left[\sum_{i=1}^2 (a_n^i c_n^i + b_n^i d_n^i) + \kappa(a_n^1 + b_n^2)(c_n^1 + d_n^2) + \kappa(b_n^1 - a_n^2)(d_n^1 - c_n^2) \right],$$

$$(3.19) \quad b_N(u, v) = \frac{2\mu\pi}{1 + 2\kappa} \sum_{n=1}^N n \left[\sum_{i=1}^2 (a_n^i c_n^i + b_n^i d_n^i) + \kappa(a_n^1 + b_n^2)(c_n^1 + d_n^2) + \kappa(b_n^1 - a_n^2)(d_n^1 - c_n^2) \right].$$

We denote Q the domain enclosed by Γ_i , and $\hat{\Omega}$ the disk bounded by Γ_e (i.e., $\hat{\Omega} = Q \cup \Omega_i \cup \Gamma_i$) (see Figure 1). Then for any $v \in V$, we define the function $v^{(0)}$ which satisfies

$$(3.20) \quad -\mu \Delta v^{(0)} - (\lambda + \mu) \operatorname{grad} \operatorname{div} v^{(0)} = 0 \quad \text{in } \Omega_i,$$

$$(3.21) \quad v^{(0)} = v \quad \text{on } \Gamma_e \cup \Gamma_i.$$

$$(3.22) \quad v^{(0)} \equiv 0 \quad \text{in } Q.$$

Then we know $v^{(0)} \in [H^1(\hat{\Omega})]^2$. We also define the function $v^{(1)} \in [H^1(\hat{\Omega})]^2$ which satisfies

$$(3.23) \quad -\mu \Delta v^{(1)} - (\lambda + \mu) \operatorname{grad} \operatorname{div} v^{(1)} = 0 \quad \text{in } \hat{\Omega},$$

$$(3.24) \quad v^{(1)} = v \quad \text{on } \Gamma_e.$$

Then $v^{(0)}|_{\Omega_i}$ minimizes the functional $\int_{\Omega_i} [\lambda(\operatorname{div} w)^2 + 2\mu|\varepsilon(w)|^2] \, dx$ among all functions $w \in [H^1(\Omega_i)]^2$ which are equal to v on $\Gamma_e \cup \Gamma_i$. Similarly, $v^{(1)}$ minimizes the functional $\int_{\hat{\Omega}} [\lambda(\operatorname{div} w)^2 + 2\mu|\varepsilon(w)|^2] \, dx$ among all functions $w \in [H^1(\hat{\Omega})]^2$ which are equal to v on Γ_e . Therefore we have that

$$(3.25) \quad \begin{aligned} a(v, v) &= \int_{\Omega_i} [\lambda(\operatorname{div} v)^2 + 2\mu|\varepsilon(v)|^2] \, dx \geq a(v^{(0)}, v^{(0)}) \\ &= \int_{\hat{\Omega}} [\lambda(\operatorname{div} v^{(0)})^2 + 2\mu|\varepsilon(v^{(0)})|^2] \, dx \\ &\geq \int_{\hat{\Omega}} [\lambda(\operatorname{div} v^{(1)})^2 + 2\mu|\varepsilon(v^{(1)})|^2] \, dx. \end{aligned}$$

Recalling that $v^{(1)}$ is a solution of the problem (3.23)-(3.24), using separation of variables and noting $v^{(1)}|_{\Gamma_e} = v^{(1)}(R, \theta) = v(R, \theta)$, we obtain

(3.26)

$$v_1^{(1)}(r, \theta) = \frac{R^2 - r^2}{2 + 4\kappa} \sum_{n=0}^{\infty} (n + 2) [(c_{n+2}^1 - d_{n+2}^2) \cos n\theta + (d_{n+2}^1 + c_{n+2}^2) \sin n\theta] \frac{r^n}{R^{n+2}} + \frac{c_0^1}{2} + \sum_{n=1}^{\infty} \frac{r^n}{R^n} (c_n^1 \cos n\theta + d_n^1 \sin n\theta) \quad 0 \leq r \leq R \quad 0 \leq \theta \leq 2\pi,$$

(3.27)

$$v_2^{(1)}(r, \theta) = \frac{R^2 - r^2}{2 + 4\kappa} \sum_{n=0}^{\infty} (n + 2) [-(d_{n+2}^1 + c_{n+2}^2) \cos n\theta + (c_{n+2}^1 - d_{n+2}^2) \sin n\theta] \frac{r^n}{R^{n+2}} + \frac{c_0^2}{2} + \sum_{n=1}^{\infty} \frac{r^n}{R^n} (c_n^2 \cos n\theta + d_n^2 \sin n\theta) \quad 0 \leq r \leq R \quad 0 \leq \theta \leq 2\pi,$$

where

(3.28)

$$c_n^i = \frac{1}{\pi} \int_0^{2\pi} v_i^{(1)}(R, \theta) \cos n\theta \, d\theta = \frac{1}{\pi} \int_0^{2\pi} v_i(R, \theta) \cos n\theta \, d\theta \quad i = 1, 2 \quad n \geq 0,$$

(3.29)

$$d_n^i = \frac{1}{\pi} \int_0^{2\pi} v_i^{(1)}(R, \theta) \sin n\theta \, d\theta = \frac{1}{\pi} \int_0^{2\pi} v_i(R, \theta) \sin n\theta \, d\theta \quad i = 1, 2 \quad n \geq 1.$$

Combining (3.25), (3.26) and (3.27) with $r = R$, (3.28), (3.29), (2.10) and (2.11) with $u = v^{(1)}$, (3.18) and (3.19) with $u = v$, and integration by parts, we obtain

(3.30)

$$\begin{aligned} a(v, v) &\geq \int_{\Omega} [\lambda(\operatorname{div} v^{(1)})^2 + 2\mu|\varepsilon(v^{(1)})|^2] \, dx = \int_{\Gamma_e} \sigma_n(v^{(1)}) \cdot v^{(1)} \, ds \\ &= \frac{2 + 2\kappa}{1 + 2\kappa} \frac{\mu}{\pi} \sum_{n=1}^{\infty} n \int_0^{2\pi} \int_0^{2\pi} [v_1^{(1)}(R, \phi)v_1^{(1)}(R, \theta) + v_2^{(1)}(R, \phi)v_2^{(1)}(R, \theta)] \cos n(\theta - \phi) \, d\theta \, d\phi \\ &\quad + \frac{2\kappa}{1 + 2\kappa} \frac{\mu}{\pi} \sum_{n=1}^{\infty} n \int_0^{2\pi} \int_0^{2\pi} [v_1^{(1)}(R, \phi)v_2^{(1)}(R, \theta) - v_2^{(1)}(R, \phi)v_1^{(1)}(R, \theta)] \sin n(\theta - \phi) \, d\theta \, d\phi \\ &= b(v^{(1)}, v^{(1)}) = b(v, v) \\ &= \frac{2\mu\pi}{1 + 2\kappa} \sum_{n=1}^{\infty} n [(c_n^1)^2 + (d_n^1)^2 + (c_n^2)^2 + (d_n^2)^2 + \kappa(c_n^1 + d_n^2)^2 + \kappa(d_n^1 - c_n^2)^2] \\ &\geq \frac{2\mu\pi}{1 + 2\kappa} \sum_{n=1}^N n [(c_n^1)^2 + (d_n^1)^2 + (c_n^2)^2 + (d_n^2)^2 + \kappa(c_n^1 + d_n^2)^2 + \kappa(d_n^1 - c_n^2)^2] \\ &= b_N(v, v) \geq 0. \end{aligned}$$

Then the desired inequality (3.13) is proved. Thus the inequality (3.14) follows from (3.13) and the Schwarz inequality immediately.

It follows immediately from (3.11), (3.12) (3.14) and (3.13) that the variational problems (VP), (VP_N) and (VP_N^h) are well posed; that is, for $f \in V'$, the dual of V , there exists a unique $u \in V$ solving (VP), a unique $u_N \in V$ solving (VP_N), a unique $u_N^h \in V^h$ solving (VP_N^h), and

$$(3.31) \quad \|u\|_* + \|u_N\|_* + \|u_N^h\|_* \leq 3\|f\|_{V'}.$$

Note that the well-posedness of (VP) implies immediately the well-posedness of the original problem (1.1)-(1.3).

Let $R_0 = \max\{|x| : x \in \text{supp } f \cup \Gamma_i\}$, $\Gamma_0 = \{(R_0, \theta) : 0 \leq \theta \leq 2\pi\}$ and $\Omega_0 = \{x \in \Omega_i : |x| < R_0\}$ and $\Gamma_r = \{(r, \theta) : 0 \leq \theta \leq 2\pi\}$. We recall an equivalent definition of Sobolev space $H^s(\Gamma_r)$ for any real number s [22]:

$$w \in H^s(\Gamma_r) \iff w(r, \theta) = \frac{p_0}{2} + \sum_{m=1}^{\infty} (p_m \cos m\theta + q_m \sin m\theta)$$

and

$$\frac{\pi p_0^2}{2} + \sum_{m=1}^{\infty} \pi(1 + m^2)^s (p_m^2 + q_m^2) < \infty.$$

Thus we use

$$(3.32) \quad |w|_{s, \Gamma_r} = \left[\sum_{m=1}^{\infty} \pi m^{2s} (p_m^2 + q_m^2) \right]^{1/2}$$

as a semi-norm of the space $H^s(\Gamma_r)$. Then we have the following estimate.

Lemma 3.2. *Suppose $u \in V$ is a solution of the exterior problem (1.1)-(1.3) and there exists an integer $k \geq 1$ such that $u|_{\Gamma_0} \in [H^{k+\frac{1}{2}}(\Gamma_0)]^2$. Then we have that*

$$(3.33) \quad |b(u, v) - b_N(u, v)| \leq \frac{C_0}{(N + 1)^{k-1}} \left(\frac{R_0}{R} \right)^{\max\{1, N-1\}} |u|_{k+\frac{1}{2}, \Gamma_0} \cdot \|v\|_* \quad \forall v \in V,$$

where C_0 is a generic constant independent of u, N, h and R .

Proof. Assume that

$$(3.34) \quad u_i(R_0, \theta) = \frac{p_0^i}{2} + \sum_{n=1}^{\infty} (p_n^i \cos n\theta + q_n^i \sin n\theta) \quad i = 1, 2,$$

$$(3.35) \quad v_i(R, \theta) = \frac{c_0^i}{2} + \sum_{n=1}^{\infty} (c_n^i \cos n\theta + d_n^i \sin n\theta) \quad i = 1, 2,$$

where c_n^i and d_n^i are defined in (3.28) and (3.29), respectively, and

$$(3.36) \quad p_n^i = \frac{1}{\pi} \int_0^{2\pi} u_i(R_0, \theta) \cos n\theta \, d\theta, \quad q_n^i = \frac{1}{\pi} \int_0^{2\pi} u_i(R_0, \theta) \sin n\theta \, d\theta, \\ i = 1, 2 \quad n \geq 0.$$

Note that u satisfies the homogeneous Navier equations (say (1.1) with $f = 0$) in the domain $\{x : |x| > R_0\}$. By separation of variables, we get

$$\begin{aligned}
 (3.37) \quad u_1(r, \theta) &= \frac{r^2 - R_0^2}{2 + 4\kappa} \sum_{n=3}^{\infty} (n - 2) [(p_{n-2}^1 - q_{n-2}^2) \cos n\theta \\
 &\quad + (q_{n-2}^1 + p_{n-2}^2) \sin n\theta] \frac{R_0^{n-2}}{r^n} \\
 &\quad + \frac{p_0^1}{2} + \sum_{n=1}^{\infty} (p_n^1 \cos n\theta + q_n^1 \sin n\theta) \frac{R_0^n}{r^n} \quad R_0 \leq r \quad 0 \leq \theta \leq 2\pi,
 \end{aligned}$$

$$\begin{aligned}
 (3.38) \quad u_2(r, \theta) &= \frac{r^2 - R_0^2}{2 + 4\kappa} \sum_{n=3}^{\infty} (n - 2) [-(q_{n-2}^1 + p_{n-2}^2) \cos n\theta \\
 &\quad + (p_{n-2}^1 - q_{n-2}^2) \sin n\theta] \frac{R_0^{n-2}}{r^n} \\
 &\quad + \frac{p_0^2}{2} + \sum_{n=1}^{\infty} (p_n^2 \cos n\theta + q_n^2 \sin n\theta) \frac{R_0^n}{r^n} \quad R_0 \leq r \quad 0 \leq \theta \leq 2\pi.
 \end{aligned}$$

Setting $r = R$ in (3.37) and (3.38), we obtain

$$(3.39) \quad u_i(R, \theta) = \frac{a_0^i}{2} + \sum_{n=1}^{\infty} (a_n^i \cos n\theta + b_n^i \sin n\theta), \quad i = 1, 2,$$

where

$$(3.40) \quad a_n^1 = \left(\frac{R_0}{R}\right)^n \begin{cases} p_n^1 & n = 0, 1, 2, \\ p_n^1 + \frac{(n-2)(R^2 - R_0^2)}{(2+4\kappa)R_0^2} (p_{n-2}^1 - q_{n-2}^2) & n \geq 3; \end{cases}$$

$$(3.41) \quad b_n^1 = \left(\frac{R_0}{R}\right)^n \begin{cases} q_n^1 & n = 1, 2, \\ q_n^1 + \frac{(n-2)(R^2 - R_0^2)}{(2+4\kappa)R_0^2} (q_{n-2}^1 + p_{n-2}^2) & n \geq 3; \end{cases}$$

$$(3.42) \quad a_n^2 = \left(\frac{R_0}{R}\right)^n \begin{cases} p_n^2 & n = 0, 1, 2, \\ p_n^2 - \frac{(n-2)(R^2 - R_0^2)}{(2+4\kappa)R_0^2} (q_{n-2}^1 + p_{n-2}^2) & n \geq 3; \end{cases}$$

$$(3.43) \quad b_n^2 = \left(\frac{R_0}{R}\right)^n \begin{cases} q_n^2 & n = 1, 2, \\ q_n^2 + \frac{(n-2)(R^2 - R_0^2)}{(2+4\kappa)R_0^2} (p_{n-2}^1 - q_{n-2}^2) & n \geq 3. \end{cases}$$

From (3.39), (3.35), (3.18) and (3.19), noting (3.40)-(3.43), (3.36), (3.14) and (3.13), we obtain

$$\begin{aligned}
 (3.44) \quad & |b(u, v) - b_N(u, v)| \\
 &= \frac{2\mu\pi}{1 + 2\kappa} \left| \sum_{n=N+1}^{\infty} n \left[\sum_{i=1}^2 (a_n^i c_n^i + b_n^i d_n^i) + \kappa(a_n^1 + b_n^2)(c_n^1 + d_n^2) \right. \right. \\
 &\qquad \qquad \qquad \left. \left. + \kappa(b_n^1 - a_n^2)(d_n^1 - c_n^2) \right] \right| \\
 &\leq \frac{2\mu\pi}{1 + 2\kappa} \left[\sum_{n=N+1}^{\infty} n((a_n^1)^2 + (b_n^1)^2 + (a_n^2)^2 + (b_n^2)^2 \right. \\
 &\qquad \qquad \qquad \left. + \kappa(a_n^1 + b_n^2)^2 + \kappa(b_n^1 - a_n^2)^2) \right]^{1/2} \\
 &\quad \cdot \left[\sum_{n=N+1}^{\infty} n((c_n^1)^2 + (d_n^1)^2 + (c_n^2)^2 + (d_n^2)^2 + \kappa(c_n^1 + d_n^2)^2 + \kappa(c_n^1 - d_n^2)^2) \right]^{1/2} \\
 &\leq C_0 \left[\sum_{n=N+1}^{\infty} n \sum_{i=1}^2 ((p_n^i)^2 + (q_n^i)^2) \frac{R_0^{2n}}{R^{2n}} \right. \\
 &\qquad \qquad \qquad \left. + \sum_{n=\max\{1, N-1\}}^{\infty} n^3 \sum_{i=1}^2 ((p_n^i)^2 + (q_n^i)^2) \frac{R_0^{2n}}{R^{2n}} \right]^{1/2} \cdot \|v\|_* \\
 &\leq \frac{C_0}{(N + 1)^{k-1}} \left(\frac{R_0}{R} \right)^{\max\{1, N-1\}} |u|_{k+\frac{1}{2}, \Gamma_0} \cdot \|v\|_*.
 \end{aligned}$$

Combining Lemmas 3.1 and 3.2, we get the following error estimate.

Theorem 3.1. *Suppose u is the solution of the problem (1.1)-(1.3) and u_N^h is the solution of the problem (VP_N^h). Suppose $f \in [L^2(\Omega_i)]^2$ and $u|_{\Gamma_0} \in [H^{k+\frac{1}{2}}(\Gamma_0)]^2$ ($k \geq 1$). Then we have the following error estimate:*

$$(3.45) \quad \|u - u_N^h\|_* \leq C_0 \left[\inf_{v^h \in V^h} \|u - v^h\|_* + \frac{1}{(N + 1)^{k-1}} \left(\frac{R_0}{R} \right)^{\max\{1, N-1\}} |u|_{k+\frac{1}{2}, \Gamma_0} \right].$$

Proof. Let $e := u - u_N^h$, $e^v := v^h - u$ and $e^h := v^h - u_N^h$. Subtracting (3.7) from (3.1), we have that

$$(3.46) \quad a(e, v^h) + b_N(e, v^h) = b(u, v^h) - b_N(u, v^h) \quad \forall v^h \in V^h.$$

From (3.12), (3.11), (3.14), (3.13), (3.46) with $v^h = e^h$ and (3.33), we have that

$$\begin{aligned}
 (3.47) \quad \|e^h\|_*^2 &= a(e^h, e^h) \leq a(e^h, e^h) + b_N(e^h, e^h) \\
 &= a(e^v, e^h) + b_N(e^v, e^h) + a(e, e^h) + b_N(e, e^h) \\
 &= a(e^v, e^h) + b_N(e^v, e^h) + b_N(u, e^h) - b(u, e^h) \\
 &\leq 2\|e^v\|_* \cdot \|e^h\|_* + \frac{C_0}{(N+1)^{k-1}} \left(\frac{R_0}{R}\right)^{\max\{1, N-1\}} |u|_{k+\frac{1}{2}, \Gamma_0} \cdot \|e^h\|_*.
 \end{aligned}$$

Thus

$$(3.48) \quad \|e^h\|_* \leq C_0 \left[\|e^v\|_* + \frac{1}{(N+1)^{k-1}} \left(\frac{R_0}{R}\right)^{\max\{1, N-1\}} |u|_{k+\frac{1}{2}, \Gamma_0} \right] \quad \forall v^h \in V^h.$$

Then the desired result (3.45) follows from (3.48) and the triangle inequality.

If we suppose $u \in [H^{p+1}(\Omega_i)]^2$, $u|_{\Gamma_0} \in [H^{p+\frac{1}{2}}(\Gamma_0)]^2$, and the interpolation error of V^h approximate to V is [2]

$$(3.49) \quad \inf_{v^h \in V^h} \|u - v^h\|_V \leq C_0 h^p |u|_{p+1, \Omega_i},$$

then combining (3.49) and (3.45), noting the Körn inequality and Poincaré inequality, we get

$$\begin{aligned}
 (3.50) \quad \|u - u_N^h\|_{1, \Omega_0} &\leq C_0 |u - u_N^h|_{1, \Omega_0} \leq C_0 \int_{\Omega_0} [\lambda |\operatorname{div}(u - u_N^h)|^2 + 2\mu |\varepsilon(u - u_N^h)|^2] dx \\
 &\leq C_0 a(u - u_N^h, u - u_N^h) \\
 &\leq C_0 \left[h^p |u|_{p+1, \Omega_i} + \frac{1}{(N+1)^{p-1}} \left(\frac{R_0}{R}\right)^{\max\{1, N-1\}} |u|_{p+\frac{1}{2}, \Gamma_0} \right].
 \end{aligned}$$

4. THE ERROR ESTIMATES FOR THE CASE OF USING HIGH-ORDER LOCAL ARTIFICIAL BOUNDARY CONDITIONS

In this section, we will present the finite element formulation of the problem (\tilde{P}_N) and provide an error estimate for the finite element approximation. To cope with the high-order local artificial boundary condition (2.31), we define

$$\tilde{V} = \{v \in [H^1(\Omega_i)]^2 \mid v|_{\Gamma_e} \in [H^N(\Gamma_e)]^2 \quad v|_{\Gamma_i} = 0\}.$$

Let

$$\begin{aligned}
 (4.1) \quad \tilde{b}_N(u, v) &= - \int_{\Gamma_e} v \cdot \tilde{T}^N(u) ds \\
 &= \frac{\mu(2+2\kappa)}{1+2\kappa} \int_0^{2\pi} \sum_{m=1}^N \alpha_m^{(N)} \left[\frac{\partial^m u_1(R, \theta)}{\partial \theta^m} \frac{\partial^m v_1(R, \theta)}{\partial \theta^m} \right. \\
 &\quad \left. + \frac{\partial^m u_2(R, \theta)}{\partial \theta^m} \frac{\partial^m v_2(R, \theta)}{\partial \theta^m} \right] d\theta \\
 &\quad + \frac{2\mu\kappa}{1+2\kappa} \int_0^{2\pi} \left[\frac{\partial u_2(R, \theta)}{\partial \theta} v_1(R, \theta) - \frac{\partial u_1(R, \theta)}{\partial \theta} v_2(R, \theta) \right] d\theta \quad \forall u, v \in \tilde{V}.
 \end{aligned}$$

Then the weak form of the problem (\tilde{P}_N) is as follows.

(\tilde{VP}_N) Find $\tilde{u}_N \in \tilde{V}$ such that

$$(4.2) \quad a(\tilde{u}_N, v) + \tilde{b}_N(\tilde{u}_N, v) = f(v) \quad \forall v \in \tilde{V}.$$

If we replace \tilde{V} by its finite dimensional subspace, $\tilde{V}^h \subset \tilde{V}$ in which h is the mesh size [2] (a family of such subspaces were introduced by Givoli et al. [10]), then the finite element approximation of the problem (\tilde{VP}_N) is as follows.

(\tilde{VP}_N^h) Find $\tilde{u}_N^h \in \tilde{V}^h$ such that

$$(4.3) \quad a(\tilde{u}_N^h, v^h) + \tilde{b}_N(\tilde{u}_N^h, v^h) = f(v^h) \quad \forall v^h \in \tilde{V}^h.$$

From (3.11) and (3.12), the well-posedness of the problems (\tilde{VP}_N) and (\tilde{VP}_N^h) depend on the property of the symmetric bilinear form $\tilde{b}_N(u, v)$. For any $u, v \in \tilde{V}$, we can also expand $u|_{\Gamma_e} = u(R, \theta)$ and $v|_{\Gamma_e} = v(R, \theta)$ in Fourier series (see (3.15) and (3.16)). Substituting (3.15) and (3.16) into (4.1) and using the orthogonality of the cosines and sines, we obtain

$$(4.4) \quad \tilde{b}_N(u, v) = \frac{2\mu}{1 + 2\kappa} \sum_{n=1}^{\infty} [\gamma_n^{(N)} (a_n^1 c_n^1 + b_n^1 d_n^1 + a_n^2 c_n^2 + b_n^2 d_n^2) + \kappa n (a_n^1 + b_n^2)(c_n^1 + d_n^2) + \kappa n (b_n^1 - a_n^2)(d_n^1 - c_n^2)] \quad \forall u, v \in \tilde{V},$$

$$(4.5) \quad \tilde{b}_N(v, v) = \frac{2\mu}{1 + 2\kappa} \sum_{n=1}^{\infty} \left[\gamma_n^{(N)} \sum_{i=1}^2 ((c_n^i)^2 + (d_n^i)^2) + \kappa n (c_n^1 + d_n^2)^2 + \kappa n (d_n^1 - c_n^2)^2 \right] \quad \forall v \in \tilde{V},$$

where

$$(4.6) \quad \gamma_n^{(N)} = (1 + \kappa) \sum_{m=1}^N n^{2m} \alpha_m^{(N)} - \kappa n \equiv (1 + \kappa) \beta_n^{(N)} - \kappa n \quad \forall n \in \mathbb{N}.$$

Thus the property of $\tilde{b}_N(u, v)$ depends on the property of $\gamma_n^{(N)}$ (or $\beta_n^{(N)}$).

Table 3 shows $\alpha_m^{(N)}$ is positive for odd m , and is negative for even $m > 0$ for $1 \leq N \leq 5$. This property can be demonstrated numerically for $1 \leq N \leq 20$ (in fact this property can be proved for any positive integer N using the method given by Sidi in [25]). Thus we have that

$$(4.7) \quad \alpha_1^{(N)} > 0, \quad \alpha_N^{(N)} = \begin{cases} > 0 & N \text{ is odd,} \\ < 0 & N \text{ is even,} \end{cases} \quad 1 \leq N \leq 20.$$

From the engineering application point of view, the parameter N in (2.31) is always less than 10. Therefore in this paper we assume $N \leq 20$ in (2.31).

Then for the $\beta_n^{(N)}$ we have

Lemma 4.1. *If $1 \leq N \leq 20$ is odd, then*

$$(4.8) \quad \beta_n^{(N)} \geq n \quad \forall n \geq 1 \quad \lim_{n \rightarrow +\infty} \frac{\beta_n^{(N)}}{n^{2N}} = \alpha_N^{(N)} > 0.$$

If $1 \leq N \leq 20$ is even, then

$$(4.9) \quad \beta_n^{(N)} < 0 \quad \text{when } n \text{ is sufficiently large} \quad \lim_{n \rightarrow +\infty} \frac{\beta_n^{(N)}}{n^{2N}} = \alpha_N^{(N)} < 0.$$

Proof. We set a polynomial function whose degree is $2N$, say

$$(4.10) \quad \eta_N(t) = \sum_{m=1}^N \alpha_m^{(N)} t^{2m} - t.$$

Since $\eta_N''(t)$ is an even polynomial function whose degree is $2N - 2$ and $\eta_N''(0) = 2\alpha_1^{(N)} > 0$ for $1 \leq N \leq 20$ by noting (4.7), we know that $\eta_N''(t) = 0$ has at most $N - 1$ nonnegative roots. Thus $\eta_N(t) = 0$ has at most $N + 1$ nonnegative roots. From (4.10) and (2.30), we know that $t = 0, 1, 2, \dots, N$ are roots of $\eta_N(t) = 0$. Thus for $1 \leq N \leq 20$, we have that

$$(4.11) \quad \eta_N(t) \neq 0 \quad \forall t > N \quad \lim_{t \rightarrow +\infty} \frac{\eta_N(t)}{t^{2N}} = \alpha_N^{(N)}.$$

Then the desired inequalities (4.8) and (4.9) follow immediately from (4.11) and (4.7).

From the above discussion we have

Lemma 4.2. *For odd $1 \leq N \leq 20$, there exist two generic positive constants $C_N^{(1)}$ and $C_N^{(2)}$ depending only on N such that*

$$(4.12) \quad |\tilde{b}_N(u, v)| \leq C_N^{(2)} |u|_{N, \Gamma_R} \cdot |v|_{N, \Gamma_R} \quad \forall u, v \in \tilde{V},$$

$$(4.13) \quad C_N^{(1)} |v|_{N, \Gamma_R}^2 \leq \tilde{b}_N(v, v) \quad \forall v \in \tilde{V}.$$

Proof. For any odd $1 \leq N \leq 20$, noting (4.6) and (4.8), we know that there exist positive constants $C_N^{(1)}$ and $C_N^{(2)}$ such that

$$(4.14) \quad C_N^{(1)} n^{2N} \leq \gamma_n^{(N)} \leq C_N^{(2)} n^{2N} \quad n = 1, 2, 3, \dots$$

From (4.14) and (4.4), noting (3.32) with $r = R$, (3.15) and (3.16), we have that

$$(4.15) \quad \begin{aligned} |\tilde{b}_N(u, v)| &\leq C_N^{(2)} \sum_{n=1}^{\infty} n^{2N} \left[|a_n^1 c_n^1 + b_n^1 d_n^1 + a_n^2 c_n^2 + b_n^2 d_n^2| \right. \\ &\quad \left. + |a_n^1 d_n^2 - b_n^1 c_n^2 - a_n^2 d_n^1 + b_n^2 c_n^1| \right] \\ &\leq C_N^{(2)} \left[\sum_{n=1}^{\infty} n^{2N} \sum_{i=1}^2 [(a_n^i)^2 + (b_n^i)^2] \right]^{1/2} \\ &\quad \cdot \left[\sum_{n=1}^{\infty} n^{2N} \sum_{i=1}^2 [(d_n^i)^2 + (c_n^i)^2] \right]^{1/2} \\ &\leq C_N^{(2)} |u|_{N, \Gamma_e} \cdot |v|_{N, \Gamma_e} \quad \forall u, v \in \tilde{V}. \end{aligned}$$

Furthermore from (4.14) and (4.5), noting (3.32) with $r = R$ and (3.16), we obtain

$$\begin{aligned}
 \tilde{b}_N(v, v) &\geq \frac{2\mu}{1 + 2\kappa} \sum_{n=1}^{\infty} \gamma_n^{(N)} ((c_n^1)^2 + (d_n^1)^2 + (c_n^2)^2 + (d_n^2)^2) \\
 (4.16) \quad &\geq C_N^{(1)} \sum_{n=1}^{\infty} n^{2N} ((c_n^1)^2 + (d_n^1)^2 + (c_n^2)^2 + (d_n^2)^2) \\
 &= C_N^{(1)} |v|_{N, \Gamma_e}^2 \quad \forall v \in \tilde{V}.
 \end{aligned}$$

Thus the desired inequalities (4.12) and (4.13) are proved.

From the discussion above, noting the Körn inequality and Poincaré inequality, we assign the following norm on \tilde{V} :

$$(4.17) \quad \|v\|_{\Delta} := [a(v, v) + |v|_{N, \Gamma_e}^2]^{1/2} \equiv [\|v\|_*^2 + |v|_{N, \Gamma_e}^2]^{1/2} \quad \forall v \in \tilde{V}.$$

It follows immediately from (3.11), (3.12), (4.12), (4.13), (4.6), (4.8) and (4.9) that the variational problems (\tilde{VP}_N) and (\tilde{VP}_N^h) are well posed in the case of odd $1 \leq N \leq 20$ or $N = 0$ and they are not well posed in the case of even $0 < N \leq 20$; that is, for $f \in \tilde{V}'$, the dual of \tilde{V} , there exists a unique $\tilde{u}_N \in \tilde{V}$ solving (\tilde{VP}_N) , a unique $\tilde{u}_N^h \in \tilde{V}^h$ solving (\tilde{VP}_N^h) , and

$$(4.18) \quad \|\tilde{u}_N\|_{\Delta} + \|\tilde{u}_N^h\|_{\Delta} \leq M_N \|f\|_{\tilde{V}'}, \quad \forall \text{ odd } 1 \leq N \leq 20,$$

where M_N is a constant.

Then we have the following estimate.

Lemma 4.3. *Suppose $u \in \tilde{V}$ is a solution of the exterior problem (1.1)-(1.3) and $u|_{\Gamma_0} \in [H^{N+1}(\Gamma_0)]^2$. Then we have the following estimate for odd $1 \leq N \leq 20$:*

$$(4.19) \quad |b(u, v) - \tilde{b}_N(u, v)| \leq C_{(N)} \left(\frac{R_0}{R}\right)^{\max\{1, N-1\}} |u|_{N+1, \Gamma_0} \cdot |v|_{N, \Gamma_e} \quad \forall v \in \tilde{V},$$

where $C_{(N)}$ is a constant independent of u , R and h .

Proof. Subtracting (3.18) from (4.4), noting (4.6), (4.8) (3.40)-(3.43) and (3.32), we obtain for odd $1 \leq N \leq 20$

$$\begin{aligned}
 (4.20) \quad & \left| b(u, v) - \tilde{b}_N(u, v) \right| \\
 &= \frac{2\mu\pi}{1 + 2\kappa} \left| \sum_{n=N+1}^{\infty} (1 + \kappa) \left(\sum_{m=1}^N n^{2m} \alpha_m^{(N)} - n \right) (a_n^1 c_n^1 + b_n^1 d_n^1 + a_n^2 c_n^2 + b_n^2 d_n^2) \right| \\
 &\leq C_{(N)} \sum_{n=N+1}^{\infty} n^{2N} |a_n^1 c_n^1 + b_n^1 d_n^1 + a_n^2 c_n^2 + b_n^2 d_n^2| \\
 &\leq C_{(N)} \left[\sum_{n=N+1}^{\infty} n^{2N} \sum_{i=1}^2 ((a_n^i)^2 + (b_n^i)^2) \right]^{1/2} \\
 &\quad \cdot \left[\sum_{n=N+1}^{\infty} n^{2N} \sum_{i=1}^2 ((c_n^i)^2 + (d_n^i)^2) \right]^{1/2} \\
 &\leq C_{(N)} \left[\sum_{n=\max\{1, N-1\}}^{\infty} n^{2N+2} ((p_n^1)^2 + (q_n^1)^2 + (p_n^2)^2 + (q_n^2)^2) \frac{R_0^{2n}}{R^{2n}} \right]^{1/2} \cdot |v|_{N, \Gamma_e} \\
 &\leq C_{(N)} \left(\frac{R_0}{R} \right)^{\max\{1, N-1\}} |u|_{N+1, \Gamma_0} \cdot |v|_{N, \Gamma_e} \quad \forall v \in \tilde{V}.
 \end{aligned}$$

Combining Lemmas 4.2 and 4.3, we get the following error estimate.

Theorem 4.1. *Let u be the solution of the problem (1.1)-(1.3) and \tilde{u}_N^h the solution of the problem $(\tilde{V}P_N^h)$. Suppose $f \in [L^2(\Omega_i)]^2$ and $u|_{\Gamma_0} \in [H^{N+1}(\Gamma_0)]^2$. Then we have the following error estimate for odd $1 \leq N \leq 20$:*

$$(4.21) \quad \|u - \tilde{u}_N^h\|_{\Delta} \leq C_{(N)} \left[\inf_{v^h \in \tilde{V}^h} \|u - v^h\|_{\Delta} + \left(\frac{R_0}{R} \right)^{\max\{1, N-1\}} |u|_{N+1, \Gamma_0} \right]$$

with constant $C_{(N)}$ independent of u, R and h .

Proof. The proof of this theorem is similarly to the proof of Theorem 3.1. It is omitted here.

If we suppose $u \in [H^{p+1}(\Omega_i)]^2$, $u|_{\Gamma_e} \in [H^{p+N}(\Gamma_e)]^2$ and the interpolation error of \tilde{V}^h approximate to \tilde{V} is

$$(4.22) \quad \inf_{v^h \in \tilde{V}^h} \|u - v^h\|_{\Delta} \leq C_0 h^p [|u|_{p+1, \Omega_i} + |u|_{p+N, \Gamma_e}]$$

(a family of this kind of finite element subspaces was proposed in [10]), then combining (4.22) and (4.21), noting the K\"orn inequality and Poincar\'e inequality, we get for odd $1 \leq N \leq 20$

$$\begin{aligned}
 (4.23) \quad & \|u - \tilde{u}_N^h\|_{1, \Omega_0} \leq C_0 |u - \tilde{u}_N^h|_{1, \Omega_0} \leq C_0 \|u - \tilde{u}_N^h\|_{\Delta} \\
 &\leq C_{(N)} \left[h^p (|u|_{p+1, \Omega_i} + |u|_{p+N, \Gamma_e}) + \left(\frac{R_0}{R} \right)^{\max\{1, N-1\}} |u|_{N+1, \Gamma_0} \right].
 \end{aligned}$$

5. NUMERICAL IMPLEMENTATION AND RESULTS

In this section we present the numerical results which demonstrate the performance of the error estimates (3.50) and (4.23). In our example, we take $\mu = 1$, $x^+ = (0, 0.25)$, and $x^- = (0, -0.25)$, and the unbounded domain $\Omega = \{x \in \mathbb{R}^2 : 0.5 < |x|\}$ is the exterior domain outside a circle $\Gamma_i = \{x \in \mathbb{R}^2 : |x| = 0.5\}$. In our computation, when dealing with nonlocal approximate artificial boundary conditions (2.20), continuous piecewise linear elements were used throughout the domain Ω_i . When dealing with high-order local artificial boundary conditions (2.31), continuous piecewise bilinear elements were used throughout the domain Ω_i , except in the single layer elements adjacent to the artificial boundary Γ_e . There, special finite elements, $C_{\Gamma_e}^{2,1}$ (which were introduced by Givoli et al., [10] and has $C^2(\Gamma_e)$ regularity at Γ_e), were used. That is to say, $p = 1$ in the interpolation errors (3.49) and (4.22) [2, 10].

Example. An exterior problem for Navier equations.

We consider the Navier equations in the planar domain outside a circular obstacle of radius $a = 0.5$ (see Figure 1). The problem is governed by the following boundary value problem:

$$(5.1) \quad -\mu \Delta u - (\lambda + \mu) \operatorname{grad} \operatorname{div} u = f = (f_1, f_2)^T \quad \text{in } \Omega = \{x : 0.5 < |x|\},$$

$$(5.2) \quad u(0.5, \theta) = g(\theta) = (g_1, g_2)^T \quad \text{on } \Gamma_i = \partial\Omega,$$

$$(5.3) \quad u \text{ is bounded when } r = \sqrt{x_1^2 + x_2^2} \rightarrow +\infty,$$

where

$$f_1(x) = \begin{cases} -8x_2[(3\lambda + 6\mu)x_1^2 + (\lambda + 4\mu)x_2^2 - (\lambda + 3\mu)] & 0.5 \leq |x| < 1.0, \\ 0 & 1.0 \leq |x|, \end{cases}$$

$$f_2(x) = \begin{cases} -8x_1[(\lambda + 4\mu)x_1^2 + (3\lambda + 6\mu)x_2^2 - (\lambda + 3\mu)] & 0.5 \leq |x| < 1.0, \\ 0 & 1.0 \leq |x|; \end{cases}$$

$$g_1(\theta) = \frac{\lambda + 3\mu}{2\mu(\lambda + 2\mu)} \ln \frac{1.25 + \sin \theta}{1.25 - \sin \theta} + \frac{\lambda + \mu}{\mu(\lambda + 2\mu)} \frac{2 \cos^3 \theta \sin \theta}{1.5625 - \sin^2 \theta} + 0.28125 \sin \theta \quad 0 \leq \theta < 2\pi,$$

$$g_2(\theta) = \frac{(\lambda + \mu) \cos \theta}{\mu(\lambda + 2\mu)} \left[\frac{\sin \theta - 0.5}{1.25 - \sin \theta} - \frac{\sin \theta + 0.5}{1.25 + \sin \theta} \right] + 0.28125 \cos \theta \quad 0 \leq \theta < 2\pi.$$

This problem has an exact solution:

$$u_1(x) = \begin{cases} \frac{\lambda + 3\mu}{\mu(\lambda + 2\mu)} \ln \frac{|x - x^-|}{|x - x^+|} + \frac{\lambda + \mu}{\mu(\lambda + \mu)} \left[\frac{x_1^2}{|x - x^+|^2} - \frac{x_1^2}{|x - x^-|^2} \right] + x_2(|x|^2 - 1)^2 & 0.5 \leq |x| < 1.0, \\ \frac{\lambda + 3\mu}{\mu(\lambda + 2\mu)} \ln \frac{|x - x^-|}{|x - x^+|} + \frac{\lambda + \mu}{\mu(\lambda + \mu)} \left[\frac{x_1^2}{|x - x^+|^2} - \frac{x_1^2}{|x - x^-|^2} \right] & 1.0 \leq |x|; \end{cases}$$




FIGURE 2. The effect of N using nonlocal artificial boundary conditions

TABLE 2. The effect of the mesh size h using nonlocal artificial boundary conditions

Mesh	$h = 0.31416$	$h = 0.15708$	$h = 0.07854$	$h = 0.03927$
$\max u - u_N^h $	1.5631E-2	4.1784E-3	1.0707E-3	2.6869E-4
$\ u - u_N^h\ _{0,\Omega_0}$	4.4819E-2	1.2438E-2	3.1956E-3	8.0471E-4
$ u - u_N^h _{1,\Omega_0}$	0.7745	0.4067	0.2060	0.1034

$$u_2(x) = \begin{cases} \frac{\lambda+\mu}{\mu(\lambda+2\mu)} \left[\frac{x_1(x_2-x_2^+)}{|x-x^+|^2} - \frac{x_1(x_2-x_2^-)}{|x-x^-|^2} \right] \\ \quad + x_1(|x|^2 - 1)^2 & 0.5 \leq |x| < 1.0, \\ \frac{\lambda+\mu}{\mu(\lambda+2\mu)} \left[\frac{x_1(x_2-x_2^+)}{|x-x^+|^2} - \frac{x_1(x_2-x_2^-)}{|x-x^-|^2} \right] & 1.0 \leq |x|. \end{cases}$$

First we test the effect of the mesh size h in the error estimates (3.50). We introduce a circular artificial boundary $\Gamma_e = \Gamma_0$ of radius $R = R_0 = 1.0$. On Γ_0 we apply the nonlocal artificial boundary condition (2.20) with $N = 0, 1, 2, \dots$ or high-order local artificial boundary condition (2.31). In the annular computational domain Ω_0 , we use four meshes. The first mesh consists of 2 radial layers of elements, with 20 quadrilateral elements in each layer. We denote it as 2×20 . The other three meshes are 4×40 , 8×80 and 16×160 . Table 2 shows the maximum errors of $u - u_N^h$ over the mesh points, $\|u - u_N^h\|_{0,\Omega_0}$ and $|u - u_N^h|_{1,\Omega_0}$ for large N (say $N = 51$).

The results show that the convergent rates of $|u - u_N^h|_{1,\Omega_0}$ and $\|u - u_N^h\|_{0,\Omega_0}$ with respect to h are 1 and 2 when using nonlocal artificial boundary conditions, respectively. Second we test the effect of N in the error estimate (3.50). Let u_∞^h denote the finite element approximation of the problem on the domain Ω_0 with




FIGURE 3. The effect of R using nonlocal artificial boundary conditions

TABLE 3. The effect of the location of Γ_R using a local artificial boundary condition ($N = 1$)

Location	$R = 1.0$	$R = 1.5$	$R = 2.0$	$R = 2.5$	$R = 3.0$
$\max u - \tilde{u}_N^h $	9.5675E-2	6.0485E-2	4.3960E-2	3.4578E-2	2.8531E-2
$\ u - \tilde{u}_N^h\ _{0,\Omega_0}$	5.9929E-2	2.2710E-2	1.5862E-2	1.0996E-2	7.8054E-3
$ u - \tilde{u}_N^h _{1,\Omega_0}$	0.6290	0.3570	0.3096	0.2740	0.2483

the mesh size h when N is very large (say $N = 51$). In this case $R_0 = R$, so the effect of R in the error estimate (3.50) disappears. Figure 2 shows the errors $E_N := \|u_\infty^h - u_N^h\|_{k,\Omega_0}$ ($k = 0, 1$) on the mesh 16×160 for different N . Third we test the effect of the location of the artificial boundary Γ_ϵ . Let $\Omega_R = \{x : 0.5 < |x| < R\}$ denotes the bounded computational domain with the artificial boundary Γ_R . We choose $R = 1.0, 1.5, 2.0, 2.5, 3.0$. The corresponding meshes we used were $8 \times 40, 16 \times 40, 24 \times 40, 32 \times 40$ and 40×40 , respectively. That is to say, each computational domain has a mesh with the fixed mesh size $h = 0.07854$. Let u_N^R denote the finite element approximation of the problem on the domain Ω_R with the corresponding mesh by using the nonlocal artificial boundary condition (2.20) on the artificial boundary Γ_R . u_∞^R corresponds to the solution when N is very large (say $N = 51$) and \tilde{u}_N^R corresponds to the solution using the high-order local artificial boundary condition (2.31) at Γ_R . Tables 3 and 4 shows the maximum errors of $u - \tilde{u}_N^R$ over the mesh points, $\|u - \tilde{u}_N^R\|_{0,\Omega_0}$ and $|u - \tilde{u}_N^R|_{1,\Omega_0}$ for $N = 1, 3$. Further, Figure 3 shows the errors $E_R := \|u_\infty^R - u_N^R\|_{1,\Omega_0}$ for different R , and Figure 4 shows the errors $E_R := \|u_\infty^R - \tilde{u}_N^R\|_{1,\Omega_0}$ for different R .

Tables 2–4 and Figures 2–4 demonstrate the performance of the error estimate (3.50) and (4.23). In practice, if one wants to use a local artificial boundary condition, we advise using the one corresponding to $N = 1$. This condition is very simple




FIGURE 4. The effect of R using local artificial boundary conditions

TABLE 4. The effect of the location of Γ_R using a local artificial boundary condition ($N = 3$)

Location	$R = 1.0$	$R = 1.5$	$R = 2.0$	$R = 2.5$	$R = 3.0$
$\max u - \tilde{u}_N^h $	0.2552	0.1910	0.1489	0.1212	0.1019
$\ u - \tilde{u}_N^h\ _{0,\Omega_0}$	0.2020	0.1136	7.0255E-2	4.6411E-2	3.2470E-2
$ u - \tilde{u}_N^h _{1,\Omega_0}$	1.3323	0.7880	0.6025	0.4889	0.4142

and easy to deal with by using the standard finite elements. From our numerical results, the local artificial boundary condition corresponding to $N = 3$ is not better than the one corresponding to $N = 1$. One reason is the constant $C_{(1)} \ll C_{(3)}$ in (4.13) and errors in the numerical integration for the part $\tilde{b}_N(u, v)$. Thus in practical computation, the cases $N \geq 3$ need further study. The analogous numerical results for the Laplacian equation can be found in [10].

REFERENCES

[1] R. A. Adams, Sobolev Spaces, Academic Press, 1975. MR 56:9247
 [2] P. G. Ciarlet, The finite element method for elliptic problems, North-Holland, 1978. MR 58:25001
 [3] B. Engquist and A. Majda, Absorbing boundary conditions for the numerical simulation of waves, Math. Comp. 31 (1977), 629-651. MR 55:9555
 [4] K. Feng, Asymptotic radiation conditions for reduced wave equations, J. Comput. Math. 2, (1984), 130-138. MR 89f:65134
 [5] D. Gilbarg and N. S. Trudinger, Elliptic partial differential equations of second order, Springer-Verlag, 1977. MR 57:13109
 [6] D. Givoli, Numerical Methods for Problems in Infinite Domains, Elsevier, Amsterdam, 1992. MR 94j:65003
 [7] D. Givoli and J. B. Keller, A finite element method for large domains, Comp. Meth. Appl. Mech. Engng., 76 (1989), 41-66. MR 90i:65196

- [8] D. Givoli and J. B. Keller, Non-reflecting boundary conditions for elastic waves, *Wave Motion*, 12 (1990), 261-279.
- [9] D. Givoli and I. Patashenko, Optimal local non-reflecting boundary conditions, *Appl. Numer. Math.*, 27 (1998), 367-384. MR **99f**:65154
- [10] D. Givoli, I. Patlashenko and J. Keller, High-order boundary conditions and finite elements for infinite domains, *Comput. Methods Appl. Mech. Engrg.* 143 (1997), 13-39. MR **97m**:65203
- [11] C. I. Goldstein, A finite element method for solving Helmholtz type equations in waveguides and other unbounded domains, *Math. Comp.* 39, (1982), 309-324. MR **84e**:65112
- [12] T. M. Hagstrom and H. B. Keller, Exact boundary conditions at artificial boundary for partial differential equations in cylinders, *SIAM J. Math. Anal.* 17, (1986), 322-341. MR **87g**:35022
- [13] T. M. Hagstrom and H. B. Keller, Asymptotic boundary conditions and numerical methods for nonlinear elliptic problems on unbounded domains, *Math. Comp.* 48, (1987), 449-470. MR **88d**:65173
- [14] L. Halpern and M. Schatzman, Artificial boundary conditions for incompressible viscous flows, *SIAM J. Math. Anal.* 20, (1989), 308-353. MR **90c**:35164
- [15] H. Han and W. Bao, An artificial boundary condition for two-dimensional incompressible viscous flows using the method of lines, *Int. J. Numer. Methods Fluids* 22, (1996), 483-493. MR **97a**:76093
- [16] H. Han and W. Bao, An artificial boundary condition for the incompressible viscous flows in a no-slip channel, *J. Comput. Math.* 13 (1995), 51-63. MR **95k**:76091
- [17] H. Han and W. Bao, The artificial boundary conditions for incompressible materials on an unbounded domain, *Numer. Math.* 77, (1997), 347-363. MR **98f**:73037
- [18] H. Han, J. Lu and W. Bao, A discrete artificial boundary condition for steady incompressible viscous flows in a no-slip channel using a fast iterative method, *J. Comput. Phys.* 114, (1994), 201-208. MR **95e**:76061
- [19] H. Han, W. Bao and T. Wang, Numerical simulation for the problem of infinite elastic foundation, *Comput. Methods Appl. Mech. Engrg.* 147, (1997), 369-385. MR **98e**:73131
- [20] H. Han and X. Wu, Approximation of infinite boundary condition and its application to finite element method, *J. Comput. Math.* 3, (1985), 179-192. MR **87k**:65134
- [21] H. Han and X. Wu, The approximation of exact boundary condition at an artificial boundary for linear elastic equation and its application, *Math. Comp.* 59, (1992), 21-37. MR **92k**:35076
- [22] J. L. Lions, *Optimal control of systems governed by partial differential equations*, Springer-Verlag, Berlin and Heidelberg, 1971. MR **42**:6395
- [23] S. G. Mikhailin, *The problem of the minimum of a quadratic functional*, Holden-Day, San Francisco, London, Amsterdam, 1965. MR **30**:1427
- [24] F. Nataf, An open boundary condition for the computation of the steady incompressible Navier-Stokes equations, *J. Comput. Phys.* 85, (1989), 104-129. MR **90i**:76015
- [25] A. Sidi, A family of matrix problems, *SIAM Review (Problems and Solutions)*, 40 (1998), 718-723.

DEPARTMENT OF APPLIED MATHEMATICS, TSINGHUA UNIVERSITY, BEIJING 100084, PEOPLE'S REPUBLIC OF CHINA

E-mail address: hhan@math.tsinghua.edu.cn

DEPARTMENT OF APPLIED MATHEMATICS, TSINGHUA UNIVERSITY, BEIJING 100084, PEOPLE'S REPUBLIC OF CHINA

E-mail address: wbao@math.tsinghua.edu.cn