

TABLE I

Coordinates and coefficients for the n -dimensional sphere angular integration rule

$$\int_0^\pi \sin^\nu \phi g(\phi) d\phi \doteq \sum_{i=1}^{2m+2} b_i g(\phi_i), \quad \nu = 1, 2, \dots, n - 2,$$

$$y = \cos \phi$$

ν	Polynomial $Q_{2m+2}^{(\nu-1)/2}$	Coordinates y_i	Coefficients b_i
$m = 0$ (3rd degree rules)			
1	$\frac{3}{2} \left(y^2 - \frac{1}{3} \right)$	$\pm \frac{1}{\sqrt{3}}$	1
2	$\frac{5}{2} \left(y^2 - \frac{1}{4} \right)$	$\cos \frac{\pi}{3}, \cos \frac{2\pi}{3}$	$\frac{\pi}{4}$
3	$\frac{15}{4} \left(y^2 - \frac{1}{5} \right)$	$\pm \frac{1}{\sqrt{5}}$	$\frac{2}{3}$
4	$\frac{21}{4} \left(y^2 - \frac{1}{6} \right)$	$\pm \frac{1}{\sqrt{6}}$	$\frac{3\pi}{16}$
5	$7 \left(y^2 - \frac{1}{7} \right)$	$\pm \frac{1}{\sqrt{7}}$	$\frac{8}{15}$
6	$9 \left(y^2 - \frac{1}{8} \right)$	$\pm \frac{1}{\sqrt{8}}$	$\frac{5\pi}{32}$
$m = 1$ (7th degree rules)			
1	$\frac{35}{8} \left(y^4 - \frac{6}{7} y^2 + \frac{3}{35} \right)$	$\pm \sqrt{\frac{15 + 2\sqrt{30}}{35}}$ $\pm \sqrt{\frac{15 - 2\sqrt{30}}{35}}$	$\frac{49}{6(18 + \sqrt{30})}$ $\frac{49}{6(18 - \sqrt{30})}$
2	$\frac{63}{8} \left(y^4 - \frac{3}{4} y^2 + \frac{1}{16} \right)$	$\cos \frac{\pi}{5}, \cos \frac{4\pi}{5}$ $\cos \frac{2\pi}{5}, \cos \frac{3\pi}{5}$	$\frac{\pi}{8} \left(1 - \frac{1}{\sqrt{5}} \right)$ $\frac{\pi}{8} \left(1 + \frac{1}{\sqrt{5}} \right)$
3	$\frac{105}{8} \left(y^4 - \frac{2}{3} y^2 + \frac{1}{21} \right)$	$\pm \sqrt{\frac{7 + \sqrt{28}}{21}}$ $\pm \sqrt{\frac{7 - \sqrt{28}}{21}}$	$\frac{6}{5} \left(\frac{1}{5 - \sqrt{7}} \right)$ $\frac{6}{5} \left(\frac{1}{5 + \sqrt{7}} \right)$

TABLE I (Continued)

ν	Polynomial $Q_{2m+2}^{(\nu-1)/2}$	Coordinates y_i	Coefficients b_i
4	$\frac{165}{8} \left(y^4 - \frac{3}{5} y^2 + \frac{3}{80} \right)$	$\pm \sqrt{\frac{6 + \sqrt{21}}{20}}$	$\frac{125\pi}{32(63 + 8\sqrt{21})}$
		$\pm \sqrt{\frac{6 - \sqrt{21}}{20}}$	$\frac{125\pi}{32(63 - 8\sqrt{21})}$
5	$\frac{495}{16} \left(y^4 - \frac{6}{11} y^2 + \frac{1}{33} \right)$	$\pm \sqrt{\frac{9 + 4\sqrt{3}}{33}}$	$\frac{242}{105(14 + 5\sqrt{3})}$
		$\pm \sqrt{\frac{9 - 4\sqrt{3}}{33}}$	$\frac{242}{105(14 - 5\sqrt{3})}$
6	$\frac{715}{16} \left(y^4 - \frac{1}{2} y^2 + \frac{1}{40} \right)$	$\pm \sqrt{\frac{5 + \sqrt{15}}{20}}$	$\frac{35\pi}{128(6 + \sqrt{15})}$
		$\pm \sqrt{\frac{5 - \sqrt{15}}{20}}$	$\frac{35\pi}{128(6 - \sqrt{15})}$

TABLE II

Coordinates and coefficients for the n -dimensional sphere radial integration rule

$$\int_R^1 r^{n-1} g(r^2) dr \doteq \sum_{j=1}^{m+1} c_j g(r_j^2)$$

$m = 0, n, R$ arbitrary

$$Q_1(r^2) = r^2 - \frac{n}{n+2} \left(\frac{1 - R^{n+2}}{1 - R^n} \right),$$

$$r_1^2 = \frac{n}{n+2} \left(\frac{1 - R^{n+2}}{1 - R^n} \right), \quad c_1 = \frac{1}{n} (1 - R^n)$$

$m = 1, R = 0, n$ arbitrary

$$r_1^2 = \frac{(n+2)(n+4) - 2\sqrt{2}(n+2)(n+4)}{(n+4)(n+6)},$$

$$c_1 = \frac{1}{2n} \left\{ 1 - \frac{(n-2)(n+4)}{(n+2)\sqrt{2}(n+2)(n+4)} \right\}$$

$$r_2^2 = \frac{(n+2)(n+4) + 2\sqrt{2}(n+2)(n+4)}{(n+4)(n+6)},$$

$$c_2 = \frac{1}{2n} \left\{ 1 + \frac{(n-2)(n+4)}{(n+2)\sqrt{2}(n+2)(n+4)} \right\}$$

TABLE II—Continued

$m = 1, n = 4$ (For $n = 3$, see the table of W. H. Peirce.)

R	$Q_2(r^2)$	r_i^2	c_i
0	$r^4 - \frac{6}{5}r^2 + \frac{3}{10}$	$r_1^2 = 0.355\ 051\ 024$ $r_2^2 = 0.844\ 948\ 977$	$c_1 = 0.090\ 979\ 310$ $c_2 = 0.159\ 020\ 691$
$\frac{1}{4}$	$r^4 - \frac{1037}{856}r^2$ $+ \frac{16\ 909}{54\ 784}$	$r_1^2 = 0.364\ 366\ 862$ $r_2^2 = 0.847\ 081\ 739$	$c_1 = 0.091\ 808\ 296$ $c_2 = 0.157\ 215\ 143$
$\frac{1}{2}$	$r^4 - \frac{29}{22}r^2$ $+ \frac{689}{1760}$	$r_1^2 = 0.451\ 910\ 920$ $r_2^2 = 0.866\ 270\ 899$	$c_1 = 0.094\ 048\ 036$ $c_2 = 0.140\ 326\ 967$
$\frac{3}{4}$	$r^4 - \frac{3\ 956\ 505}{2\ 505\ 272}r^2$ $+ \frac{487\ 302\ 501}{801\ 687\ 040}$	$r_1^2 = 0.664\ 422\ 219$ $r_2^2 = 0.914\ 849\ 411$	$c_1 = 0.077\ 239\ 032$ $c_2 = 0.093\ 659\ 411$

$m = 1, n = 5$

0	$r^4 - \frac{14}{11}r^2 + \frac{35}{99}$	$r_1^2 = 0.409\ 596\ 534$ $r_2^2 = 0.863\ 130\ 739$	$c_1 = 0.065\ 637\ 839$ $c_2 = 0.134\ 362\ 161$
$\frac{1}{4}$	$r^4 - \frac{49 \times 227\ 879}{8 \times 1092\ 923}r^2$ $+ \frac{35 \times 31\ 387\ 427}{2816 \times 1\ 092\ 923}$	$r_1^2 = 0.413\ 170\ 409$ $r_2^2 = 0.863\ 917\ 619$	$c_1 = 0.066\ 037\ 827$ $c_2 = 0.133\ 766\ 869$
$\frac{1}{2}$	$r^4 - \frac{7 \times 53\ 557}{22 \times 12\ 601}r^2$ $+ \frac{35 \times 237\ 793}{1584 \times 12\ 601}$	$r_1^2 = 0.475\ 583\ 690$ $r_2^2 = 0.876\ 758\ 471$	$c_1 = 0.070\ 121\ 263$ $c_2 = 0.123\ 628\ 742$
$\frac{3}{4}$	$r^4 - \frac{7 \times 8\ 334\ 140\ 936}{704 \times 52\ 193\ 401}r^2$ $+ \frac{35 \times 92\ 932\ 900\ 132}{99 \times 1024 \times 52\ 193\ 401}$	$r_1^2 = 0.669\ 472\ 413$ $r_2^2 = 0.918\ 235\ 562$	$c_1 = 0.065\ 438\ 853$ $c_2 = 0.087\ 100\ 212$

Let

$$(1.4) \quad f(x_1, x_2, \dots, x_n) = f_1(\theta_1, \theta_2, \dots, \theta_{n-1}, r).$$

The Jacobian of the transformation is

$$(1.5) \quad J(\theta_1, \theta_2, \dots, \theta_{n-1}, r) = r^{n-1} \sin \theta_2 \sin^2 \theta_3 \dots \sin^{n-2} \theta_{n-1}$$

D is now defined by:

$$(1.6) \quad \begin{aligned} 0 \leq \theta_1 \leq 2\pi; \quad 0 \leq \theta_i \leq \pi \quad (i = 2, 3, \dots, n - 1), \text{ and} \\ R \leq r \leq 1. \end{aligned}$$

The rule to be constructed has the form:

$$(1.7) \quad \int_R^1 \int_0^\pi \dots \int_0^\pi \int_0^{2\pi} Jf_1(\theta_1, \theta_2, \dots, \theta_{n-1}, r) d\theta_1 d\theta_2 \dots d\theta_{n-1} dr$$

$$\doteq \sum_{g,h,\dots,i,j} \alpha_{gh\dots ij} f_1(\theta_{1g}, \theta_{2h}, \dots, \theta_{n-1,i}, r_j).$$

The main result is the following theorem:

2. Theorem. *If it be required that*

- (a) rule (1.7) have precision $k = 4m + 3$ ($m = 0, 1, 2, \dots$) in x_1, x_2, \dots, x_n ;
- (b) the evaluation points be taken at all intersection points of the surfaces

$$(2.1) \quad \begin{aligned} \theta_1 &= \theta_{1g} & (g = 1, 2, \dots, p_1), \\ \theta_2 &= \theta_{2h} & (h = 1, 2, \dots, p_2), \\ &\dots\dots\dots \\ \theta_{n-1} &= \theta_{n-1,i} & (i = 1, 2, \dots, p_{n-1}), \\ r &= r_j & (j = 1, 2, \dots, p_n); \end{aligned}$$

- (c) the number of evaluation points in this configuration be a minimum; then it is both necessary and sufficient for the existence of a unique set of real weight coefficients, $\alpha_{gh\dots ij}$, that

$$(2.2)(a) \quad \theta_{1g} = \frac{2\pi g}{k + 1}, \quad g = 0, 1, 2, \dots, k;$$

- (b) the $\cos \theta_{li}$ ($i = 1, 2, \dots, p_i$) be the $2m + 2$ zeros of the orthogonal polynomials $Q_{2m+2}^{l/2-1}(y)$ ($l = 2, \dots, n - 1$), where

$$(2.3) \quad Q_p^\alpha(y) \equiv \binom{p + \alpha}{p} G_p(2\alpha + 1, \alpha + 1, \frac{1}{2}(1 - y)),$$

where the $G_p(a, b, z)$ are the Jacobi polynomials;

- (c) the r_j^2 be the $m + 1$ zeros of the polynomial $Q_{m+1}(r^2)$ of degree $m + 1$ in r^2 which satisfies

$$(2.4) \quad \int_R^1 r^{n-2} Q_{m+1}(r^2) T_m(r^2) dr^2 = 0,$$

where $T_m(r^2)$ is an arbitrary polynomial in r^2 of degree m or less.

The unique weights are

$$(2.5) \quad \alpha_{gh\dots ij} = a_g b_{2,h} \cdots b_{n-1,i} c_j,$$

where

$$(2.6) \quad a_g = \frac{2\pi}{k+1}, \quad g = 1, 2, \dots, k+1,$$

$$(2.7) \quad b_{l,i} = \frac{((2m+1+\frac{1}{2}l)!)^2}{(2m+2)!(2m+l)!} \cdot \frac{2^{l-1}}{(1-y_i^2)(Q'_{2m+2}(y_i))^2},$$

$$l = 2, \dots, n-1, \quad i = 1, 2, \dots, 2m+2,$$

$$(2.8) \quad c_j = \frac{1}{Q'_{m+1}(r_j^2)} \int_R \frac{r^{n-1} Q_{m+1}(r^2) dr}{r^2 - r_j^2}.$$

Proof. Because the evaluation points of the rule (1.7) are arranged in a product lattice the integral can be written

$$(2.9) \quad I \equiv \int_D f(\mathbf{x}) d\tau = \int_R r^{n-1} \int_0^\pi \sin^{n-2} \theta_{n-1} \int_0^\pi \cdots \int_0^\pi \sin \theta_2$$

$$\cdot \int_0^{2\pi} f_1(\theta_1, \theta_2, \dots, \theta_{n-1}, r) d\theta_1 d\theta_2 \cdots d\theta_{n-1} dr$$

Define functions f_2, \dots, f_n as follows:

$$(2.10) \quad f_2(\theta_2, \dots, \theta_{n-1}, r) = \int_0^{2\pi} f_1(\theta_1, \theta_2, \dots, \theta_{n-1}, r) d\theta_1,$$

$$(2.11) \quad f_l(\theta_l, \dots, \theta_{n-1}, r) = \int_0^\pi \sin^{l-2} \theta_{l-1} f_{l-1}(\theta_{l-1}, \dots, \theta_{n-1}, r) d\theta_{l-1}$$

$$(l = 3, 4, \dots, n-1),$$

$$(2.12) \quad f_n(r) = \int_0^\pi \sin^{n-2} \theta_{n-1} f_{n-1}(\theta_{n-1}, r) d\theta_{n-1};$$

then

$$(2.13) \quad I = \int_R r^{n-1} f_n(r) dr.$$

We look for integration rules:

$$(2.14) \quad \int_0^{2\pi} f_1 d\theta_1 \doteq \sum_{g=1}^{p_1} a_g f_1(\theta_{1g}, \theta_2, \dots, \theta_{n-1}, r),$$

$$(2.15) \quad \int_0^\pi \sin^{l-1} \theta_l f_l d\theta_l \doteq \sum_{i=1}^{p_l} b_{li} f_l(\theta_{l,i}, \theta_{l+1}, \dots, \theta_{n-1}, r)$$

$$(l = 2, \dots, n-1),$$

and

$$(2.16) \quad \int_R r^{n-1} f_n(r) dr \doteq \sum_{j=1}^{p_n} c_j f_n(r_j).$$

The first of these, (2.14), must be exact whenever f_1 is a polynomial of degree k in

$$\begin{aligned} & r \sin \theta_1 \sin \theta_2 \cdots \sin \theta_{n-1}, \\ & r \cos \theta_1 \sin \theta_2 \cdots \sin \theta_{n-1}, \\ & \dots\dots\dots, \\ & r \cos \theta_{n-1} \end{aligned}$$

for $\theta_2, \theta_3, \dots, \theta_{n-1}, r$ parameters; i.e., whenever f_1 is of degree k in $\sin \theta_1$ and $\cos \theta_1$ and, hence, whenever $f_1 = 1, \cos \theta_1, \sin \theta_1, \cos 2\theta_1, \sin 2\theta_1, \dots, \cos k\theta_1$ or $\sin k\theta_1$. This requirement leads to the set of $2k + 1$ equations:

$$(2.17) \quad \left. \begin{aligned} \sum_{g=1}^{p_1} a_g &= \int_0^{2\pi} d\theta_1 = 2\pi, \\ \sum_{g=1}^{p_1} a_g \cos t\theta_{1g} &= \int_0^{2\pi} \cos t\theta_1 d\theta_1 = 0, \\ \sum_{g=1}^{p_1} a_g \sin t\theta_{1g} &= \int_0^{2\pi} \sin t\theta_1 d\theta_1 = 0. \end{aligned} \right\} \quad (t = 1, 2, \dots, k)$$

One solution with minimum p_1 for given k is

$$(2.18) \quad \begin{aligned} p_1 &= k + 1 = 4(m + 1), \\ a_g &= \frac{2\pi}{k + 1}, \quad \theta_{1g} = \frac{2\pi g}{k + 1}, \quad g = 1, 2, \dots, k + 1. \end{aligned}$$

This is the only solution, apart from a single arbitrary additive constant to θ_{1g} , for which the a_g are real so the above, (2.18), is a necessary and sufficient condition for the first rule, (2.14), to have the required precision with a minimum number of points.

Put, temporarily,

$$(2.19) \quad \begin{aligned} \theta_l &= \phi, \quad b_{li} = b_i, \quad p_l = p, \quad l = \nu + 1, \quad \text{and} \\ & f_l(\theta_l, \dots, \theta_{n-1}, r) = g(\phi), \end{aligned}$$

so a typical member of the second set of rules, (2.15), is now

$$(2.20) \quad \int_0^\pi \sin^\nu \phi g(\phi) d\phi \doteq \sum_{i=1}^p b_i g(\phi_i), \quad \nu = 1, 2, \dots, n - 2,$$

and is to be exact when the function $g(\phi)$ is a polynomial in $\sin \phi$ and $\cos \phi$ of degree at most k . The integral over the centrally symmetric n -spherical shell of any monomial $x_1^{t_1} x_2^{t_2} \cdots x_n^{t_n}$ with an odd power t_s is zero and is computed exactly by any rule centrally symmetric about the origin. So, provided rule (2.20) is symmetric, it needs to be exact only for all even polynomials in $\sin \phi$ and $\cos \phi$ of degree at most k or, equivalently, for all polynomials in $\cos \phi$ alone of degree at most k . This requirement leads to the $k + 1$ equations:

$$(2.21) \quad \sum_{i=1}^p b_i \cos^t \phi_i = \int_0^\pi \sin^\nu \phi \cos^t \phi d\phi, \quad t = 0, 1, \dots, k.$$

Change the variable to y by the equations

$$(2.22) \quad y = \cos \phi \quad \text{and} \quad y_j = \cos \phi_j,$$

and (2.21) becomes

$$(2.23) \quad \sum_{i=1}^p b_i y_i^t = \int_{-1}^1 (1 - y^2)^{(\nu-1)/2} y^t dy, \quad t = 0, 1, \dots, k.$$

These are the equations determining a rule

$$(2.24) \quad \int_{-1}^1 (1 - y^2)^{(\nu-1)/2} h(y) dy \doteq \sum_{i=1}^p b_i h(y_i)$$

of precision k in y and with weight function $(1 - y^2)^{(\nu-1)/2}$. Now the unique rule of precision $2p - 1$ using a minimum number of points for the weight function $(1 - y)^\alpha (1 + y)^\beta$ over the interval $(-1, 1)$ is the p -point Gauss-Mehler rule, which has abscissae at the zeros of the orthogonal polynomials

$$(2.25) \quad Q_p^{(\alpha, \beta)}(y) \equiv \binom{p + \alpha}{p} G_p(\alpha + \beta + 1, \alpha + 1, \frac{1}{2}(1 - y)),$$

where $G_p(a, b, z)$ are the Jacobi polynomials defined by

$$(2.26) \quad G_p(a, b, z) \equiv F(-p, p + a; b, z),$$

where $F(a, b; c, z)$ is the ordinary hypergeometric series (terminating in this case) and which has weights [2]

$$(2.27) \quad \begin{aligned} b_i &= \frac{1}{Q_p^{(\alpha, \beta)}(y_i)} \int_{-1}^1 \frac{Q_p^{(\alpha, \beta)}(y) dy}{y - y_i} \\ &= \frac{(p + \alpha)!(p + \beta)!}{p!(p + \alpha + \beta)!} \cdot \frac{2^{\alpha+\beta+1}}{(1 - y_i^2)(Q_p^{(\alpha, \beta)}(y_i))^2}. \end{aligned}$$

For the rule (2.24)

$$\alpha = \beta = (\nu - 1)/2 \quad (\nu = 1, 2, \dots, n - 2),$$

and for brevity write

$$(2.28) \quad Q_p^{(\alpha, \alpha)}(y) = Q_p^\alpha(y).$$

It follows that for the rule (2.24) to be of precision $4m + 3$ with a minimum number of points, p , that $p = 2m + 2$, the abscissae y_i are the $2m + 2$ zeros of

$$(2.29) \quad Q_{2m+2}^{(\nu-1, 2)}(y) = \binom{2m + 2 + (\nu - 1)/2}{2m + 2} G_{2m+2}\left(\nu, \frac{\nu + 1}{2}, \frac{1 - y}{2}\right),$$

and the weights are

$$(2.30) \quad b_i = \frac{\left(\left(2m + 2 + \frac{\nu - 1}{2}\right)!\right)^2}{(2m + 2)!(2m + \nu + 1)!} \cdot \frac{2^\nu}{(1 - y_i^2)(Q_{2m+2}^{(\nu-1)/2}(y_i))^2}.$$

On replacing ν by $l - 1$, y by $\cos \theta$, etc., according to (2.19) one gets the result in the second part of the theorem.

Because of the symmetry of the angular integration rules all the monomials with an odd power of some coordinate x_i have their integrals (zero) computed exactly, independently of rule (2.16), so rule (2.16) need be exact only for all even polynomials in r of degree at most $4m + 3$, i.e., one must have

$$(2.31) \quad \int_R^1 r^{n-1} g(r^2) dr \doteq \sum_{j=1}^q c_j g(r_j^2)$$

exact whenever $g(r^2)$ is a polynomial in r^2 of degree at most $2m + 1$ and with q a minimum. Put $r^2 = t$ and this becomes

$$(2.32) \quad \int_{R^2}^1 t^{n/2-1} g(t) dt \doteq \sum_{j=1}^q 2c_j g(t_j).$$

The rule (2.32) of precision $2m + 1$ and minimum number of points, q , is unique [3] and has $q = m + 1$. The t_j are the $m + 1$ zeros of the polynomial $Q_{m+1}(t)$ in t of degree $m + 1$ that is orthogonal with respect to the weight function $t^{n/2-1}$ over $(R^2, 1)$ to all polynomials in t of lower degree; i.e.,

$$(2.33) \quad \int_{R^2}^1 t^{n/2-1} Q_{m+1}(t) T_m(t) dt = 0,$$

where $T_m(t)$ is an arbitrary polynomial of degree at most m , and the weights $2c_j$ are given by:

$$(2.34) \quad 2c_j = \frac{1}{Q'_{m+1}(t_j)} \int_{R^2}^1 \frac{t^{n/2-1} Q_{m+1}(t) dt}{t - t_j}.$$

On replacing t by r^2 , (2.33) and (2.34) become (2.4) and (2.8), respectively, in the last part of the statement of the theorem. This completes the proof.

3. Discussion.

COROLLARY 1. *The weights $\beta_{gh\dots i} = a_g b_{2,h} \cdots b_{n-1,i}$ are positive and their sum is*

$$\frac{2\pi^{n/2}}{\left(\frac{n}{2} - 1\right)!},$$

the area of the surface of the unit n -sphere.

COROLLARY 2. *The weights $\alpha_{gh\dots ij} = a_g b_{2,h} \cdots b_{n-1,ij}$ are positive and their sum is*

$$\frac{1 - R^n}{n} \cdot \frac{2\pi^{n/2}}{\left(\frac{n}{2} - 1\right)!},$$

the volume of the n -spherical shell.

COROLLARY 3. *All evaluation points of the rule lie between the inner and outer n -spherical surfaces of the shell.*

COROLLARY 4. *Integration rule (2.14) uses $4(m + 1)$ points; the $n - 2$ rules (2.15) each use $2(m + 1)$ points and rule (2.16) uses $m + 1$ points so the number of points used by the complete rule (1.7) is*

$$(4(m + 1))(2(m + 1))^{n-2}(m + 1) \quad \text{or} \quad 2^n(m + 1)^n.$$

Peirce's result of $8(m + 1)^3$ points for the three-dimensional shell is recovered by taking $n = 3$.

COROLLARY 5. *When, in the integration rule (2.24), ν is odd:*

$$\nu = 2l + 1 \quad l = 0, 1, 2, \dots,$$

then in (2.25) $\alpha = \beta = l$, an integer, and the rule becomes a Gauss-Christoffel rule for which the $Q_p^{(l)}(y)$ can be written in terms of the Legendre polynomials $P_n(y)$:

$$(3.1) \quad (1 - y^2)^l Q_p^{(l)}(y) = \begin{vmatrix} P_p(y) & P_{p+2}(y) & \dots & P_{p+2l}(y) \\ P_p(1) & P_{p+2}(1) & \dots & P_{p+2l}(1) \\ P_p'(1) & P_{p+2}'(1) & \dots & P_{p+2l}'(1) \\ \dots & \dots & \dots & \dots \\ P_p^{(l-1)}(1) & P_{p+2}^{(l-1)}(1) & \dots & P_{p+2l}^{(l-1)}(1) \end{vmatrix},$$

where [2], [3]

$$P_n^{(l)}(y) \equiv \frac{d^l}{dy^l} P_n(y).$$

When $\nu = 1$ the rule is the classical Gauss' one with $p = 2m + 2$, $y_i = \cos \phi_i$, the zeros of the Legendre polynomial $P_{2m+2}(y)$ and

$$(3.2) \quad b_i = \frac{2}{(1 - y_i^2)(P'_{2m+2}(y_i))^2}.$$

For the integral over the three-dimensional shell ν takes just the single value 1. This particular case is the one given by Peirce [1].

When $\nu = 3$ one gets from (3.1) and the properties of $P_n(y)$

$$(3.3) \quad Q_p^1(y) = \frac{2p + 3}{(p + 1)(p + 2)} \cdot P'_{p+1}(y)$$

so the zeros of $Q_p^1(y)$ are the same as the zeros of $P'_{p+1}(y)$.

COROLLARY 6. *When ν is even put*

$$\nu = 2\lambda, \quad \lambda = 1, 2, \dots,$$

then

$$(3.4) \quad \alpha = \beta = \lambda - \frac{1}{2}$$

and

$$(3.5) \quad Q_p^\alpha(y) = Q_p^{\lambda-1/2}(y) = \binom{p + \lambda - \frac{1}{2}}{p} G_p(2\lambda, \lambda + \frac{1}{2}, \frac{1}{2}(1 + y)).$$

For the case $\lambda = 1$ the polynomial is a particularly simple one and has the explicit form for all p [2]:

$$(3.6) \quad Q_p^{1/2}(\cos \phi) = 2 \frac{1 \cdot 3 \cdot 5 \cdots (2p + 1)}{2 \cdot 4 \cdot 6 \cdots (2p + 2)} \cdot \frac{\sin(p + 1)\phi}{\sin \phi}$$

which has zeros:

$$(3.7) \quad \phi_i = \frac{\pi i}{p + 1}.$$

The associated weights are

$$(3.8) \quad b_i = \frac{\pi(1 - \cos^2 \phi_i)}{p + 1}.$$

The values for ϕ_i have the great practical advantage, as have those of the θ_{1q} , of being equally spaced.

The ordinary hypergeometric function $F(a, b; c, z)$ satisfies the relation

$$(3.9) \quad \frac{d}{dz} F(a, b; c, z) = \frac{ab}{c} F(a + 1, b + 1; c + 1, z),$$

so from the definition (2.26) of the Jacobi polynomials, $G_p(a, b, z)$,

$$(3.10) \quad \frac{d}{dz} G_{p+1}(a, b, z) = \frac{-(p + 1)(p + 1 + a)}{b} G_p(a + 2, b + 1, z).$$

From (3.10):

$$(3.11) \quad \begin{aligned} G_p(2(\alpha + \frac{3}{2}), \alpha + 2, z) \\ = \frac{-(\alpha + 1)}{(p + 1)(p + 2(\alpha + 1))} \cdot \frac{d}{dz} G_{p+1}(2(\alpha + \frac{1}{2}), \alpha + 1, z). \end{aligned}$$

Now, using (2.25), (2.28) and (3.11), one finds

$$Q_p^{\alpha+1}(y) = \binom{p + \alpha + 1}{p} \frac{-(\alpha + 1)}{(p + 1)(p + 2(\alpha + 1))} \cdot \frac{d}{dz} G_{p+1}(2(\alpha + \frac{1}{2}), \alpha + 1, z)$$

where $z = (1 - y)/2$ so

$$Q_p^{\alpha+1}(y) = \frac{-1}{(p + 2(\alpha + 1))} \cdot \frac{d}{dz} Q_{p+1}^{\alpha}(y).$$

From this

$$(3.12) \quad Q_p^{\alpha+1}(y) = \frac{2}{p + 2(\alpha + 1)} \cdot \frac{d}{dy} Q_{p+1}^{\alpha}(y),$$

a useful differential recursion relation.

One can use this relation to find $Q_p^{3/2}$ from $Q_{p+1}^{1/2}$:

$$(3.13) \quad Q_p^{3/2} = \frac{2}{p + 3} \frac{d}{dy} Q_{p+1}^{1/2}(y)$$

where $y = \cos \phi$ so, by (3.6),

$$(3.14) \quad Q_p^{3/2} = \frac{2}{p + 3} \cdot 2 \cdot \frac{1 \cdot 3 \cdot 5 \cdots (2p + 3)}{2 \cdot 4 \cdot 6 \cdots (2p + 4)} \cdot \frac{d}{d\phi} \left(\frac{\sin(p + 2)\phi}{\sin \phi} \right) \cdot \frac{d\phi}{dy}.$$

Now

$$(3.15) \quad \begin{aligned} \frac{d}{d\phi} \left(\frac{\sin(p + 2)\phi}{\sin \phi} \right) \cdot \frac{d\phi}{dy} \\ = \frac{2}{2 \sin^3 \phi} ((p + 3) \sin(p + 1)\phi - (p + 1) \sin(p + 3)\phi) \end{aligned}$$

so the zeros of $Q_p^{3/2}$ are the roots (in $(0, \pi)$) of

$$(3.16) \quad (p + 3) \sin (p + 1)\phi - (p + 1) \sin (p + 3)\phi = 0.$$

This does not have a simple explicit solution for all the ϕ ; and for all the relevant p so it is unlikely that any $Q_p^{\lambda-1/2}$ where $\lambda > 1$ has such a solution. For each value of p a separate solution has to be found. A few are given in Table I.

COROLLARY 7. *When $R = 0$ the radial rule (2.32) becomes*

$$(3.17) \quad \int_0^1 t^{n/2-1} g(t) dt \doteq \sum_{j=1}^{m+1} 2c_j g(t_j),$$

and the polynomials $Q_{m+1}(t)$ of equations (2.4) and (2.8) are simply the Jacobi polynomials $G_{m+1}(n/2, n/2, t)$.

For $n = 3$ Peirce [1] has tabulated the $Q_{m+1}(t)$, t_j and c_j for $m = 0$ and 1 (also for $R = \frac{1}{4}, \frac{1}{2}$ and $\frac{3}{4}$). For n an even number, $n/2 - 1$ is an integer and (3.17) is a Gauss-Christoffel rule for which the $Q_{m+1}(t)$ can be obtained either from the Legendre polynomials (but with argument $2t - 1$) as in (3.1) or as a Jacobi polynomial above. The tables of Fishman [4] give normalised $Q_{m+1}(t)$ and values of t_j and $2c_j$ above for $n = 2, 4, 6, \dots, 12$ and for $m = 0, 1, 2, \dots, 7$. (His n is my $n/2 - 1$ and his m is my $m + 1$.)

Table II contains some values of r_j^2 and c_j for $m = 0, 1$. (Note: for n an odd number Fishman's tables *could* be used *directly*—treating the sphere as a cone with a spherical base—to give an integration rule but the number of points used in such a rule would not be a minimum since it would unnecessarily compute exactly integrals of certain odd powers of r .)

COROLLARY 8. *If the integration rule (1.2) is modified to include the weight function r^s (s an integer), all the foregoing is still applicable except that in (2.4), (2.8), etc., $n + s$ must be substituted for n so, for example, the table of c_j and r_j for the five-dimensional shell can be used for the three-dimensional shell with weight function r^2 .*

The author is grateful to the referee for drawing his attention to the work in References 5 and 6.

School of Mathematics
University of New South Wales
Sydney, New South Wales,
Australia

1. W. H. PEIRCE, "Numerical integration over the spherical shell," *MTAC*, v. 11, 1957, p. 244-249.
2. Z. KOPAL, *Numerical Analysis*, Wiley, New York, 1955, p. 367-386.
3. G. SZEGÖ, *Orthogonal Polynomials*, Amer. Math. Soc. Colloq. Publ. Vol. 23, Amer. Math. Soc., Providence, R. I., 1939.
4. H. FISHMAN, "Numerical integration constants," *MTAC*, v. 11, 1957, p. 1-9.
5. A. H. STROUD & D. SECREST, "Approximate integration formulas for certain spherically symmetric regions," *Math. Comp.*, v. 17, 1963, p. 105-135.
6. I. P. MYSOVSKIY, "Cubature formulas for evaluating integrals over a sphere," *Dokl. Akad. Nauk SSSR*, v. 147, No. 3, 1962, p. 552-555. (Russian)