

when $n = 15$, for .69379239, read .69378268; when $n = 19$, for .39464231, read .39465249. The remaining entries require additive last-place corrections as follows:

n	Correction	n	Correction
2	+5	11	+11
3	3	12	7
4	6	13	6
5	18	14	37
7	25	16	24
8	8	17	21
9	4	18	19
10	19	20	19

Furthermore, the exponents in the floating-point values of S_n^0 when $n = 18$, 19, 20 should each be decreased by unity.

J. A. ROBERTS

North Carolina State of the University of North Carolina
Raleigh, North Carolina

EDITORIAL NOTE: The exponents corresponding to $n = 14$, 18, 19, 20 in the values of S_n^0 appearing in the same table should also be decreased by unity.

378.—JEAN PETERS, *Eight-Place Tables of Trigonometric Functions for Every Second of Arc*, Chelsea Publishing Company, New York, 1965.

On p. 503, the value of $\cos 25^\circ 5' 53''$ should read 0.90558319, instead of 0.90538319. This typographical error appears also in the first printing (1963), in which other errors have been previously found [1].

ROBERT C. MILLER

Walsh and Company
Albuquerque, New Mexico

1. *Math. Comp.*, v. 19, 1965, p. 174, MTE 363. See also *Math. Comp.*, v. 18, 1964, p. 509, RMT 65.

CORRIGENDUM

D. TEICHROEW, "Use of continued fractions in high speed computing," *MTAC*, v. 6, 1952, pp. 127-133.

On p. 129, under method III, the denominator of the formula for $1 + \rho_i$ should read $1 + r_i(1 + \rho_{i-1})$, instead of $1 + r_i(1 - \rho_{i-1})$.

HENRY E. FETTIS

Aerospace Research Laboratories
Wright-Patterson Air Force Base, Ohio