

5. E. C. TITCHMARSH, *The Theory of Functions*, Oxford Univ. Press, London, 1939.
 6. E. C. TITCHMARSH, *Eigenfunction Expansions Associated With Second-Order Differential Equations*, Oxford, at the Clarendon Press, 1946; Part I, Oxford Univ. Press, London, 1962. MR 8, 458.
 7. G. N. WATSON, *A Treatise on the Theory of Bessel Functions*, 2nd ed., Cambridge Univ. Press and Macmillan, New York, 1944. MR 6, 64.
 8. W. H. YOUNG, "On series of Bessel functions," *Proc. London Math. Soc.*, (2), v. 18, 1920, pp. 163-200.

On a Numerical Solution of an Integral Equation with Singularities

By Robert G. Voigt

1. **Introduction.** Annular airfoil theory gives rise to Fredholm integral equations of the second kind in the following form:

$$(1) \quad f(x) = g(x) + \int_0^1 G(x, y)f(y) dy, \quad 0 \leq x \leq 1,$$

where the kernel $G(x, y)$ has the form

$$G(x, y) = \int_0^1 \frac{q(y, z)}{z - x} dz,$$

and $g(x)$ is a continuous function; in particular it may be of the form

$$(2) \quad g(x) = \int_0^1 \frac{r(x, z)}{z - x} dz.$$

For what follows, we will assume that $q(y, z)$ and $r(x, z)$ are continuous functions as they would be in most physical problems; however, the results are valid for more general functions. By using a Fourier series technique given in Collatz [1], we are able to neatly evaluate the singular integrals involved, but as will be seen, this is not the only advantage of the technique. We also obtain a kernel function of degenerate type; that is

$$G(x, y) = \sum_{i=1}^n m_i(x)M_i(y).$$

Then the integral equation may be solved using a method applicable to degenerate kernels such as the simple one given in Mikhlin [2].

An example of the method applied to an integral equation arising in annular airfoil theory is included at the end of this paper.

2. **Handling the Singularities.** The first step in handling the singularities is to apply the changes of variables suggested by Collatz [1]: Let

$$\begin{aligned} x &= \frac{1}{2}(1 + \cos \theta), \\ y &= \frac{1}{2}(1 + \cos \psi), \\ z &= \frac{1}{2}(1 + \cos \varphi). \end{aligned}$$

Equation (1) may then be written in the form

$$f^*(\theta) = g^*(\theta) + \frac{1}{2} \int_0^\pi G^*(\theta, \psi) f^*(\psi) \sin \psi \, d\psi,$$

where

$$G^*(\theta, \psi) = \int_0^\pi \frac{q^*(\psi, \varphi) \sin \varphi}{\cos \varphi - \cos \theta} \, d\varphi.$$

The * is used to simplify notation, e.g.,

$$f(x) = f[\frac{1}{2}(1 + \cos \theta)] \equiv f^*(\theta).$$

Now we may expand $q^*(\psi, \varphi) \sin \varphi$ in a Fourier cosine series obtaining

$$G^*(\theta, \psi) = \int_0^\pi \frac{\sum_{n=0}^{\infty} a_n(\psi) \cos n\varphi}{\cos \varphi - \cos \theta} \, d\varphi.$$

Since $q^*(\psi, \varphi) \sin \varphi$ is a continuous function of φ , the series is uniformly convergent. Thus we may write

$$G^*(\theta, \psi) = \sum_{n=0}^{\infty} a_n(\psi) \int_0^\pi \frac{\cos n\varphi}{\cos \varphi - \cos \theta} \, d\varphi.$$

If we now consider the Cauchy Principal Value of the integral, we have that

$$G^*(\theta, \psi) = \pi \sum_{n=1}^{\infty} \frac{a_n(\psi) \sin n\theta}{\sin \theta}.$$

Since the infinite sum is convergent, we may approximate it to a prescribed degree of accuracy with a finite sum. This yields a kernel of degenerate type.

If $g(x)$ is given by Equation (2) it may be evaluated in the same way that $G(x, y)$ was evaluated. Thus we obtain Equation (1) in a form free of singularities.

3. Solution of the Equation. Equation (1) now has the form

$$(3) \quad f^*(\theta) \doteq g^*(\theta) + \frac{\pi}{2} \int_0^\pi \sum_{n=1}^N \frac{a_n(\psi) \sin n\theta}{\sin \theta} [\sin \psi f^*(\psi)] \, d\psi.$$

Proceeding with the method of solution found in Mikhlin [2] we write Equation (3) in the form

$$(4) \quad f^*(\theta) \doteq g^*(\theta) + \frac{\pi}{2} \sum_{n=1}^N \frac{\sin n\theta}{\sin \theta} C_n,$$

where

$$C_n = \int_0^\pi a_n(\psi) \sin \psi f^*(\psi) \, d\psi.$$

Using Equation (4) in the expression for C_n we may write

$$C_n - \int_0^\pi a_n(\psi) \sin \psi \left[g^*(\psi) + \frac{\pi}{2} \sum_{m=1}^N \frac{\sin m\psi}{\sin \psi} C_m \right] \, d\psi \doteq 0.$$

Since this is true for $n = 1, 2, \dots, N$, we have the following linear system:

$$(5) \quad C_n - \frac{\pi}{2} \sum_{m=1}^N \alpha_{n,m} C_m \doteq \beta_n, \quad n = 1, 2, \dots, N,$$

where

$$\beta_n = \int_0^\pi a_n(\psi) \sin \psi g^*(\psi) d\psi,$$

and

$$\alpha_{n,m} = \int_0^\pi a_n(\psi) \sin m\psi d\psi.$$

After evaluating β_n and $\alpha_{n,m}$, the system (5) may be solved for C_n . Substitution of C_n values into Equation (4) then gives a solution to (1).

4. Example. The following equation arises as part of the solution of the problem of finding the shape of an annular airfoil from a specified pressure distribution on the inner and outer surfaces of the airfoil, $p^-(x)$ and $p^+(x)$, respectively [3]. The annular airfoil has been nondimensionalized to have length 1 with the trailing edge at 0, thus $x \in [0, 1]$. The equation is in the form of Equation (1) with $g(x)$ in the form of Equation (2):

$$f(x) = \frac{1}{\pi^2 \sqrt{x(1-x)}} \left\{ - \int_0^1 \frac{r(x,z)}{z-x} dz + \int_0^1 \left[\int_0^1 \frac{q(y,z)}{z-x} dz \right] f(y) dy \right\},$$

$$r(x,z) = \sqrt{z(1-z)} \left\{ \frac{h}{2} \int_0^1 u(t)k(z,t)[K[k(z,t)] - E[k(z,t)]] dt + \pi w(z) \right\},$$

$$q(y,z) = \frac{\sqrt{z(1-z)}}{z-y} \{1 - k(y,z)E[k(y,z)]\}.$$

The apparent square root singularities at $x = 0$ and $x = 1$ are removed when the changes of variables are made since $f(x)$ represents a first derivative; i.e., the slope of the thickness distribution. Thus for this particular example

$$f(x) = -\frac{f^*(\theta) \sin \theta}{2}.$$

$K[k(z,t)]$ and $E[k(z,t)]$ are complete elliptic integrals of the first and second kind, respectively, with modulus $k(z,t) = 1/\sqrt{(h^2(z-t)^2 + 1)}$. The quantity h is a specified constant, and $u(t)$ and $w(z)$, which are given, depend on p^- and p^+ ; a complete discussion is given in Reference [3].

The above example has been run on the IBM-7090 at the David Taylor Model Basin. The following table illustrates the effect of the number of coefficients on the stability of the solution. N is the number of coefficients used in the evaluation of the kernel function G ; the next column is the number of significant places in the integral term of Equation (1) unaffected by increasing N ; M is the number of co-

efficients used in the evaluation of g ; and the last column is the number of significant places in g unaffected by increasing M .

N		M	
5	4	5	2
10	5	10	2
15	5	15	2
20	6	20	2
25	6	25	2
30	6	30	2
35	6	35	3
40	6	40	3
		45	4
		50	4
		55	4

This table indicates that a better method should be investigated for the evaluation of g ; however, in evaluating the kernel, the Fourier series technique seems to be satisfactory, and it is in this part of the problem that the two-fold advantage of the technique is realized. The present program requires less than two minutes on the 7090 for $N = 20$ and $M = 45$.

5. Acknowledgment. This paper covers research initiated by the Hydromechanics Laboratory, David W. Taylor Model Basin, Washington, D. C., under Subproject No. S-R011-0101, Task 0401.

David Taylor Model Basin
Hydromechanics Laboratory
Washington, D. C.

1. L. COLLATZ, *The Numerical Treatment of Differential Equations*, 3rd ed. of English transl., Die Grundlehren der Mathematischen Wissenschaften, Bd. 60, Springer, Berlin, 1960, pp. 505-506. MR **22** #322.

2. S. G. MIKHLIN, *Integral Equations*, Macmillan, New York, 1957, pp. 19-21.

3. W. B. MORGAN, "Theory of the annular airfoil and ducted propeller," Fourth Symposium on Naval Hydrodynamics, ACR-92, U. S. Government Printing Office, Washington, D. C., 1962, pp. 151-197.