

# A New Function Associated with the Prime Factors of $\binom{n}{k}$

By E. F. Ecklund, Jr., P. Erdős and J. L. Selfridge

**Abstract.** Let  $g(k)$  denote the least integer  $> k + 1$  so that all the prime factors of  $\binom{g(k)}{k}$  are greater than  $k$ . The irregular behavior of  $g(k)$  is studied, obtaining the following bounds:

$$k^{1+\epsilon} < g(k) < \exp(k(1 + o(1))).$$

Numerical values obtained for  $g(k)$  with  $k \leq 52$  are listed.

The prime factors of  $\binom{n}{k}$  have been studied a great deal. In a recent paper, Erdős [2] stated several results and unsolved problems on this subject. In this paper, we discuss one of the problems stated there: Denote by  $g(k)$  the least integer  $> k + 1^*$  so that all prime factors of  $\binom{g(k)}{k}$  are greater than  $k$ . Determine or estimate  $g(k)$ .

The behavior of  $g(k)$  is surprisingly irregular. We searched for values of  $g(k) \leq 2500000$  for  $2 \leq k \leq 100$ ; the results of this search are reported in Table 1. In reviewing Table 1, we noticed the surprising example  $g(28) = 284$ . This motivated a second search for other such examples with  $g(k) \leq 100000$  and  $101 \leq k \leq 500$ ; none were found.

TABLE 1. Values of  $g(k) \leq 2500000$  for  $2 \leq k \leq 100$

| $k$ | $g(k)$ | $k$ | $g(k)$ | $k$ | $g(k)$ | $k$ | $g(k)$ | $k$ | $g(k)$   |
|-----|--------|-----|--------|-----|--------|-----|--------|-----|----------|
|     |        | 11  | 47     | 21  | 14871  | 31  | 341087 | 41  | <i>B</i> |
| 2   | 6      | 12  | 174    | 22  | 19574  | 32  | 371942 | 42  | 96622    |
| 3   | 7      | 13  | 2239   | 23  | 35423  | 33  | 6459   | 43/ | <i>B</i> |
| 4   | 7      | 14  | 239    | 24  | 193049 | 34  | 69614  | 45  |          |
| 5   | 23     | 15  | 719    | 25  | 2105   | 35  | 37619  | 46  | 692222   |
| 6   | 62     | 16  | 241    | 26  | 36287  | 36  | 152188 | 47/ | <i>B</i> |
| 7   | 143    | 17  | 5849   | 27  | 1119   | 37  | 152189 | 51  |          |
| 8   | 44     | 18  | 2098   | 28  | 284    | 38  | 487343 | 52  | 366847   |
| 9   | 159    | 19  | 2099   | 29  | 240479 | 39  | 767919 | 53/ | <i>B</i> |
| 10  | 46     | 20  | 43196  | 30  | 58782  | 40  | 85741  | 100 |          |

*B:  $g(k)$  exceeds the search bound of 2500000*

The following conjectures on  $g(k)$  all seem certainly true, and perhaps some of them will not be difficult to prove. First, we conjecture

- (1)  $\limsup_{k \rightarrow \infty} g(k + 1)/g(k) = \infty$  and
- (2)  $\liminf_{k \rightarrow \infty} g(k + 1)/g(k) = 0$ .

Received May 7, 1973.

AMS (MOS) subject classifications (1970). Primary 10H15.

\* The condition  $g(k) > k + 1$  was inserted to avoid the special case  $k + 1 = p$ , a prime.

Copyright © 1974, American Mathematical Society

Also, it seems that  $g(k)$  is not of polynomial growth—in other words, for every  $n$  and  $k > k_0(n)$ ,

$$(3) \quad g(k) > k^n.$$

On the other hand,

$$(4) \quad \lim_{k \rightarrow \infty} g(k)^{1/k} = 1$$

certainly seems to hold, and we expect that

$$(5) \quad g(k) < \exp(c_1 \pi(k))$$

is true.

We now give lower and upper bounds for  $g(k)$ . For a lower bound, we show there is an absolute constant  $c > 0$  such that

$$(6) \quad g(k) > k^{1+c}.$$

We first show that  $g(k) > 2k$  (for  $k > 4$ ) always holds. By definition,  $g(k) > k + 1$ , and  $g(k) \neq 2k$  since  $\binom{2k}{k}$  is always even. Suppose  $g(k) = k + t$  with  $1 < t < k$ . We have  $\binom{k+t}{k} = \binom{k+t}{t}$ . Ecklund [1] showed that  $\binom{k+t}{t}$  has a prime factor not exceeding  $(k + t)/2 < k$ , the only exception being  $\binom{3}{2}$  which corresponds to the case  $k = 4$ ,  $t = 3$ . Erdős and Selfridge [2, p. 406] proved that if  $m \geq 2k$ , then  $\binom{m}{k}$  always has a prime factor  $< m/k^c$ , for some absolute constant  $c > 0$ . This immediately implies (6).

Next, we give a very crude upper bound on  $g(k)$ . Denote by  $L_k$  the least common multiple of the integers  $1, 2, \dots, k$  and put  $P_l = \prod_{p \leq l} p$ . Let  $N(k, l) = L_k P_l$ . If  $n + 1$  is any multiple of  $N(k, l)$ , then

$$\binom{n}{k} = \left( \frac{mN(k, l)}{1} - 1 \right) \left( \frac{mN(k, l)}{2} - 1 \right) \dots \left( \frac{mN(k, l)}{k} - 1 \right)$$

has no prime factors less than  $l$ . Thus,

$$(7) \quad g(k) < N(k, k) = \prod_{p \leq k} p^{\alpha_p+1},$$

where  $\alpha_p = [\log_p k]$ . For  $k > k_0$ , this upper bound can be improved a bit. We show

$$(8) \quad g(k) < k^2 L_k P_l \quad \text{with } l = [6k/\log k].$$

To prove (8), consider the integers  $tL_k P_l - 1$  for  $1 \leq t \leq k^2$ . We show that, for at least one of these values of  $t$ ,

$$(9) \quad p \nmid \binom{tL_k P_l - 1}{k} \quad \text{for every } p \leq k.$$

For  $p \leq l$ , (9) holds as before. If  $l < p \leq k$ ,

$$p \mid \binom{tL_k P_l - 1}{k}$$

can only hold if there is a  $j$ ,  $1 \leq j \leq k$ , for which

$$(10) \quad tL_k P_l \equiv j \pmod{p^{\alpha_p+1}}.$$

The number of integers  $t$  with  $1 \leq t \leq k^2$ , for which (10) holds, is at most

$$(11) \quad k(\lfloor k^2/p^2 \rfloor + 1), \quad \text{since } \alpha_p = 1 \text{ for } p > l.$$

Thus, by (10) and (11), the number of integers  $t$ ,  $1 \leq t \leq k$ , for which (10) holds for some prime  $p$ ,  $l < p \leq k$ , is at most

$$(12) \quad \sum_{l < p \leq k} k(\lfloor k^2/p^2 \rfloor + 1) < k^3 \sum_{p > l} 1/p^2 + k\pi(k).$$

It easily follows from the prime number theorem that, for  $k > k_0$ ,

$$(13) \quad \sum_{p > l} 1/p^2 < \frac{2}{l \log l} < \frac{1}{2k}.$$

From (12) and (13), for  $k > k_0$ , the number of integers  $t$ ,  $1 \leq t \leq k$ , for which (10) holds, is less than  $k^2/2 + k\pi(k) < k^2$ . Thus, there is a  $t \leq k^2$  with (9) holding for every  $p \leq k$ . Thus,  $g(k) < k^2 L_k P_l$  as stated. The value 6 could be replaced by a smaller constant, but we cannot prove  $g(k) < L_k$ , which seems to hold for all  $k$ .

It is well known that  $L_k < \exp(k(1 + o(1)))$  and  $k^2 P_l < \exp(o(k))$ . Thus,  $g(k) < \exp(k(1 + o(1)))$ . So  $g(k) < L_k$  should be achievable.

Department of Mathematical Sciences  
Northern Illinois University  
DeKalb, Illinois 60115

1. E. F. ECKLUND, JR., "On prime divisors of the binomial coefficient," *Pacific J. Math.*, v. 29, 1969, pp. 267–270. MR 39 #5465.

2. P. ERDÖS, "Some problems in number theory," in *Computers in Number Theory*, Academic Press, London, 1971, pp. 405–414.