

The Second Largest Prime Factor of an Odd Perfect Number

By Carl Pomerance

Abstract. Recently Hagis and McDaniel have studied the largest prime factor of an odd perfect number. Using their results, we begin the study here of the second largest prime factor. We show it is at least 139. We apply this result to show that any odd perfect number not divisible by eight distinct primes must be divisible by 5 or 7.

1. Introduction. Suppose $n = p_1^{a_1} p_2^{a_2} \cdots p_t^{a_t}$ is an odd perfect number (opn) where a_1, a_2, \dots, a_t are positive integers and $p_1 > p_2 > \cdots > p_t$ are primes. In [7], Kanold proved that $p_1 \geq 61$. Recently, Hagis and McDaniel [5], [6] have succeeded in showing that

$$(1) \quad p_1 \geq 100129.$$

It is the purpose of this paper to study the second largest prime factor of an opn. In Section 2 we develop a method of attack. In Section 3 we prove that

$$(2) \quad p_2 \geq 139.$$

The proof of (2) makes only marginal direct use of computers (we used some computer factorizations in the construction of Table 3). But the proof of (2) does depend strongly on (1), and (1) could not have been accomplished without electronic assistance.

In Section 4 we illustrate the value of the seemingly weak (2). Indeed, in Pomerance [9] and Robbins [10] it is shown that any opn is divisible by at least 7 distinct primes. Sylvester [11] showed that if an opn n is divisible by precisely 7 distinct primes, then $3|n$. In Section 4, using (1) and (2), we prove that if an opn n is divisible by precisely 7 distinct primes, then either $5|n$ or $7|n$. This result would be very difficult to establish without the use of (2).

Before we proceed, it should be pointed out that there is an effective (but not practical) procedure for deciding the following:

Problem. Given any k, N , either find an opn with k th largest prime factor $p_k < N$, or prove no such opn exists.

Indeed, Dickson [2] and Gradštein [4] proved that for any given m there are only finitely many opn's divisible by at most m distinct primes, and it is clear, at least from Dickson's proof, that these opn's are effectively computable. Hence, to resolve the above problem, one need only examine the finite set of opn's divisible by at most $k + \pi(N) - 2$ distinct primes (where $\pi(N)$ denotes the number of primes less than N). Indeed, if the k th largest prime p_k of the opn n satisfies $p_k < N$, then n is divisible by

Received July 10, 1974.

AMS (MOS) subject classifications (1970). Primary 10A20.

Copyright © 1975, American Mathematical Society

at most $k - 1$ distinct primes not less than N . Together with the $\pi(N) - 1$ odd primes less than N , we see that n is divisible by at most $k + \pi(N) - 2$ distinct primes.

Using this procedure to prove (2), one would “merely” have to enumerate the set of opn 's divisible by at most 33 distinct primes. However, this would be an enormous undertaking and hardly practical. Since we do know that every opn is divisible by at least 7 distinct primes, this method gives the result $p_2 \geq 17$. Using the additional fact that no opn is divisible by $3 \cdot 5 \cdot 7$ (Sylvester [12]), we can get $p_2 \geq 19$.

By the above discussion, for any N there are at most a finite number of opn 's with second largest prime factor $p_2 < N$. We denote this set by $\mathcal{P}(N)$. Hence, given any prime p , the set $E(p, N) = \{a > 0: p^a \parallel n \text{ for some } n \in \mathcal{P}(N)\}$ is finite (we write $x \parallel y$ if $x|y$ and $(x, y/x) = 1$). The main goal of Section 2 is to develop procedures for proving $E(p, N) = \emptyset$ or perhaps $E(p, N) \subset S$ where S is some small, explicitly determined set of positive integers.

I wish to acknowledge the expert assistance of David E. Penney on the computer work used for constructing Table 3.

2. Notation, Preliminaries, and a Theorem. If p is a prime and m is a natural number, we write $v_p(m)$ for the exponent (possibly 0) on p in the prime factorization of m . If $p \nmid m$, we write $\text{ord}_p(m)$ for the least positive integer h for which $p|(m^h - 1)$. We write

$$a_p(m) = 0, \quad \text{if } \text{ord}_p(m) = 1;$$

$$a_p(m) = v_p(m^h - 1), \quad \text{if } h = \text{ord}_p(m) > 1.$$

LEMMA 1. *If p is a prime and $m = q^c > 1$ where q, c are natural numbers, then $a_p(q) \leq a_p(m) + v_p(m - 1)$.*

We denote by $\sigma(m)$ the sum of the positive divisors of m . We note that if q is a prime, then $\sigma(q^c) = (q^{c+1} - 1)/(q - 1)$ for any natural number c . The following lemma is a corollary of Theorems 94 and 95 in Nagell [8].

LEMMA 2. *If p, q are odd primes and a, c are natural numbers, then $p^a \parallel \sigma(q^c)$ if and only if $\text{ord}_p(q)|(c + 1)$ and $v_p(c + 1) = a - a_p(q)$.*

LEMMA 3. *If p, q are odd primes, a, b are natural numbers, and $\sigma(p^a) = q^b$, then $a_p(q) = 0$ or 1.*

Proof. We have $q^b - 1 = p + p^2 + \dots + p^a$, so that $p|(q^b - 1)$. Let $h = \text{ord}_p(q)$. Then $h|b$ and $p|(q^h - 1)$.

We state now a result found often in the literature; it is originally due to Bang [1]: Given arbitrary integers $a \geq 2, b \geq 2$, there is a prime p with $\text{ord}_p(a) = b$, unless (i) $a = 2$ and $b = 6$, or (ii) $a = 2^k - 1$ for some k and $b = 2$.

LEMMA 4. *If q is an odd prime, c is a natural number, and $4 \nmid \sigma(q^c)$, then for each divisor $d > 1$ of $c + 1$, there is a prime $p|\sigma(q^c)$ with $\text{ord}_p(q) = d$.*

Proof. We apply the theorem quoted above to the natural numbers q, d , noting that $q \neq 2$ so (i) does not apply, and since $4 \nmid \sigma(q^c)$ we have $c + 1$ odd if $q = 2^k - 1$ for some k , so (ii) does not apply. Hence there is a prime p with $\text{ord}_p(q) = d$. Also, $p|(q^d - 1)/(q - 1)|(q^{c+1} - 1)/(q - 1) = \sigma(q^c)$.

Suppose now $N < 100129$ and p is an odd prime. We denote by $A(p, N)$ the set

of all q^c , q an odd prime, c a natural number, with

- (i) $q < N$;
- (ii) $p \mid \sigma(q^c)$;
- (iii) $4 \nmid \sigma(q^c)$;
- (iv) if $c' < c$, then $v_p(\sigma(q^{c'})) < v_p(\sigma(q^c))$; and either
- (v) every prime factor of $\sigma(q^c)$ is less than N ; or
- (vi) $\sigma(q^c)$ has precisely one prime factor $r \geq N$ and $r \geq 100129$.

We remark that given any q, p, N , there are at most finitely many c with $q^c \in A(p, N)$. Indeed, Lemma 2 and condition (iv) imply that if $c + 1 \neq \text{ord}_p(q)$, then $p \mid (c + 1)$. But if $c + 1 = pm$ where $m \geq N$, then Lemma 4 implies there are primes r_1, r_2 with $r_1 r_2 \mid \sigma(q^c)$, $\text{ord}_{r_1}(q) = m$, and $\text{ord}_{r_2}(q) = pm$. Then $r_1 \geq m + 1 > N$, $r_2 \geq pm + 1 > N$, so that neither (v) nor (vi) is satisfied. We conclude that the set $A(p, N)$ is effectively computable.

We now define several subsets of $A(p, N)$. Let $A_1(p, N)$ denote the subset of those q^c for which $2 \mid \sigma(q^c)$; let $A_2(p, N)$ denote the subset of those q^c for which $2 \nmid \sigma(q^c)$ and (v) holds; and let $A_3(p, N)$ denote the subset of those q^c for which $2 \nmid \sigma(q^c)$ and (vi) holds. Let

$$a_1(p, N) = \max \{v_p(\sigma(q^c)): q^c \in A_1(p, N)\},$$

$$a_2(p, N) = a_1(p, N) + \sum_q \max_c \{v_p(\sigma(q^c)): q^c \in A_2(p, N)\},$$

$$a_3(p, N) = a_1(p, N) + \sum_q \max_c \{v_p(\sigma(q^c)): q^c \in A_2(p, N) \cup A_3(p, N)\}.$$

Let

$$Q(p, N) = \{r \geq 100129: r \text{ prime, } r \mid \sigma(q^c) \text{ for some } q^c \in A_3(p, N)\},$$

$$b(p, N) = \max \{v_p(q - 1): q \text{ prime, } q < N\}.$$

We recall now the definitions of $P(N)$ and $E(p, N)$ from Section 1.

LEMMA 5. Suppose $n \in P(N)$ where $N < 100129$, p_1 is the largest prime factor of n , p is any prime factor of n , $p_1^{a_1} \parallel n$, and $p^a \parallel n$. Then $v_p(\sigma(p_1^{a_1})) \geq a - a_3(p, N)$. If $p_1 \notin Q(p, N)$, then $v_p(\sigma(p_1^{a_1})) \geq a - a_2(p, N)$.

Proof. Let the prime factorization of n be written $p_1^{a_1} p_2^{a_2} \cdots p_t^{a_t}$ where $p_1 \geq 100129 > N > p_2 > \cdots > p_t$. Now $\sigma(n) = 2n$, so $a = \sum_{i=1}^t v_p(\sigma(p_i^{a_i}))$. But $\sum_{i=2}^t v_p(\sigma(p_i^{a_i})) \leq a_3(p, N)$, and if $p_1 \notin Q(p, N)$, then $\sum_{i=2}^t v_p(\sigma(p_i^{a_i})) \leq a_2(p, N)$.

THEOREM 1. Let $N < 100129$ and let p denote an odd prime. Each of the following conditions implies $a \notin E(p, N)$:

- (i) $4 \mid \sigma(p^a)$;
- (ii) there is a prime $q \mid \sigma(p^a)$ with $N \leq q < 100129$;
- (iii) there are primes q_1, q_2 with $q_1 q_2 \mid \sigma(p^a)$ and $q_1 > q_2 \geq N$;
- (iv) $\sigma(p^a)$ is divisible by a prime $q \geq N$ and $a > a_3(p, N) + b(p, N) + a_p(q)$;
- (v) $\sigma(p^a)$ is divisible by a prime $q \geq N$, $q \notin Q(p, N)$, and $a > a_2(p, N) + b(p, N) + a_p(q)$;
- (vi) every prime divisor of $\sigma(p^a)$ is at least N and $a > a_3(p, N) + b(p, N) + 1$;

(vii) $\sigma(p^a) = m_1 m_2$ where every prime divisor of m_1 is less than N , every prime divisor of m_2 is at least N , $m_2 > 1$, and $a > a_3(p, N) + b(p, N) + a_p(m_2) + v_p(m_2 - 1)$;

(viii) $p < N$, $a + 1$ is prime, and $a + 1 \geq \max\{\frac{1}{2}(N - 1), a_3(p, N) + b(p, N) + 3\}$;

(ix) $p^{a'}$ satisfies one of the above and $(a' + 1)|(a + 1)$.

Proof. Note that (i), (ii), and (iii) are obvious.

Suppose (iv) holds and suppose $n \in \mathcal{P}(N)$ with $p^a \| n$. Then $q = p_1$, the largest prime factor of n . Say $p_1^{a_1} \| n$. Lemma 5 implies $v_p(\sigma(p_1^{a_1})) \geq a - a_3(p, N) > b(p, N) + a_p(p_1)$. Hence Lemma 2 implies $p^{1+b(p, N)} | (a_1 + 1)$. By Lemma 4, there is a prime $r | \sigma(p_1^{a_1})$ with $\text{ord}_r(p_1) = p^{1+b(p, N)}$, so that $v_p(r - 1) \geq 1 + b(p, N)$. Then $r \geq N$. But $r \neq p_1$, contradicting $n \in \mathcal{P}(N)$. Hence $a \notin E(p, N)$.

To show that (v) implies $a \notin E(p, N)$ we proceed as with the proof of (iv), except that we note the condition $p_1 = q \notin Q(p, N)$ implies by Lemma 5 that $v_p(\sigma(p_1^{a_1})) \geq a - a_2(p, N)$.

Now assume (vi) holds. Then we may assume $\sigma(p^a) = q^b$ where q is a prime and $q \geq N$. Lemma 3 implies $a_p(q) = 0$ or 1. Hence $a \notin E(p, N)$ by (iv).

If (vii) holds, we may assume $m_2 = q^b$ where q is a prime and $q \geq N$. Then Lemma 1 and (iv) imply $a \notin E(p, N)$.

Suppose (viii) holds. Since $a + 1$ is prime, if q is a prime divisor of $\sigma(p^a)$, then $\text{ord}_q(p) = a + 1$ or 1 by Lemma 2. In the former case, $q \equiv 1 \pmod{a + 1}$, so $q \geq 2(a + 1) + 1 \geq N$ (since $a + 1 \geq 3$ and $a + 1 \geq \frac{1}{2}(N - 1)$). Suppose $\text{ord}_q(p) = 1$, so that $p \equiv 1 \pmod{q}$. Then $\sigma(p^a) = 1 + p + \dots + p^a \equiv a + 1 \pmod{q}$, so that $a + 1 = q$. Hence $p \geq 2(a + 1) + 1 \geq N$, a contradiction. Thus every prime divisor of $\sigma(p^a)$ is at least N , and since $a > a_3(p, N) + b(p, N) + 1$, (vi) implies $a \notin E(p, N)$.

Finally, suppose (ix) holds. Since $\sigma(p^{a'}) | \sigma(p^a)$, we have $a \notin E(p, N)$ due to our above proofs for (i)–(viii).

Remark. We note that Lemma 4 together with (iii), (viii), and (ix) of Theorem 1 imply that if $a \in E(p, N)$ and $p < N$, then every prime divisor r of $a + 1$ satisfies $r < \max\{\frac{1}{2}(N - 1), a_3(p, N) + b(p, N) + 3\}$ and $v_r(a + 1) \leq b(r, N) + 1$. Hence Theorem 1 provides an effective means for examining the finite set $E(p, N)$.

3. The Proof of (2). In Section 2 we remarked that the sets $A(p, N)$ (and hence the subsidiary notions $A_1(p, N)$, $A_2(p, N)$, $A_3(p, N)$, $Q(p, N)$, $a_1(p, N)$, $a_2(p, N)$, and $a_3(p, N)$) are effectively computable. We have performed these computations for $N = 139$ and $p = 3, 7, 11, 13, 19, 31, 61, 97$, and 127. The fruits of this labor may be found in Tables 1, 2, and 3. Also, $b(p, 139)$ for the above p may be found in Table 1. Making use of this numerical information and Theorem 1, we are able to conclude:

$$\begin{aligned}
 E(3, 139) &\subset \{2, 4\}, & E(31, 139) &= \emptyset, \\
 E(7, 139) &\subset \{2\}, & E(61, 139) &\subset \{1, 2\}, \\
 (3) \quad E(11, 139) &\subset \{2\}, & E(97, 139) &\subset \{1\}, \\
 E(13, 139) &\subset \{1, 2\}, & E(127, 139) &= \emptyset, \\
 E(19, 139) &\subset \{2\}, & &
 \end{aligned}$$

Below we give the details of the proof that $E(3, 139) \subset \{2, 4\}$. These details are fairly representative of the techniques used in establishing the remainder of (3), the proof of which we omit.

Since $a_3(3, 139) + b(3, 139) +_v 3 = 19 < 69 = \frac{1}{2}(139 - 1)$, (viii) and (ix) of Theorem 1 imply that if $a \in E(3, 139)$, then every prime divisor of $a + 1$ is less than 69. Since every prime divisor of $\sigma(3^a)$ is at least 139 for $a = 30, 36, 42, 46, 58, 60, 66$, (vi) and (ix) of Theorem 1 imply that if $a \in E(3, 139)$, then $(a + 1, 31 \cdot 37 \cdot 43 \cdot 47 \cdot 59 \cdot 61 \cdot 67) = 1$. Since $1093|\sigma(3^6)$, $3851|\sigma(3^{10})$, $1871|\sigma(3^{16})$, $1597|\sigma(3^{18})$, $28537|\sigma(3^{28})$, (ii) and (ix) of Theorem 1 imply that if $a \in E(3, 139)$, then $(a + 1, 7 \cdot 11 \cdot 17 \cdot 19 \cdot 29) = 1$. Let $m_2 = \sigma(3^{52})/107$, $\sigma(3^{40})/83$, or $\sigma(3^{22})/47$. Then in each case m_2 is an integer, every prime divisor of m_2 is at least 139, and $a_3(m_2) + v_3(m_2 - 1) = 1$. Then (vii) and (ix) of Theorem 1 imply that if $a \in E(3, 139)$, then $(a + 1, 23 \cdot 41 \cdot 53) = 1$. Since $4|\sigma(3)$, (i) and (ix) of Theorem 1 imply that if $a \in E(3, 139)$, then $(a + 1, 2) = 1$.

Hence if $a \in E(3, 139)$, then every prime divisor of $a + 1$ is found in the set $\{3, 5, 13\}$. Suppose $n \in \mathcal{P}(139)$, $3^a || n$, and $13|(a + 1)$. Then $p_1 = 797161 = \sigma(3^{12})$ is the largest prime divisor of n . Say $p_1^{\alpha_1} || n$. Now $797161 \notin Q(3, 139)$, so Lemma 5 implies $v_3(\sigma(797161^{\alpha_1})) \geq a - a_2(3, 139) = a - 10 \geq 2$. Then Lemma 2 implies $9|(a_1 + 1)$. Hence $151|\sigma(797161^2)|\sigma(797161^{\alpha_1})|2n$, contradicting $n \in \mathcal{P}(139)$. Hence if $a \in E(3, 139)$, then $(a + 1, 13) = 1$. Since $757|\sigma(3^8)$, $4561|\sigma(3^{14})$, and $8951|\sigma(3^{24})$, (ii) and (ix) of Theorem 1 imply that if $a \in E(3, 139)$, then $9 \nmid (a + 1)$, $15 \nmid (a + 1)$, $25 \nmid (a + 1)$. We conclude that $E(3, 139) \subset \{2, 4\}$.

We now use (3) to prove (2). Suppose $n \in \mathcal{P}(139)$. Then (3) implies $31 \nmid n$ and $127 \nmid n$. Now, if $19|n$, (3) implies $19^2 || n$. But $\sigma(19^2) = 3 \cdot 127$ and $127 \nmid n$. Hence $19 \nmid n$. If either $7|n$ or $11|n$, then $7^2 || n$ or $11^2 || n$, respectively. But $\sigma(7^2) = 3 \cdot 19$ and $\sigma(11^2) = 7 \cdot 19$, and $19 \nmid n$. Hence $7 \nmid n$ and $11 \nmid n$. If $97|n$, then $97 || n$. But $\sigma(97) = 2 \cdot 7^2$ and $7 \nmid n$. Hence $97 \nmid n$. If $61|n$, then either $61 || n$ or $61^2 || n$. But $\sigma(61) = 2 \cdot 31$, $\sigma(61^2) = 3 \cdot 13 \cdot 97$, and $31 \nmid n, 97 \nmid n$. Hence $61 \nmid n$. If $13|n$, then $13 || n$ or $13^2 || n$. But $\sigma(13) = 2 \cdot 7$, $\sigma(13^2) = 3 \cdot 61$, and $7 \nmid n, 61 \nmid n$. Hence $13 \nmid n$. Finally, if $3|n$, then $3^2 || n$ or $3^4 || n$. But $\sigma(3^2) = 13$, $\sigma(3^4) = 11^2$, and $11 \nmid n, 13 \nmid n$. Hence $3 \nmid n$.

Summing up, if $n \in \mathcal{P}(139)$, then $(n, 3 \cdot 7 \cdot 11 \cdot 13 \cdot 19 \cdot 31 \cdot 61 \cdot 97 \cdot 127) = 1$. Say the prime factorization of n is $p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_t^{\alpha_t}$ where $p_1 \geq 100129 > 139 > p_2 > \cdots > p_t$. Then

$$2 = \frac{\sigma(n)}{n} = \prod_{i=1}^t \frac{\sigma(p_i^{\alpha_i})}{p_i^{\alpha_i}}$$

$$= \prod_{i=1}^t \frac{p_i^{\alpha_i+1} - 1}{p_i^{\alpha_i}(p_i - 1)} < \prod_{i=1}^t \frac{p_i}{p_i - 1} \leq \frac{100129}{100128} P < 1.9,$$

where P is the product of all $p/(p - 1)$ as p ranges over all primes less than 139 and not equal to 2, 3, 7, 11, 13, 19, 31, 61, 97, 127. This contradiction shows that $\mathcal{P}(139) = \emptyset$; that is, if an opn exists, its second largest prime factor is at least 139.

4. **An Application.** It was noted by Euler [3], that if n is an *opn*, then in n 's prime factorization, every exponent is even except for one exponent which is $\equiv 1 \pmod{4}$ as is the corresponding prime. We also note that if p is a prime, then $\sigma(p^a)/p^a$ is an increasing function of a and $\lim_{a \rightarrow \infty} \sigma(p^a)/p^a = p/(p-1)$. We are now in a position to prove:

THEOREM 2. *If $n = p_1^{a_1} p_2^{a_2} \cdots p_7^{a_7}$ is an *opn* where $p_1 > p_2 > \cdots > p_7$ are primes and a_1, a_2, \dots, a_7 are positive integers, then $p_7 = 3$ and $p_6 = 5$ or 7 .*

Proof. As we remarked in Section 1, Sylvester [11] proved that $p_7 = 3$.

Suppose $p_6 \geq 11$. Then $p_6 = 11, p_5 = 13, p_4 = 17$, and $p_3 = 19$. Indeed, if not, using (1) and (2) we have

$$\begin{aligned} 2 &= \frac{\sigma(n)}{n} = \prod_{i=1}^7 \frac{\sigma(p_i^{a_i})}{p_i^{a_i}} < \prod_{i=1}^7 \frac{p_i}{p_i - 1} \\ &\leq \frac{3}{2} \cdot \frac{11}{10} \cdot \frac{13}{12} \cdot \frac{17}{16} \cdot \frac{23}{22} \cdot \frac{139}{138} \cdot \frac{100129}{100128} < 2, \end{aligned}$$

a contradiction. We next note that $a_7 \geq 4, a_6 \geq 4, a_5 \geq 4$, and $a_3 \geq 4$. Indeed, if $a_7 = 2$, then

$$\frac{\sigma(n)}{n} < \frac{\sigma(3^2)}{3^2} \cdot \frac{11}{10} \cdot \frac{13}{12} \cdot \frac{17}{16} \cdot \frac{19}{18} \cdot \frac{139}{138} \cdot \frac{100129}{100128} < 2.$$

If $a_6 = 2$, then $7|\sigma(11^2)|n$. If $a_5 = 1$, then $7|\sigma(13)|2n$. If $a_5 = 2$, then $61|\sigma(13^2)|n$. If $a_3 = 2$, then $127|\sigma(19^2)|n$.

We now show that $a_4 \geq 4$. First, suppose $a_4 = 2$. Then $p_2 = 307 = \sigma(17^2)$. Then

$$\frac{\sigma(n)}{n} > \frac{\sigma(3^4)}{3^4} \cdot \frac{\sigma(11^4)}{11^4} \cdot \frac{\sigma(13^4)}{13^4} \cdot \frac{\sigma(17^2)}{17^2} \cdot \frac{\sigma(19^4)}{19^4} \cdot \frac{\sigma(307^2)}{307^2} > 2,$$

a contradiction. Next suppose $a_4 = 1$. Then from 1.13 in [9], we have $p_1 \equiv p_2 \equiv 1 \pmod{17}$. But $p_2 \geq 139$, so $p_2 \geq 239$. Then since

$$\frac{\sigma(3^4)}{3^4} \cdot \frac{11}{10} \cdot \frac{13}{12} \cdot \frac{\sigma(17)}{17} \cdot \frac{19}{18} \cdot \frac{239}{238} \cdot \frac{100129}{100128} < 2,$$

we have $a_7 \geq 6$. But $1093|\sigma(3^6)$ and $1093 \not\equiv 1 \pmod{17}$, so $a_7 \geq 8$. Now

$$\frac{\sigma(3^8)}{3^8} \cdot \frac{\sigma(11^4)}{11^4} \cdot \frac{\sigma(13^4)}{13^4} \cdot \frac{\sigma(17)}{17} \cdot \frac{\sigma(19^4)}{19^4} \cdot \frac{\sigma(647^2)}{647^2} > 2,$$

so $p_2 > 647$. Then since $p_2 \equiv 1 \pmod{17}$, we have $p_2 \geq 919$. Then

$$\frac{\sigma(n)}{n} < \frac{3}{2} \cdot \frac{11}{10} \cdot \frac{13}{12} \cdot \frac{\sigma(17)}{17} \cdot \frac{19}{18} \cdot \frac{919}{918} \cdot \frac{100129}{100128} < 2,$$

a contradiction. Hence $a_4 \geq 4$.

Also since

$$\frac{\sigma(3^6)}{3^6} \cdot \frac{\sigma(11^4)}{11^4} \cdot \frac{\sigma(13^4)}{13^4} \cdot \frac{\sigma(17^4)}{17^4} \cdot \frac{\sigma(19^4)}{19^4} > 2,$$

we have $a_7 = 4$.

Using the natural generalization of 1.11 from [9], since $a_4 \geq 4$, we have for either $p = p_1$ or $p = p_2$ that $p \equiv 1 \pmod{4}$ and $p \equiv -1 \pmod{17^4}$. Then $p \equiv 167041 \pmod{334084}$. But 167041 is not prime, so $p > 500000$. Now, if $p_2 \geq 569$, we have

$$\frac{\sigma(n)}{n} < \frac{\sigma(3^4)}{3^4} \cdot \frac{11}{10} \cdot \frac{13}{12} \cdot \frac{17}{16} \cdot \frac{19}{18} \cdot \frac{569}{568} \cdot \frac{500001}{500000} < 2.$$

Hence $p_2 \leq 563$. But then

$$\frac{\sigma(n)}{n} > \frac{\sigma(3^4)}{3^4} \cdot \frac{\sigma(11^4)}{11^4} \cdot \frac{\sigma(13^4)}{13^4} \cdot \frac{\sigma(17^4)}{17^4} \cdot \frac{\sigma(19^4)}{19^4} \cdot \frac{\sigma(563)}{563} > 2,$$

a contradiction.

TABLE 1

p	$b(p, 139)$	$a_1(p, 139)$	$A_1(p, 139)$
3	3	3	5, 5 ⁵ , 17, 29, 41, 53, 89, 101, 113, 137
7	1	2	5 ⁵ , 13, 41, 97
11	1	1	109
13	1	0	∅
19	0	1	37, 113
31	0	1	37 ⁵ , 61
61	0	0	∅
97	0	0	∅
127	0	0	∅

TABLE 2

p	$a_2(p, 139)$	$A_2(p, 139)$
3	10	7 ² , 13 ² , 19 ² , 37 ² , 61 ² , 67 ² , 79 ²
7	11	11 ² , 23 ² , 37 ² , 67 ² , 79 ² , 107 ² , 137 ²
11	4	3 ⁴ , 5 ⁴
13	4	3 ² , 29 ² , 61 ² , 107 ²
19	3	7 ² , 11 ²
31	3	5 ² , 67 ²
61	2	13 ² , 47 ²
97	1	61 ²
127	2	19 ² , 107 ²

TABLE 3

p	$a_3(p, 139)$	$A_3(p, 139)$	$Q(p, 139)$
3	13	$13^8, 61^8, 67^8$	1609669, 903870199, 30152894311
7	12	29^6	88009573
11	8	$23^{10}, 97^4, 103^4, 137^4$	262321, 319411, 10332211, 3937230404603
13	4	\emptyset	\emptyset
19	6	$23^8, 47^8, 61^8$	7792003, 567332587, 903870199
31	4	97^4	262321
61	3	131^4	973001
97	1	\emptyset	\emptyset
127	2	\emptyset	\emptyset

Department of Mathematics
University of Georgia
Athens, Georgia 30602

1. A. S. BANG, "Taltheoretiske Undersøgelser," *Tidsskrift Math.*, v. 5, 1886, pp. 70–80, 130–137.
2. L. E. DICKSON, "Finiteness of the odd perfect and primitive abundant numbers with n distinct prime factors," *Amer. J. Math.*, v. 35, 1913, pp. 413–422.
3. L. EULER, "Tractatus de numerorum doctrina," *Commentationes Arithmeticae Collectae*, v. 2, 1849, p. 514.
4. I. S. GRADŠTEIN, "O nečetnych soveršenných číslah," *Mat. Sb.*, v. 32, 1925, pp. 476–510.
5. P. HAGIS, JR. & W. L. McDANIEL, "On the largest prime divisor of an odd perfect number," *Math. Comp.*, v. 27, 1973, pp. 955–957. MR 48 #3855.
6. P. HAGIS, JR. & W. L. McDANIEL, "On the largest prime divisor of an odd perfect number. II," *Math. Comp.*, v. 29, 1975, pp. 922–924.
7. H.-J. KANOLD, "Folgerungen aus dem Vorkommen einer Gaussischen Primzahl in der Primfaktorenzerlegung einer ungeraden vollkommenen Zahl," *J. Reine Angew. Math.*, v. 186, 1944, pp. 25–29. MR 6, 255.
8. T. NAGELL, *Introduction to Number Theory*, 2nd ed., Chelsea, New York, 1964. MR 30 #4714.
9. C. POMERANCE, "Odd perfect numbers are divisible by at least seven distinct primes," *Acta Arith.*, v. 25, 1974, pp. 265–300.
10. N. ROBBINS, *The Non-Existence of Odd Perfect Numbers With Less Than Seven Distinct Prime Factors*, Doctoral Dissertation, Polytechnic Institute of Brooklyn, June 1972.
11. J. J. SYLVESTER, "Sur une classe spéciale des diviseurs de la somme d'une série géométrique," *Comptes Rendus*, v. CVI, 1888, pp. 446–450.
12. J. J. SYLVESTER, "Sur les nombres parfaits," *Comptes Rendus*, v. CVI, 1888, pp. 403–405.