

## Odd Perfect Numbers Not Divisible by 3 Are Divisible by at Least Ten Distinct Primes

By Masao Kishore

**Abstract.** Hagis and McDaniel have shown that the largest prime factor of an odd perfect number  $N$  is at least 100111, and Pomerance has shown that the second largest prime factor is at least 139. Using these facts together with the method we develop, we show that if  $3 \nmid N$ ,  $N$  is divisible by at least ten distinct primes.

**1. Introduction.** A positive integer  $N$  is called perfect if  $\sigma(N) = 2N$ ,  $\sigma(N)$  being the sum of positive divisors of  $N$ . No odd perfect (OP) numbers are known; however, it has been proved that if  $N$  is OP and  $\omega(N)$  denotes the number of distinct prime factors of  $N$ , then  $\omega(N) \geq 5$  by Sylvester (1888), Dickson (1913) and Kanold (1949);  $\omega(N) \geq 6$  by Gradstein (1925), Kühnel (1949) and Weber (1951);  $\omega(N) \geq 7$  by Pomerance (1972, [1]) and Robbins (1972);  $\omega(N) \geq 8$  by Hagis (1975, [3]); and that if  $N$  is OP and  $3 \nmid N$ , then  $\omega(N) \geq 8$  by Sylvester (1888), and  $\omega(N) \geq 9$  by Kanold (1949, [6]). Also, it has been proved that if  $N$  is OP, then  $N > 10^{200}$  by Buxton and Elmore (1976, [5]), the largest prime factor of  $N > 100110$  by Hagis and McDaniel (1975, [4]), and the second largest prime factor of  $N \geq 139$  by Pomerance (1975, [2]).

In this paper we prove

**THEOREM.** *If  $N$  is OP and  $3 \nmid N$ ,  $\omega(N) \geq 10$ .*

**2. Preliminary results.** Throughout this paper let

$$N = \prod_{i=1}^r p_i^{a_i},$$

where  $p_1 < p_2 < \dots < p_r$  are odd primes and  $a_1, \dots, a_r$  are positive integers. We call  $p_i^{a_i}$  a component of  $N$  and write  $V_{p_i}(N)$  for  $a_i$ .

Euler proved that if  $N$  is OP, then for some  $j$ ,  $p_j \equiv a_j \equiv 1 \pmod{4}$  and for  $i \neq j$ ,  $a_i \equiv 0 \pmod{2}$ .  $p_j$  is called the special prime denoted by  $\Pi$ .

**LEMMA 1.** *Suppose  $N$  is OP,  $3 \nmid N$ , and  $p^a$  is a component of  $N$ . If  $p \equiv 2 \pmod{3}$ , then  $p \neq \Pi$ , and if  $p \equiv 1 \pmod{3}$ , then  $a \not\equiv 2 \pmod{3}$ .*

*Proof.* If  $p \equiv 2 \pmod{3}$  and  $p = \Pi$ , then  $\sigma(p^a) \equiv 0 \pmod{3}$  because  $a$  is odd, while if  $p \equiv 1 \pmod{3}$  and  $a \equiv 2 \pmod{3}$ , then  $\sigma(p^a) \equiv 0 \pmod{3}$ , both of which contradict the fact that  $3 \nmid N$ . Q.E.D.

From Euler's Theorem and Lemma 1 we have

**COROLLARY 1.** *Suppose  $N$  is OP,  $3 \nmid N$ , and  $p^a$  is a component of  $N$ . If  $p \equiv 1 \pmod{4}$  and  $p \equiv 1 \pmod{3}$ , then  $a = 1, 4, 6, 9, 10, 12, \dots$ ; if  $p \equiv 1 \pmod{4}$  and  $p \equiv 2 \pmod{3}$ ,*

---

Received December 4, 1975; revised June 2, 1976.

AMS (MOS) subject classifications (1970). Primary 10A20.

Copyright © 1977, American Mathematical Society

then  $a = 2, 4, 6, 8, 10, 12, \dots$ ; if  $p \not\equiv 1 \pmod{4}$  and  $p \equiv 1 \pmod{3}$ , then  $a = 4, 6, 10, 12, \dots$ ; if  $p \not\equiv 1 \pmod{4}$  and  $p \equiv 2 \pmod{3}$ , then  $a = 2, 4, 6, 8, 10, 12, \dots$ .

We are interested in finding  $p_i^{a_i}$ 's for which

$$\prod_{i=1}^r S(p_i^{a_i}) = 2, \quad \text{where } S(p^a) = \sigma(p^a)/p^a.$$

Since the accuracy of the computer is limited, we use the inequality

$$(1) \quad 0.693147180 < \log 2 = \sum_{i=1}^r \log S(p_i^{a_i}) < 0.693147181.$$

With nine-digit figures we have sufficient accuracy, and with log we can easily control computational errors involved.

Suppose  $N$  is OP,  $3 \nmid N$ , and  $p^a$  is a component of  $N$ . We define

$$a(p) = \text{minimum } \{b \mid b > 1 \text{ is an allowable power of } p \text{ as determined by Corollary 1 and } p^{b+1} > 10^9\}$$

and

$$L(p^a) = \begin{cases} [10^9 \log S(p^a)] 10^{-9} & \text{if } a < a(p), \\ \left[10^9 \log \frac{p}{p-1}\right] 10^{-9} & \text{if } a \geq a(p), \end{cases}$$

where  $[ ]$  is the greatest integer function.

We note that if  $p$  and  $q$  are odd primes with  $p < q$ , then for any positive integers  $a$  and  $b$

$$S(q^a) < \frac{q}{q-1} < \frac{p+1}{p} \leq S(p^b),$$

and so  $L(q^a) \leq L(p^b)$ .

LEMMA 2. Suppose

$$N = \prod_{i=1}^r p_i^{a_i}$$

is OP and  $3 \nmid N$ . Then

$$(2) \quad S_r < \sum_{i=1}^r L(p_i^{b_i}) < T_r,$$

where  $S_r = 0.693147180 - r \cdot 10^{-9}$ ,  $T_r = 0.693147181 + r \cdot 10^{-9}$ ,  $b_i = a_i$  if  $a_i < a(p_i)$ , and  $b_i = a(p_i)$  if  $a_i \geq a(p_i)$ .

*Proof.* Since  $N$  is OP, (1) holds. Suppose  $p^a$  is a component of  $N$ . If  $a < a(p)$ , then

$$|\log S(p^a) - L(p^a)| < 10^{-9}.$$

If  $a \geq a(p)$ , then

$$\begin{aligned} 10^{-9} &\geq \log \frac{p}{p-1} - L(p^a) > \log S(p^a) - L(p^{a(p)}) \\ &\geq \log \frac{p^{a+1} - 1}{p^{a+1} - p^a} - \log \frac{p}{p-1} = \log \left(1 - \frac{1}{p^{a+1}}\right) = -\sum_{i=1}^{\infty} \frac{1}{i(p^{a+1})^i} \\ &> -\sum_{i=1}^{\infty} \frac{1}{(p^{a+1})^i} = \frac{-1}{p^{a+1} - 1} \geq -10^{-9}, \end{aligned}$$

and so

$$|\log S(p^a) - L(p^a(p))| < 10^{-9}.$$

Hence,

$$(3) \quad \left| \sum_{i=1}^r \log S(p_i^{a_i}) - \sum_{i=1}^r L(p_i^{b_i}) \right| < r \cdot 10^{-9},$$

and (2) follows from (1) and (3). Q.E.D.

We also need the following lemmas, which were proved in [1, pp. 269–271]:

LEMMA 3. *If  $q$  is a prime for which  $q - 1$  is a power of 2,  $N$  is OP, and if  $p^a$  is a component of  $N$ , then*

$$V_q(\sigma(p^a)) = \begin{cases} V_q(a + 1) & \text{if } p \equiv 1 \pmod{q}, \\ V_q(p + 1) + V_q(a + 1) & \text{if } p \equiv -1 \pmod{q} \text{ and } p = \Pi, \\ 0 & \text{otherwise.} \end{cases}$$

LEMMA 4. *If  $N$  is OP,  $p^a$  is a component of  $N$ , and if  $q$  is a prime and  $q^b | a + 1$ , then  $N$  is divisible by at least  $b$  distinct primes  $\equiv 1 \pmod{q}$  other than  $p$ .*

LEMMA 5. *If  $n$  is OP,  $17^a$  is a component of  $N$ , and if  $17^a \nmid \Pi + 1$ , then  $N$  is divisible by at least two primes  $\equiv 1 \pmod{17}$ .*

3. **Proof of the Theorem.** In this section, we shall prove that if  $3 \nmid N$  and  $\omega(N) = 9$ , then  $N$  is not OP.

LEMMA 6. *If  $N$  is OP,  $3 \nmid N$ , and if  $\omega(N) = 9$ , then*

$$p_1 = 5, \quad p_2 = 7, \quad p_3 = 11, \quad p_4 = 13, \quad p_5 \leq 19, \quad p_6 \leq 23, \quad p_7 \leq 53, \\ p_8 \geq 139 \quad \text{and} \quad p_9 > 100110.$$

*Proof.* By [4]  $p_9 > 100110$ , and by [2]  $p_8 \geq 139$ . Others follow from

$$\frac{5}{4} \frac{7}{6} \frac{11}{10} \frac{17}{16} \frac{19}{18} \frac{23}{22} \frac{29}{28} \frac{139}{138} \frac{100111}{100110} < 2,$$

$$\frac{5}{4} \frac{7}{6} \frac{11}{10} \frac{13}{12} \frac{23}{22} \frac{29}{28} \frac{31}{30} \frac{139}{138} \frac{100111}{100110} < 2,$$

$$\frac{5}{4} \frac{7}{6} \frac{11}{10} \frac{13}{12} \frac{17}{16} \frac{29}{28} \frac{31}{30} \frac{139}{138} \frac{100111}{100110} < 2,$$

and

$$\frac{5}{4} \frac{7}{6} \frac{11}{10} \frac{13}{12} \frac{17}{16} \frac{19}{18} \frac{59}{58} \frac{139}{138} \frac{100111}{100110} < 2. \quad \text{Q.E.D.}$$

LEMMA 7.  $p_5 = 17$  in Lemma 6.

*Proof.* Suppose  $p_5 = 19$ . Then  $p_6 = 23, p_7 = 29$  and  $p_8 = 139$  because

$$\frac{5}{4} \frac{7}{6} \frac{11}{10} \frac{13}{12} \frac{19}{18} \frac{23}{22} \frac{31}{30} \frac{139}{138} \frac{100111}{100110} < 2$$

and

$$\frac{5}{4} \frac{7}{6} \frac{11}{10} \frac{13}{12} \frac{19}{18} \frac{23}{22} \frac{29}{28} \frac{149}{148} \frac{100111}{100110} < 2.$$

Hence

$$N = 5^{a_1} 7^{a_2} 11^{a_3} 13^{a_4} 19^{a_5} 23^{a_6} 29^{a_7} 139^{a_8} p_9^{a_9}.$$

Since

$$\frac{5}{4} \frac{7}{6} S(11^2) \frac{13}{12} \frac{19}{18} \frac{23}{22} \frac{29}{28} \frac{139}{138} \frac{100111}{100110} < 2$$

and

$$\frac{5}{4} \frac{7}{6} \frac{11}{10} S(13^1) \frac{19}{18} \frac{23}{22} \frac{29}{28} \frac{139}{138} \frac{100111}{100110} < 2,$$

$a_3 \neq 2$  and  $a_4 \neq 1$ . Also,  $a_2, a_4, a_5, a_8 \neq 2$  by Corollary 1. Since every odd prime factor of  $\sigma(p_i^{a_i})$  is a factor of  $N$ ,  $a_1, a_6, a_7 \neq 2$  and  $a_1, a_2 \neq 4$  because  $31 \mid \sigma(5^2)$ ,  $71 \mid \sigma(5^4)$ ,  $2801 \mid \sigma(7^4)$ ,  $79 \mid \sigma(23^2)$  and  $67 \mid \sigma(29^2)$ . Hence for  $1 \leq i \leq 2$ ,  $a_i \geq 6$  and for  $3 \leq i \leq 8$ ,  $a_i \geq 4$ . Then  $N$  is not OP because

$$S(N) > \prod_{i=1}^8 S(p_i^{a_i}) > 2. \quad \text{Q.E.D.}$$

LEMMA 8.  $17^{a_5} \mid \Pi + 1$  and  $\Pi > 100110$  in Lemma 6.

*Proof.* Suppose  $17^{a_5} \nmid \Pi + 1$ . Since  $p_i \neq \pm 1$  (17) for  $1 \leq i \leq 7$ ,  $p_8 \equiv p_9 \equiv 1$  (17) by Lemma 5. If  $17^2 \mid \sigma(p_j^{a_j})$  for  $j = 8$  or  $9$ , then by Lemma 3,  $17^2 \mid a_j + 1$ , and by Lemma 4  $N$  would be divisible by at least two primes  $\equiv 1$  (17) other than  $p_j$ . Hence  $17^2 \nmid \sigma(p_j^{a_j})$ . Since  $17 \nmid \sigma(p_i^{a_i})$  for  $1 \leq i \leq 7$ , we conclude that  $a_5 = 2$ ,  $17 \mid \sigma(p_8^{a_8})$  and  $17 \mid \sigma(p_9^{a_9})$ . Then  $p_8 = \sigma(17^2) = 307$ , and for  $j = 8, 9$ ,  $a_j = 16$ ,  $p_j \neq \Pi$ ,  $5 \nmid \sigma(p_j^{a_j})$ , and so for some  $1 \leq i \leq 7$ ,  $5 \mid \sigma(p_i^{a_i})$ . By Lemma 3 and Corollary 1,  $p_i = 11, 31$ , or  $41$ , and  $\sigma(p_i^4) \mid \sigma(p_i^{a_i})$  because  $5 \mid a_i + 1$ ; however,  $3221 \mid \sigma(11^4)$ ,  $17351 \mid \sigma(31^4)$ ,  $579281 \mid \sigma(41^4)$ , and none of these primes  $\equiv 1$  (17). Hence  $p_i \neq 11, 31, 41$ , a contradiction, and  $17^{a_5} \mid \Pi + 1$ .

If  $a_5 \geq 4$ ,  $\Pi \geq 2 \cdot 17^4 - 1 = 167041$ , while if  $a_5 = 2$ ,  $\Pi = p_9 > 100110$  because  $p_8 = 307$ . Q.E.D.

LEMMA 9. If  $3 \nmid N$ ,  $\omega(N) = 9$ , and if  $p_8 > 1000$ ,  $N$  is not OP.

*Proof.* Suppose  $N$  is OP. Then by Lemma 2

$$S_9 < \sum_{i=1}^9 L(p_i^{b_i}) < T_9.$$

If  $a_i < a(p_i)$ ,  $b_i = a_i$ , and so every prime factor of  $\sigma(p_i^{b_i})$  is a factor of  $N$  except when  $p_i = \Pi$ . Hence if

$$M = \left( \prod_{i=1}^7 p_i \right) \left( \prod_{i=1; b_i < a(p_i)}^7 \sigma(p_i^{b_i}) \right),$$

we have

(4)  $\omega(M) = 7,$

(5)  $\omega(M) = 8, \text{ or}$

(6)  $\omega(M) = 9.$

Suppose (4) holds. Since  $p_8 > 1000$  and  $p_9 > 100110$ ,

$$(7) \quad S_9 < \sum_{i=1}^7 L(p_i^{b_i}) + \log \frac{1009}{1008} + \log \frac{100111}{100110} \quad \text{and} \quad \sum_{i=1}^7 L(p_i^{b_i}) < T_7.$$

Suppose (5) holds, and let  $p$  be the prime factor of  $M$  other than  $p_i, 1 \leq i \leq 7$ . Then

$$(8) \quad \begin{aligned} &\text{if } 1000 < p < 100110, S_9 < \sum_{i=1}^7 L(p_i^{b_i}) + L(p^b) + \log \frac{100111}{100110}, \\ &\text{if } p > 100110, S_9 < \sum_{i=1}^7 L(p_i^{b_i}) + \log \frac{1009}{1008} + L(p^b), \text{ and} \\ &\sum_{i=1}^7 L(p_i^{b_i}) + L(p^b) < T_8, \end{aligned}$$

where  $b \leq a(p)$  is an allowable power of  $p$ . Suppose (6) holds. Then the two prime factors of  $M$  other than  $p_i, 1 \leq i \leq 7$ , are  $p_8$  and  $p_9$ , and

$$(9) \quad p_8 > 1000, \quad p_9 > 100110 \quad \text{and} \quad S_9 < \sum_{i=1}^9 L(p_i^{b_i}) < T_9.$$

Computer was used to find  $\prod_{i=1}^7 p_i^{b_i}$  satisfying

- (A) (4) and (7),
- (B) (5) and (8), or
- (C) (6) and (9),

with the following results:

$$\begin{aligned} &5^{12}7^{10}11^813^917^823^629^6, \quad 5^{12}7^{10}11^813^917^823^629^4, \\ &5^{12}7^{10}11^813^917^823^429^6, \quad 5^{12}7^{10}11^813^917^623^629^6 \\ &5^{12}7^{10}11^813^617^823^629^6, \quad \text{or} \quad 5^{10}7^{10}11^813^917^823^629^6. \end{aligned}$$

In every case  $p_8 \geq 3011$  because

$$S(5^{10}7^{10}11^813^617^623^429^43001^1) > 2.$$

Then  $N$  is not OP because  $p_9 \geq \Pi > 17^6 - 1$  and

$$S(N) < \frac{5}{4} \frac{7}{6} \frac{11}{10} \frac{13}{12} \frac{17}{16} \frac{23}{22} \frac{29}{28} \frac{3011}{3010} \frac{p_9}{p_9 - 1} < 2. \quad \text{Q.E.D.}$$

LEMMA 10. *If  $3 \nmid N, \omega(N) = 9$ , and if  $p_8 < 1000, N$  is not OP.*

*Proof.* Suppose  $N$  is OP. Then by Lemma 2

$$S_9 < \sum_{i=1}^9 L(p_i^{b_i}) < T_9.$$

If

$$M = \left( \sum_{i=1}^8 p_i \right) \left( \prod_{i=1; b_i < a(p_i)}^7 \sigma(p_i^{b_i}) \right),$$

then

(10)  $\omega(M) = 8,$  or

(11)  $\omega(M) = 9.$

Suppose (10) holds. Then

(12) 
$$p_8 < 1000, S_9 < \sum_{i=1}^8 L(p_i^{b_i}) + \log \frac{100111}{100110}, \text{ and}$$

$$\sum_{i=1}^8 L(p_i^{b_i}) < T_8.$$

Suppose (11) holds. Then the prime factor of  $M$  other than  $p_i, 1 \leq i \leq 8,$  is  $p_9,$  and

(13) 
$$p_8 < 1000, p_9 > 100110 \text{ and } S_9 < \sum_{i=1}^9 L(p_i^{b_i}) < T_9.$$

Computer was used to find  $\prod_{i=1}^8 p_i^{b_i}$  satisfying

(A) (10) and (12), or

(B) (11) and (13),

with the following results:

$$5^{12}7^{10}11^813^917^819^647^6233^4, \quad 5^{12}7^{10}11^813^917^819^647^6233^2,$$

$$5^{12}7^{10}11^213^917^819^643^6331^4, \quad 5^{27}7^{10}11^813^917^819^631^6953^4,$$

$$5^{27}7^{10}11^213^917^819^631^6557^4, \text{ or } 5^{27}7^{10}11^213^917^819^631^6557^2.$$

Then  $N$  is not OP because in every case  $p_9 = \Pi \geq 2 \cdot 17^8 - 1$  and  $S(N) < 2.$  Q.E.D.  
 Lemmas 9 and 10 prove our theorem.

Computer (PDP 11 at the University of Toledo) program run time for Lemmas 9 and 10 was three minutes.

I would like to thank Professor J. Chidambaraswamy for his help in preparing this paper. Also, I would like to thank the referee who suggested the use of [2], which cut down the computer time.

Department of Mathematics  
 University of Toledo  
 Toledo, Ohio 43606

1. C. POMERANCE, "Odd perfect numbers are divisible by at least seven distinct primes," *Acta Arith.*, v. 25, 1973/74, pp. 265-300. MR 49 #4925.
2. C. POMERANCE, "The second largest prime factor of an odd perfect number," *Math. Comp.*, v. 29, 1975, pp. 914-921. MR 51 #8018.
3. P. HAGIS, JR., "Every odd perfect number has at least eight prime factors," *Notices Amer. Math. Soc.*, v. 22, 1975, p. A-60. Abstract #720-10-14.
4. P. HAGIS, JR. & W. L. McDANIEL, "On the largest prime divisor of an odd perfect number. II," *Math. Comp.*, v. 29, 1975, pp. 922-924. MR 51 #8021.
5. M. BUXTON & S. ELMORE, "An extension of lower bounds for odd perfect numbers," *Notices Amer. Math. Soc.*, v. 23, 1976, p. A-55. Abstract #731-10-40.
6. H.-J. KANOLD, "Folgerungen aus dem Vorkommen einer Gauss'schen Primzahl in der Primfaktorenzerlegung einer ungeraden Vollkommenen Zahl," *J. Reine Angew. Math.*, v. 186, 1944, pp. 25-29. MR 6, 255.