

# On Maximal Finite Irreducible Subgroups of $GL(n, \mathbf{Z})$

## IV. Remarks on Even Dimensions with Applications to $n = 8$

By Wilhelm Plesken and Michael Pohst

**Abstract.** The general methods for the determination of maximal finite absolutely irreducible subgroups of  $GL(n, \mathbf{Z})$  developed in Part I of this series of papers [6] are refined for even  $n$ . Applications are made to  $n = 8$  in view of Part V [7], where a complete classification is obtained.

**1. Introduction.** The general procedure for the determination of the  $\mathbf{Z}$ -classes of maximal finite irreducible (i.e.  $\mathbf{C}$ -irreducible) subgroups of  $GL(n, \mathbf{Z})$  suggested in Part I [6] consists of three steps: finding representatives of the  $\mathbf{Q}$ -classes of the minimal irreducible finite subgroups of  $GL(n, \mathbf{Z})$ , calculating the  $\mathbf{Z}$ -classes of these groups by the centering algorithm, and computing the  $\mathbf{Z}$ -automorphism groups of the quadratic forms fixed by the minimal irreducible finite subgroups of  $GL(n, \mathbf{Z})$ . These methods turned out to be very effective for odd dimensions such as  $n = 5, 7, 9$ , where we had to consider only two, respectively three,  $\mathbf{Q}$ -classes of minimal irreducible subgroups of  $GL(n, \mathbf{Z})$ . On the other hand, for  $n = 6$  this number is already 33 and a cautious estimate yields more than a hundred  $\mathbf{Q}$ -classes for  $n = 8$ . The main reason for these big numbers is that there exist many possibilities for the decomposition scheme of normal abelian subgroups of irreducible matrix groups in  $GL(n, \mathbf{Q})$ , if  $n$  has many even divisors. Moreover, if  $n$  is a power of two, a lot of 2-groups occur. Therefore, it is desirable to have a method which allows us to avoid the determination of all minimal irreducible finite subgroups of  $GL(n, \mathbf{Z})$  in case  $n = 2r$ ,  $r \in \mathbf{N}$ . In Section 2 we describe a method which provides all quadratic forms fixed by a finite irreducible subgroup  $G$  of  $GL(2r, \mathbf{Z})$ , where  $G$  has a  $\mathbf{Q}$ -reducible subgroup of index 2. Note that these groups include all 2-groups in case  $2r$  is a power of 2. The method requires information about the finite irreducible subgroups of  $GL(r, \mathbf{Z})$ .

In Section 3 we carry out the computations for  $2r = 8$  and obtain 17 of 26 primitive positive definite integral quadratic forms the automorphism groups of which are the maximal finite irreducible subgroups of  $GL(8, \mathbf{Z})$ . The remaining discussions for  $n = 8$  and a complete description of the results for dimensions less than 10 appear in Part V [7].

---

Received November 27, 1978.

AMS (MOS) subject classifications (1970). Primary 20C10, 20H15.

Key words and phrases. Integral matrix groups.

**2. Irreducible Subgroups of  $GL(2r, \mathbf{Z})$  Derived from Subgroups of  $GL(r, \mathbf{Z})$  and Associated Forms.** As we already mentioned in the introduction it is desirable to avoid the computation of all  $\mathbf{Q}$ -classes of minimal irreducible finite subgroups of  $GL(n, \mathbf{Z})$ ,  $n = 2r$ . Therefore, we discuss the following two types of these groups  $G$  separately.

Type ( $\alpha$ ):  $G$  has a  $\mathbf{Q}$ -reducible subgroup of index two.

Type ( $\beta$ ):  $G$  has no  $\mathbf{Q}$ -reducible subgroup of index two.

For  $n = 8$  the majority of the groups belongs to Type ( $\alpha$ ), for instance all those groups  $G$  the order of which is a power of two. In the cases already treated we computed all centerings of  $G$  and the corresponding quadratic forms. Since only the forms are used to determine the maximal finite subgroups of  $GL(n, \mathbf{Z})$ , we shall develop a method for groups of Type ( $\alpha$ ) to find the forms without computing all centerings. Moreover, the method allows us to treat many groups simultaneously.


Let  $G$  be an irreducible finite subgroup of  $GL(2r, \mathbf{Z})$  ( $r \in \mathbf{N}$ ) with a  $\mathbf{Q}$ -reducible subgroup  $N$  of index two. Since  $N$  is normal in  $G$ , the restriction  $\Delta|_N$  of the natural representation  $\Delta$  of  $G$  to  $N$  can be assumed to be of the form  $\Delta|_N = \Gamma_1 \dot{+} \Gamma_2$ , where  $\Gamma_1, \Gamma_2$  are rationally inequivalent irreducible integral representations of  $N$  with  $\Gamma_1(N) = \Gamma_2(N)$  (Corollary (6.19) in [3] and Theorem (3.1) in [6]).

A short calculation shows that  $G$  can be chosen as  $G = N \dot{\cup} \begin{pmatrix} 0 & h \\ I_r & 0 \end{pmatrix} N$ , where  $N$  consists of block diagonal matrices and  $h \in \Gamma_1(N) = \Gamma_2(N)$ . If  $X$  denotes the integral positive definite matrix representing a form fixed by  $\Gamma_1(N)$ , then  $I_2 \otimes X$  represents the corresponding form of  $G$ . We are interested in the integral forms which are induced on the centerings of  $G$ . Let  $M = \mathbf{Z}^{2r \times 1}$  be the natural representation module of  $G$  and  $L$  a centering of  $M$ . We use a description of  $M$  as  $\mathbf{Z}N$ -module given in [5].  $M$  splits into a direct sum, say  $M = M_1 \oplus M_2$  with associated projections  $\pi_1, \pi_2$

$$\left( M_1 := \left\{ \begin{pmatrix} l_1 \\ 0 \end{pmatrix} \in M \mid l_1 \in \mathbf{Z}^r \times 1 \right\}, M_2 := \left\{ \begin{pmatrix} 0 \\ l_2 \end{pmatrix} \in M \mid l_2 \in \mathbf{Z}^r \times 1 \right\} \right).$$

We define  $N$ -centerings of  $M_i$ :  $L_i := \pi_i(L)$ ,  $L'_i := M_i \cap L_i$  ( $i = 1, 2$ ) and the finite  $\mathbf{Z}N$ -module  $A := (L_1 \oplus L_2)/L$ . Then  $A \cong L_i/L'_i$  ( $i = 1, 2$ ) holds, and there exist  $\mathbf{Z}N$ -epimorphisms  $\mu_i: L_i \rightarrow A$  such that the kernel of  $\mu_1 \oplus \mu_2: L_1 \oplus L_2 \rightarrow A: (l_1, l_2) \rightarrow \mu_1(l_1) + \mu_2(l_2)$  is equal to  $L$ . Furthermore,  $L$  is not only an  $N$ -centering but also a  $G$ -centering. Therefore,  $\begin{pmatrix} 0 & h \\ I_r & 0 \end{pmatrix} L = L$  and  $\begin{pmatrix} 0 & h \\ I_r & 0 \end{pmatrix} L_1 = L_2$  holds. Hence,  $\mu_1, \mu_2$  can be chosen in such a way that  $\mu_1 \oplus \mu_2$  is a  $\mathbf{Z}G$ -epimorphism. Now, let us assume  $L_1 = M_1$ .\* Then  $L_2 = M_2$  follows. Clearly,  $A$  can be generated by  $m \leq r$  elements. Therefore, homomorphisms of  $L_i$  ( $i = 1, 2$ ) into  $A$ , respectively endomorphisms of  $A$ , can be described by  $m \times r$ , respectively  $m \times m$ , matrices over  $\mathbf{Z}/a\mathbf{Z}$  with  $a := \exp(A)$ . The matrix of such a mapping  $\beta$  is denoted by  $\bar{\beta}$ . Since  $A$  is a  $\mathbf{Z}G$ -module there is a homomorphism  $\alpha: G \rightarrow \text{Aut}_{\mathbf{Z}}(A)$  with  $(\mu_1 \oplus \mu_2)g = \alpha(g)(\mu_1 \oplus \mu_2)$  for all  $g \in G$ . In particular, we obtain  $(\mu_1 \oplus \mu_2)\begin{pmatrix} 0 & h \\ I_r & 0 \end{pmatrix} = \chi(\mu_1 \oplus \mu_2)$  with  $\chi = \alpha\left(\begin{pmatrix} 0 & h \\ I_r & 0 \end{pmatrix}\right)$ . Hence,  $\bar{\mu}_2 = \bar{\chi}\bar{\mu}_1$  and  $\bar{\mu}_1 h = \bar{\chi}\bar{\mu}_2$ . Note that  $\chi^2 \in \alpha(N)$  because of  $\begin{pmatrix} 0 & h \\ I_r & 0 \end{pmatrix}^2 \in N$ .

\*Compare discussion in the second paragraph following (2.1).



In our later computations we want to deal with all groups  $G$  just discussed which simultaneously fix the same quadratic form. This can be done because of the considerations of this paragraph and especially because of the following lemma.

(2.1) LEMMA. *Let  $G, \mu_1, \mu_2, X$  as above and  $k_1, k_2 \in \text{Aut}_{\mathbb{Z}}(X)$ . Then the quadratic form belonging to the centering  $L = \ker(\mu_1 \oplus \mu_2)$  is equal to the one induced by  $X$  on  $\ker(\mu_1 k_1 \oplus \mu_2 k_2)$ .*

*Proof.* Let  $S$  be the matrix of the basis transformation from  $M$  to  $L$ . The form belonging to  $M$  is  $S^t \text{diag}(X, X)S$ . Then  $\text{diag}(k_1^{-1}, k_2^{-2})S$  is the matrix of a basis transformation from  $M$  to  $\ker(\mu_1 k_1 \oplus \mu_2 k_2)$ . Therefore, the form induced by  $\text{diag}(X, X)$  on  $\ker(\mu_1 k_1 \oplus \mu_2 k_2)$  is given by

$$(\text{diag}(k_1^{-1}, k_2^{-1})S)^t \text{diag}(X, X)(\text{diag}(k_1^{-1}, k_2^{-1})S)$$

which is equal to  $S^t \text{diag}(X, X)S$ , since  $k_1^{-1}, k_2^{-1}$  are elements of  $\text{Aut}_{\mathbb{Z}}(X)$ . Q.E.D.

We explicitly describe the form induced by  $I_2 \otimes X$  on  $L = \ker(\mu_1 \oplus \mu_2)$ . Let  $B \in \mathbb{Z}^r \times r$  be the matrix of the basis transformation from  $L_1$  to  $L'_1$ . Since  $\mu_1$  is an epimorphism there is a matrix  $C \in \mathbb{Z}^r \times r$  which satisfies  $\bar{\mu}_1 C + \bar{\chi} \bar{\mu}_1 = 0$ . Then  $S := \begin{pmatrix} B & C \\ 0 & I_r \end{pmatrix}$  is the matrix of the basis transformation from  $M$  to  $L$ . The form induced on  $M$  is  $S^t \text{diag}(X, X)S$ .

For our discussion we assumed  $L_1 = M_1$  so far. If  $L_1$  is a proper  $N$ -centering of  $M_1$ , we transform  $G$  by  $\text{diag}(D, D)$ , where  $D$  is the matrix of a basis transformation from  $M_1$  to  $L_1$ . Note that the new group has the same “block pattern” as  $G$ . However,  $\Gamma_1(N)$  is replaced by a rationally equivalent integral group. This leads us to associate the quadratic forms of the centerings of  $G$  with the  $N$ -centerings of  $M_1$ . Therefore, we proceed in our computation as follows.

Let  $X \in \mathbb{Z}^r \times r$  be the matrix of a positive definite quadratic form with irreducible  $\text{Aut}_{\mathbb{Z}}(X)$ , and let  $K_1, \dots, K_l$  be all minimal irreducible subgroups of  $\text{Aut}_{\mathbb{Z}}(X)$  up to  $\mathbb{Z}$ -equivalence. By  $M = \mathbb{Z}^r \times 1$  we denote the natural representation module of  $\text{Aut}_{\mathbb{Z}}(X)$ . For each submodule  $M'$  of  $M$  which is a centering with respect to one of the  $K_i, i \in \{1, \dots, l\}$ , we define  $K(M')$  to be the biggest subgroup of  $\text{Aut}_{\mathbb{Z}}(X)$  leaving  $M'$  invariant

and  $\alpha'$  to be the homomorphism of  $K(M')$  into  $\text{Aut}_{\mathbf{Z}}(M/M')$  which describes the action of  $K(M')$  on  $M/M'$ .

There are two possibilities for  $M'$  to yield a group  $G$  as considered above for which  $\Gamma_1, \Gamma_2$  are inequivalent. Either  $K(M')$  does not act faithfully on  $M/M'$  or  $\alpha'$  is injective and there exists an irreducible subgroup  $H$  of  $K(M')$  with an outer automorphism  $c$  subject to the following three properties:  $c^2$  is an inner automorphism,  $c$  is induced by the normalizer of  $\alpha'(H)$  in  $\text{Aut}_{\mathbf{Z}}(M/M')$  corresponding to  $H$ , and  $c$  is not induced by the normalizer of  $H$  in  $GL(r, \mathbf{Q})$ .

Let  $\mathcal{N}(M')$  be the set of all  $\chi \in \text{Aut}_{\mathbf{Z}}(M/M')$  for which an irreducible subgroup  $H$  of  $K(M')$  exists which is normalized by  $\chi$  and for which  $\chi^2 \in \alpha'(H)$ . In case  $\alpha'$  is injective  $\chi$  must correspond to an automorphism  $c$  described above. The forms we are interested in are induced by  $I_2 \otimes X$  on  $\ker(\mu_1 \oplus \mu_2)$  with  $\bar{\mu}_2 = \bar{\chi}\bar{\mu}_1$  for  $\chi \in \mathcal{N}(M')$ . However, we know from Lemma (2.1) that the  $\chi$  lying in the same coset of  $\text{Aut}_{\mathbf{Z}}(M/M')$  modulo  $\alpha'(K(M'))$  provide the same form. Hence, it suffices to pick one  $\chi$  out of each coset. If  $M'$  runs through a set of representatives of the  $\text{Aut}_{\mathbf{Z}}(X)$ -orbits of the centerings of  $M$  as discussed above we get all forms derived from  $I_2 \otimes X$ . Moreover, if  $X$  runs through a set of representatives of integral positive definite primitive  $r$ -ary forms with an irreducible automorphism group, we obtain all integral  $2r$ -ary forms an automorphism group of which is irreducible and has a  $\mathbf{Q}$ -reducible subgroup of index 2. This procedure is performed in the next paragraph for  $r = 4$ .

There can be made further simplifications the underlying ideas of which are demonstrated by the following example.

(2.2) LEMMA. *If  $\mathcal{N}(2M)$  is contained in the subgroup induced by  $\text{Aut}_{\mathbf{Z}}(X)$  in  $\text{Aut}_{\mathbf{Z}}(M/2M)$ , each centering  $M' \subseteq 2M$  provides only multiples of quadratic forms which can already be obtained by centerings  $M''$  which are not contained in  $2M$ .*

*Proof.* For  $M' = 2M$  we have  $\bar{\mu}_1 = \bar{\mu}_2 = I_n \in \mathbf{Z}_2^{r \times r}$ . A matrix for the basis transformation is  $S = \begin{pmatrix} I_r & I_r \\ I_r & -I_r \end{pmatrix}$ . Because of  $S^t(I_2 \otimes X)S = 2(I_2 \otimes X)$  the result follows. Q.E.D.

**3. Irreducible Subgroups of  $GL(4, \mathbf{Z})$  and Derived Octonary Forms.** There are—up to  $\mathbf{Z}$ -equivalence—six quaternary integral primitive quadratic forms admitting an irreducible automorphism group [1], [2]. They are represented by the following matrices:

$$Q_1 = I_4, \quad Q_2 = \begin{pmatrix} 2 & 1 & 0 & 0 \\ 1 & 2 & 1 & 1 \\ 0 & 1 & 2 & 0 \\ 0 & 1 & 0 & 2 \end{pmatrix}, \quad Q_3 = I_2 \otimes \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix},$$

$$Q_4 = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix} \otimes \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}, \quad Q_5 = I_4 + J_4, \quad Q_6 = 5I_4 - J_4,$$

where all entries of  $J_4$  are 1.


For each of these forms we have to proceed as described in the last paragraph in

order to obtain the forms belonging to the centerings of the irreducible subgroups of  $\text{Aut}_{\mathbf{Z}}(I_2 \otimes Q_i)$  ( $i = 1, \dots, 6$ ). For this we can make use of the list of the finite subgroups of  $GL(4, \mathbf{Z})$  in [1].

Ad  $Q_1$ . The automorphism group of  $Q_1$  is the full monomial group  $H_4$  of order  $2^4 4!$ . From Theorem (3.2) in [6] one sees immediately that the minimal irreducible subgroups of  $H_4$  are 2-groups. Hence we only have to consider 2-centerings of the natural representation module  $L = \mathbf{Z}^4 \times^1$ . For instance, the extraspecial 2-group

$$\left\langle g \otimes h | g, h \in \left\langle \left( \begin{matrix} -1 & 0 \\ 0 & 1 \end{matrix} \right), \left( \begin{matrix} 0 & 1 \\ 1 & 0 \end{matrix} \right) \right\rangle \right\rangle$$

of order 32 has the following lattice of centerings which contains the centerings of all other irreducible subgroups of  $H_4$ .



The orbits under the action of  $H_4$  are  $\{M_1\}$ ,  $\{M_2\}$ ,  $\{M_3, M_4, M_5\}$ ,  $\{M_6\}$ ,  $\{M_7, M_8\}$ ,  $\{2M_1\}$ ,  $\dots$ . The corresponding bases are expressed in the basis of  $M_1$  via the transformation matrices:

$$(i) B(M_1) = I_4, \quad (ii) B(M_2) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & -1 \end{pmatrix}, \quad (iii) B(M_3) = I_2 \otimes \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix},$$

$$(iv) B(M_6) = \begin{pmatrix} 1 & -1 & 1 & 1 \\ 1 & 1 & -1 & 1 \\ 1 & 1 & 1 & -1 \\ 1 & 1 & 1 & 1 \end{pmatrix}, \quad (v) B(2M_1) = 2I_4,$$

$$(vi) \quad B(M_7) = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}.$$

(3.1) LEMMA. *Only the forms induced by  $I_2 \otimes Q_1 = I_8$  on  $\ker(\mu_1 \oplus \mu_2)$  need to be considered:*

(i)  $\bar{\mu}_1 = \bar{\mu}_2 = (0 \ 0 \ 0 \ 0) \in \mathbf{Z}_2^{1 \times 4},$

(ii)  $\bar{\mu}_1 = \bar{\mu}_2 = (1 \ 1 \ 1 \ 1) \in \mathbf{Z}_2^{1 \times 4},$

(iii)  $\bar{\mu}_1 = \bar{\mu}_2 = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 \end{pmatrix} \in \mathbf{Z}_2^{2 \times 4},$

(iv)  $\bar{\mu}_1 = \bar{\mu}_2 = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \end{pmatrix} \in \mathbf{Z}_2^{3 \times 4},$

(v)  $\bar{\mu}_1, \bar{\mu}_2 \in \left\{ \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{pmatrix} \right\} \subseteq \mathbf{Z}_2^{4 \times 4}.$

(3.2) PROPOSITION. *The forms obtained from Lemma (3.1) are:*

(i)  $F_1 = I_8, \det F_1 = 1^8;$

(ii)  $F_2 = \begin{pmatrix} 2 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 2 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 2 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 2 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 2 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 2 \end{pmatrix}, \det F_2 = 1^6 2^2;$

(iii)  $F_3 = I_2 \otimes \begin{pmatrix} 2 & 1 & 0 & 0 \\ 1 & 2 & 1 & 1 \\ 0 & 1 & 2 & 0 \\ 0 & 1 & 0 & 2 \end{pmatrix}, \det F_3 = 1^4 2^4;$

(iv)  $F_4 = \begin{pmatrix} 2 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 2 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 2 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 2 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 2 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 2 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 2 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 4 \end{pmatrix}, \det F_4 = 1^2 2^6;$

$$(v) \ 2F_1; \ 2F_5 \text{ with } F_5 = \begin{pmatrix} 2 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & -1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 2 & -1 & 0 & 0 & 0 & 0 \\ 0 & -1 & -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 2 \end{pmatrix}, \det F_5 = 1^8; \ 2F_1.$$

*Proof of (3.1).* An easy calculation yields that  $K(M')$  acts faithfully on  $M_1/M'$  except for those centerings  $M'$  containing  $2M_1$ . We consider these exceptions first. In the cases (i) and (ii) we have  $\bar{\mu}_1 = \bar{\mu}_2$  since the automorphism group of  $M_1/M_2$  is trivial. In case (iii)  $K(M_3)$  is a subgroup of index 3 of  $H_4$  and its image in  $\text{Aut}_{\mathbb{Z}}(M_1/M_3)$  is isomorphic to  $C_2$ . Since the image of  $K(M_3)$  in  $\text{Aut}_{\mathbb{Z}}(M_1/M_3)$  is self normal, we again obtain  $\bar{\mu}_1 = \bar{\mu}_2$ . In case (iv) we see that  $K(M_6)$  is equal to  $H_4$  and the subgroup induced by  $K(M_6)$  in  $\text{Aut}_{\mathbb{Z}}(M_1/M_6)$  is isomorphic to the symmetric group  $S_4$  which is the biggest subgroup of  $\text{Aut}_{\mathbb{Z}}(M_1/M_6)$  transforming  $M_2/M_6$  into itself. Since  $M_2$  is the only centering of  $M_1$  of index 2 with respect to an irreducible subgroup of  $H_4$  any element of  $\text{Aut}_{\mathbb{Z}}(M_1/M_6)$  normalizing the image of an irreducible subgroup of  $H_4$  leaves  $M_2/M_6$  invariant and, hence, lies in the image of  $H_4$ . Therefore,  $\bar{\mu}_1 = \bar{\mu}_2$  holds. Also, in case (v)  $K(2M_1)$  is equal to  $H_4$  and the subgroup induced in  $\text{Aut}_{\mathbb{Z}}(M_1/M_2)$  consists of all permutation matrices of degree 4 with respect to the standard basis. Clearly the normalizer of this group is the direct product of the group itself and its centralizer which is easily seen to be generated by  $J_4 + I_4$  (over  $\mathbb{Z}_2$ ). The images of the irreducible subgroups of  $H_4$  are the transitive groups of  $4 \times 4$ -permutation matrices. Some standard arguments show that their normalizers are contained in the normalizer of the group of all permutation matrices. Therefore, we end up with two possibilities for  $\bar{\mu}_2$ .

It remains to discuss the cases in which  $K(M')$  acts faithfully on  $M_1/M'$ . For  $M' = M_7$  (or  $M' = M_8$ )  $K(M')$  is a subgroup of  $H_4$  of index 2 and its image in  $\text{Aut}_{\mathbb{Z}}(M_1/M_7)$  is the biggest automorphism group leaving  $M_2/M_7$  invariant. (Note  $M_6/M_7$  is the Frattini-subgroup of  $M_1/M_7$ .) The same argument as in case (iv) shows that no  $\mu_2$  can exist.

For  $M' = 2M_2$  the module  $\ker(\mu_1 \oplus \mu_2)$  is contained in


$$L := \langle 2(M_1 \oplus M_1), \ker(\mu_1 \oplus \mu_2) \rangle$$

of index 2. Clearly, there are two possible forms which can be induced on  $L$ , namely  $2F_1$  or  $2F_5$ ; compare (3.2). In the first case we can only obtain a form which already occurred in (ii). In the second we observe that the automorphism group of  $F_5$  is the Weyl-group  $W(E_8)$  of the root system  $E_8$ . The  $2^8 - 1$  subgroups of index 2 of the lattice on which  $W(E_8)$  acts fall into two orbits of length 135 and 120. The stabilizer of a lattice in the first orbit is irreducible of order  $2^7 8!$  and the corresponding form is  $F_2$ . The stabilizers of lattices in the second orbit are reducible

since they permute all roots which are orthogonal to a given root. Hence, we can only obtain the form  $F_2$ .

If  $M'$  is a centering properly contained in  $2M_2$ , then there must exist two centerings  $L_1, L_2$  between  $M_1 \oplus M_1$  and  $\ker(\mu_1 \oplus \mu_2)$  which correspond to  $M_6$  and  $2M_2$  and are of index  $2^3$ , respectively  $2^5$ , in  $M_1 \oplus M_1$ . Clearly, we have  $L_1 \supset L_2$ ; and there must be at least one more centering between  $L_1$  and  $L_2$ . The forms induced on the subgroups  $S$  with  $L_1 \supset S \supset L_2$  are easily seen to be equivalent to  $2F_1$  in one case and  $2F_5$  in two cases. Therefore, the subgroup with the form  $2F_1$  must be a centering and a multiple of the form induced on  $\ker(\mu_1 \oplus \mu_2)$  has already been obtained. Q.E.D.

Ad  $Q_2$ . The automorphism group of  $Q_2$  is the Weyl group of the root system  $F_4$ . It is of order  $2^7 3^2 = 1152$  and has a subgroup  $\tilde{H}_4$  of index 3 which is rationally equivalent to  $H_4$ . First, we consider the minimal irreducible subgroups which are rationally equivalent to a subgroup of  $H_4$ . They yield the following centerings:



The orbits under the action of  $\text{Aut}_{\mathbb{Z}}(Q_2)$  are  $\{M_1\}, \{M_2, M_3, M_4\}, \{M_5\}, \{M_6, M_7, M_8\}, \{2M_1\}, \dots$ . Transformation matrices of the corresponding bases:

(i)  $B(M_1) = I_4,$

(ii)  $B(M_2) = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & -2 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{pmatrix},$  (iii)  $B(M_5) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 1 & -1 & 1 & 1 \\ 0 & -1 & 1 & -1 \end{pmatrix},$

(iv)  $B(M_6) = \begin{pmatrix} 2 & 0 & 0 & 0 \\ -2 & 2 & 0 & 0 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & 1 & 1 \end{pmatrix},$  (v)  $B(2M_1) = 2I_4.$



(3.3) LEMMA. Only the forms induced by  $I_2 \otimes Q_2 = F_3$  on  $\ker(\mu_1 \oplus \mu_2)$  need to be considered:

(i)  $\bar{\mu}_1 = \bar{\mu}_2 = (0 \ 0 \ 0 \ 0) \in \mathbb{Z}_2^{1 \times 4}$ ,

(ii)  $\bar{\mu}_1 = \bar{\mu}_2 = (0 \ 1 \ 0 \ 0) \in \mathbb{Z}_2^{1 \times 4}$ ,

(iii)  $\bar{\mu}_1 = \bar{\mu}_2 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 1 \end{pmatrix} \in \mathbb{Z}_2^{2 \times 4}$ ,

(iv)  $\bar{\mu}_1 = \bar{\mu}_2 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 1 \end{pmatrix} \in \mathbb{Z}_2^{3 \times 4}$ .

(3.4) PROPOSITION. The forms obtained from Lemma (6.3) are:

- (i)  $F_3$ ; (ii)  $F_4$ ; (iii)  $2F_5$ ; (iv)  $2F_2$ .

They occurred already in (3.2).


*Proof of (3.3).* The centerings  $M'$  for which  $K(M')$  acts faithfully on  $M_1/M'$  are properly contained in  $2M_1$ . In the cases (i), (ii) and (iii)  $K(M')$  induces the full automorphism group of  $M_1/M'$ . Therefore,  $\bar{\mu}_1 = \bar{\mu}_2$  holds. The argument in the cases (iv) and (v) is exactly the same as in (3.1), case (iv). The rest follows from Lemma (2.2). Q.E.D.

Next, we discuss the minimal irreducible subgroups of  $\text{Aut}_{\mathbb{Z}}(Q_2)$  which are not rationally equivalent to a subgroup of  $H_4$ .

(3.5) LEMMA. If  $U$  is an absolutely irreducible subgroup of  $\text{Aut}_{\mathbb{Z}}(Q_2)$  which has more 2-centerings than  $\text{Aut}_{\mathbb{Z}}(Q_2)$  itself, then  $U$  is rationally equivalent to a subgroup of  $H_4$ .

*Proof.* The 2-centerings of  $\text{Aut}_{\mathbb{Z}}(Q_2)$  are  $M_1, M_5, 2M_1, \dots$ . If  $M_1/M_5$  or  $M_5/2M_1$  become reducible as  $\mathbb{Z}_2 U$ -modules, the statement is obvious, since the forms induced on  $M_2, M_3, M_4, M_6, M_7, M_8$  are multiples of  $Q_1$ . If  $M_1/M_5$  and  $M_5/2M_1$  stay irreducible, then the lattice of 2-centerings with respect to  $U$  is not linearly ordered and the order of  $U$  divides 72. Since  $2^2$  does not divide  $72/4$  this yields a contradiction to Corollary 3.6 in [4]. Q.E.D.

We obtain the following centerings and transformation matrices:



$$(vi) \quad B(M_9) = \begin{pmatrix} 2 & 0 & 0 & -1 \\ -1 & 2 & 1 & 3 \\ 1 & 0 & 0 & -2 \\ 1 & -1 & -2 & -1 \end{pmatrix},$$

$$(vii) \quad B(M_{10}) = \begin{pmatrix} 6 & 0 & -1 & 3 \\ 0 & 6 & -2 & -2 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

$$(viii) \quad B(3M_1) = 3I_4.$$

(3.6) LEMMA. We have only to consider forms which are induced by  $I_2 \otimes Q_2$  on  $\ker(\mu_1 \oplus \mu_2)$ :

$$(vi) \quad \bar{\mu}_1 = \bar{\mu}_2 = \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & -1 \end{pmatrix} \in \mathbf{Z}_3^{2 \times 4},$$

$$(vii) \quad \bar{\mu}_1 = \begin{pmatrix} 2 & 3 & 2 & 0 \\ 3 & 2 & 1 & 1 \end{pmatrix}, \quad \bar{\mu}_2 = \begin{pmatrix} -1 & 0 & -1 & 3 \\ 0 & -1 & -2 & -2 \end{pmatrix} \in \mathbf{Z}_6^{2 \times 4}.$$

(3.7) PROPOSITION. From (3.6) we obtain the forms

$$(vi) \quad F_6 = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix} \otimes \begin{pmatrix} 2 & 1 & 0 & 0 \\ 1 & 2 & 1 & 1 \\ 0 & 1 & 2 & 0 \\ 0 & 1 & 0 & 2 \end{pmatrix}, \quad \det F_6 = 1^4 6^4;$$

$$(vii) \quad F_7 = I_4 \otimes \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}, \quad \det F_7 = 1^4 3^4.$$


*Proof of (3.6).*  $M_9$  is the only centering  $M'$  for which  $K(M')$  acts faithfully on  $M_1/M'$  which we have not yet discussed.  $K(M_9)$  has order 144 and induces the full automorphism group of  $M_1/M_9$ . Hence, in case (iv) we have  $\bar{\mu}_1 = \bar{\mu}_2$ . As for case (vii) we have  $K(M_{10}) = K(M_9)$  and  $K(M_{10})$  acts faithfully on  $M_1/M_{10}$ . The automorphism group  $\overline{K(M_{10})}$  of  $M_1/M_{10}$  induced by  $K(M_{10})$  is of index two in  $\text{Aut}_{\mathbf{Z}}(M_1/M_{10})$ . Hence  $\bar{\mu}_2 = \epsilon \bar{\mu}_1$ , where  $\epsilon$  lies in  $\text{Aut}_{\mathbf{Z}}(M_1/M_{10}) \setminus \overline{K(M_{10})}$ . The centerings contained in  $2M_1$  need not be considered because of Lemma (2.2). The centerings contained in  $3M_1$  need not be considered, since  $\text{Aut}_{\mathbf{Z}}(Q_2)$  acts faithfully on  $M_1/3M_1$  and different representations  $\Delta$  of the irreducible subgroups  $H$  of  $\text{Aut}_{\mathbf{Z}}(Q_2)$  with  $\Delta(H) = H$  can be distinguished by the signs of their character values which can already be determined from the action on  $M_1/3M_1$ . (All character tables are listed in 1.) Q.E.D.

Ad  $Q_3$ . The automorphism group of  $Q_3$  is the wreath product  $\text{Aut}_{\mathbf{Z}}(\begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}) \sim C_2 \wr D_{12} \sim C_2$  of order  $12^2 2$ . As in the case of the form  $Q_2$ , one recognizes that the

subgroups of  $\text{Aut}_{\mathbb{Z}}(Q_3)$  are of two kinds. Either their nontrivial centerings are 3-centerings, or they have just two  $\prec$ -maximal 3-centerings and nontrivial 2-centerings. If  $G$  is a subgroup of the first type, the centerings of  $G$  are also centerings of

$$\left\langle \text{diag} \left( \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \text{diag} \left( \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & I_2 \\ I_2 & 0 \end{pmatrix} \right) \right\rangle.$$

These are given by:



The orbits under the action of  $\text{Aut}_{\mathbb{Z}}(Q_3)$  are  $\{M_1\}$ ,  $\{M_2, M_3\}$ ,  $\{M_4\}$ ,  $\{M_5, M_6\}$ ,  $\{3M_1\}$ ,  $\dots$ .

The corresponding bases are given by the transformation matrices:

$$(i) B(M_1) = I_4, \quad (ii) B(M_2) = \begin{pmatrix} 1 & 1 & -1 & 0 \\ -1 & 2 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad (iii) B(M_4) = I_2 \otimes \begin{pmatrix} 1 & 1 \\ -1 & 2 \end{pmatrix},$$

$$(iv) B(M_5) = \begin{pmatrix} 3 & 0 & -1 & -1 \\ 0 & 3 & 1 & -2 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & -1 & 2 \end{pmatrix}, \quad (v) B(3M_1) = 3I_4.$$

(3.8) LEMMA. Only those forms have to be considered which are induced by

$I_2 \otimes Q_3$  on  $\ker(\mu_1 + \mu_2)$ :

$$(i) \bar{\mu}_1 = \bar{\mu}_2 = (0 \ 0 \ 0 \ 0) \in \mathbb{Z}_3^{1 \times 4},$$

$$(ii) \bar{\mu}_1 = \bar{\mu}_2 = (1 \ 1 \ 1 \ 1) \in \mathbb{Z}_3^{1 \times 4},$$

$$(iii) \bar{\mu}_1, \bar{\mu}_2 \in \left\{ \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & -1 & -1 \\ 1 & 1 & 1 & 1 \end{pmatrix} \right\} \subset \mathbb{Z}_3^{2 \times 4},$$

$$(iv) \bar{\mu}_1 = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 \end{pmatrix}, \quad \bar{\mu}_2 = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 0 & 1 & 0 & 1 \end{pmatrix} \in \mathbb{Z}_3^{3 \times 4}.$$

(3.9) PROPOSITION. From (3.8) we obtain the following forms

$$(i) F_7 = I_4 \otimes \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix} \text{ (occurred already in (3.7));}$$

$$(ii) \quad F_8 = \begin{pmatrix} 4 & -2 & 0 & 0 & -2 & 1 & 1 & 1 \\ -2 & 4 & 0 & 0 & 1 & -2 & -2 & 1 \\ 0 & 0 & 4 & -2 & -1 & 2 & -1 & -1 \\ 0 & 0 & -2 & 4 & -1 & -1 & 2 & -1 \\ -2 & 1 & -1 & -1 & 4 & -2 & -2 & 1 \\ 1 & -2 & 2 & -1 & -2 & 4 & 1 & -2 \\ 1 & -2 & -1 & 2 & -2 & 1 & 4 & -2 \\ 1 & 1 & -1 & -1 & 1 & -2 & -2 & 4 \end{pmatrix}, \quad \det F_8 = 1^3 3^4 9;$$

(iii)  $F_9 = I_2 \otimes \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix} \otimes \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}, \det F_9 = 1^2 3^4 9^2; 3F_5; F_9;$

(iv)  $3F_{10}$  with  $F_{10} = I_8 + J_8, \det F_{10} = 1^7 9.$

*Proof of (3.8).*  $K(M')$  acts faithfully on  $M_1/M'$  exactly for those centerings which are properly contained in  $M_4$ . In the cases (i) and (ii)  $K(M')$  clearly induces the full automorphism group of  $M_1/M'$ , hence  $\bar{\mu}_1 = \bar{\mu}_2$ .


In case (iii) we obtain  $K(M_4) = \text{Aut}_{\mathbb{Z}}(Q_3)$  and the induced subgroup of  $\text{Aut}_{\mathbb{Z}}(M_1/M_4)$  is isomorphic to the dihedral group of order 8. Because of  $|\text{Aut}_{\mathbb{Z}}(M_1/M_4)| = 48$  there are two possibilities for  $\bar{\mu}_2$ .

In case (iv)  $K(M_5)$  acts faithfully on  $M_1/M_5$  and is of order 144. If  $\overline{K(M_5)}$  denotes the induced subgroup of  $\text{Aut}_{\mathbb{Z}}(M_1/M_5)$ , then  $M_4/M_5$  has to be invariant under the normalizer of  $\overline{K(M_5)}$ . Hence, the normalizer is contained in a subgroup of order  $3^2 \cdot 2 \cdot 48$ . The index is 6, and there is at most one possibility for a relevant outer automorphism. We end up with  $\bar{\mu}_1, \bar{\mu}_2$  as given. Finally, if  $M'$  is contained in  $3M_1$ , the same argument as at the end of the proof of (3.6) applies. Q.E.D.

If  $G$  is a subgroup of  $\text{Aut}(Q_3)$  with nontrivial 2-centerings, those centerings are also centerings of the group

$$\left\langle \text{diag} \left( \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix}, \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix} \right), \text{diag} \left( \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right), \text{diag}(-I_2, I_2), \begin{pmatrix} 0 & I_2 \\ I_2 & 0 \end{pmatrix} \right\rangle.$$

They are given by:



The bases are described by the following transformation matrices:

$$(vi) \ B(M_7) = \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & -1 \end{pmatrix}, \quad (vii) \ B(M_8) = \begin{pmatrix} 6 & 0 & 2 & 3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 6 & 3 & 2 \end{pmatrix},$$

$$(viii) \ B(2M_1) = 2I_4, \quad (ix) \ B(2M_4) = 2I_2 \otimes \begin{pmatrix} 1 & 1 \\ -1 & 2 \end{pmatrix}.$$

(3.10) LEMMA. *We have to consider only forms which are induced by  $I_2 \otimes Q_3$  on  $\ker(\mu_1 \oplus \mu_2)$ :*

$$(vi) \quad \bar{\mu}_1 = \bar{\mu}_2 = \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{pmatrix} \in \mathbb{Z}_2^{2 \times 4},$$

$$(vii) \quad \bar{\mu}_1 = \begin{pmatrix} 1 & -2 & 3 & 0 \\ 0 & 3 & -2 & 1 \end{pmatrix}, \quad \bar{\mu}_2 = \begin{pmatrix} 1 & -2 & -1 & 2 \\ -2 & 1 & -2 & 1 \end{pmatrix} \in \mathbb{Z}_6^{2 \times 4}.$$

(3.11) PROPOSITION. *Lemma (3.10) provides the forms*

- (vi)  $F_6$  (occurred already in (3.7)),
- (vii)  $3F_3$  ( $F_3$  occurred already in (3.2)).


*Proof of (3.10).*  $K(M')$  acts faithfully on  $M_1/M'$  except for  $M' \in \{M_4, M_7, 2M_1\}$ . The case  $M' = M_4$  was already treated in (3.8). In case (v) the group  $K(M_7)$  has order 48 and induces the full automorphism group of  $M_1/M_7$ . Hence,  $\bar{\mu}_1 = \bar{\mu}_2$  holds. In case (vi) we have  $K(M_8) = K(M_7)$ , and  $K(M_8)$  induces a subgroup of index 6 in  $\text{Aut}_{\mathbb{Z}}(M_1/M_7)$ . Therefore, we have at most one possibility for a relevant outer automorphism yielding  $\bar{\mu}_2$ .

For  $M' \subseteq 2M_1$  we can apply Lemma (2.2), since  $\text{Aut}_{\mathbb{Z}}(Q_3)$  induces a maximal imprimitive and, hence, self-normal subgroup of  $\text{Aut}_{\mathbb{Z}}(M_1/2M_1)$ . The cases  $M' \subseteq 3M_1$  were already discussed in (3.8). Q.E.D.

Ad  $Q_4$ . The automorphism group of  $Q_4$  is rationally equivalent to a subgroup of order 144 of  $\text{Aut}_{\mathbb{Z}}(Q_3)$ . There is no irreducible subgroup of  $\text{Aut}_{\mathbb{Z}}(Q_4)$  which admits nontrivial 2-centerings. All possible 3-centerings already occur as centerings of

$$\left\langle \left( \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix} \right) \right\rangle.$$

We obtain the following lattice of centerings:



The orbits under the action of  $\text{Aut}_{\mathbb{Z}}(Q_4)$  are  $\{M_1\}, \{M_2\}, \{M_3, M_4\}, \{M_5\}, \{M_6\}, \{3M_1\}, \dots$

Corresponding bases:

$$(i) B(M_1) = I_4, \quad (ii) B(M_2) = \begin{pmatrix} -1 & -1 & -1 & 2 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix},$$

$$(iii) B(M_3) = I_2 \otimes \begin{pmatrix} -1 & 2 \\ 1 & 1 \end{pmatrix}, \quad (iv) B(M_5) = \begin{pmatrix} 1 & 0 & 0 & 3 \\ -1 & 3 & 0 & 0 \\ -1 & 0 & 3 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix},$$

$$(v) B(M_6) = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} \otimes \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}.$$

(3.12) LEMMA. *Only forms induced by  $I_2 \otimes Q_4$  on  $\ker(\mu_1 \oplus \mu_2)$  need to be considered:*

$$(i) \bar{\mu}_1 = \bar{\mu}_2 = (0 \ 0 \ 0 \ 0) \in \mathbf{Z}_3^1 \times^4,$$

$$(ii) \bar{\mu}_1 = \bar{\mu}_2 = (1 \ 1 \ 1 \ 1) \in \mathbf{Z}_3^1 \times^4,$$

$$(iii) \bar{\mu}_1 = \bar{\mu}_2 = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix} \in \mathbf{Z}_3^2 \times^4,$$

$$(iv) \bar{\mu}_1 = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & -1 \end{pmatrix}, \quad \bar{\mu}_2 = \begin{pmatrix} 1 & 1 & 1 & 1 \\ -1 & 1 & 0 & -1 \\ 1 & 1 & 0 & 0 \end{pmatrix} \in \mathbf{Z}_3^3 \times^4.$$

(3.13) PROPOSITION. *From Lemma (3.12) we obtain the forms:*

(i)  $F_9$  (occurred already in (3.9));

$$(ii) F_{11} = \begin{pmatrix} 6 & 0 & 0 & 0 & 0 & 3 & 3 & 0 \\ 0 & 6 & 0 & 3 & 0 & 0 & -3 & 0 \\ 0 & 0 & 6 & 0 & -3 & 0 & 0 & 3 \\ 0 & 3 & 0 & 6 & 0 & 0 & -3 & -3 \\ 0 & 0 & -3 & 0 & 6 & 0 & 3 & 0 \\ 3 & 0 & 0 & 0 & 0 & 6 & 3 & 3 \\ 3 & -3 & 0 & -3 & 3 & 3 & 8 & 4 \\ 0 & 0 & 3 & -3 & 0 & 3 & 4 & 8 \end{pmatrix}, \quad \det F_{11} = 1 \cdot 3^4 \cdot 9^3;$$

$$(iii) F_{12} = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix} \otimes \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix} \otimes \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}, \quad \det F_{12} = 1 \cdot 3^3 \cdot 9^3 27;$$

$$(iv) F_{13} = 9I_8 - J_8, \quad \det F_{13} = 1 \cdot 9^7.$$

*Proof of (3.12).*  $K(M')$  acts faithfully on  $M_1/M'$  except for  $M' \in \{M_1, M_2, M_3, M_4, 2M_1\}$ . In cases (i), (ii) we clearly have  $\bar{\mu}_1 = \bar{\mu}_2$ . In case (iii)

$K(M_3)$  has order 72 and induces a subgroup isomorphic to  $C_2 \times S_3$  in  $\text{Aut}_{\mathbb{Z}}(M_1/M_3)$  which is already the biggest subgroup leaving  $M_2/M_3$  invariant. Hence this subgroup is self-normalizing ( $M_2$  is an  $\text{Aut}_{\mathbb{Z}}(Q_4)$ -centering!), and we obtain  $\bar{\mu}_1 = \bar{\mu}_2$ . In case (iv)  $K(M_5)$  acts faithfully on  $M_1/M_5$  and is equal to  $\text{Aut}_{\mathbb{Z}}(Q_4)$ . Since the index of the induced subgroup in  $\text{Aut}_{\mathbb{Z}}(M_1/M_5)$  in the biggest subgroup leaving  $M_2/M_5$  invariant is 6, there is at most one possibility for  $\bar{\mu}_2$  (compare (3.10)).

A similar argument shows that  $M_6$  can yield at most one form. This form would necessarily be  $9F_5$ , since the automorphism group of the unique form obtained from  $M_5$ , namely  $9I_8 - J_8$ , has a unique centering of index 3. The form provided by this centering is  $9F_5$  (for a similar argument compare the proof of (3.1)).

The cases  $M' \subseteq 3M'$  and  $M' \subseteq 2M'$  are treated as in (3.10). Q.E.D.

Ad  $Q_5$ . The automorphism group of  $Q_5$  is isomorphic to  $C_2 \times S_5$  of order 240. The orders of the irreducible subgroups are all divisible by 5, hence we have only non-trivial 5-centerings. They already occur as centerings of

$$\left\langle \left\langle \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & -1 & -1 & -1 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} \right\rangle \right\rangle.$$

The lattice of 5-centerings is linearly ordered:  $M_1 \supset M_2 \supset M_3 \supset M_4 \supset 5M_1 \supset \dots$ .

Bases for  $M_i$  are:

$$B(M_i) = \left( \left( \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & -1 & -1 & -1 \end{pmatrix} - I_4 \right)^{i-1} \right) \quad (i = 1, 2, 3, 4).$$

(3.13) LEMMA. *We have to consider only forms which are induced by  $I_2 \otimes Q_5$  on  $\ker(\mu_1 \oplus \mu_2)$ :*

- (i)  $\bar{\mu}_1 = \bar{\mu}_2 = (0 \ 0 \ 0 \ 0) \in \mathbb{Z}_5^{1 \times 4}$ ,
- (ii)  $\bar{\mu}_1 = \bar{\mu}_2 = (1 \ 2 \ 3 \ 4) \in \mathbb{Z}_5^{1 \times 4}$ .

(3.14) PROPOSITION. (3.13) provides the forms:

- (i)  $F_{14} = I_2 \otimes (I_4 + J_4)$ ,  $\det F_{14} = 1^6 5^2$ ;

(ii)  $F_{15} = \begin{pmatrix} 4 & -2 & 1 & 1 & -1 & -2 & 0 & 2 \\ -2 & 4 & -1 & 0 & -1 & 1 & 1 & -1 \\ 1 & -1 & 4 & -1 & 1 & -2 & -2 & 0 \\ 1 & 0 & -1 & 4 & 1 & 1 & 2 & 1 \\ -1 & -1 & 1 & 1 & 4 & 1 & -1 & -2 \\ -2 & 1 & -2 & 1 & 1 & 4 & 2 & 0 \\ 0 & 1 & -2 & 2 & -1 & 2 & 4 & 1 \\ 2 & -1 & 0 & 1 & -2 & 0 & 1 & 4 \end{pmatrix}, \det F_{15} = 1^4 5^4.$

*Proof of (3.13).*  $K(M')$  acts faithfully on  $M_1/M'$  except for  $M' \in \{M_1, M_2, 2M_1\}$ . The order of  $K(M_2)$  is 40 and  $K(M_2)$  already induces the full automorphism group of  $M_1/M_2$ . Hence  $\bar{\mu}_1 = \bar{\mu}_2$  holds in case (ii). Because of  $K(M_2) = K(M_3)$  one easily recognizes that there is just one representation  $\Delta$  of  $K(M_3)$  with  $\Delta(K(M_3)) = K(M_3)$ . Hence  $M_3$  need not be considered, since  $K(M_3)$  acts faithfully on  $M_1/M_3$ . Similar arguments work for all  $M'$  contained in  $M_3$ . If  $M'$  is contained in  $2M_1$ , Lemma (2.2) can be applied because the outer automorphism group of  $S_5$  is trivial. Q.E.D.

Ad  $Q_6$ . The automorphism group of  $Q_6$  is rationally equivalent to the automorphism group of  $Q_5$ . Hence, we are only concerned with 5-centerings which already occur as centerings of

$$\left\langle \left( \begin{pmatrix} 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & -1 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -1 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \end{pmatrix} \right) \right\rangle.$$

Again the lattice of 5-centerings is linearly ordered:  $M_1 \supset M_2 \supset M_3 \supset M_4 \supset 5M_1 \supset \dots$ . Bases for  $M_i$  are:

$$B(M_i) = \left( \left( \begin{pmatrix} 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & -1 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -1 \end{pmatrix} - I_4 \right)^{i-1} \right) \quad (i = 1, 2, 3, 4).$$

(3.15) LEMMA. *Only the forms induced by  $I_2 \otimes Q_6$  on  $\ker(\mu_1 \oplus \mu_2)$  need to be considered:*

- (i)  $\bar{\mu}_1 = \bar{\mu}_2 = (0 \ 0 \ 0 \ 0) \in \mathbf{Z}_5^1 \times^4$ ,
- (ii)  $\bar{\mu}_1, \bar{\mu}_2 \in \{(1 \ 1 \ 1 \ 1), (2 \ 2 \ 2 \ 2)\} \subseteq \mathbf{Z}_5^1 \times^4$ .

(3.16) PROPOSITION. (3.15) yields the forms:

- (i)  $F_{16} = I_2 \otimes (5I_4 - J_4)$ ,  $\det F_{16} = 1^2 5^6$ ;

$$(ii) \quad F_{17} = \begin{pmatrix} 8 & 3 & 3 & 3 & -3 & -3 & -2 & -2 \\ 3 & 8 & 3 & 3 & 2 & 2 & -2 & -2 \\ 3 & 3 & 8 & 3 & 2 & 2 & -2 & -2 \\ 3 & 3 & 3 & 8 & 2 & 2 & -2 & 3 \\ -3 & 2 & 2 & 2 & 8 & 3 & 2 & 2 \\ -3 & 2 & 2 & 2 & 3 & 8 & 2 & 2 \\ -2 & -2 & -2 & -2 & 2 & 2 & 8 & 3 \\ -2 & -2 & -2 & 3 & 2 & 2 & 3 & 8 \end{pmatrix}, \quad \det F_{17} = 1 \cdot 5^6 25; 5F_5.$$

*Proof of (3.15).* The proof is analogous to the one of (3.13) the main difference being  $K(M_2) = \text{Aut}_{\mathbf{Z}}(Q_6)$ . Q.E.D.

The automorphism groups of the forms  $F_1, \dots, F_{17}$  obtained in this paragraph are discussed in Part V [7]. It turns out that all these groups are irreducible which is



not completely clear from Section 2. In Part V we also determine the remaining forms of degree 8, i.e. those forms  $F$  for which there is no subgroup  $H$  in  $\text{Aut}_{\mathbf{Z}}(F)$  having a  $\mathbf{Q}$ -reducible subgroup of index 2. We finally obtain 26  $\mathbf{Z}$ -classes of maximal finite irreducible subgroups of  $GL(8, \mathbf{Z})$ .

RWTH Aachen  
Lehrstuhl D für Mathematik  
Templergraben 64  
5100 Aachen, West Germany

Mathematisches Institut der Universität zu Köln  
Weyertal 86–90  
5000 Köln 41, West Germany

1. H. BROWN, R. BÜLOW, J. NEUBÜSER, H. WONDRATSCHEK & H. ZASSENHAUS, *Crystallographic Groups of Four-Dimensional Space*, Wiley, New York, 1978.
2. E. C. DADE, "The maximal finite groups of  $4 \times 4$  integral matrices," *Illinois J. Math.*, v. 9, 1965, pp. 99–122.
3. I. M. ISAACS, *Character Theory of Finite Groups*, Academic Press, New York, 1976.
4. W. PLESKEN, "On absolutely irreducible representations of orders," in *Number Theory and Algebra* (H. Zassenhaus, Editor), Academic Press, New York, 1977.
5. W. PLESKEN, "On reducible and decomposable representations of orders," *J. Reine Angew. Math.*, v. 297, 1978, pp. 188–210.
6. W. PLESKEN & M. POHST, "On maximal finite irreducible subgroups of  $GL(n, \mathbf{Z})$ . I. The five and seven dimensional cases," *Math. Comp.*, v. 31, 1977, pp. 536–551.
7. W. PLESKEN & M. POHST, "On maximal finite irreducible subgroups of  $GL(n, \mathbf{Z})$ . V. The eight dimensional case and a complete description of dimensions less than ten," *Math. Comp.*, v. 34, 1980, pp. 277–301.