

Eigenvalue Problems on Infinite Intervals*

By Peter A. Markowich

Abstract. This paper is concerned with eigenvalue problems for boundary value problems of ordinary differential equations posed on an infinite interval. Problems of that kind occur for example in fluid mechanics when the stability of laminar flows is investigated. Characterizations of eigenvalues and spectral subspaces are given, and the convergence of approximating problems, which are derived by reducing the infinite interval to a finite but large one and by imposing additional boundary conditions at the far end, is proved. Exponential convergence is shown for a large class of problems.

1. Introduction. This paper deals with eigenvalue problems of the form

$$(1.1) \quad y' - t^\alpha A(t)y = \lambda t^\alpha G(t)y, \quad 1 \leq t < \infty, \alpha > -1,$$

$$(1.2) \quad By(1) = 0,$$

$$(1.3) \quad y \in C([1, \infty]): \Leftrightarrow y \in C([1, \infty)) \text{ and } \lim_{t \rightarrow \infty} y(t) \text{ exists,}$$

where the $n \times n$ matrices $A, G \in C([1, \infty))$ and $A(\infty) \neq 0$. A theory for inhomogeneous boundary value problems on infinite intervals has been developed (see Lentini and Keller [11], de Hoog and Weiss [6], [7], Markowich [12], [13], [14]) but not much attention has been paid to eigenvalue problems with a singularity of the second kind. de Hoog and Weiss [5] established a theory for eigenvalue problems in the case that the differential equation has a singularity of the first kind ($\alpha = -1$) and that $G(\infty) = 0$. They showed that the spectrum has no finite limit point and that the spectral subspaces associated with a particular eigenvalue are finite dimensional. They also considered difference schemes for problems which have been transformed to a finite interval, and they derived convergence results for eigenvalues and spectral subspaces using the collective compactness of the difference schemes. For the problems (1.1), (1.2), (1.3) de Hoog and Weiss [6] showed that all eigenvalues λ , for which $A(\infty) + \lambda G(\infty)$ has no eigenvalue on the imaginary axis, are isolated and the spectral subspaces are finite dimensional. Their proofs hinge on the Fredholm property of the differential operator.

The goal of this paper is twofold. First to derive properties of the spectrum and the generalized eigenvectors of (1.1), (1.2), (1.3) and second to consider the approximating eigenvalue problems

$$(1.4) \quad x'_T - t^\alpha A(t)x_T = \lambda_T t^\alpha G(t)x_T, \quad 1 \leq t \leq T, T \geq 1,$$

Received December 19, 1980; revised October 2, 1981.

1980 *Mathematics Subject Classification.* Primary 34B05, 34B25, 34C11, 34E05, 34A45.

Key words and phrases. Boundary value problems of linear equations, spectral theory of boundary value problems, boundedness of solutions, asymptotic expansion, theoretical approximation of solutions.

*Sponsored by the United States Army under Contract No. DAAG29-80-C-0041. This material is based upon work supported by the National Science Foundation under Grant No. MCS-7927062.

$$(1.5) \quad Bx_T(1) = 0,$$

$$(1.6) \quad S(T)x_T(T) = 0.$$

These problems, with a suitably chosen matrix $S(T)$, are ‘regular’ two-point boundary eigenvalue problems which can be solved by any appropriate code, for example by collocation; see de Boor and Swartz [1]. A class of matrices $S(T)$, for which the eigenvalues and spectral subspaces of (1.4), (1.5), (1.6) converge to those of (1.1), (1.2), (1.3), will be defined and the order of convergence, which turns out to be exponential in the most important cases, will be estimated.

This paper is organized as follows. In Section 2 the case where $A(\infty)$ has no eigenvalue on the imaginary axis and where $G(\infty) = 0$ is treated. In Section 3 the assumption $G(\infty) = 0$ is eliminated. In Section 4 no assumption on the eigenvalues of $A(\infty)$ are made, but a certain order of convergence of $G(t)$ to 0 is required. In Section 5 the Orr-Sommerfeld equation, a fluid dynamical problem posed as an eigenvalue problem on an infinite interval, is dealt with, and appropriate approximating problems are devised.

It is of particular interest that the approximation theory in the case $G(\infty) \neq 0$ is treated by using Grigorieff’s [3] ‘discrete’ approximation theory for eigenvalue problems, which allows the approximating operators to be defined on different spaces which—in some sense—converge to the space on which the eigenvalue problem is posed. This approach simplifies the analysis essentially.

2. $G(\infty) = 0$: The ‘Compact’ Case. We assume that $A(\infty) := \lim_{t \rightarrow \infty} A(t)$ has no eigenvalue with real part zero and that $G(\infty) := \lim_{t \rightarrow \infty} G(t) = 0$. We transform $A(\infty)$ to its Jordan canonical form $J(\infty)$

$$(2.1) \quad A(\infty) = FJ(\infty)F^{-1}$$

and assume that $J(\infty)$ has the block structure

$$(2.2) \quad J(\infty) = \text{diag}(J_\infty^+, J_\infty^-),$$

where J_∞^+ contains all Jordan blocks which have eigenvalues with real part larger than zero and J_∞^- contains all Jordan blocks with eigenvalues with negative real part. Let J_∞^+ be a $r_+ \times r_+$ matrix and J_∞^- a $r_- \times r_-$ matrix, and let D_+ and D_- be the projections onto the sum of invariant subspaces associated with the eigenvalues of J_∞^+ and J_∞^- , respectively. We define a solution operator H of the problem

$$(2.3) \quad z' = t^\alpha J(\infty)z + t^\alpha g(t)$$

for all real $\alpha > -1$ as follows:

$$(2.4) \quad (Hg)(t) = \phi(t) \int_\delta^t D_+ \phi^{-1}(s) s^\alpha g(s) ds + \phi(t) \int_\delta^t D_- \phi^{-1}(s) s^\alpha g(s) ds,$$

where $\delta \geq 1$ and

$$(2.5) \quad \phi(t) = \exp\left(\frac{J(\infty)}{\alpha + 1} t^{\alpha+1}\right).$$

This operator has been used by de Hoog and Weiss [6], [7] and they showed that $H: C([\delta, \infty]) \rightarrow C([\delta, \infty])$ and that

$$(2.6) \quad (Hg)(\infty) = -J(\infty)^{-1} g(\infty)$$

holds. As norm in $C([\delta, \infty])$ we take $\|\cdot\|_{[\delta, \infty]}$ which denotes the max-norm on the interval $[\delta, \infty]$. Then $H: C([\delta, \infty]) \rightarrow C([\delta, \infty])$ is bounded uniformly in $\delta \geq 1$, and

$$(2.7) \quad \|(Hg)(t)\| \leq \text{const} \left(\|g\|_{[t/2, \infty]} + \left\| \exp\left(\frac{J_\infty^-}{\alpha + 1} \left(\frac{2^{\alpha+1} - 1}{2^{\alpha+1}}\right) t^{\alpha+1}\right) \right\| \|g\|_{[1, \infty]} \right)$$

holds for $t \geq 2$.

Now we investigate the problem

$$(2.8) \quad y' - t^\alpha A(t)y = t^\alpha G(t)f(t), \quad 1 \leq t < \infty,$$

$$(2.9) \quad By(1) = 0,$$

$$(2.10) \quad y \in C([1, \infty]),$$

where $f \in C([1, \infty])$. Substituting $u = F^{-1}y$, the general solution of the transformed problem (2.8), (2.10) is

$$(2.11)(a) \quad u(t) = \phi(t)G_- \xi + (H(J - J(\infty))u)(t) + (HF^{-1}Gf)(t), \quad \xi \in C^{r_-},$$

where $J(t) = F^{-1}A(t)F$ and the $n \times r_-$ matrix G_- is obtained from D_- by cancelling all columns which have only zero entries.

Obviously, the operator

$$(2.11)(b) \quad H(J - J(\infty)): C([\delta, \infty]) \rightarrow C([\delta, \infty])$$

fulfills $\|H(J - J(\infty))\|_{[\delta, \infty]} < 1$ for δ sufficiently large. Therefore u in (2.11)(a) is defined uniquely on $[\delta, \infty]$ and can be extended uniquely to $[1, \infty]$.

Defining

$$(2.12) \quad \psi_-(t) = ((I - H(J - J(\infty)))^{-1} \phi G_-)(t),$$

$$(2.13) \quad \psi(Gf)(t) = ((I - H(J - J(\infty)))^{-1} HF^{-1}Gf)(t),$$

we write the general solution of (2.8), (2.9), (2.10) as

$$(2.14) \quad y(t) = F\psi_-(t)\xi + F\psi(Gf)(t), \quad t \in [1, \infty], \xi \in C^{r_-}.$$

So (2.8), (2.9), (2.10) is uniquely soluble for every $f \in C([1, \infty])$ if and only if the $r_- \times r_-$ matrix

$$(2.15) \quad BF\psi_-(1) \text{ is nonsingular.}$$

B is assumed to be an $r_- \times n$ matrix. (2.6) and (2.11)(a) imply that $y(\infty) = 0$.

We define the operator V as follows:

$$(2.16) \quad V: \begin{cases} C([1, \infty]) \rightarrow C([1, \infty]), \\ f \rightarrow Vf = y, \end{cases}$$

where y is the solution of (2.8), (2.9), (2.10). V is defined properly if and only if (2.15) holds. This is no restriction because if $\lambda = 0$ is an eigenvalue of (1.1), (1.2), (1.3), we substitute $\lambda = \bar{\lambda} + \gamma$, so that the problem with $A(t)$ replaced by $A(t) + \bar{\lambda}G(t)$ has not $\gamma = 0$ as eigenvalue.

(2.7) and (2.14) imply that V is bounded.

Obviously the eigenvalue problem (1.1), (1.2), (1.3) is equivalent to

$$(2.17) \quad Vf = \mu f,$$

with

$$(2.18) \quad \mu = 1/\lambda.$$

Our goal is to show that V is compact. We need the following

LEMMA 2.1. Assume $\sigma \in C([\delta, \infty])$, $\sigma(t) \rightarrow 0$ as $t \rightarrow \infty$, $\sigma \geq 0$, and $\delta \geq 1$. Then the set A_σ , defined by

$A_\sigma := \{f \in C([1, \infty]) \mid \|f\|_{[1, \infty]} \leq C_1, \|f(t)\| \leq C_2\sigma(t) \text{ for } t \geq \delta, \|f'(t)\| \leq C_3t^\alpha\}$
for $\alpha \in \mathbb{R}$, is conditionally compact in $C([1, \infty])$.

Proof. Given $\varepsilon > 0$, we choose $T = T(\varepsilon) \geq \delta$ so large that $\sigma(t) < \varepsilon/C_2$ for $t \geq T$. Obviously there is a finite collection of intervals I_i , for $i = 1(1)(N(\varepsilon) - 1)$, whose conjunction is $[1, T]$, and there are points t_i in I_i so that

$$\sup_{f \in A_\sigma} \sup_{t \in I_i} \|f(t_i) - f(t)\| < \varepsilon, \quad i = 1(1)(N(\varepsilon) - 1).$$

This is fulfilled if $|I_i| < \varepsilon/(C_3T^\alpha)$ with t_i arbitrary in I_i .

Setting $t_{N(\varepsilon)} = \infty$, Theorem 5, Chapter IV in Dunford and Schwartz [2] is satisfied and the lemma follows.

From (2.11) and (2.7) we conclude that

$$(2.19) \quad \|(Vf)(t)\| \leq \text{const} \left(\|\phi(t)G_- \| + \|J(t) - J(\infty)\|_{[t/2, \infty]} \right. \\ \left. + \|G(t)\|_{[t/2, \infty]} + \left\| \exp \left(\frac{J_\infty^-}{\alpha + 1} \left(\frac{2^{\alpha+1} - 1}{2^{\alpha+1}} \right) t^{\alpha+1} \right) \right\| \right) \|f\|_{[1, \infty]}, \\ t \geq \delta > 2,$$

holds because ξ in (2.11)(a) equals $-(BE\psi_-(1))^{-1}E\psi(f)(1)$. Setting

$$(2.20) \quad \sigma(t) = \|\phi(t)G_- \| + \|J(t) - J(\infty)\|_{[t/2, \infty]} + \|G(t)\|_{[t/2, \infty]} \\ + \left\| \exp \left(\frac{J_\infty^-}{\alpha + 1} \left(\frac{2^{\alpha+1} - 1}{2^{\alpha+1}} \right) t^{\alpha+1} \right) \right\|,$$

we notice that $\sigma(t) \downarrow 0$ as $t \rightarrow \infty$ and therefore

$$(2.21) \quad \{Vf \mid f \in C([1, \infty]), \|f\|_{[1, \infty]} \leq 1\} \in A_\sigma$$

for some constants C_1, C_2, C_3 . So V is a compact operator on $C([1, \infty])$.

Therefore, the spectrum $\sigma(V)$ consists of a countable set of eigenvalues $\mu \neq 0$ of finite algebraic multiplicities, and $\mu = 0 \in \sigma(V)$ is the only possible accumulation point of the μ 's. The spectrum of compact operators is described in Dunford and Schwartz [2, Chapter VII, Theorem 5].

Let $\mu (\neq 0)$ be a fixed eigenvalue of V . We want to investigate the spectral subspace associated with μ . The spectral projection is given by

$$(2.22) \quad E = E(\mu) = \frac{1}{2\pi i} \int_\Gamma (z - V)^{-1} dz: C([1, \infty]) \rightarrow C([1, \infty]),$$

where Γ is a circle centered at μ which contains no other eigenvalue of V . Moreover,

$$(2.23) \quad \text{rank}(E(\mu)) = m,$$

where m is the algebraic multiplicity of μ . Let

$$(2.24) \quad \text{Range}(E) = \text{span}\{\varphi_1, \dots, \varphi_m\} = N((\mu - V)^\beta)$$

hold, where the φ_i are generalized eigenfunctions of V . N denotes the null space and β the ascent of $\mu - V$.

As the range of E is invariant under V , we get

$$(2.25) \quad \varphi'_i - t^\alpha A(t)\varphi_i = \sum_{j=1}^m a_{i,j} t^\alpha G(t)\varphi_j, \quad B\varphi_i(1) = 0, \quad \varphi_i \in C([1, \infty)).$$

The $m \times m$ matrix (a_{ij}) can be assumed to be in Jordan canonical form with the only eigenvalue $\lambda = 1/\mu$. This can always be achieved by a basis transformation. So every element φ_k is contained in a finite chain $\varphi_{r_1}, \dots, \varphi_{r_i}$ which fulfills

$$(2.26) \quad \varphi'_{r_1} - t^\alpha(A(t) + \lambda G(t))\varphi_{r_1} = 0, \quad B\varphi_{r_1}(1) = 0, \quad \varphi_{r_1} \in C([1, \infty)),$$

$$(2.27) \quad \varphi'_{r_2} - t^\alpha(A(t) + \lambda G(t))\varphi_{r_2} = t^\alpha G(t)\varphi_{r_1}, \quad B\varphi_{r_2}(1) = 0, \quad \varphi_{r_2} \in C([1, \infty)),$$

$$(2.28) \quad \begin{matrix} \vdots & & \vdots & & \vdots \\ \varphi'_{r_i} - t^\alpha(A(t) + \lambda G(t))\varphi_{r_i} = t^\alpha G(t)\varphi_{r_{i-1}}, & B\varphi_{r_i}(1) = 0, & \varphi_{r_i} \in C([1, \infty)). \end{matrix}$$

Using the properties of the spectrum $\sigma(V)$ and applying the estimates defined in Markowich [13, Sections 2, 3] to (2.26), (2.27), (2.28), we obtain

THEOREM 2.1. *The spectrum of (1.1), (1.2), (1.3) consists of countably many eigenvalues λ which have no finite accumulation point, and every (generalized) eigenfunction φ satisfies*

$$(2.29) \quad \|\varphi(t)\| \leq \text{const} \exp\left(\frac{\nu_- + \varepsilon}{\alpha + 1} t^{\alpha+1}\right), \quad t \geq \delta,$$

where ν_- is the largest (in modulus) negative real part of the eigenvalues of $A(\infty)$, and $\varepsilon = \varepsilon(\delta) > 0$ fulfills $\nu_- + \varepsilon < 0$ and $\varepsilon(\delta) \rightarrow 0$ as $\delta \rightarrow \infty$.

Now we want to investigate the convergence of the eigenvalue and generalized eigenvectors of the approximating problems (1.4), (1.5), (1.6). As a notion of the distance of closed subspaces we use the ‘gap’ (see Osborn [16]) which is defined as follows

$$(2.30) \quad \text{gap}(M, N) = \max\left(\sup_{\substack{x \in M \\ \|x\|=1}} \text{dist}(x, N), \sup_{\substack{y \in N \\ \|y\|=1}} \text{dist}(M, y)\right),$$

where M, N are closed subspaces of a Banach space $(X, \|\cdot\|)$ and dist is defined as

$$(2.31) \quad \text{dist}(x, N) = \inf_{y \in N} \|x - y\|.$$

We define the operators V_T for T sufficiently large by

$$(2.32) \quad V_T: \begin{cases} C([1, \infty)) \rightarrow C([1, \infty)), \\ f \rightarrow V_T f = x_T, \end{cases}$$

where x_T satisfies

$$(2.33) \quad x'_T - t^\alpha A(t)x_T = t^\alpha G(t)f(t), \quad 1 \leq t \leq T,$$

$$(2.34) \quad Bx_T(1) = 0,$$

$$(2.35) \quad S(T)x_T(T) = 0,$$

and

$$(2.36) \quad x_T(t) = x_T(T) \quad \text{for } t \geq T.$$

This definition makes sense if and only if (2.33), (2.34), (2.35) is soluble for every $f \in C([1, \infty])$ and T sufficiently large. De Hoog and Weiss [7] have shown that this is the case if (2.15) holds and the $r_+ \times n$ matrix $S(T)$ satisfies

$$(2.37) \quad \|S(T)\| \leq \text{const} \quad \text{as } T \rightarrow \infty,$$

$$(2.38) \quad \|(S(T)FG_+)^{-1}\| \leq \text{const} \quad \text{as } T \rightarrow \infty,$$

where the $n \times r_+$ matrix G_+ is obtained by cancelling all columns of D_+ which have only zero entries. Moreover the stability estimate

$$(2.39) \quad \|x_T\|_{[1,T]} \leq \text{const}(\|f\|_{[1,T]} + \|\gamma(T)\|)$$

holds for problems of the form (2.33), (2.34) and

$$(2.40) \quad S(T)x_T(T) = \gamma(T)$$

instead of the homogeneous boundary condition (2.35). de Hoog and Weiss [7] have also shown that (2.38) is necessary if (2.37) holds, and they constructed matrices $S(T)$ fulfilling (2.37), (2.38) explicitly. Obviously, the estimate (2.39) with $\gamma(T) \equiv 0$ and the definition of V_T imply

$$(2.41) \quad \|V_T\|_{[1,\infty]} \leq \text{const}.$$

Every operator V_T is compact because

$$(2.42) \quad \|(V_T f)\|_{[1,T]} \leq \text{const } T^\alpha \|f\|_{[1,T]}$$

holds and $V_T f$ is constant on $[T, \infty]$. By adding the identity $S(T)y(T) = S(T)y(T)$ to (2.8), (2.9), (2.10) and by subtracting from (2.33), (2.34), (2.35), we get the problem

$$(2.43) \quad (x_T - y)' - t^\alpha A(t)(x_T - y) = 0, \quad 1 \leq t \leq T,$$

$$(2.44) \quad B(x_T - y)(1) = 0,$$

$$(2.45) \quad S(T)(x_T - y)(T) = -S(T)y(T).$$

Applying estimate (2.39) implies

$$(2.46) \quad \|V_T f - Vf\|_{[1,T]} \leq \text{const} \|y(T)\| \leq \text{const } \sigma(T) \|f\|_{[1,\infty]},$$

where σ is defined in (2.20). Also, we get

$$(2.47) \quad \|V_T f - Vf\|_{[1,\infty]} \leq 2\|V_T f - Vf\|_{[1,T]} + 2\|y\|_{[T,\infty]}$$

because of (2.36). Therefore V_T converges to V (in the norm) and

$$(2.48) \quad \|V_T - V\|_{[1,\infty]} \leq \text{const } \sigma(T)$$

holds.

It should be noticed that $G(t) \rightarrow 0$ as $t \rightarrow \infty$ is absolutely crucial for the norm convergence.

The eigenvalue problem (1.4), (1.5), (1.6) is equivalent to

$$(2.49) \quad V_T f = \mu_T f$$

with

$$(2.50) \quad \mu_T = 1/\lambda_T.$$

The generalized eigenfunctions of (1.4), (1.5), (1.6) are obtained by restricting the generalized eigenfunction of (2.49) to $[1, T]$.

Because of the compactness of V_T , there is a countable set of eigenvalues $\mu_T \neq 0$ possibly accumulating at 0. The compactness and norm convergence (2.48) allows us to apply Osborn's [16] result. From this and Theorem 2.1 we derive

THEOREM 2.2. *Let λ be an eigenvalue of (1.1), (1.2), (1.3) with algebraic multiplicity m . Then, for T sufficiently large, there are exactly m eigenvalues $\lambda_T^1, \dots, \lambda_T^m$, and these satisfy*

$$(2.51) \quad |\lambda - \lambda_T^i| \leq \text{const} \exp\left(\frac{\nu_- + \varepsilon}{(\alpha + 1)\beta} T^{\alpha+1}\right), \quad i = 1(1)m,$$

$$(2.52) \quad \left| \lambda - \frac{1}{\hat{\mu}_T} \right| \leq \text{const} \exp\left(\frac{\nu_- + \varepsilon}{(\alpha + 1)\beta} T^{\alpha+1}\right),$$

where $\hat{\mu}_T = (1/m)\sum_{i=1}^m 1/\lambda_T^i$ and β is defined in (2.24). Moreover, the spectral projections

$$(2.53) \quad E_T = \frac{1}{2\pi i} \int_{\Gamma} (z - V_T)^{-1} dz$$

satisfy $\text{rank}(E_T) = m$ and

$$(2.54) \quad \text{gap}(\text{Range}(E), \text{Range}(E_T)) \leq \text{const} \exp\left(\frac{\nu_- + \varepsilon}{\alpha + 1} T^{\alpha+1}\right)$$

holds.

The constants in (2.51), (2.52), and (2.54) are independent of T but may very well depend on λ . Sharper estimates will be proven in Section 3.

A possible choice for $S(T)$ is

$$(2.55) \quad S(T) \equiv S \equiv (G_+)^T F^{-1},$$

where the superscript T denotes transposition. The condition (2.38) is satisfied because

$$(2.56) \quad SFG_+ = I_{r_+}$$

holds for the choice (2.55), which has been used by de Hoog and Weiss [7] for the solution of inhomogeneous boundary value problems on infinite intervals.

3. The Case $G(\infty) \neq 0$. Again we consider the problem (1.1), (1.2), (1.3), but we drop the restriction $G(\infty) = 0$. We again assume that B is an $r_- \times n$ matrix.

The following assumption will be needed.

(I) The problem

$$(3.1) \quad y'_h - t^\alpha A(t)y_h = 0, \quad 1 \leq t < \infty,$$

$$(3.2) \quad By_h(1) = 0,$$

$$(3.3) \quad y_h \in C([1, \infty])$$

has the unique solution $y_h = 0$. This guarantees that the $r_- \times r_-$ matrix

$$(3.4) \quad BF\psi_-(1) \text{ is nonsingular,}$$

and therefore the inhomogeneous problem

$$(3.5) \quad y' - t^\alpha A(t)y = t^\alpha G(t)f(t), \quad 1 \leq t < \infty,$$

with the boundary conditions (3.2) and (3.3) has a unique solution for every $f \in C([1, \infty])$. Moreover, we restrict the eigenparameter λ to an open and connected set $\Omega \subset \mathbb{C}$ with $0 \in \Omega$, so that the matrix $A(\infty) + \lambda G(\infty)$ for $\lambda \in \Omega$ has no eigenvalue $\nu(\lambda)$ on the imaginary axis, and therefore the matrices G_+ and G_- are constant for $\lambda \in \Omega$.

De Hoog and Weiss [6] proved that all eigenvalues λ of (1.1), (1.2), (1.3) which fulfill $\lambda \in \Omega$ are isolated and that the associated spectral subspaces are finite dimensional. Each (generalized) eigenfunction y associated with an eigenvalue $\lambda \in \Omega$ satisfied $y(\infty) = 0$. The spectrum of (1.1), (1.2), (1.3) has no finite limit point in Ω .

Of course this settles the case $G(\infty) = 0$ completely because then $\Omega = \mathbb{C}$ holds, but the compactness arguments in Section 2 were included because they will be used in Section 4 where imaginary eigenvalues of $A(\infty)$ will be admitted.

We define the operator V differently than in Section 2:

$$(3.6) \quad V: \begin{cases} C_0([1, \infty]) \rightarrow C_0([1, \infty]), \\ f \rightarrow Vf = y, \end{cases}$$

where $C_0([1, \infty])$ is the Banach space of all functions $f \in C([1, \infty])$ which satisfy $f(\infty) = 0$ and y is the solution of the problem (2.8), (2.9), (2.10). Assumption (I) makes V well-defined on $C_0([1, \infty])$, and (2.6), (2.11) guarantee that $y(\infty) = 0$ if $f(\infty) = 0$.

The eigenvalue problem (1.1), (1.2), (1.3) is equivalent to

$$(3.7) \quad Vf = \mu f, \quad f \in C_0([1, \infty]),$$

with

$$(3.8) \quad \mu = 1/\lambda, \quad \lambda \in \Omega,$$

because all generalized eigenfunctions associated with $\lambda \in \Omega$ are in $C_0([1, \infty])$ and because $\lambda = 0$ is no eigenvalue.

Now let us consider a fixed eigenvalue $\mu = 1/\lambda, \lambda \in \Omega$, with algebraic multiplicity m and ascent β . The spectral projection is again given by

$$(3.9) \quad E = E(\mu) = \frac{1}{2\pi i} \int_{\Gamma} (z - V)^{-1} dz: C_0([1, \infty]) \rightarrow C_0([1, \infty]),$$

where the circle Γ centered at μ contains no other eigenvalue of (3.7) and the image of Γ under the mapping $\lambda = 1/\mu$ (denoted by $(1/\mu)(\Gamma)$) is in Ω . E satisfies (2.24), (2.25).

We want to approximate the generalized eigenpair $(\lambda, \text{Range}(E(\mu)))$ by a sequence of nearby eigenpairs of (1.4), (1.5), (1.6).

Therefore we define the operators V_T for T sufficiently large

$$(3.10) \quad V_T: \begin{cases} C([1, T]) \rightarrow C([1, T]), \\ f_T \rightarrow V_T f_T = x_T, \end{cases}$$

where x_T solves (2.33), (2.34), (2.35), $S(T)$ is independent of λ and satisfies (2.37), (2.38). Then the V_T 's are defined properly and satisfy

$$(3.11) \quad \|V_T f_T\|_{[1, T]} \leq \text{const} \|f_T\|_{[1, T]},$$

$$(3.12) \quad \|(V_T f_T)'\|_{[1, T]} \leq \text{const } T^\alpha \|f_T\|_{[1, T]}.$$

Therefore each V_T is compact and has a countable set of eigenvalues $\mu_T \neq 0$ which may only accumulate at 0. The associated spectral subspaces are finite dimensional.

It is therefore clear that the finite interval problems (1.4), (1.5), (1.6) cannot be used to approximate continuous parts of the spectrum of (1.1), (1.2), (1.3) which may very well exist outside of Ω .

We define the restriction operator

$$(3.13) \quad r_T: \begin{cases} C_0([1, \infty]) \rightarrow C([1, T]), \\ f \rightarrow r_T f = f|_{[1, T]}. \end{cases}$$

Then, for every sequence $T_n \rightarrow \infty$, the sequence of spaces $C([1, T_n])$ forms a discrete approximation $A(C_0([1, \infty]), \prod_n C([1, T_n]), r_{T_n})$ for the space $C_0([1, \infty])$ in the sense of Stummel [17].

A sequence $f_{T_n} \in C([1, T])$ is said to converge to an element $f \in C_0([1, \infty])$, denoted by $f_{T_n} \rightarrow f$, if

$$(3.14) \quad \|f_{T_n} - r_{T_n} f\|_{[1, T_n]} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

A sequence of bounded operators A_{T_n} on $C([1, T_n])$ is said to converge to a bounded operator A on $C_0([1, \infty])$, again denoted by $A_{T_n} \rightarrow A$, if for every $f \in C_0([1, \infty])$ and for every sequence f_{T_n} ,

$$(3.15) \quad f_{T_n} \rightarrow f \quad \text{implies} \quad A_{T_n} f_{T_n} \rightarrow Af.$$

We will drop the subscript n mostly.

Taking a fixed $z \neq 0$ in the resolvent set of V and in $(1/\lambda)(\Omega)$, we investigate $(z - V_T)^{-1}$. Setting $u_T = (z - V_T)^{-1} f_T$ for an arbitrary $f_T \in C([1, T])$, we easily find that

$$(3.16) \quad u_T = \frac{1}{z}(y_T + f_T),$$

where y_T solves

$$(3.17) \quad y'_T - t^\alpha \left(A(t) + \frac{1}{z} G(t) \right) y_T = \frac{1}{z} t^\alpha G(t) f_T,$$

$$(3.18) \quad B y_T(1) = 0,$$

$$(3.19) \quad S(T) y_T(T) = 0.$$

Defining $F(1/z)$ as the matrix which transforms $A(\infty) + G(\infty)/z$ to its Jordan canonical form (which is assumed to be partitioned as in (2.2) for $1/z \in \Omega$), we derive from de Hoog and Weiss [7] that (3.17), (3.18), (3.19) is uniquely soluble for T sufficiently large if

$$(3.20) \quad \left\| \left(S(T) F \left(\frac{1}{z} \right) G_+ \right)^{-1} \right\| \leq \text{const} \quad \text{as } T \rightarrow \infty$$

and the estimate

$$(3.21) \quad \|(z - V_T)^{-1}\|_{[1, T]} \leq \text{const}(z)$$

follows if (3.20) holds. This bound is uniform in $z \in K_1$, where K_1 is compact, $0 \notin K_1$ and $(1/\mu)(K_1) \subset \Omega$ (see Kreiss [10]).

This analysis also shows that

$$(3.22) \quad (z - V_T)^{-1} \rightarrow (z - V)^{-1}$$

uniformly for $z \in K_1$. Therefore (3.20) guarantees that

$$(3.23) \quad \inf_{\mu} |\mu - \mu_T| \rightarrow 0 \quad \text{for } t \rightarrow \infty,$$

where $\mu \in (1/\lambda)(\Omega)$ are the eigenvalues of V and μ_T are the eigenvalues of V_T . Moreover the spectral projections satisfy

$$(3.24) \quad E_T(\mu) = \frac{1}{2\pi i} \int_{\Gamma} (z - V_T)^{-1} dz \rightarrow E(\mu),$$

$$(3.25) \quad \lim_{T \rightarrow \infty} \text{rank}(E_T(\mu)) \geq \text{rank}(E(\mu)).$$

The sets $\text{Range}(E_T(\mu))$ form a discrete approximation

$$A\left(\text{Range}(E(\mu)), \prod_T \text{Range}(E_T(\mu)), r_T\right)$$

for $\text{Range}(E(\mu))$; see Grigorieff [3].

In order to make sure that $\text{rank}(E_T(\mu)) = \text{rank}(E(\mu))$ for T sufficiently large, it is sufficient to show that the sequence $E_T(\mu)$ is discretely compact (see Stummel [18]) because $E(\mu)$ has finite rank.

We recall that the sequence of bounded operators A_{T_n} in $C([1, T_n])$ is discretely compact if for every bounded sequence $f_{T_n} \in C([1, T_n])$, there is a subsequence $f_{T_{n_k}}$ so that $A_{T_{n_k}} f_{T_{n_k}}$ is convergent to an element in $C_0([1, \infty])$.

We write

$$(3.26) \quad E_T r_T = r_T E + (E_T r_T - r_T E): C_0([1, \infty]) \rightarrow C([1, T]).$$

Obviously,

$$(3.27) \quad E_T r_T - r_T E = \frac{1}{2\pi i} \int_{\Gamma} (z - V_T)^{-1} (V_T r_T - r_T V) (z - V)^{-1} dz$$

holds.

For an arbitrary $f \in C_0([1, \infty])$, the function $e_T = (V_T r_T - r_T V)f \in C([1, T])$ is the solution of the problem

$$(3.28) \quad e'_T - t^\alpha A(t) e_T = 0, \quad 1 \leq t \leq T,$$

$$(3.29) \quad B e_T(1) = 0,$$

$$(3.30) \quad S(T) e_T(T) = -S(T)(Vf)(T).$$

Proceeding similarly to de Hoog and Weiss [7] we can express e_T explicitly.

We substitute $F\tilde{e}_T = e_T$, where F is as in (2.1), and get the problem

$$(3.31) \quad \tilde{e}'_T = t^\alpha J(\infty)\tilde{e}_T + t^\alpha (J(t) - J(\infty))\tilde{e}_T.$$

$J(t), J(\infty)$ are as in (2.11), (2.2). Now we write

$$(3.32) \quad \tilde{e}_T = \tilde{e}_T^+ \xi_1^T + \tilde{e}_T^- \xi_2^T, \quad \xi_1^T \in C^{r^+}, \xi_2^T \in C^{r^-},$$

where $\tilde{e}_T^+, \tilde{e}_T^-$ satisfy

$$(3.33) \quad \tilde{e}_T^+(t) = \begin{bmatrix} \exp\left(\frac{J_\infty^+}{\alpha + 1}(t^{\alpha+1} - T^{\alpha+1})\right) \\ 0 \end{bmatrix} + (H_T(J - J(\infty))\tilde{e}_T^+)(t),$$

$$(3.34) \quad \tilde{e}_T^-(t) = \begin{bmatrix} 0 \\ \exp\left(\frac{J_\infty^-}{\alpha + 1}t^{\alpha+1}\right) \end{bmatrix} + (H_T(J - J(\infty))\tilde{e}_T^-)(t).$$

H_T is a suitable solution operator of the problem

$$(3.35) \quad z' = t^\alpha J(t)z + t^\alpha g(t), \quad 1 \leq t \leq T, \quad g \in C([1, T]).$$

We choose

$$(3.36) \quad (H_T g)(t) = (H\hat{g})(t), \quad 1 \leq t \leq T,$$

with H defined in (2.4) where \hat{g} has been set to

$$(3.37) \quad \hat{g}(t) = \begin{cases} g(t), & 1 \leq t \leq T, \\ g(T), & T \leq t \leq \infty. \end{cases}$$

Because H is bounded on $[\delta, \infty]$ independently of δ , we get

$$(3.38) \quad \|H_T(J - J(\infty))\|_{[\delta, T]} \leq \text{const}\|J - J(\infty)\|_{[\delta, T]} \leq \frac{1}{2}$$

for δ, T sufficiently large. The operator

$$(3.39) \quad I - H_T(J - J(\infty)): C([\delta, T]) \rightarrow C([\delta, T])$$

is invertible and $\tilde{e}_T^+, \tilde{e}_T^- \in C([\delta, T])$ are uniquely defined and can be continued to $[1, T]$.

Inserting (3.32) into the boundary conditions (3.29), (3.30) gives

$$(3.40) \quad \begin{bmatrix} BF\tilde{e}_T^+(1) & BF\tilde{e}_T^-(1) \\ S(T)F\tilde{e}_T^+(T) & S(T)F\tilde{e}_T^-(T) \end{bmatrix} \begin{bmatrix} \xi_1^T \\ \xi_2^T \end{bmatrix} = \begin{pmatrix} 0 \\ -S(T)(Vf)(T) \end{pmatrix}.$$

De Hoog and Weiss [6] showed that

$$(3.41) \quad (a) \quad \lim_{T \rightarrow \infty} \tilde{e}_T^+(T) = G_+, \quad (b) \quad \|\tilde{e}_T^- - r_T\psi_-\|_{[1, T]} \rightarrow 0 \quad \text{as } T \rightarrow \infty$$

hold, where ψ_- is defined in (2.12).

A block system of the form

$$(3.42) \quad \begin{bmatrix} \bar{A} & \bar{B} \\ \bar{C} & \bar{D} \end{bmatrix} \begin{pmatrix} \xi_1 \\ \xi_2 \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix},$$

where \bar{B}, \bar{C} are quadratic matrices, is uniquely soluble if and only if $\bar{B}, (\bar{C} - \bar{D}\bar{B}^{-1}\bar{A})$ are invertible and the solution is

$$(3.43) \quad \begin{pmatrix} \xi_1 \\ \xi_2 \end{pmatrix} = \begin{bmatrix} -(\bar{C} - \bar{D}\bar{B}^{-1}\bar{A})^{-1}\bar{D}\bar{B}^{-1} & (\bar{C} - \bar{D}\bar{B}^{-1}\bar{A})^{-1} \\ \bar{B}^{-1} + \bar{B}^{-1}\bar{A}(\bar{C} - \bar{D}\bar{B}^{-1}\bar{A})^{-1}\bar{D}\bar{B}^{-1} & -\bar{B}^{-1}\bar{A}(\bar{C} - \bar{D}\bar{B}^{-1}\bar{A})^{-1} \end{bmatrix} \begin{pmatrix} a \\ b \end{pmatrix}.$$

The off-diagonal matrices in (3.40) are invertible, their inverses are bounded as $T \rightarrow \infty$, the matrix in the (2, 2) position converges to 0 as $T \rightarrow \infty$, and the matrix in (1, 1) position is bounded, and therefore the system is invertible for T sufficiently

large. Moreover,

$$(3.44) \quad \lim_{T \rightarrow \infty} \tilde{e}_T^+(1) = 0$$

holds because we get from the series expansion of (3.33):

$$(3.44)(a) \quad \left\| \tilde{e}_T^+ - \begin{bmatrix} \exp\left(\frac{J_\infty^+}{\alpha + 1}(\omega - T^{\alpha+1})\right) \\ 0 \end{bmatrix} \right\|_{[\delta, T]} \leq C \left\| (J - J(\infty)) \begin{bmatrix} \exp\left(\frac{J_\infty^+}{\alpha + 1}(\omega - T^{\alpha+1})\right) \\ 0 \end{bmatrix} \right\|_{[\delta, T]},$$

where $\omega(t) = t^{\alpha+1}$ has been set. The right-hand side of this inequality can be estimated by

$$(3.44)(b) \quad \bar{\sigma}(\kappa, T) = C \max_{i=1(1)k} \max_{t \in [\delta, T]} \left(\|J(t) - J(\infty)\| \exp\left(\frac{\kappa}{\alpha + 1}(t^{\alpha+1} - T^{\alpha+1})\right) \times (T^{\alpha+1} - t^{\alpha+1})^i \right),$$

where κ is the smallest (positive) real part of the eigenvalues of J_∞^+ and k is the dimension of the largest Jordan block with eigenvalue with real part κ . Obviously $\bar{\sigma}(\kappa, T) \rightarrow 0$ as $T \rightarrow \infty$, and (3.44) follows by continuation to $[1, T]$.

We get from (3.40), (3.41), (3.43), (3.44) that

$$(3.45)(a) \quad \xi_1^T = - \left((S(T)FG_+)^{-1} + o(T) \right) S(T)(Vf)(T),$$

$$(3.45)(b) \quad \xi_2^T = o(T)S(T)(Vf)(T)$$

holds. For $g \in C_0([1, \infty])$ we therefore obtain

$$(3.45)(c) \quad (V_T r_T - r_T V)(z - V)^{-1} g = \left(-F\tilde{e}_T^+ (S(T)FG_+)^{-1} + F\tilde{e}_T^- o(T) + o(T) \right) S(T)(V(z - V)^{-1} g)(T).$$

Obviously $h = V(z - V)^{-1} g$ is the solution to the problem

$$(3.46) \quad h' - t^\alpha \left(A(t) + \frac{1}{z} G(t) \right) h = \frac{1}{z} t^\alpha G(t) g(t),$$

$$(3.47) \quad Bh(1) = 0,$$

$$(3.48) \quad h \in C_0([1, \infty]).$$

We define

$$(3.49) \quad A(t, \lambda) = A(t) + \lambda G(t) \quad \text{for } t \in [1, \infty],$$

and the family of operators $\hat{H}(\lambda): C([\delta, \infty]) \rightarrow C([\delta, \infty])$

$$(3.50) \quad \begin{aligned} (\hat{H}(\lambda)g)(t) &= \hat{\phi}(t, \lambda) \int_\infty^t P_+(\lambda) \hat{\phi}^{-1}(s, \lambda) s^\alpha g(s) ds \\ &+ \hat{\phi}(t, \lambda) \int_\delta^t P_-(\lambda) \hat{\phi}^{-1}(s, \lambda) s^\alpha g(s) ds, \end{aligned}$$

for $\delta > 1$, where

$$(3.51) \quad \hat{\phi}(t, \lambda) = \exp\left(A(\infty, \lambda) \frac{t^{\alpha+1}}{\alpha + 1} \right),$$

$$(3.52) \quad (a) \quad P_+(\lambda) = F(\lambda)D_+F^{-1}(\lambda), \quad (b) \quad P_-(\lambda) = F(\lambda)D_-F^{-1}(\lambda)$$

hold. Obviously $\hat{H}(0) = FHF^{-1}$ for H as in (2.4). The projections $P_+(\lambda), P_-(\lambda)$ are holomorphic in Ω ; see Kato [8].

Using the techniques of de Hoog and Weiss [6] and Markowich [13], we conclude that

$$(3.53) \quad \|\hat{H}(\lambda)\|_{[\delta, \infty]} \leq C(\lambda),$$

where $C(\lambda)$ is independent of δ and bounded on a compact set $K \subset \Omega$. Moreover it is an easy exercise to show that $(H(\lambda)g(\cdot, \lambda))(t)$ is holomorphic in \mathring{K} for all $t \in [\delta, \infty]$ if $g(t, \lambda)$ is continuous for $t \in [1, \infty]$ and holomorphic (for all t) in $\lambda \in \mathring{K}$.

Proceeding as in Section 2, we rewrite (3.46)

$$(3.54) \quad h' = t^\alpha A\left(\infty, \frac{1}{z}\right)h + t^\alpha \underbrace{\left(A\left(t, \frac{1}{z}\right) - A\left(\infty, \frac{1}{z}\right) \right)}_{B\left(t, \frac{1}{z}\right)} h + \frac{1}{z} t^\alpha G(t)g(t)$$

and get

$$(3.55) \quad h(t) = \hat{\phi}\left(t, \frac{1}{z}\right)W_-\left(\frac{1}{z}\right)\zeta + \left(\hat{H}\left(\frac{1}{z}\right)B\left(\cdot, \frac{1}{z}\right)h\right)(t) + \frac{1}{z}\left(\hat{H}\left(\frac{1}{z}\right)Gg\right)(t),$$

where the columns of the holomorphic $n \times r_-$ matrix $W_-(1/z)$ span the range of $P_-(1/z)$ and $\zeta \in C^{r_-}$. Because of (3.53) there is a fixed δ so that

$$(3.56) \quad \left\| \hat{H}\left(\frac{1}{z}\right)B\left(\cdot, \frac{1}{z}\right) \right\|_{[\delta, \infty]} \leq \frac{1}{2} \quad \text{for all } z \in \bar{S}_\mu,$$

where \bar{S}_μ is the closed disk contoured by Γ .

Setting

$$(3.57)(a) \quad \hat{\psi}_-(\cdot, \lambda) = (I - \hat{H}(\lambda)B(\cdot, \lambda))^{-1}\hat{\phi}(\cdot, \lambda)W_-(\lambda),$$

$$(3.57)(b) \quad \hat{\psi}(Gf, \lambda) = (I - \hat{H}(\lambda)B(\cdot, \lambda))^{-1}\hat{H}(\lambda)Gg,$$

the general solution of (3.46), (3.48) on $[\delta, \infty]$ is

$$(3.58) \quad h(t) = \hat{\psi}_-\left(t, \frac{1}{z}\right)\zeta + \frac{1}{z}\hat{\psi}\left(Gg, \frac{1}{z}\right)(t).$$

$\hat{\psi}_-, \hat{\psi}(Gg, \lambda)$ can be uniquely extended to $[1, \infty]$.

Proceeding as in (2.19) using (3.53), we get

$$(3.59)(a) \quad \left\| \hat{\psi}_-\left(T, \frac{1}{z}\right) \right\| = o(T) \quad \text{uniformly for } z \in \Gamma,$$

and

$$(3.59)(b) \quad \|\zeta\| = O(\|g\|_{[1, \infty]}) \quad \text{uniformly for } z \in \Gamma,$$

can be concluded as in Section 2 using (3.56).

(3.56) and the uniform convergence of the series expansion of (3.57) assure the analyticity of $\hat{\psi}(Gg, 1/z)(T)$ for $z \in S_\mu$ and continuity in \bar{S}_μ . Therefore

$$(3.60) \quad \begin{aligned} (V_T r_T - r_T V)(z - V)^{-1} g \\ = \left(-F\bar{e}_T^+ (S(T)FG_+)^{-1} + F\bar{e}_T^- o(T) + o(T) \right) \\ \times \left(\frac{1}{z} S(T) \hat{\psi} \left(Gg, \frac{1}{z} \right) (T) + o(T) \right). \end{aligned}$$

$u_T = (z - V_T)^{-1} F\bar{e}_T^+$ is the solution to the problem

$$(3.61)(a) \quad u_T' - t^\alpha A \left(t, \frac{1}{z} \right) u_T = 0,$$

$$(3.61)(b) \quad Bu_T(1) = \frac{1}{z} BF\bar{e}_T^+(1),$$

$$(3.61)(c) \quad S(T)u_T(T) = \frac{1}{z} S(T)F\bar{e}_T^+(T).$$

We set similarly to (3.32)

$$(3.62)(a) \quad u_T(t) = e_T^+ \left(t, \frac{1}{z} \right) \xi_1^T + e_T^- \left(t, \frac{1}{z} \right) \xi_2^T,$$

where

$$(3.62)(b) \quad \begin{aligned} e_T^+(t, \lambda) &= \hat{\phi}(t, \lambda) \hat{\phi}^{-1}(T, \lambda) W_+(\lambda) \\ &+ (\hat{H}(\lambda, T) B(\cdot, \lambda) e_T^+(\cdot, \lambda))(t), \end{aligned}$$

$$(3.62)(c) \quad e_T^-(t, \lambda) = \hat{\phi}(t, \lambda) W_-(\lambda) + (\hat{H}(\lambda, T) B(\cdot, \lambda) e_T^-(\cdot, \lambda))(t)$$

hold. The columns of the holomorphic matrix $W_+(\lambda)$ span the range of $P_+(\lambda)$ and $\hat{H}(\lambda, T): C([\delta, T]) \rightarrow C([\delta, T])$ so that

$$(3.62)(d) \quad \hat{H}(\lambda, T)g := \hat{H}(\lambda)\hat{g}, \quad \hat{g}(t) = \begin{cases} g(t), & \delta \leq t \leq T, \\ g(T), & t \geq T, \end{cases}$$

holds. Because of (3.53) the equations (3.62)(b), (c) are uniquely soluble for all $z \in \bar{S}_\mu$ if δ is sufficiently large, and the analyticity of $e_T^\pm(t, 1/z)$ for all $t \in [1, T]$ follows by the above argument and by continuation from $[\delta, T]$ to $[1, T]$.

Moreover, we derive as in (3.41) from de Hoog and Weiss [6]

$$(3.63)(a) \quad \lim_{T \rightarrow \infty} e_T^+ \left(T, \frac{1}{z} \right) = W_+ \left(\frac{1}{z} \right),$$

$$(3.63)(b) \quad \left\| e_T^- \left(\cdot, \frac{1}{z} \right) - r_T \hat{\psi}_- \left(\cdot, \frac{1}{z} \right) \right\|_{[1, T]} \rightarrow 0$$

and as in (3.44)

$$(3.63)(c) \quad \lim_{T \rightarrow \infty} e_T^+ \left(1, \frac{1}{z} \right) = 0$$

uniformly for $z \in \Gamma$.

Inserting into the boundary condition (3.61)(b), (c) results in a block system of the form (3.42) and using (3.43) gives

$$(3.64)(a) \quad \begin{aligned} u_T = e_T^+ \left(\cdot, \frac{1}{z} \right) & \left(-\frac{1}{z} \left(S(T)W_+ \left(\frac{1}{z} \right) \right)^{-1} S(T)FG_+ + o(T) \right) \\ & + e_T^- \left(\cdot, \frac{1}{z} \right) o(T) \end{aligned}$$

uniformly for $z \in \Gamma$. The solvability of the problem (3.61) follows from the invertibility of the (analytic) matrix $S(T)W_+(1/z)$ which is a direct consequence of Assumption (3.20). Combining all gives

$$(3.64)(b) \quad \begin{aligned} & ((z - V_T)^{-1}(V_T r_T - r_T V)(z - V)^{-1}g)(t) \\ & = \frac{1}{z^2} e_T^+ \left(t, \frac{1}{z} \right) \left(S(T)W_+ \left(\frac{1}{z} \right) \right)^{-1} S(T)\psi \left(Gg, \frac{1}{z} \right)(T) \\ & \quad + e_T^- \left(t, \frac{1}{z} \right) A_1(T, z) + (z - V_T)^{-1} F\bar{e}_T^-(t)A_2(T, z) + A_3(T, z), \end{aligned}$$

where

$$(3.65) \quad \|A_i(t, z)\| \leq o(T)\|g\|_{[1, \infty]} \quad \text{for } i = 1, 2, 3,$$

uniformly for $z \in \Gamma$.

The first summand on the right-hand side of (3.64)(b) is holomorphic in S_μ and continuous in \bar{S}_μ . Therefore its contour integral along Γ vanishes.

Now we define the imbedding operators

$$(3.66)(a) \quad i_T: C([1, T]) \rightarrow C_0([1, \infty]),$$

$$(3.66)(b) \quad (i_T f_T)(t) = \begin{cases} f_T(t), & 1 \leq t \leq T, \\ f_T(T) \frac{T}{t}, & t \geq T. \end{cases}$$

Obviously $r_T i_T f_T = f_T$ and $\|i_T f_T\|_{[1, \infty]} = \|f_T\|_{[1, T]}$ hold.

Because of (3.64), (3.65) the operator

$$(3.67) \quad (z - V_T)^{-1}(V_T r_T - r_T V)(z - V)^{-1} i_T: C([1, T]) \rightarrow C([1, T])$$

is discretely compact for every $z \in \Gamma$, and the contour-integral-operator (3.27) is also discretely compact, see Grigorieff [3]. Because $\text{Range}(E)$ is finite dimensional, $r_T E i_T$ is discretely compact and so is $E_T r_T i_T = E_T$ and

$$(3.68) \quad \text{rank}(E_T(\mu)) = \text{rank}(E(\mu)).$$

Therefore it is guaranteed that the eigenvalue $\mu = 1/\lambda$ is stable with regard to the V_T 's (see Grigorieff [3]), so that there are exactly $m = \text{rank}(E(\mu))$ eigenvalues μ^1_T, \dots, μ^m_T of V_T which converge to μ , and the estimates

$$(3.69) \quad \text{gap}(\text{Range}(E_T), r_T(\text{Range}(E))) \leq \text{const} \|(V_T r_T - r_T V)|_{\text{Range}(E)}\|_{[1, T]}$$

and

$$(3.70) \quad \max \left(\left| \frac{1}{\hat{\mu}_T} - \lambda \right|, \max_i |\lambda'_T - \lambda|^\beta \right) \leq \text{const} \|(V_T r_T - r_T V)|_{\text{Range}(E)}\|_{[1, T]}$$

hold; see Grigorieff [3].

However, a stronger estimate can be derived by proceeding as Osborn [16] did but without carrying out the last estimates which lead to his Theorems 1, 2, 3. In the same way the estimates given by Grigorieff [3] can be changed. We get

$$(3.71)(a) \quad \text{gap}(\text{Range}(E_T), r_T(\text{Range}(E))) \leq \text{const} \| (E_T r_T - r_T E) |_{\text{Range}(E)} \|,$$

$$(3.71)(b) \quad \max \left(\left| \frac{1}{\hat{\mu}_T} - \lambda \right|, \max_{i=1(1)m} |\lambda_T^i - \lambda|^\beta \right) \leq \text{const} \| (E_T (V_T r_T - r_T V)) |_{\text{Range}(E)} \|.$$

An estimate for the right-hand side of (3.71)(a) can be obtained by using (3.64)(b) with $g \in \text{Range}(E)$ and (3.65). We obtain, collecting the results,

THEOREM 3.1. *If the problem (1.1), (1.2), (1.3) satisfies (3.4) and if (3.20) holds, then to every eigenvalue $\lambda \in \Omega$ of (1.1), (1.2), (1.3) with algebraic multiplicity m there are exactly m eigenvalues λ_T^i of (1.4), (1.5), (1.6) in a sufficiently small neighborhood of λ and they fulfill*

$$(3.72)(a) \quad |\lambda - \lambda_T^i| \leq \text{const } o(T)^{1/\beta} \exp \left(\frac{\nu_-(\lambda) + \varepsilon}{(\alpha + 1)\beta} T^{\alpha+1} \right), \quad T \rightarrow \infty,$$

$$(3.72)(b) \quad \left| \lambda - \frac{1}{\hat{\mu}_T} \right| < \text{const } o(T) \exp \left(\frac{\nu_-(\lambda) + \varepsilon}{\alpha + 1} T^{\alpha+1} \right), \quad T \rightarrow \infty,$$

where $\hat{\mu}_T$ and β are defined as in Theorem 2.2 and $\nu_-(\lambda)$ is the largest (in modulus) negative real part of the eigenvalues of $A(\infty) + \lambda G(\infty)$. The spectral projections satisfy $\text{rank}(E_T) = m$ and

$$(3.72)(c) \quad \text{gap}(\text{Range}(E), r_T(\text{Range}(E_T))) \leq \text{const } o(T) \exp \left(\frac{\nu_-(\lambda) + \varepsilon}{\alpha + 1} T^{\alpha+1} \right)$$

as $T \rightarrow \infty$.

Therefore the standard error estimates are not sharp for this problem. Tracing the history of the $o(T)$, we get

$$(3.73) \quad o(T) \leq \max(\|A(T) - A(\infty)\|, \|G(T) - G(\infty)\|, \bar{\sigma}(\kappa + \varepsilon_1, T)),$$

where $\bar{\sigma}$ is defined as in (3.44)(b) and ε_1 is small when the radius of Γ is sufficiently small.

$S(T)$ can be chosen independently of λ for a large class of problems, for example, if $G(\infty)$ is regular. In this case we can set $G(\infty) = I$ because this always can be achieved by a linear transformation. Then $F(\lambda) \equiv F$ and (3.20) is equal to (2.38). Ω is then the strip $\nu_- < \lambda < \nu_+$ where ν_- is the largest negative and ν_+ is the smallest positive real part of eigenvalues of $A(\infty)$. In this case the asymptotic boundary condition (2.55) can be used.

In the case that $G(\infty) \neq 0$ is not regular, $S(T)$ can be chosen independently of λ if we know a sufficiently close approximation $\tilde{\lambda} \in \Omega$ to an eigenvalue λ of (1.1), (1.2), (1.3). Then we rewrite (1.1) as

$$(3.74) \quad \begin{aligned} y' - t^\alpha (A(t) + \tilde{\lambda} G(t)) y &= \gamma t^\alpha G(t) y, & \gamma &= \lambda - \tilde{\lambda}. \\ \tilde{A}(t) &= A(t) + \tilde{\lambda} G(t). \end{aligned}$$

The isolatedness of the eigenvalues $\lambda \in \Omega$ guarantees that the problem

$$(3.75)(a) \quad y'_h - t^\alpha \tilde{A}(t)y_h = 0,$$

$$(3.75)(b) \quad By_h(1) = 0,$$

$$(3.75)(c) \quad y_h \in C([1, \infty])$$

has only the trivial solution $y_h \equiv 0$ if $\tilde{\lambda}$ is sufficiently close to λ . Therefore the above theory can be used for the eigenvalue problem (3.74), (1.2), (1.3) (with γ as eigenparameter). The change consists of taking $\tilde{\lambda}$ instead of 0 as reference point.

We set

$$(3.76) \quad S \equiv S(T) = (G_+)^T F^{-1}(\tilde{\lambda}),$$

where the superscript T denotes transposition and $F(\tilde{\lambda})$ transforms $A(\infty) + \tilde{\lambda}G(\infty) = \tilde{A}(\infty)$ to its Jordan canonical form.

Defining $F = F(\tilde{\lambda})$, (2.40), (2.41) follows immediately.

From the analysis for the approximating problems (1.4), (1.5), (1.6) it follows that it is sufficient to require that (3.20) holds locally if the particular eigenvalue λ is to be calculated. ‘‘Locally’’ means in this context the closed set bounded by the contour $(1/\mu)(\Gamma)$ defined in (3.9).

Since the family of projections $F(\lambda)D_+ F^{-1}(\lambda)$ is holomorphic in Ω , there is a nonsingular $r_+ \times r_+$ matrix $T(\lambda)$ so that

$$(3.77) \quad W_+(\lambda) = F(\lambda)G_+ T(\lambda) \text{ is holomorphic in } \Omega.$$

Therefore

$$\begin{aligned} SF(\lambda)G_+ T(\lambda) &= (G_+)^T F^{-1}(\tilde{\lambda})(F(\tilde{\lambda})G_+ T(\tilde{\lambda}) + O(|\lambda - \tilde{\lambda}|)) \\ &= T(\tilde{\lambda}) + O(|\lambda - \tilde{\lambda}|) \end{aligned}$$

holds, and

$$(3.78) \quad (SF(\lambda)G_+)^{-1} = T(\lambda)T(\tilde{\lambda})^{-1} + O(|\lambda - \tilde{\lambda}|)$$

for λ sufficiently close to $\tilde{\lambda}$. So (3.20) holds locally for λ close to $\tilde{\lambda}$, and the asymptotic boundary condition (3.76) can be used for the calculation of λ .

This analysis leads to the idea to use asymptotic boundary conditions which depend on the eigenparameter λ . This leads, even in the case that the ‘infinite’ problem is a linear eigenvalue problem, to nonlinear approximating ‘finite’ eigenvalue problems which, suggested by Keller [9], have been successfully used in computation (see Ng and Reid [15]), and their analysis will be presented in a subsequent paper.

However, for many important fluid-dynamical problems it is possible to choose simpler asymptotic boundary conditions. An example is presented in Section 5.

4. Imaginary Eigenvalues of $A(\infty)$. We are now going to neglect the crucial restriction that all eigenvalues of $A(\infty)$ have a nonzero real part, but we will require a sufficiently fast convergence of $G(t)$ to 0 which puts us back into the compactness argument of Section 2.

We assume that

$$(4.1) \quad A\left(\frac{1}{\delta}\right) \in C^{(\alpha+1)\max(k_0, k_-)+1}\left(\left[0, \frac{1}{\delta}\right]\right), \quad \delta \geq 1,$$

$$(4.2) \quad \|G(t)\| \leq \text{const } t^{-(\alpha+1)k_0-\epsilon}, \quad \epsilon > 0,$$

where k_0 is the largest algebraic multiplicity of an eigenvalue of $A(\infty)$ with real part zero and k_- is the largest algebraic multiplicity of an eigenvalue of $A(\infty)$ with negative real part.

Markowich [12], [13] has shown that there is a solution operator \bar{H} of the inhomogeneous problem (2.8) which fulfills

$$(4.3) \quad \|(\bar{H}f)(t)\| \leq \text{const } t^{-\epsilon}(\ln t)^j \|f\|_{[1,\infty]}, \quad 1 \leq j \leq n,$$

if (4.2) holds.

Therefore, if the homogeneous problem (2.8), (2.9), (2.10) has only the trivial solution $y \equiv 0$, then the operator V (see (2.16)) is well defined, and as the sum of a degenerate and a compact operator it is compact and the same consideration as in Section 2 holds for the eigenvalues and the generalized eigenvectors, except the decay statements, because the eigenvalues with real part zero may produce solutions which are asymptotically constant or which decay algebraically. An algorithm which determines the nature of the basic solutions under the assumption (4.1) is given in Markowich [13, Sections 3 and 4].

The construction of the supplementary boundary condition $S(T)x_T(T) = 0$ for the approximating problems (1.4), (1.5), (1.6) now relies heavily on the asymptotic nature of the basic solutions and is explained in Markowich [14, Sections 3 and 4]. The matrix $S(T)$ constructed in the mentioned paper takes care that the basic solutions, which are in $C([1, \infty])$ but which do not decay sufficiently fast, are damped by the multiplication with $S(T)$ so that norm convergence of the operators V_T , defined as in (2.32), to V results; see Markowich [14, Section 4].

Exponential convergence of eigenvalues and spectral subspaces holds if all (basic) solutions of the problem

$$(4.4) \quad y' - t^\alpha A(t)y = 0,$$

$$(4.5) \quad y \in C([1, \infty]),$$

decay exponentially.

5. The Orr-Sommerfeld Equation. The Orr-Sommerfeld equation (see Ng and Reid [15]) governs the stability of laminar boundary layers in the parallel flow approximation:

$$(5.1) \quad \frac{1}{iR\alpha} \left(\frac{d^2}{dz^2} - \alpha^2 \right)^2 \phi - \left[(U(z) - \lambda) \left(\frac{d^2}{dz^2} - \alpha^2 \right) \phi - U'''(z)\phi \right] = 0, \quad 0 \leq z < \infty,$$

$\alpha \in R, \alpha > 0$. $\phi(z)e^{i\alpha(x-\lambda t)}$ is the disturbance stream function, $R > 0$ is the Reynolds number, and $U(z)$ is the velocity distribution fulfilling

$$(5.2) \quad U \in C^2([0, \infty]), \quad U(\infty) = 1, \quad U''(\infty) = 0.$$

The boundary conditions for the Orr-Sommerfeld problem at $z = 0$ and $z = \infty$ are

$$(5.3) \quad \phi(0) = \phi'(0) = 0,$$

$$(5.4) \quad \phi(\infty) = \phi'(\infty) = 0.$$

This problem is of singular perturbation type for R large, but we disregard this computational difficulty and just derive appropriate asymptotic boundary conditions.

We substitute

$$(5.5) \quad y = (\phi, \phi', \phi'', \phi''')^T$$

and get the problem

$$(5.6) \quad y' - \overbrace{\begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ f_1(z) & 0 & f_2(z) & 0 \end{bmatrix}}^{A(z)} y = \lambda \overbrace{\begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ a & 0 & b & 0 \end{bmatrix}}^G y, \quad 0 \leq z < \infty,$$

$$(5.7) \quad \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix} y(0) = 0,$$

$$(5.8) \quad y \in C([0, \infty)),$$

where

$$(5.9) \quad \begin{aligned} (a) \quad & f_1(z) = -(\alpha^4 + i\alpha R(\alpha^2 U(z) + U''(z))), \\ (b) \quad & f_2(z) = 2\alpha^2 + i\alpha R U(z), \end{aligned}$$

and

$$(5.10) \quad (a) \quad a = i\alpha^3 R, \quad (b) \quad b = -i\alpha R.$$

The eigenvalues of $A(\infty)$ are

$$(5.11) \quad \nu_1 = \alpha, \quad \nu_2 = (\alpha^2 + i\alpha R)^{1/2}, \quad \nu_3 = -\alpha, \quad \nu_4 = -(\alpha^2 + i\alpha R)^{1/2},$$

so that $\text{Re } \nu_1, \text{Re } \nu_2 > 0; \text{Re } \nu_3, \text{Re } \nu_4 < 0$ and the eigenvalues of $A(\infty) + \lambda G$ are

$$(5.12) \quad \begin{aligned} \nu_1(\lambda) &= \alpha, \quad \nu_2(\lambda) = (\alpha^2 + i\alpha R(1 - \lambda))^{1/2}, \\ \nu_3(\lambda) &= -\alpha, \quad \nu_4(\lambda) = -(\alpha^2 + i\alpha R(1 - \lambda))^{1/2}, \end{aligned}$$

so that $\text{Re } \nu_1(\lambda), \text{Re } \nu_2(\lambda) > 0; \text{Re } \nu_3(\lambda), \text{Re } \nu_4(\lambda) < 0$ for all $\lambda \in \Omega = C - \{\lambda \mid \text{Re } \lambda = 1, \text{Im } \lambda \leq -\alpha/R\}$ holds. We get

$$(5.13)(a) \quad J(\infty) = \text{diag}(\nu_1, \nu_2, \nu_3, \nu_4) \quad \text{for } \lambda \in \Omega - \{1\},$$

and

$$(5.13)(b) \quad J(\infty) = \begin{bmatrix} \nu_1 & 1 & & 0 \\ & \nu_1 & & \\ & & \nu_2 & 1 \\ 0 & & & \nu_2 \end{bmatrix} \quad \text{for } \lambda = 1,$$

so that

$$(5.14) \quad G_- = \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad G_+ = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$$

holds. We calculate

$$(5.15)(a) \quad F(\lambda) = \begin{bmatrix} v_i^1(\lambda) \\ v_i^2(\lambda) \\ v_i^3(\lambda) \end{bmatrix}_{i=1(1)4} \quad \text{for } \lambda \in \Omega - \{1\}, F(0) = F,$$

$$(5.15)(b) \quad F(1) = \begin{bmatrix} 1 & 0 & 1 & 0 \\ \alpha & 1 & -\alpha & 1 \\ \alpha^2 & 2\alpha & \alpha^2 & 0 \\ \alpha^3 & 3\alpha^2 & -\alpha^3 & \alpha^2 \end{bmatrix}.$$

All eigenvalues $\lambda \in \Omega$ are isolated and have finite algebraic multiplicities.

We choose the matrix $S(Z)$ which sets up the asymptotic boundary condition $S(Z)y_Z(Z) = 0, Z > 0$, independent of Z , so that

$$(5.16) \quad S(Z) \equiv S = (S_{ij})_{i=1,2; j=1,2,3,4}$$

holds. Then the regularity condition (3.20) reduces to:

$$(5.17) \quad \det \begin{bmatrix} \sum_{j=1}^4 S_{1j} \alpha^{j-1} & \sum_{j=1}^4 S_{1j} v_2(\lambda)^{j-1} \\ \sum_{j=1}^4 S_{2j} \alpha^{j-1} & \sum_{j=1}^4 S_{2j} v_2(\lambda)^{j-1} \end{bmatrix} \neq 0 \quad \text{for } \lambda \in \Omega - \{1\},$$

and

$$(5.18) \quad \det \begin{bmatrix} \sum_{j=1}^4 S_{1j} \alpha^{j-1} & \sum_{j=1}^4 S_{1j} (j-1) \alpha^{j-2} \\ \sum_{j=1}^4 S_{2j} \alpha^{j-1} & \sum_{j=1}^4 S_{2j} (j-1) \alpha^{j-2} \end{bmatrix} \neq 0 \quad \text{for } \lambda = 1.$$

For example, the ‘natural’ asymptotic boundary condition

$$(5.19) \quad S = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}$$

satisfies (5.17) and (5.18).

The order of convergence for eigenvalues and spectral subspaces of the approximating problems (1.4), (1.5), (1.6), where S fulfills (5.17), (5.18), at a particular eigenvalue $\lambda \in \Omega$ of (5.6), (5.7), (5.8), can be estimated by

$$(5.20) \quad o(Z) \exp((\max(-\alpha, \operatorname{Re} v_4(\lambda)) + \epsilon)Z/\beta),$$

where

$$(5.21) \quad \operatorname{Re} \lambda_4(\lambda) = - \left(\frac{\alpha^2 + \alpha R \operatorname{Im} \lambda}{2} + \left(\frac{(\alpha^2 + \alpha R \operatorname{Im} \lambda)^2}{4} + \frac{\alpha^2 R^2 (1 - \operatorname{Re} \lambda)^2}{4} \right)^{1/2} \right)^{1/2}.$$

holds and β is the ascent of λ .

Computation of the Orr-Sommerfeld problem using the boundary conditions set up by (5.19) can be found in Grosch and Orszag [4]. They used the Blasius velocity profile $U(z) = 1 + O(e^{-wz^2})$, $w > 0$.

Their numerical experiments indicate that the order of convergence is $e^{-2\alpha Z}$ in the case that $\alpha < |\operatorname{Re} \nu_4(\lambda)| < 1$ and λ has ascent 1. Checking our order formula (3.73), (3.79) gives

$$(5.22) \quad o(Z) \leq \max \left(e^{-wZ^2}, \max_{z \in [\delta, Z]} e^{-wZ^2 + (\alpha - \varepsilon)(z - Z)} \right) = \text{const } e^{(-\alpha + \varepsilon)Z},$$

and the order of convergence the theory predicts is $e^{-2(\alpha - \varepsilon)Z}$ for eigenvalues and spectral subspaces at eigenvalues with ascent 1.

Acknowledgement. The author is grateful to Carl de Boor (Mathematics Research Center, Madison, Wisconsin) and Richard Weiss (Technical University of Vienna, Institute for Applied Mathematics) for stimulating discussions.

Department of Mathematics
The University of Texas at Austin
RLM 8-100
Austin, Texas 78712

1. C. DE BOOR & B. SWARTZ, *Collocation Approximation to Eigenvalues of an Ordinary Differential Equation. The Principle of the Thing*, MRC Tech. Report #1937, Madison, Wisc., 1980.
2. N. DUNFORD & J. T. SCHWARTZ, *Linear Operators*. I, Interscience, New York, 1957; Vol. VII in *Pure and Appl. Math.* (To appear.)
3. R. D. GRIGORIEFF, "Diskrete Approximation von Eigenwertproblemen. I, II," *Numer. Math.*, v. 24, 1975, pp. 355–374, 415–433.
4. CH. E. GROSCH & S. A. ORSZAG, "Numerical solution of problems in unbounded regions: Coordinate transforms," *J. Comput. Phys.*, v. 25, 1977, pp. 273–296.
5. F. R. DE HOOG & R. WEISS, "Difference methods for boundary value problems with a singularity of the first kind," *SIAM J. Numer. Anal.*, v. 13, 1976, pp. 775–813.
6. F. R. DE HOOG & R. WEISS, "On the boundary value problem for systems of ordinary differential equations with a singularity of the second kind," *SIAM J. Math. Anal.*, v. 11, 1980, pp. 46–61.
7. F. R. DE HOOG & R. WEISS, "An approximation method for boundary value problems on infinite intervals," *Computing*, v. 24, 1980, pp. 227–239.
8. T. KATO, *Perturbation Theory for Linear Operators*, Springer-Verlag, Berlin and New York, 1966.
9. H. KELLER, "Numerical solution of two point boundary value problems," *Regional Conf. Series in Appl. Math.*, No. 24, SIAM, Philadelphia, Pa., 1976.
10. H. O. KREISS, "Difference approximations for boundary and eigenvalue problems for ordinary differential equations," *Math. Comp.*, v. 26, 1972, pp. 649–686.
11. M. LENTINI & H. B. KELLER, "Boundary value problems on semi-infinite intervals and their numerical solution," *SIAM J. Numer. Anal.*, v. 13, 1980, pp. 577–604.
12. P. A. MARKOWICH, *Randwertprobleme auf Unendlichen Intervallen*, Dissertation, TU Wien, 1980.
13. A. MARKOWICH, "Analysis of boundary value problems on infinite intervals," *SIAM J. Appl. Math.*, v. 42, 1982, pp. 549–557.
14. P. A. MARKOWICH, "A theory for the approximation of solutions of boundary value problems on infinite intervals," *SIAM J. Math. Anal.*, v. 13, 1982, pp. 484–513.
15. B. S. NG & W. H. REID, "On the numerical solution of the Orr-Sommerfeld problem. Asymptotic initial conditions for shooting methods," *J. Comput. Phys.*, v. 38, 1980, pp. 275–293.
16. J. E. OSBORN, "Spectral approximation for compact operators," *Math. Comp.*, v. 28, 1975, pp. 712–725.
17. F. STUMMEL, "Diskrete Konvergenz linearer Operatoren. I," *Math. Ann.*, v. 190, 1970/71, pp. 45–92.
18. F. STUMMEL, "Diskrete Konvergenz linearer operatoren. II," *Math. Z.*, v. 120, 1971, pp. 231–264.