

## Further Inequalities for the Gamma Function\*

By **Andrea Laforgia**

**Abstract.** For  $\lambda > 0$  and  $k \geq 0$  we present a method which permits us to obtain inequalities of the type  $(k + \alpha)^{\lambda-1} < \Gamma(k + \lambda)/\Gamma(k + 1) < (k + \beta)^{\lambda-1}$ , with the usual notation for the gamma function, where  $\alpha$  and  $\beta$  are independent of  $k$ . Some examples are also given which improve well-known inequalities. Finally, we are also able to show in some cases that the values  $\alpha$  and  $\beta$  in the inequalities that we obtain cannot be improved.

**1. Introduction.** In this paper we are concerned with some inequalities for the function  $\Gamma(k + \lambda)/\Gamma(k + 1)$ , where  $k \geq 0$  and  $\lambda > 0$  is independent of  $k$ .

Many authors have studied inequalities for this function. For example, Gautschi [2, (7)] has proved that

$$(1.1) \quad \frac{1}{(k + 1)^{1-\lambda}} < \frac{\Gamma(k + \lambda)}{\Gamma(k + 1)} < \frac{1}{k^{1-\lambda}}, \quad 0 < \lambda < 1, k = 1, 2, \dots,$$

and, in the particular case  $\lambda = 1/2$ , Watson [5] has given the lower bound

$$(1.2) \quad \frac{\Gamma(k + 1/2)}{\Gamma(k + 1)} > \frac{1}{(k + 4\pi^{-1} - 1)^{1/2}}, \quad k > 1,$$

where  $k$  is real.

Recently Lorch [4] has given some useful improvements of the bounds in (1.1) and has used his results to obtain a very interesting inequality for ultraspherical polynomials.

Here we show that Lorch's method can be sharpened somewhat, so as to obtain new inequalities and improvements upon known inequalities.

Moreover our results refer to the general case of real and positive  $k$  and not only integer  $k$  as in Lorch's results.

**2. The Function  $A_k(\lambda; \alpha)$ .** From the relation [1, p. 257, (6.1.46)]

$$\lim_{k \rightarrow \infty} k^{b-a} \frac{\Gamma(k + a)}{\Gamma(k + b)} = 1$$

we see that

$$\lim_{k \rightarrow \infty} f_k = 1,$$

where

$$f_k \equiv \frac{\Gamma(k + \lambda)}{\Gamma(k + 1)} (k + \alpha)^{1-\lambda}, \quad k \geq 0, \quad \lambda, \alpha > 0.$$

Received April 18, 1983.

1980 *Mathematics Subject Classification*. Primary 33A15.

\*Work sponsored by the Consiglio Nazionale delle Ricerche-Italy.

Thus, if for some fixed  $\alpha$  we can show that  $f_k$  ultimately increases (or decreases) to 1 for  $k \rightarrow \infty$ , we can conclude that  $f_k < 1$  (or  $f_k > 1$ ) for  $k \geq k_0$ , and we obtain the inequalities

$$\frac{\Gamma(k + \lambda)}{\Gamma(k + 1)} < (k + \alpha)^{\lambda-1} \quad \text{or} \quad \frac{\Gamma(k + \lambda)}{\Gamma(k + 1)} > (k + \alpha)^{\lambda-1}, \quad k \geq k_0,$$

for the cases  $f_k < 1$  and  $f_k > 1$ , respectively. This suggests studying the function  $g_k \equiv f_{k+1}/f_k$  which, using the functional relation  $\Gamma(z + 1) = z\Gamma(z)$ , can be written in the form

$$g_k \equiv \frac{f_{k+1}}{f_k} = \frac{k + \lambda}{k + 1} \left( \frac{k + \alpha + 1}{k + \alpha} \right)^{1-\lambda}$$

to see for what values of  $\alpha$ ,  $\lambda$  and  $k$  one has  $g_k > 1$  or  $g_k < 1$ . Lorch in [4] has already used this method, but only for some particular values of  $\alpha$ . Since  $\lim_{k \rightarrow \infty} g_k = 1$ , we are interested in the monotonicity of  $g_k$  with respect to  $k$ . Letting  $k > 0$  be a continuous variable, we have

$$(k + \alpha)^{2-\lambda}(k + 1)^2(k + \alpha + 1)^\lambda g'_k = A_k(\lambda; \alpha),$$

where

$$(2.1) \quad A_k(\lambda; \alpha) = (1 - \lambda)(-\lambda k + 2\alpha k - \lambda + \alpha^2 + \alpha).$$

Thus we need to study the sign of the function  $A_k(\lambda; \alpha)$ .

When  $\alpha = \lambda/2$  we have

$$A_k(\lambda; \lambda/2) \equiv \frac{\lambda}{2}(1 - \lambda) \left( \frac{\lambda}{2} - 1 \right)$$

and

$$\begin{aligned} A_k(\lambda; \lambda/2) &> 0, & 1 < \lambda < 2, k \geq 0, \\ A_k(\lambda; \lambda/2) &< 0, & 0 < \lambda < 1 \text{ or } \lambda > 2, k \geq 0. \end{aligned}$$

We thus obtain

$$(2.2) \quad \frac{\Gamma(k + \lambda)}{\Gamma(k + 1)} < (k + \lambda/2)^{\lambda-1}, \quad 0 < \lambda < 1 \text{ or } \lambda > 2, k \geq 0,$$

and

$$(2.3) \quad \left( k + \frac{\lambda}{2} \right)^{\lambda-1} < \frac{\Gamma(k + \lambda)}{\Gamma(k + 1)}, \quad 1 < \lambda < 2, k \geq 0,$$

which for integer  $k$  are the inequalities obtained by Lorch [4]. For the case  $0 < \lambda < 1$  and real  $k$  the inequality (2.2) has also been proved independently by Kershaw [3]. We observe that the term  $\lambda/2$  in the bounds (2.2) and (2.3) cannot be improved upon. For example, if  $\lambda > 2$  and  $\alpha < \lambda/2$ , (2.1) shows that there exists a  $k_0$  such that for  $k \geq k_0$  the function  $A_k(\lambda; \alpha)$  becomes, and remains, positive.

This proves that, for (2.2) to be valid for any nonnegative real  $k$ , the term  $\lambda/2$  in (2.2) is best possible. Similarly, the term  $\lambda/2$  in (2.3) and in (2.2), for  $0 < \lambda < 1$ , cannot be replaced by  $\alpha > \lambda/2$ .

**3. New Inequalities.** The study of the sign of the function  $A_k(\lambda; \alpha)$  permits one to obtain new inequalities for  $\Gamma(k + \lambda)/\Gamma(k + 1)$ . For example, for  $\alpha = \frac{2}{3}\lambda$ , (2.1) shows that

$$A_k\left(\lambda; \frac{2}{3}\lambda\right) \equiv \frac{\lambda}{3}(1 - \lambda)\left(k + \frac{4}{3}\lambda - 1\right)$$

is positive for  $0 < \lambda < 1$  and  $k \geq 1$ , thus yielding the inequality

$$(3.1) \quad \left(k + \frac{2}{3}\lambda\right)^{\lambda-1} < \frac{\Gamma(k + \lambda)}{\Gamma(k + 1)}, \quad 0 < \lambda < 1, k \geq 1.$$

This lower bound is more precise than the one in Gautschi's inequality (1.1).

It is clear that (3.1) can be further improved if we confine ourselves to values  $k \geq k_0$  for some  $k_0$ . For example, if  $k \geq 5$ , it is sufficient to choose  $\alpha$  such that  $A_5(\lambda; \alpha) \geq 0$ , that is,  $\alpha \geq (-11 + \sqrt{121 + 24\lambda})/2$ . Since for these values of  $\alpha$  the condition  $2\alpha > \lambda$  is satisfied, we get  $A_k(\lambda; \alpha) > 0$  for  $k > 5$ , and the inequality

$$\left(k + \frac{-11 + \sqrt{121 + 24\lambda}}{2}\right)^{\lambda-1} < \frac{\Gamma(k + \lambda)}{\Gamma(k + 1)}, \quad 0 < \lambda < 1, k \geq 5,$$

holds.

This lower bound is more precise than the one in (1.2), which is valid only for  $\lambda = 1/2$ . In fact for  $\lambda = 1/2$  the term  $4\pi^{-1} - 1 = 0.273 \dots$  in (1.2) is replaced by  $0.266 \dots$  in the new inequality. Generally, we have that for every  $\epsilon > 0$  there exists a  $k_0(\epsilon)$  such that the inequality

$$\left(k + \frac{\lambda + \epsilon}{2}\right)^{\lambda-1} < \frac{\Gamma(k + \lambda)}{\Gamma(k + 1)}, \quad 0 < \lambda < 1, k \geq k_0,$$

holds.

Other useful inequalities can be obtained for the case  $1 < \lambda < 2$ . If we put  $\alpha = \lambda/2 + 1/8$  we find  $A_k(\lambda; \lambda/2 + 1/8) < 0$  for  $k \geq 0$ , and this proves the inequality

$$\frac{\Gamma(k + \lambda)}{\Gamma(k + 1)} < \left(k + \frac{\lambda}{2} + \frac{1}{8}\right)^{\lambda-1}, \quad 1 < \lambda < 2, k \geq 0.$$

Similarly, if  $\alpha = \frac{\lambda}{2} + \frac{1}{10}$ , one has  $A_k(\lambda; \lambda/2 + 1/10) < 0$  for  $k \geq 1$ , and we obtain

$$\frac{\Gamma(k + \lambda)}{\Gamma(k + 1)} < \left(k + \frac{\lambda}{2} + \frac{1}{10}\right)^{\lambda-1}, \quad 1 < \lambda < 2, k \geq 1.$$

Our calculations show that the last upper bounds hold in the general case  $\lambda > 1$ , but our interest is for  $1 < \lambda < 2$ , because (2.2) is more precise for  $\lambda > 2$ .

It is clear that it is possible to obtain many other inequalities of a similar type.

**Acknowledgments.** The author is grateful to Professor Gian Carlo Rota for the hospitality extended by the Department of Mathematics of M.I.T. (Massachusetts Institute of Technology) where the author was a visitor when this paper was written. He also is indebted to Professor Walter Gautschi for pointing out the reference [3].

Department of Mathematics  
 University of Torino  
 Via Carlo Alberto, 10  
 10123 Torino, Italy

1. M. ABRAMOWITZ & I. A. STEGUN, Editors, *Handbook of Mathematical Functions*, Appl. Math. Series No. 55, National Bureau of Standards, Washington, D.C., 1964.
2. W. GAUTSCHI, "Some elementary inequalities relating to the gamma and incomplete gamma function," *J. Math. Phys.*, v. 38, 1959, pp. 77–81.
3. D. KERSHAW, "Some extensions of Gautschi's inequalities for the gamma function," *Math. Comp.*, v. 41, 1983, pp. 607–611.
4. L. LORCH, "Inequalities for ultraspherical polynomials and the gamma function," *J. Approx. Theory*. (To appear.)
5. G. N. WATSON, "A note on gamma functions," *Proc. Edinburgh Math. Soc.* (2), v. 11, 1958/59, Edinburgh Math. Notes No. 41, 1959, pp. 7–9.