

## $L_\infty$ -Boundedness of $L_2$ -Projections on Splines for a Multiple Geometric Mesh\*

By Rong-Qing Jia\*\*

**Abstract.** This paper concerns the  $L_2$ -projectors from  $L_\infty$  to the normed linear space of polynomial splines. It is shown that for the multiple geometric meshes the  $L_\infty$  norms of the corresponding  $L_2$ -projectors are bounded independently of the mesh ratio.

**0. Introduction.** Let us begin with some notation. Let  $k$  be a positive integer, and  $\mathbf{x} = (x_i)_{i \in \mathbf{Z}}$  a real nondecreasing knot sequence with  $x_i < x_{i+k}$ , all  $i$ . Set

$$x_{-\infty} := \lim_{i \rightarrow -\infty} x_i, \quad x_\infty := \lim_{i \rightarrow \infty} x_i, \quad I := (x_{-\infty}, x_\infty).$$

For  $i \in \mathbf{Z}$  and  $x \in I$ , define

$$M_{i,k}(x) := k[x_i, \dots, x_{i+k}](\cdot - x)_+^{k-1}, \quad N_{i,k}(x) := (x_{i+k} - x_i)M_{i,k}(x)/k,$$

where  $[\rho_0, \dots, \rho_r]f$  denotes the  $r$ th divided difference of the function  $f$  at the points  $\rho_0, \dots, \rho_r$ . Then  $N_{i,k}$  is called the  $i$ th  $B$ -spline of order  $k$  on the knot sequence  $\mathbf{x}$ .

For  $\mathbf{a} \in \mathbf{R}^{\mathbf{Z}}$ , the rule

$$f(x) := \sum_i \mathbf{a}(i)N_{i,k}(x)$$

defines a function on  $I$  if we take the sum to be pointwise. Every such function is called a polynomial spline of order  $k$  with the knot sequence  $\mathbf{x}$ , and their collection is denoted by  $\mathbb{S}_{k,\mathbf{x}}$ . Further, let  $\mathbb{S}$  denote the normed linear space of bounded polynomial splines of order  $k$  with the knot sequence  $\mathbf{x}$  and norm

$$\|f\| := \sup_{x \in I} |f(x)|.$$

We shall be concerned with  $P_{\mathbb{S}}$ , the orthogonal projector onto  $\mathbb{S}$  with respect to the ordinary inner product

$$(f, g) = \int_I f(x)g(x) dx,$$

but restricted to  $L_\infty(I)$ . We want to bound its norm

$$\|P_{\mathbb{S}}\| := \sup_{f \in L_\infty(I)} \|P_{\mathbb{S}}f\| / \|f\|_\infty.$$

Received March 26, 1984; revised December 16, 1985.

1980 *Mathematics Subject Classification*. Primary 41A15, 15A60.

\*This is a part of the author's thesis, which was submitted to the graduate school, University of Wisconsin-Madison, in partial fulfillment of the requirement for Ph.D. degree.

\*\**Present address*: Department of Mathematics, Zhejiang University, Hangzhou, Zhejiang, The People's Republic of China.

In 1973, de Boor [1] made the conjecture that

$$\sup_x \|P_{\mathbb{S}}\| \leq \text{const}_k < \infty.$$

This conjecture has been verified only for  $k = 2$  (Ciesielski [8]),  $k = 3$  and  $4$  (de Boor [5]). Moreover, de Boor [4] showed that  $P_{\mathbb{S}}$  is bounded in terms of the global mesh ratio. For geometric meshes, Höllig [12] proved that  $P_{\mathbb{S}}$  is bounded independently of the mesh ratio. Later on, Feng and Kozak [9] reproved this result. For a tri-multiple geometric mesh, Mityagin [15] established the uniform boundedness of  $P_{\mathbb{S}}$  for  $k = 6$ .

In this paper we shall be concerned with multiple geometric knot sequences  $(x_i)_{i \in \mathbb{Z}}$ ; that is,

$$x_{li} = x_{li+1} = \dots = x_{li+l-1} = q^i, \quad \text{all } i, 0 < q < \infty,$$

where  $l \in \mathbb{N}$  is the multiplicity of the knots. Our main result is

$$\sup_x \|P_{\mathbb{S}}\| \leq \text{const}_k < \infty$$

where  $\mathbf{x}$  runs through all the multiple geometric knot sequences. This result extends the results of Höllig and Mityagin.

Let  $A = A_q$  be the  $\mathbb{Z} \times \mathbb{Z}$  matrix given by

$$A(i, j) := \int M_{i,k} N_{j,k} \quad \text{for } i, j \in \mathbb{Z}.$$

It is shown by de Boor ([1], also [2]), that

$$D_k^{-2} \|A^{-1}\|_{\infty} \leq \|P_{\mathbb{S}}\|_{\infty} \leq \|A^{-1}\|_{\infty},$$

where  $D_k$  is a constant depending only on  $k$ . Thus, bounding  $P_{\mathbb{S}}$  is equivalent to bounding  $A^{-1}$ .

Now since the mesh  $(x_i)_{i \in \mathbb{Z}}$  is an  $l$ -multiple geometric one, we have

$$\begin{aligned} A(i + l, j + l) &= \int M_{i+l}(x) N_{j+l}(x) dx \\ &= \int M_{i+l}(x) N_j(q^{-1}x) dx = \int q M_{i+l}(qx) N_j(x) dx \\ &= \int M_i(x) N_j(x) dx = A(i, j). \end{aligned}$$

This shows that  $A$  is an  $l$ -block Toeplitz matrix. Moreover,  $A$  is totally positive (see [13, Chapter 10], also [2]). This motivates us to investigate totally positive block Toeplitz matrices.

**1. Totally Positive Block Toeplitz Matrices.** Let  $A$  be a bi-infinite  $N$ -block Toeplitz matrix. Then there exists a sequence of  $N \times N$  matrices  $A_j$  ( $j \in \mathbb{Z}$ ) such that  $A$  has the following form:

$$A = \begin{bmatrix} \cdots & \cdot & \cdot & \cdot & \cdots \\ \cdots & A_0 & A_1 & A_2 & \cdots \\ \cdots & A_{-1} & A_0 & A_1 & \cdots \\ \cdots & A_{-2} & A_{-1} & A_0 & \cdots \\ \cdots & \cdot & \cdot & \cdot & \cdots \end{bmatrix}.$$

Now

$$\hat{A}(z) := \sum_{n=-\infty}^{\infty} A_n z^n, \quad z \in \mathbf{C},$$

is a formal power series with matrix coefficients. It is called the symbol of  $A$ . If all the rows of  $A$  are in  $l_1$ , i.e.,

$$(1.1) \quad \sum_{j=-\infty}^{\infty} |a_{ij}| < \infty \quad \text{for all } i \in \mathbf{Z},$$

then  $\hat{A}(z)$  makes sense for  $|z| \leq 1$ . In this case,  $A$  determines a bounded linear operator from  $l_\infty$  to  $l_\infty$ . This operator is also denoted by  $A$ . A basic question is when  $A$  is boundedly invertible. In the case where  $A$  is also totally positive, the following theorem gives an answer. In its statement and further on, we use the abbreviation

$$\mathbf{1}' := \left( (-1)^{j-1} \right)_{j \in J}$$

for the vector on the index set  $J$  whose entries are alternately 1 and  $-1$ . Typically,  $J = (1, \dots, N)$  or else  $J = \mathbf{Z}$ . We also use

$$\omega := (-1)^N.$$

**THEOREM 1.1.** *Let  $A$  be a bi-infinite totally positive  $N$ -block Toeplitz matrix with all the rows in  $l_1$ . Then  $A$  is boundedly invertible if and only if*

$$\det \hat{A}(\omega) \neq 0.$$

Moreover,  $\det \hat{A}(\omega) \neq 0$  if and only if there exists  $b \in \mathbf{R}^N$  such that

$$(1.2) \quad \hat{A}(\omega)\mathbf{b} = \mathbf{1}',$$

and, for such  $\mathbf{b}$ ,  $\|A^{-1}\| = \|\mathbf{b}\|_\infty$ .

*Remark.* When  $A$  is a banded matrix, the first part of this theorem is already obtained in [7]. The above theorem removes the restriction of bandedness.

To prove Theorem 1.1, we need a lemma.

**LEMMA 1.1.** *Let  $\varphi$  be a real number. If  $\varphi/(2\pi)$  is not an integer, then for any positive integer  $n_0$  there exist integers  $k, l$  such that  $l > k \geq n_0$  and*

$$(1.3) \quad |\cos k\varphi| \geq \frac{1}{2}, \quad |\cos l\varphi| \geq \frac{1}{2} \quad \text{and} \quad \cos k\varphi \cos l\varphi < 0.$$

Furthermore, if  $\varphi/\pi$  is not an integer, then  $k$  and  $l$  can be chosen to be even numbers.

*Proof.* Suppose first that  $\varphi/2\pi$  is not an integer. Without loss of generality we may assume  $0 < \varphi < 2\pi$ . We shall argue case by case.

1°.  $0 < \varphi \leq 2\pi/3$ . By  $[x]$  we denote the integer part of  $x$ , i.e., the largest integer  $\leq x$ . Let

$$k := [(2n_0\pi + \pi/3)/\varphi], \quad l := [((2n_0 + 1)\pi + \pi/3)/\varphi].$$

Then

$$\begin{aligned} k\varphi &\leq 2n_0\pi + \pi/3 \leq (k+1)\varphi, \\ k\varphi &\geq 2n_0\pi + \pi/3 - \varphi \geq 2n_0\pi - \pi/3; \end{aligned}$$

hence,

$$k\varphi \in [2n_0\pi - \pi/3, 2n_0\pi + \pi/3].$$

Similarly,

$$l\varphi \in [(2n_0 + 1)\pi - \pi/3, (2n_0 + 1)\pi + \pi/3].$$

Thus (1.3) is fulfilled.

2°.  $4\pi/3 \leq \varphi < 2\pi$ . In this case we set

$$k := [2n_0\pi/(2\pi - \varphi)] \quad \text{and} \quad l := [(2n_0 + 1)\pi/(2\pi - \varphi)].$$

3°.  $2\pi/3 < \varphi < \pi$ . Since  $4\pi/3 < 2\varphi < 2\pi$ , this case reduces to 2°.

4°.  $\varphi = \pi$ . This case is trivial.

5°.  $\pi < \varphi < 4\pi/3$ . Since  $2\pi < 2\varphi < 8\pi/3$ , this case reduces to 1°.

Assume now that  $\varphi/\pi$  is not an integer. Then  $2\varphi/2\pi$  is not an integer and we obtain (1.3) with  $\varphi$  replaced by  $2\varphi$ ; hence we obtain (1.3) itself with both  $k$  and  $l$  even. This ends the proof of Lemma 1.1.  $\square$

*Proof of Theorem 1.1.* It is known (see [11]) that  $A$  is boundedly invertible if and only if  $\det \hat{A}(e^{i\theta}) \neq 0$  for all real  $\theta$ . Thus, to prove the theorem, it is sufficient to show that the solvability of (1.2) for  $\mathbf{b}$  implies that  $\det \hat{A}(e^{i\theta}) \neq 0$  for all  $\theta$ . We prove this implication by contradiction. Suppose that  $\det \hat{A}(e^{i\theta}) = 0$  for some real  $\theta$ . Then there exists a nonzero  $\mathbf{y} \in \mathbf{C}^N$  such that  $\hat{A}(e^{i\theta})\mathbf{y} = 0$ . Without loss of generality we may assume  $y_1 \neq 0$ , and, after multiplying by a suitable complex number, we may assume further that  $y_1$  is a positive real number. Let

$$(1.4) \quad \begin{aligned} y_{kN+j} &:= (e^{i\theta})^k y_j, \quad k \in \mathbf{Z}; \quad j = 1, 2, \dots, N, \\ \tilde{\mathbf{y}} &:= (\dots, y_{-1}, y_0, y_1, y_2, \dots). \end{aligned}$$

Then  $A\tilde{\mathbf{y}} = 0$ . Suppose that  $\mathbf{b}$  is a solution of (1.2). Let

$$\begin{aligned} b_{kN+j} &:= \omega^k b_j, \quad k \in \mathbf{Z}; \quad j = 1, 2, \dots, N, \\ \tilde{\mathbf{b}} &:= (\dots, b_{-1}, b_0, b_1, b_2, \dots). \end{aligned}$$

Then  $A\tilde{\mathbf{b}} = \mathbf{1}'$ .

We consider the two possibilities:  $e^{i\theta} \neq \omega$  or  $e^{i\theta} = \omega$  separately. First we consider the case  $e^{i\theta} \neq \omega$ . Fix  $\varepsilon > 0$  such that  $y_1/2 > \varepsilon|b_1|$  and let  $\mathbf{u} := \text{Re } \tilde{\mathbf{y}} + \varepsilon\tilde{\mathbf{b}}$ . Then  $\mathbf{u} \in l_\infty$  and  $A\mathbf{u} = \varepsilon\mathbf{1}'$ . By (1.1) there exists a positive integer  $m$  such that

$$(1.5) \quad \sum_{|k-j|>m} |a_{jk}| \|\mathbf{u}\|_\infty < \varepsilon.$$

In the following we shall argue separately in terms of  $N$  even or odd.

(i)  $N$  is even. In this case,  $e^{i\theta} \neq \omega = 1$ ; hence  $\theta/2\pi$  is not an integer. By Lemma 1.1, there exist positive integers  $k_i, l_i$  ( $i = 0, \dots, 2m$ ) such that

$$(1.6) \quad k_0 < l_0 < k_1 < l_1 < \dots < k_{2m} < l_{2m},$$

$$(1.7) \quad |\cos k_i\theta| \geq \frac{1}{2}, \quad |\cos l_i\theta| \geq \frac{1}{2}, \quad \text{and} \quad \cos k_i\theta \cos l_i\theta < 0.$$

Since  $\varepsilon|b_1| < y_1/2$ , we have

$$u_{k_iN+1}u_{l_iN+1} = (y_1 \cos k_i\theta + \varepsilon b_1)(y_1 \cos l_i\theta + \varepsilon b_1) < 0.$$

But  $(l_iN + 1) - (k_iN + 1)$  is an even number; therefore

$$(1.8) \quad S^-(u_{k_iN+1}, \dots, u_{l_iN+1}) \leq (l_iN - k_iN) - 1, \quad i = 0, 1, \dots, 2m,$$

where by  $S^-$  we denote the number of strong sign changes in a sequence (see [13, Chapter 5]). Now set

$$\mathbf{v} := (u_{-m+k_0N+1}, \dots, u_{m+l_{2m}N+1}),$$

$$B := (a_{ij})_{k_0N+1 \leq i \leq l_{2m}N+1, -m+k_0N+1 \leq j \leq m+l_{2m}N+1}.$$

Then  $B$  is a totally positive  $((l_{2m} - k_0)N + 1) \times ((l_{2m} - k_0)N + 2m + 1)$ -matrix. The fact that  $A\mathbf{u} = \epsilon\mathbf{1}'$ , together with (1.5), implies that

$$S^-(B\mathbf{v}) = (l_{2m} - k_0)N.$$

On the other hand, (1.8) gives

$$S^-(\mathbf{v}) \leq (l_{2m} - k_0)N - (2m + 1) + 2m = (l_{2m} - k_0)N - 1.$$

Since  $B$  is totally positive, it is variation-diminishing, namely

$$S^-(B\mathbf{v}) \leq S^-(\mathbf{v})$$

(see [13, Theorem 5.1.4]). This is a contradiction.

(ii)  $N$  is odd. In this case,  $e^{i\theta} \neq \omega = -1$ . If  $e^{i\theta} \neq 1$  also, then  $\theta/\pi$  is not an integer. By Lemma 1.1 there exist even positive integers  $k_i, l_i$  ( $i = 0, \dots, 2m$ ) such that (1.6) and (1.7) hold. Thus the argument for case (i) is also valid for this case.

Now, assume  $e^{i\theta} = 1$ . Let

$$k_i = i, \quad l_i = (i + 1), \quad i = 0, 1, \dots, 2m.$$

Then  $(l_iN + 1) - (k_iN + 1)$  is an odd number, but  $\cos k_i\theta \cos l_i\theta > 0$ ; hence (1.8) is still true. Following the argument for case (i), we obtain the desired result.

It remains to treat the case  $e^{i\theta} = \omega$ . In this case, (1.4) becomes

$$y_{kN+j} = \omega^k y_j.$$

Thus we may assume that  $\tilde{\mathbf{y}}$  is a real sequence. For any  $\lambda \in \mathbf{R}$ ,  $\mathbf{c} := \tilde{\mathbf{b}} + \lambda\tilde{\mathbf{y}}$  is a real sequence and satisfies  $A\mathbf{c} = \mathbf{1}'$ . Thus we have

$$c_j c_{j+1} < 0 \quad \text{for all } j.$$

(This can be proved, as done before, by truncating  $A$  to a finite matrix, invoking the variation-diminishing property for it and using the periodicity of  $\mathbf{c}$ .) But that is impossible unless  $\tilde{\mathbf{y}} = \mathbf{0}$ .

Finally, since  $A\tilde{\mathbf{b}} = \mathbf{1}'$ , we have  $\|A^{-1}\| = \|\tilde{\mathbf{b}}\|_\infty$  (see [6]). Therefore

$$\|A^{-1}\| = \|\mathbf{b}\|_\infty.$$

The proof of Theorem 1.1 is complete.

## 2. Exponential Splines.

*Definition 2.1.*  $S \in \mathcal{S}_{2k}$  is called an exponential spline, if for some constant  $\lambda \in \mathbf{R}$ ,

$$S(qx) = \lambda S(x) \quad \text{for all } x \in I.$$

“The exponential splines”, said Schoenberg in his elegant monograph [17], “will be used as thoroughly as the American Indians utilized the buffalo, to the last bone.” The exponential splines also play an important role in this paper. By the above definition, every exponential spline is uniquely determined by its polynomial component in  $[1, q]$ . For reasons that will become clear later, we are particularly interested in the exponential splines

$$\varphi_j, \psi_j \quad (j = 0, 1, \dots, l - 1)$$

given by

$$\varphi_0(x) := x^k,$$

$$\varphi_j(x) := (-1)^{k-j+1} q^k [0, 1, \dots, k-1, k+1, \dots, 2k-j]_z \frac{x^z}{q^z - q^k}$$

for  $x \in [1, q]$  and  $j = 1, \dots, l-1$ ,

$$\psi_j(x) := (-1)^{k-j+1} q^k [0, 1, \dots, 2k-j-1]_z \frac{x^z}{q^z + q^k}$$

for  $x \in [1, q]$  and  $j = 0, \dots, l-1$

(cf. [14]). Extend the domain of  $\varphi_j$  and  $\psi_j$  to  $(0, \infty)$  by the following rule:

$$\varphi_j(q^m x) = q^{mk} \varphi_j(x), \quad m \in \mathbf{Z}, x \in [1, q],$$

$$\psi_j(q^m x) = (-q^k)^m \psi_j(x), \quad m \in \mathbf{Z}, x \in [1, q].$$

We have to verify that  $\varphi_j$  and  $\psi_j$  are in  $\mathbb{S}_{2k}$ . Obviously,  $\varphi_j$  and  $\psi_j$  are polynomials on each interval  $[q^m, q^{m+1}]$ . Thus it suffices to show that

$$(2.1) \quad q^p \varphi_j^{(p)}(q) = q^k \varphi_j^{(p)}(1), \quad p = 0, 1, \dots, 2k-2,$$

$$(2.2) \quad q^p \psi_j^{(p)}(q) = -q^k \psi_j^{(p)}(1), \quad p = 0, 1, \dots, 2k-2.$$

Indeed, for  $j = 0$ , (2.1) is trivial. For  $1 \leq j \leq l-1$ , we have

$$\begin{aligned} q^p \varphi_j^{(p)}(q) &= q^p (-1)^{k-j+1} q^k [0, 1, \dots, k-1, k+1, \dots, 2k-j]_z \\ &\quad \cdot \frac{z(z-1) \cdots (z-p+1) q^{z-p}}{q^z - q^k} \\ &= (-1)^{k-j+1} q^k [0, 1, \dots, k-1, k+1, \dots, 2k-j]_z \\ &\quad \left\{ z(z-1) \cdots (z-p+1) \left( 1 + q^k \frac{1}{q^z - q^k} \right) \right\} \\ &= q^k \varphi_j^{(p)}(1). \end{aligned}$$

One can prove (2.2) in the same fashion.

We want to investigate some properties of these exponential splines. In this section, we mean by  $i$  the imaginary unit:  $i = \sqrt{-1}$ . Also we make the abbreviation  $t := \log q$ . We notice that the case  $q \leq 1$  is symmetric to the case  $q \geq 1$ , so we need to treat the case  $q \geq 1$  only. We assume also that  $k \geq l+1$ .

In the following, we use the abbreviations

$$h_{m,j}(z) := \prod_{\nu=1}^{k-m} 1/(v+iz) \prod_{\nu=1}^{k-j} 1/(v-iz), \quad z_r := r\pi/t.$$

**THEOREM 2.1.** (i) For  $x \in [1, q]$ ,  $m = 0, 1, \dots, l-1$ , and  $j = 1, \dots, l-1$ ,

$$\varphi_j^{(m)}(x) = \sum_{n \in \mathbf{Z}} \frac{1}{t} e^{(k-m+2n\pi i/t) \log x} h_{m,j}(z_{2n}).$$

(ii) For  $x \in [1, q]$  and  $m, j = 0, 1, \dots, l-1$ ,

$$\psi_j^{(m)}(x) = \sum_{n \in \mathbf{Z}} \frac{1}{itz_{2n+1}} e^{(k-m+(2n+1)\pi i/t) \log x} h_{m,j+1}(z_{2n+1}).$$

*Proof.* Suppose that  $\Omega$  is a simply connected domain in the complex plane, and that  $C$  is a rectifiable Jordan closed curve in  $\Omega$ . If  $f$  is analytic in the domain  $\Omega$ , and if  $x$  is inside the curve  $C$ , then Cauchy's formula holds:

$$(2.3) \quad [x]f = \frac{1}{2\pi i} \int_C \frac{f(z)}{z-x} dz.$$

More generally, we have the following well-known formula for divided differences (see [10, Chapter 1]):

$$(2.4) \quad [x_0, x_1, \dots, x_n]f = \frac{1}{2\pi i} \int_C \frac{f(z)}{(z-x_0) \cdots (z-x_n)} dz,$$

as long as  $x_0, \dots, x_n$  are all inside  $C$ . This result may be generalized to an arbitrary domain (not necessarily simply connected). Let  $\Omega$  be a domain,  $C$  a cycle in  $\Omega$  (see [16, Chapter 10]). Thus  $C$  is a sum of closed paths,

$$C = \gamma_1 + \cdots + \gamma_n.$$

Each path is understood to be a continuous map from the unit circle to  $\Omega$ . The image of a path  $\gamma$  is denoted by  $\gamma^*$ . Correspondingly,

$$C^* := \gamma_1^* + \cdots + \gamma_n^*.$$

For any  $x \notin C^*$ , the index of  $x$  with respect to  $C$  is defined by

$$\text{Ind}_C(x) := \frac{1}{2\pi i} \int_C \frac{dz}{z-x}.$$

Suppose that  $\text{Ind}_C(\zeta) = 0$  for any  $\zeta$  not in  $\Omega$ . If  $f$  is analytic in  $\Omega$ , and if  $\text{Ind}_C(x) = 1$ , then Cauchy's formula (2.3) is still valid. More generally, if

$$\text{Ind}_C(x_j) = 1, \quad j = 0, \dots, n,$$

then (2.4) is still true.

Now we want to use (2.4) to prove Theorem 2.1. Fix  $x \in [1, q]$ . Consider the function  $f$  defined by

$$f(z) := x^z / (q^z - q^k) = e^{z \log x} / (e^{zt} - e^{kt}).$$

Then  $f$  is an analytic function of  $z$  in  $\mathbf{C} \setminus \{k + 2n\pi i/t; n \in \mathbf{Z}\}$ . Let  $\gamma_n$  be the path defined by

$$\gamma_n: e^{i\theta} \mapsto k + 2n\pi i/t + e^{i\theta}/t, \quad 0 \leq \theta \leq 2\pi \quad (n \in \mathbf{Z}),$$

and  $\delta_N$  the path given by

$$\delta_N: e^{i\theta} \mapsto k + ((2N + 1)\pi/t)e^{i\theta}, \quad 0 \leq \theta \leq 2\pi.$$

Let

$$C_N := \delta_N - \sum_{n=-N}^N \gamma_n.$$

Then by what has been proved, we have

$$(2.5) \quad [0, 1, \dots, k-1, k+1, \dots, 2k-j]f = \frac{1}{2\pi i} \int_{C_N} g(z) dz,$$

where

$$g(z) = f(z) \prod_{\substack{0 \leq \nu \leq 2k-j \\ \nu \neq k}} \frac{1}{z-\nu}.$$

We claim

$$(2.6) \quad \lim_{N \rightarrow \infty} \int_{\delta_N} g(z) dz = 0.$$

First, we observe that

$$\left| \prod_{\substack{0 \leq \nu \leq 2k-j \\ \nu \neq k}} \frac{1}{z - \nu} \right| \leq \left( \frac{1}{N - 2k} \right)^k \quad \text{for } z \in \delta_N^*.$$

Next,  $f$  is bounded on  $\delta_N^*$ . This will be proved case by case. Let

$$z = k + e^{i\theta}(2N + 1)\pi/t.$$

Then

$$(z - k)t = e^{i\theta}(2N + 1)\pi = (2N + 1)\pi(\cos \theta + i \sin \theta).$$

Case 1.  $\cos \theta \geq 1/(4N + 2)$ . In this case, since

$$|e^{(z-k)t}| = e^{(2N+1)\pi \cos \theta} \geq 2,$$

one has

$$|e^{(z-k)t} - 1| \geq e^{(2N+1)\pi \cos \theta} / 2.$$

It follows that

$$|f(z)| \leq e^{k \log x + (2N+1)\pi \cos \theta (\log x/t)} / (e^{kt + (2N+1)\pi \cos \theta} / 2) \leq 2,$$

noting that  $\log x/t \leq 1$ .

Case 2.  $\cos \theta \leq -1/(4N + 2)$ . In this case,

$$|e^{(z-k)t} - 1| \geq 1 - e^{-\pi/2},$$

and

$$|e^{z \log x}| \leq e^{kt}.$$

Case 3.  $|\cos \theta| \leq 1/(4N + 2)$ . In this case,

$$|\sin \theta| \geq 1 - |\cos \theta| \geq 1 - 1/(4N + 2),$$

hence

$$\text{Im}((z - k)t) \in [2N\pi + \pi/2, 2N\pi + \pi] \text{ or } [-2N\pi - \pi, -2N\pi - \pi/2].$$

It follows that

$$|e^{(z-k)t} - 1| \geq 1.$$

Moreover,

$$|e^{z \log x}| \leq e^{(k+\pi/2)t}.$$

This finishes the proof of (2.6).

Now let  $N \rightarrow \infty$  in (2.5), and take account of (2.6). We obtain

$$[0, 1, \dots, k - 1, k + 1, \dots, 2k - j]f = -\frac{1}{2\pi i} \sum_{n=-\infty}^{\infty} \int_{\gamma_n} g(z) dz.$$

Each integral  $\int_{\gamma_n} g(z) dz$  can be computed by residue calculus. The function  $g$  has only one pole  $z = k + 2n\pi i/t$  inside  $\gamma_n$ . The residue of  $g$  at this point is

$$\frac{1}{q^{kt}} e^{(k+2n\pi i/t)\log x} \prod_{\substack{0 \leq \nu \leq 2k-j \\ \nu \neq k}} \frac{1}{k + 2n\pi i/t - \nu}.$$



Finally, we obtain

$$\begin{aligned} \varphi_j(x) &= (-1)^{k-j+1} q^k [0, 1, \dots, k-1, k+1, \dots, 2k-j]_z \frac{x^z}{q^z - q^k} \\ &= (-1)^{k-j} \sum_{n \in \mathbf{Z}} \frac{1}{t} e^{(k+2n\pi i/t)\log x} \prod_{\nu=0}^{k-1} \frac{1}{k-\nu+2n\pi i/t} \prod_{\nu=k+1}^{2k-j} \frac{1}{k-\nu+2n\pi i/t} \\ &= \sum_{n \in \mathbf{Z}} \frac{1}{t} e^{(k+2n\pi i/t)\log x} \prod_{\nu=1}^k \frac{1}{\nu+2n\pi i/t} \prod_{\nu=1}^{k-j} \frac{1}{\nu-2n\pi i/t}. \end{aligned}$$

Now (i) comes from this formula by differentiating  $m$  times. The proof of (ii) is similar.  $\square$

As a consequence of Theorem 2.1, the following relations hold:

$$\begin{aligned} \varphi_j^{(m)}(1) &= \sum_{n \in \mathbf{Z}} \frac{1}{t} h_{m,j}(z_{2n}), \\ \psi_j^{(m)}(1) &= \sum_{n \in \mathbf{Z}} \frac{1}{it z_{2n+1}} h_{m,j+1}(z_{2n+1}). \end{aligned}$$

It follows from these two formulae that

$$\begin{aligned} (2.7) \quad \lim_{q \rightarrow \infty} \varphi_j^{(m)}(1) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} h_{m,j}(z) dz, \\ \lim_{q \rightarrow \infty} \psi_j^{(m)}(1) &= \frac{1}{2\pi} \text{pr.v.} \int_{-\infty}^{\infty} h_{m,j+1}(z)/(iz) dz. \end{aligned}$$

(Here, pr.v. means the Cauchy principal value.) Indeed, since  $(1/2\pi) \int_{-\infty}^{\infty} |h_{m,j}(z)| dz < \infty$ , by the very definition of integration we get

$$\begin{aligned} \frac{1}{2\pi} \int_{-\infty}^{\infty} h_{m,j}(z) dz &= \lim_{t \rightarrow \infty} \frac{1}{2\pi} \sum_{n \in \mathbf{Z}} h_{m,j}(z_{2n})(z_{2n} - z_{2n-2}) \\ &= \lim_{t \rightarrow \infty} \sum_{n \in \mathbf{Z}} h_{m,j}(z_{2n})/t = \lim_{q \rightarrow \infty} \varphi_j^{(m)}(1). \end{aligned}$$

The proof of the second relation in (2.7) is similar.

Now set

$$\Phi(q) := \left(\varphi_j^{(m)}(1)\right)_{m,j=0}^{l-1} \quad \text{and} \quad \Psi(q) := \left(\psi_j^{(m)}(1)\right)_{m,j=0}^{l-1}.$$

**THEOREM 2.2.** *The matrices  $\Phi(q)$  and  $\Psi(q)$  are invertible for  $1 < q < \infty$ . Moreover,*

$$\Phi(\infty) := \lim_{q \rightarrow \infty} \Phi(q), \quad \Psi(\infty) := \lim_{q \rightarrow \infty} \Psi(q)$$

*both exist and are invertible.*

*Proof.* We observe that

$$\varphi_0^{(m+1)}(1)/\varphi_0^{(m)}(1) = k - m, \quad h_{m+1,j}(z)/h_{m,j}(z) = k - m + zi/t.$$

Thus we subtract  $(k - m)$  times row  $m$  from row  $m + 1$  in  $\Phi(q)$  ( $m = l - 2, \dots, 0$ ). We obtain

$$\det \Phi = \begin{vmatrix} 1 & \varphi_1 & \dots & \varphi_{l-1} \\ 0 & \varphi'_1 - k\varphi_1 & \dots & \varphi'_{l-1} - k\varphi_{l-1} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & \varphi_1^{(l-1)} - (k-l+2)\varphi_1^{(l-2)} & \dots & \varphi_{l-1}^{(l-1)} - (k-l+2)\varphi_{l-1}^{(l-2)} \end{vmatrix},$$

all functions evaluated at 1. We have

$$\begin{aligned} \varphi_j^{(m+1)}(1) - (k - m)\varphi_j^{(m)}(1) &= \sum_{n \in \mathbf{Z}} [h_{m+1,j}(z_{2n}) - h_{m,j}(z_{2n})]/t \\ &= \sum_{n \in \mathbf{Z}} iz_{2n}h_{m,j}(z_{2n})/t. \end{aligned}$$

We notice that  $\Phi$  is a real matrix. If  $\Phi$  is singular, then there exist *real* numbers  $\beta_1, \dots, \beta_{l-1}$ , not all zero, such that

$$\sum_{j=1}^{l-1} \beta_j \sum_{n \in \mathbf{Z}} iz_{2n}h_{m,j}(z_{2n})/t = 0, \quad m = 1, \dots, l - 1;$$

or

$$(2.8) \quad \sum_{n \in \mathbf{Z}} iz_{2n}h_{0,1}(z_{2n}) \left[ \sum_{j=1}^{l-1} \beta_j \prod_{\nu=k-j+1}^{k-1} (\nu - iz_{2n}) \right] \prod_{\nu=k-m+2}^k (\nu + iz_{2n}) = 0.$$

Let

$$(2.9) \quad \begin{aligned} p(u) &:= \sum_{j=1}^{l-1} \beta_j \prod_{\nu=k-j+1}^{k-1} (\nu + u), \\ g_m(u) &:= \prod_{\nu=k-m+2}^k (\nu + u), \quad m = 1, \dots, l - 1. \end{aligned}$$

Then  $p$  and  $g_m$  are all real polynomials, and (2.8) becomes

$$(2.10) \quad \sum_{n \in \mathbf{Z}} iz_{2n}h_{0,1}(z_{2n})p(-iz_{2n})g_m(iz_{2n}) = 0.$$

We denote by  $\mathbf{P}_n$  the linear space of all polynomials of degree less than  $n$ . Since  $g_m$  has the leading term  $u^{m-1}$  ( $m = 1, \dots, l$ ),  $\{g_1, \dots, g_{l-1}\}$  forms a basis for  $\mathbf{P}_{l-1}$ . Since  $p \in \mathbf{P}_{l-1}$ , we can find *real* numbers  $\alpha_1, \dots, \alpha_{l-1}$  such that

$$(2.11) \quad p = \sum_{m=1}^{l-1} \alpha_m g_m.$$

It follows from (2.10) that

$$\sum_{m=1}^{l-1} \alpha_m \sum_{n \in \mathbf{Z}} iz_{2n}h_{0,1}(z_{2n})p(-iz_{2n})g_m(iz_{2n}) = 0,$$

or, by interchanging the order of summation,

$$(2.12) \quad \sum_{n \in \mathbf{Z}} iz_{2n}h_{0,1}(z_{2n})p(-iz_{2n}) \sum_{m=1}^{l-1} \alpha_m g_m(iz_{2n}) = 0.$$

Now (2.11) tells us that

$$p(-iz_{2n}) \sum_{m=1}^{l-1} \alpha_m g_m(iz_{2n}) = p(-iz_{2n})p(iz_{2n}) = |p(iz_{2n})|^2.$$

Moreover,

$$iz_{2n}h_{0,1}(z_{2n}) = iz_{2n}(k - iz_{2n})h_{0,0}(z_{2n}),$$

and

$$\operatorname{Re}(iz_{2n}(k - iz_{2n})) = |z_{2n}|^2,$$

while

$$h_{0,0}(z_{2n}) = \prod_{\nu=1}^k 1/(\nu^2 + z_{2n}^2) > 0.$$

Thus (2.12) yields

$$\sum_{n \in \mathbf{Z}} |p(iz_{2n})|^2 |z_{2n}|^2 h_{0,0}(z_{2n}) = 0,$$

and, therefore,

$$p(iz_{2n}) = 0 \quad \text{for all } n \in \mathbf{Z}.$$

Since  $p$  is a polynomial, this implies that  $p = 0$ . Recalling (2.9), we conclude that

$$\beta_j = 0 \quad \text{for all } j = 1, \dots, l - 1.$$

This shows that  $\Phi$  is invertible. We can prove in the same fashion, and even more easily, that  $\Psi$  is also invertible.

As to  $\Phi(\infty)$  and  $\Psi(\infty)$ , the series appearing in the definition of  $\varphi_j$  and  $\psi_j$  are replaced by integrals, but the above proof is readily translated to this case. The proof of Theorem 2.2 is complete.

**3. Divided Differences and Derivatives.** For simplicity, we write

$$\varphi_j := \psi_{j-l}, \quad j = l, \dots, 2l - 1.$$

Let  $S$  be the span of  $\varphi_0, \dots, \varphi_{2l-1}$ . Then, by (2.1) and (2.2), we have

$$f(q^2x) = q^{2k}f(x) \quad \text{all } f \in S.$$

Recall that  $(x_j)_{j \in \mathbf{Z}}$  is an  $l$ -multiple geometric mesh. Hence,  $x_{j+2l} = q^2x_j$ .

We use the following abbreviation,

$$\lambda_{j,m} := [x_j, \dots, x_{j+m}],$$

for the  $m$ th divided difference linear functional. According to the standard convention,

$$\lambda_{j,m} = [x_j]D^m \quad \text{in case } x_j = \dots = x_{j+m},$$

where  $D = d/dx$  is differentiation. By standard properties of divided differences,

$$(3.1) \quad \lambda_{j,m+1} = (\lambda_{j+1,m} - \lambda_{j,m}) / (x_{j+1+m} - x_j) \quad \text{if } x_{j+1+m} > x_j,$$

$$(3.2) \quad \lambda_{j+1,m} = \lambda_{j,m} \quad \text{if } x_{j+1+m} = x_j.$$

Now we restrict  $\lambda_{j,m}$  to the space  $S$ . Then one verifies that

$$(3.3) \quad \lambda_{2l,m} = q^{2k-2m}\lambda_{0,m}.$$

Let  $\Lambda_m$  be the cone generated by the  $\lambda_{j,m}$ , i.e.,

$$\Lambda_m := \left\{ \sum_{j=0}^{2l-1} a_j \lambda_{j,m}; a_j \geq 0 \right\}.$$

We claim that

$$(3.4) \quad \Lambda_m \subset \Lambda_{m+1}.$$

Indeed, (3.1) and (3.2) tell us that

$$\lambda_{j+1,m} - \lambda_{j,m} \in \Lambda_{m+1}, \quad j = 0, \dots, 2l - 1.$$

Therefore,

$$\lambda_{2l,m} - \lambda_{0,m} = \sum_{j=0}^{2l-1} (\lambda_{j+1,m} - \lambda_{j,m}) \in \Lambda_{m+1}.$$

From (3.3), we have  $\lambda_{2l,m} = q^{2k-2m}\lambda_{0,m}$ , hence  $(q^{2k-2m} - 1)\lambda_{0,m} \in \Lambda_{m+1}$ . This shows that  $\lambda_{0,m} \in \Lambda_{m+1}$ , and therefore

$$\lambda_{j,m} = \lambda_{0,m} + \sum_{i=0}^{j-1} (\lambda_{i+1,m} - \lambda_{i,m}) \in \Lambda_{m+1}.$$

Thus (3.4) is proved. Set

$$(3.5) \quad \begin{aligned} \mu_i &:= \lambda_{0,i} \\ \mu_{i+l} &:= \lambda_{l,i}/q^{k-i} \end{aligned} \quad \text{for } 0 \leq i < l.$$

Then, as a consequence of (3.4), we have

$$\mu_i \in \Lambda_k \quad \text{for } i = 0, \dots, 2l - 1.$$

We have already proved the first part of the following theorem.

**THEOREM 3.1.** *There exists a nonnegative matrix*

$$C := (c_{ij})_{i,j=0}^{2l-1}$$

such that

$$\mu_i = \sum_{j=0}^{2l-1} c_{ij} \lambda_{j,k}.$$

Moreover,  $C$  is invertible, and  $C$  is bounded independently of  $q$ ,

$$|c_{ij}| \leq \text{const}_k \quad \text{for all } i, j.$$

*Proof.* To prove the second part of this theorem, we let  $\mu_i$  act on the function  $\varphi_0 \in S$ . Then

$$\mu_i \varphi_0 = \sum_j c_{ij} (\lambda_{j,k} \varphi_0).$$

We observe that

$$\lambda_{j,k} \varphi_0 = 1 \quad \text{for all } j$$

while

$$\mu_i \varphi_0 = \mu_{i+l} \varphi_0 = k(k-1) \cdots (k-l+1) \quad \text{for } i = 0, \dots, l-1.$$

Therefore,

$$\sum_j c_{ij} \leq \text{const}_k.$$

Since each  $c_{ij}$  is nonnegative, this implies that

$$c_{ij} \leq \text{const}_k \quad \text{for all } i, j.$$

We observe in the next section (see (4.7)) that  $(\mu_i, \varphi_j)$  is invertible. This shows that  $(\mu_i)$  is linearly independent over  $S$ , hence  $C$  is invertible. With this, the proof of Theorem 3.1 is complete.

**4.  $L$ -Boundedness of  $L_2$ -Projections on Splines for a Multiple Geometric Mesh.**  
We are now in a position to prove the main result of this paper.

**THEOREM 4.1.** *Let  $\mathbf{x} = (x_i)_{-\infty}^{\infty}$  be the mesh given by*

$$x_{li} = x_{li+1} = \dots = x_{li+l-1} = q^i \quad \text{all } i, 0 < q < \infty,$$

where  $l \in \mathbf{N}$  is the multiplicity of the mesh. Let

$$A := \left( \int M_{i,k} N_{j,k} \right)_{i,j \in \mathbf{Z}}.$$

Then

$$\sup_{0 < q < \infty} \|A^{-1}\|_{\infty} \leq \text{const}_k.$$

*Proof.* When  $l = k$ ,  $S_k$  becomes the space of all piecewise polynomials with breakpoints  $x_i$ ,  $i \in \mathbf{Z}$ , so this case is trivial. Thus we may assume  $l < k$ .

We view  $A$  as a  $(2l \times 2l)$ -block Toeplitz matrix. Set

$$v_{p,j} := \sum_{m \in \mathbf{Z}} A(p, j + 2ml), \quad p, j = 0, 1, \dots, 2l - 1,$$

and

$$V := (v_{p,j})_{p,j=0}^{2l-1}.$$

Then  $V$  is just  $\hat{A}(1)$ . If  $\mathbf{b} \in R^{2l-1}$  is a solution to the equation

$$(4.1) \quad V\mathbf{b} = -\mathbf{1}',$$

then, by Theorem 1.1,

$$(4.2) \quad \|A^{-1}\|_{\infty} = \|\mathbf{b}\|_{\infty}.$$

Here is our scheme for the proof of this theorem: As mentioned before, we need only to consider the case  $1 \leq q < \infty$ . First, we show that (4.1) has a solution for any  $q \in (1, \infty)$ . Next, we investigate the uniform boundedness of  $\mathbf{b}$  as  $q \rightarrow \infty$ . Finally, we consider the behavior of  $\mathbf{b}$  near  $q = 1$ .

We need the following lemma.

**LEMMA 4.1.** *If  $f \in S$  satisfies*

$$(4.3) \quad [x_i, \dots, x_{i+k}]f = (-1)^i/k! \quad \text{for } i = 0, \dots, 2l - 1,$$

then (4.1) has a solution  $\mathbf{b}$  satisfying

$$\|\mathbf{b}\|_{\infty} \leq \text{const}_k \|f^{(k)}\|_{\infty}.$$

(Recall that  $S$  is the span of  $\varphi_0, \dots, \varphi_{2l-1}$ .)

*Proof.* By Peano's Theorem,

$$(4.4) \quad \int M_i f^{(k)} = k! [x_i, \dots, x_{i+k}]f$$

(see [2]). Since  $f^{(k)} \in S_k$ , it can be uniquely expanded in a  $B$ -spline series:

$$f^{(k)} = \sum_{j \in \mathbf{Z}} b_j N_{j,k}$$

with coefficient vector  $\tilde{\mathbf{b}} := (b_j)_{j \in \mathbf{Z}}$ . By (4.3) and (4.4),

$$(4.5) \quad A\tilde{\mathbf{b}} = -\mathbf{1}'.$$

We claim that

$$b_{j+2l} = b_j \quad \text{for any } j \in \mathbf{Z}.$$

Since  $f \in S$ ,  $f(q^2x) = q^{2k}f(x)$ , and therefore  $f^{(k)}(q^2x) = f^{(k)}(x)$ . This fact gives

$$\sum b_j N_j(q^2x) = \sum b_j N_j.$$

But  $N_j(q^2x) = N_{j-2l}(x)$ , hence

$$\sum b_j N_{j-2l} = \sum b_j N_j \quad \text{or} \quad \sum b_{j+2l} N_j = \sum b_j N_j.$$

This implies that

$$b_{j+2l} = b_j \quad \text{all } j,$$

by the linear independence of  $\{N_j; j \in \mathbf{Z}\}$  (see [2]). Therefore, with (4.5), the vector  $\mathbf{b} := (b_j)_{j=0}^{2l-1}$  solves (4.1). Moreover,  $\|\mathbf{b}\|_\infty \leq \text{const}_k \|f^{(k)}\|_\infty$  (see [2]). The proof of Lemma 4.1 is complete.  $\square$

Back to the proof of Theorem 4.1. We want to find  $f \in S$  such that (4.3) holds. Since  $S$  is the span of  $\varphi_0, \dots, \varphi_{2l-1}$ , there exists  $\mathbf{a} = (a_0, \dots, a_{2l-1})$  such that  $f = \sum a_j \varphi_j$ . Hence,

$$\mu_i f = \sum (\mu_i \varphi_j) a_j.$$

On the other hand, Theorem 3.1 tells us that

$$\mu_i f = \sum_j c_{ij} \lambda_{j,k} f.$$

Therefore,

$$\sum_j (\mu_i \varphi_j) a_j = \sum_j c_{ij} (\lambda_{j,k} f).$$

Since  $C = (c_{ij})$  is an invertible matrix, (4.3) is equivalent to

$$(4.6) \quad \sum_j (\mu_i \varphi_j) a_j = \sum_j c_{ij} (-1)^j / k!.$$

We have to take a closer look at the matrix  $(\mu_i \varphi_j)$ . By (3.5), we have

$$\mu_i \varphi_j = \begin{cases} \varphi_j^{(i)}(1) & \text{for } 0 \leq i, j < l, \\ \psi_j^{(i)}(1) & \text{for } 0 \leq i < l \leq j < 2l, \\ \varphi_j^{(i-l)}(1) & \text{for } l \leq i < 2l; 0 \leq j < l, \\ -\psi_j^{(i-l)}(1) & \text{for } l \leq i, j < 2l. \end{cases}$$

Therefore, recalling the definition of  $\Phi$  and  $\Psi$ , we obtain

$$(4.7) \quad (\mu_i \varphi_j) = \begin{bmatrix} \Phi & \Psi \\ \Phi & -\Psi \end{bmatrix}.$$

Since  $\Phi$  and  $\Psi$  are invertible, so is the matrix

$$\begin{bmatrix} \Phi & \Psi \\ \Phi & -\Psi \end{bmatrix} = \begin{bmatrix} I & 0 \\ I & I \end{bmatrix} \begin{bmatrix} \Phi & \Psi \\ 0 & -2\Psi \end{bmatrix}.$$

(Here,  $I$  is the  $l \times l$  identity matrix.) Moreover, since  $\Phi(\infty)$  and  $\Psi(\infty)$  are invertible,

$$(4.8) \quad \limsup_{q \rightarrow \infty} \|(\mu_i \varphi_j)^{-1}\|_\infty \leq \text{const}_k.$$

Since the matrix  $(\mu_i \varphi_j)$  is invertible, we can find  $\mathbf{a}$  so that (4.6) holds. Moreover, using Theorem 3.1,

$$(4.9) \quad \|\mathbf{a}\|_\infty \leq \text{const}_k \|(\mu_i \varphi_j)^{-1}\|_\infty.$$

For such  $\mathbf{a}$ , the function  $f := \sum_j a_j \varphi_j$  satisfies (4.3). Thus, by Lemma 4.1, we have already proved that (4.1) has a solution  $\mathbf{b}$  for any  $q \in (1, \infty)$ .

Now we want to prove that

$$(4.10) \quad \limsup_{q \rightarrow \infty} \|f^{(k)}\|_\infty \leq \text{const}_k.$$

By (4.8) and (4.9),

$$\limsup_{q \rightarrow \infty} \|\mathbf{a}\|_\infty \leq \text{const}_k.$$

Hence (4.10) is true, once we show that

$$(4.11) \quad \limsup_{q \rightarrow \infty} \|\varphi_j^{(k)}\|_\infty \leq \text{const}_k.$$

Appealing to Theorem 2.1, we have

$$\|\varphi_j^{(k)}\|_\infty \leq \sum_{n \in \mathbf{Z}} \left[ \prod_{\nu=1}^{k-j} (\nu^2 + 4n^2\pi^2/t^2)^{-1/2} \right] / t \quad \text{for } 1 \leq j \leq l-1.$$

Recall the abbreviation  $t = \log q$ . Since  $t \rightarrow \infty$  when  $q \rightarrow \infty$ , we have, by the very definition of the Riemann integral, that

$$\begin{aligned} \lim_{q \rightarrow \infty} \sum_{n \in \mathbf{Z}} \left[ \prod_{\nu=1}^{k-j} (\nu^2 + 4n^2\pi^2/t^2)^{-1/2} \right] / t \\ = \frac{1}{2\pi} \int_{-\infty}^{\infty} \prod_{\nu=1}^{k-j} (\nu^2 + z^2)^{-1/2} dz \leq \text{const}_k. \end{aligned}$$

This proves (4.11) for  $j = 1, \dots, l-1$ . For  $l \leq j \leq 2l-1$ , the proof is similar. The case  $j = 0$  is trivial. With this, it follows from (4.2), (4.4), and (4.10) that

$$(4.12) \quad \limsup_{q \rightarrow \infty} \|A^{-1}\|_\infty \leq \text{const}_k.$$

Finally, we consider the behavior of  $\mathbf{b}$  near  $q = 1$ . It is known (see [1]) that, for  $q = 1$ ,  $A^{-1}$  is bounded. Therefore,  $V$  is invertible for  $q = 1$ . Since each entry of  $V$  is a continuous function of  $q$ ,  $\mathbf{b}$  is continuous in  $q$  near  $q = 1$ . This shows that

$$\lim_{q \rightarrow 1} \|\mathbf{b}\|_\infty \leq \text{const}_k.$$

Invoking (4.2) once again, we obtain

$$\lim_{q \rightarrow 1} \|A^{-1}\|_\infty \leq \text{const}_k.$$

The proof of Theorem 4.1 is now complete.

**Acknowledgment.** The author wishes to thank his adviser, Professor Carl de Boor, for his valuable guidance.

Mathematics Research Center  
University of Wisconsin  
Madison, Wisconsin 53705

1. C. DE BOOR, "The quasi-interpolant as a tool in elementary polynomial spline theory," in *Approximation Theory* (G. G. Lorentz, ed.), Academic Press, New York, 1973, pp. 269–276.
2. C. DE BOOR, "Splines as linear combinations of  $B$ -splines. A survey," in *Approximation Theory II* (G. G. Lorentz, ed.), Academic Press, New York, 1976, pp. 1–47.
3. C. DE BOOR, "Total positivity of the spline collocation matrix," *Indiana Univ. Math. J.*, v. 25, 1976, pp. 541–551.
4. C. DE BOOR, "A bound on the  $L_\infty$ -norm of the  $L_2$ -approximation by splines in terms of a global mesh ratio," *Math. Comp.*, v. 30, 1976, pp. 767–771.
5. C. DE BOOR, "On a max norm bound for the least-squares spline approximation," in *Approximation and Function Spaces* (Z. Ciesielski, ed.), North-Holland, New York, 1981, pp. 163–175.
6. C. DE BOOR, S. FRIEDLAND & A. PINKUS, "Inverses of infinite sign regular matrices," *Trans. Amer. Math. Soc.*, v. 274, 1982, pp. 59–68.
7. A. S. CAVARETTA, W. DAHMEN, C. A. MICCHELLI & P. W. SMITH, "On the solvability of certain systems of linear difference equations," *SIAM J. Math. Anal.*, v. 12, 1981, pp. 833–841.
8. Z. CIESIELSKI, "Properties of the orthonormal Franklin system," *Studia Math.*, v. 23, 1963, pp. 141–157.
9. Y. Y. FENG & J. KOZAK, "On the generalized Euler-Frobenius polynomial," *J. Approx. Theory*, v. 32, 1981, pp. 327–338.
10. A. O. GELFAND, *Calculus of Finite Differences*, Authorized English translation of the 3rd Russian ed., Delhi, Hindustan Publishing Corp., 1971.
11. I. I. HIRSCHMAN, "Matrix-valued Toeplitz operators," *Duke Math. J.*, v. 34, 1967, pp. 403–415.
12. K. HÖLLIG, " $L_\infty$ -boundedness of  $L_2$ -projections on splines for a geometric mesh," *J. Approx. Theory*, v. 33, 1981, pp. 318–333.
13. S. KARLIN, *Total Positivity*, Vol. 1, Stanford University Press, Stanford, California, 1968.
14. C. A. MICCHELLI, "Cardinal  $L$ -splines," in *Studies in Spline Functions and Approximation Theory*, Academic Press, New York, 1976, pp. 203–250.
15. B. MITYAGIN, "Quadratic pencils and least-squares piecewise polynomial approximation," *Math. Comp.*, v. 40, 1983, pp. 283–300.
16. W. RUDIN, *Real and Complex Analysis*, McGraw-Hill, New York, 1974.
17. I. J. SCHOENBERG, *Cardinal Spline Interpolation*, Regional Conference in Applied Mathematics No. 12, SIAM, Philadelphia, Pa., 1973.