

## SOME ZEROS OF THE TITCHMARSH COUNTEREXAMPLE

ROBERT SPIRA

**ABSTRACT.** Zeros on and off the critical line are found for Titchmarsh's function  $f(s)$ .

Let  $s = \sigma + it$ . E. C. Titchmarsh [1, pp. 240-244] introduced the function

$$\begin{aligned} f(s) &= \frac{1}{2} \sec \theta \{e^{-i\theta} L_1(s) + e^{i\theta} L_2(s)\} \\ &= \frac{1}{1^s} + \frac{\tan \theta}{2^s} - \frac{\tan \theta}{3^s} - \frac{1}{4^s} + \frac{1}{6^s} + \cdots \\ &= 5^{-s} \{\zeta(s, 1/5) + \tan \theta \zeta(s, 2/5) - \tan \theta \zeta(s, 3/5) - \zeta(s, 4/5)\}, \end{aligned}$$

where

$$\tan \theta = \frac{\sqrt{10 - 2\sqrt{5}} - 2}{\sqrt{5} - 1} = .28407\ 90438\ 40412\ 296\dots$$

and  $L_1(s) = \sum_{n=1}^{\infty} \chi_1(n)n^{-s}$ ,  $L_2(s) = \sum_{n=1}^{\infty} \chi_2(n)n^{-s}$  are Dirichlet  $L$ -functions mod 5 with  $\chi_1$  and  $\chi_2$  the Dirichlet characters determined by  $\chi_1(2) = i$  and  $\chi_2(2) = -i$ .

Titchmarsh showed that though  $f(s)$  satisfies a functional equation identical to that of a Dirichlet  $L$ -function:

$$f(s) = 5^{1/2-s} 2(2\pi)^{s-1} \Gamma(1-s) \cos(\frac{1}{2}s\pi) f(1-s),$$

it has zeros with  $\sigma > 1$  (together with infinitely many zeros on the line  $\sigma = \frac{1}{2}$ ). According to a theorem of Voronin [2],  $f(s)$  has zeros in the critical strip off the critical line. Titchmarsh gave the equation  $\sin 2\theta = 2 \cos(2\pi/5)$ , but  $\sin 2\theta$  should be  $\tan 2\theta$ . This minor error was carried over to Voronin [2] and the review MR 86g:11048 in *Mathematical Reviews*.

With the help of programs for computing  $L$  and  $L'$  (Spira [3]), an exploratory computation of  $f(s)$  in the critical strip for  $0 \leq t \leq 200$  revealed the following zeros off the critical line:

$$\begin{aligned} &.808517 + 85.699348i \\ &.650830 + 114.163343i \\ &.574356 + 166.479306i \\ &.724258 + 176.702461i \end{aligned}$$

---

Received by the editor January 13, 1993 and, in revised form, December 7, 1993.

1991 *Mathematics Subject Classification*. Primary 11M26.

*Key words and phrases*. Riemann zeta function.

The first few zeros on the line have  $t$ -coordinates: 5.094160, 8.939914, 12.133545, 14.404003, 17.130239, 19.308800, 22.159708, 23.345370, 26.094967, 27.923799, 30.159418, 31.964500, 33.699862, 35.890855, 37.455462, 40.162578, 40.682953, 43.081265, 44.947134, 46.456355, 48.477787, 50.240086.

It was also found that  $f(0) = .6568158$ ,  $f(\frac{1}{2}) = .8253830$ . The program reproduced check values of an  $L$ -function mod 5 and its derivative, and then new values for the character were inserted simply and easily to calculate  $f(s)$ . A further check was to calculate the zeros off the line reflected in  $\sigma = \frac{1}{2}$ .

By Rolle's Theorem for a zero  $\sigma_0 + it_0$  off the line, there is a  $\sigma_1$  between  $\sigma_0$  and  $1 - \sigma_0$  such that  $|f(\sigma_1 + it_0)|$  is a maximum, or

$$(\operatorname{Re} f \cdot \operatorname{Re} f' + \operatorname{Im} f \cdot \operatorname{Im} f')(\sigma_1 + it_0) = 0.$$

For the first zero at least,  $\sigma_1 < \frac{1}{2}$  and  $f'(\sigma_1 + it_0) \neq 0$ , so the vectors  $(\operatorname{Re} f, \operatorname{Im} f)$  and  $(\operatorname{Re} f', \operatorname{Im} f')$  are orthogonal. In Spira [4] it was conjectured that for  $|t| > 6.3$ ,  $(\operatorname{Re} \zeta \cdot \operatorname{Re} \zeta' + \operatorname{Im} \zeta \cdot \operatorname{Im} \zeta') < 0$  in the left half of the critical strip, which is stronger than the Riemann hypothesis.

No zeros of  $f'(s)$  were found with  $\sigma < \frac{1}{2}$  for  $t \leq 200$ , nor any zeros of  $f(s)$  with  $\sigma > 1$  for  $t \leq 200$ , though this last is not unusual since Titchmarsh's proof relies on methods which ordinarily require a very large  $t$ . If one multiplies  $\zeta(s)$  by the four linear factors  $(s - \frac{1}{2} \pm \frac{1}{4} \pm i)$  one obtains a function with a functional equation which is not zero for  $\sigma \geq 1$ , but vanishes off  $\sigma = \frac{1}{2}$ .

Karatsuba and Voronin [5, Chapter VI, §5, pp. 212–240] is devoted to a study of zeros of  $f(s)$  in the critical strip and on  $\sigma = \frac{1}{2}$ .

#### BIBLIOGRAPHY

1. E. C. Titchmarsh, *The theory of the Riemann zeta function*, Oxford Univ. Press, London, 1951.
2. S. M. Voronin, *On the distribution of zeros of some Dirichlet series*, Proc. Steklov Inst. Math. (1985), no. 4, 89–92.
3. R. Spira, *Calculation of Dirichlet  $L$ -functions*, Math. Comp. **23** (1969), 489–498.
4. ———, *On the Riemann zeta function*, J. London Math. Soc. **44** (1969), 325–328.
5. A. A. Karatsuba and S. M. Voronin, *The Riemann zeta-function*, de Gruyter, Berlin, 1992.