

## RANDOM FIBONACCI SEQUENCES AND THE NUMBER 1.13198824...

DIVAKAR VISWANATH

ABSTRACT. For the familiar Fibonacci sequence (defined by  $f_1 = f_2 = 1$ , and  $f_n = f_{n-1} + f_{n-2}$  for  $n > 2$ ),  $f_n$  increases exponentially with  $n$  at a rate given by the golden ratio  $(1 + \sqrt{5})/2 = 1.61803398\dots$ . But for a simple modification with both additions and subtractions — the *random* Fibonacci sequences defined by  $t_1 = t_2 = 1$ , and for  $n > 2$ ,  $t_n = \pm t_{n-1} \pm t_{n-2}$ , where each  $\pm$  sign is independent and either  $+$  or  $-$  with probability  $1/2$  — it is not even obvious if  $|t_n|$  should increase with  $n$ . Our main result is that

$$\sqrt[n]{|t_n|} \rightarrow 1.13198824\dots \text{ as } n \rightarrow \infty$$

with probability 1. Finding the number 1.13198824... involves the theory of random matrix products, Stern-Brocot division of the real line, a fractal measure, a computer calculation, and a rounding error analysis to validate the computer calculation.

### 1. INTRODUCTION

The Fibonacci numbers defined by  $f_1 = f_2 = 1$  and  $f_n = f_{n-1} + f_{n-2}$  for  $n > 2$  are widely known. It is equally well-known that  $|f_n|$  increases exponentially with  $n$  at the rate  $(1 + \sqrt{5})/2$ . Consider random Fibonacci sequences defined by the random recurrence  $t_1 = 1$ ,  $t_2 = 1$ , and for  $n > 2$ ,  $t_n = \pm t_{n-1} \pm t_{n-2}$ , where each  $\pm$  sign is independent and either  $+$  or  $-$  with probability  $1/2$ . Do the random Fibonacci sequences level off because of the subtractions? Or do the random Fibonacci sequences increase exponentially with  $n$  like the Fibonacci sequence? If so, at what rate? The answer to these questions brings Stern-Brocot sequences, a beautiful way to divide the real number line that was first discovered in the 19th century, and fractals and random matrix products into play. The final answer is obtained from a computer calculation, raising questions about computer assisted theorems and proofs.

Below are three possible runs of the random Fibonacci recurrence:

1, 1, -2, -3, -1, 4, -3, 7, -4, 11, -15, 4, -19, 23, -4, ...

1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 134, 223, 357, 580, ...

1, 1, -2, 1, 1, -2, 1, 1, -2, 1, 1, -2, 1, 1, -2, ...

The first of the runs above was randomly generated on a computer. The second run is the familiar Fibonacci sequence. The last of the three runs above is a sequence

---

Received by the editor November 17, 1997 and, in revised form, July 14, 1998.

1991 *Mathematics Subject Classification*. Primary 15A52, 28A80, 60J05, 65G05.

This work was supported in part by NSF Grant DMS-9500975CS and DOE Grant DE-FG02-94ER25199 to Lloyd N. Trefethen.




FIGURE 1. (a) A semilog plot of  $|t_n|$  vs.  $n$  for a computer generated random Fibonacci sequence  $t_n$  showing a clear exponential trend. The dashed line is  $1.132^n$ . (b) Plot of  $\sqrt[n]{|t_n|}$  vs.  $n$ . As  $n$  increases to a million,  $\sqrt[n]{|t_n|}$  seems to settle down to a constant close to 1.132.

that remains bounded as  $n \rightarrow \infty$ ; but such runs with no exponential growth occur with probability 0. For longer, typical runs see Figure 1. Numerical experiments in Figure 1 illustrate our main result (Theorem 4.2), that

$$\sqrt[n]{|t_n|} \rightarrow 1.13198824\dots \text{ as } n \rightarrow \infty$$

with probability 1. Thus 1.13198824... gives the exponential rate of increase of  $|t_n|$  with  $n$  for random Fibonacci sequences, just as the golden ratio  $(1 + \sqrt{5})/2$  gives the exponential rate of increase of the Fibonacci numbers.

For the random Fibonacci recurrence  $t_n = \pm t_{n-1} \pm t_{n-2}$  as well as the recurrence  $t_n = \pm t_{n-1} + t_{n-2}$  with each  $\pm$  independent and  $+$  or  $-$  with probability  $1/2$ ,  $|t_n|$  is either  $|t_{n-1}| + |t_{n-2}|$  or  $||t_{n-1}| - |t_{n-2}||$  with probability  $1/2$ . As our interest is in  $|t_n|$  vs.  $n$  as  $n \rightarrow \infty$ , we restrict focus to  $t_n = \pm t_{n-1} + t_{n-2}$  and call it the random Fibonacci recurrence. As a result, the presentation becomes briefer, especially in Section 3.

The next step is to rewrite the random Fibonacci recurrence using matrices. In matrix form the random Fibonacci recurrence is  $\begin{pmatrix} t_{n-1} \\ t_n \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & \pm 1 \end{pmatrix} \begin{pmatrix} t_{n-2} \\ t_{n-1} \end{pmatrix}$ , with one of the two matrices

$$(1.1) \quad A = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 1 \\ 1 & -1 \end{pmatrix}$$

picked independently with probability  $1/2$  at each step. Let  $\mu_f$  denote the distribution that picks  $A$  or  $B$  with probability  $1/2$ . Then the random matrix  $M_n$  chosen at the  $n$ th step is  $\mu_f$ -distributed and independent of  $M_i$  for  $i \neq n$ . Moreover,

$$\begin{pmatrix} t_{n-1} \\ t_n \end{pmatrix} = M_{n-2} \dots M_1 \begin{pmatrix} 1 \\ 1 \end{pmatrix},$$

where  $M_{n-2} \dots M_1$  is a product of independent, identically distributed random matrices.

Known results from the theory of random matrix products imply that

$$(1.2) \quad \frac{\log \|M_n \dots M_1\|}{n} \rightarrow \gamma_f \text{ as } n \rightarrow \infty,$$

$$(1.3) \quad \sqrt[n]{|t_n|} \rightarrow e^{\gamma_f} \text{ as } n \rightarrow \infty,$$

for a constant  $\gamma_f$  with probability 1 [7, p. 11, p. 157]. About  $\gamma_f$  itself known theory can only say that  $\gamma_f > 0$  [7, p. 30]. Our aim is to determine  $\gamma_f$  or  $e^{\gamma_f}$  exactly. Theorem 4.2 realizes this aim by showing that  $e^{\gamma_f} = 1.13198824\dots$ . The limit in (1.3) is the same  $\gamma_f$  for any norm over 2-dimensional matrices, because all norms over a finite dimensional vector space are equivalent. In the rest of this paper, all norms are 2-norms, and all matrices and vectors are real and 2-dimensional except when stated otherwise. Thus, for a vector  $x$ ,  $\|x\|$  is its Euclidean length in the real plane, and for a matrix  $M$ ,  $\|M\| = \sup_{x \neq 0} \frac{\|Mx\|}{\|x\|}$ .

The limit (1.2) for  $M_i$  independent but identically distributed over  $d$ -dimensional matrices has been a central concern of the theory of random matrix products. Furstenberg and Kesten [19, 1960] have shown that the limit (1.2) exists under very general conditions. When it exists, that limit is usually denoted by  $\gamma$  and called the upper Lyapunov exponent. Furstenberg [18, 1963] has shown that when the normalizing condition  $|\det M_i| = 1$  holds, as it does for  $\mu_f$ , “usually”  $\gamma > 0$ . Furstenberg’s theorem implies, for example, that  $\gamma_f > 0$ , and hence, that  $|t_n|$  increases exponentially with  $n$  with probability 1.

In spite of the importance of the upper Lyapunov exponent  $\gamma$ ,  $\gamma$  is known exactly for very few examples. Kingman, one of the pioneers of subadditive ergodic theory, of which the theory of random matrix products is a special case, wrote [26, 1973]:

Pride of place among the unsolved problems of subadditive ergodic theory must go to the calculation of the constant  $\gamma$  ( . . . ). In none of the applications described here is there an obvious mechanism for obtaining an exact numerical value, and indeed this usually seems to be a problem of some depth.

One of the applications Kingman refers to is the general problem of finding  $\gamma$  for random matrix products. For this and other applications, Kingman’s problem is still unsolved. Bougerol [7, p. 33] and Lima and Rahibe [31] calculate  $\gamma$  for some examples. The work of Chassaing, Letac and Mora [11] is closer to our determination of  $\gamma_f$ . But in all their examples, matrices, unlike  $B$  in (1.1), have only non-negative entries. In our opinion, the random Fibonacci recurrence is more natural than these examples. In fact, the random Fibonacci recurrence in a more general form appears as a motivating example in the very first paragraph of Furstenberg’s famous paper [18].

In Section 2, we present a formula for  $\gamma_f$  due to Furstenberg that forms the basis for this paper. The matrices  $A$  and  $B$  map a direction in the real plane of slope  $m$  to directions of slope  $1 + 1/m$  and  $-1 + 1/m$ , respectively. Since  $\mu_f$  picks  $A$  or  $B$  with probability 1/2, it induces the random walk which sends a direction of slope  $m$  to a direction of slope  $1 + 1/m$  or  $-1 + 1/m$  with probability 1/2. The invariant probability measure for this random walk is central to Furstenberg’s formula.

In Section 3, we find that invariant probability measure, denoted by  $\nu_f$ , using the Stern-Brocot division of the real line. See Figures 3, 4. The measure  $\nu_f$  gives a probability measure over the real line  $R$  because the slope  $m$  can be any real number. Since the backward maps for the random walk are  $m \rightarrow 1/(\pm 1 + m)$ , the

invariance condition requires

$$\nu_f([a, b]) = \frac{1}{2}\nu_f\left(\left[\frac{1}{-1+b}, \frac{1}{-1+a}\right]\right) + \frac{1}{2}\nu_f\left(\left[\frac{1}{1+b}, \frac{1}{1+a}\right]\right),$$

for any interval  $[a, b]$  with  $\pm 1 \notin (a, b)$ . Since the slopes of the backward maps vary in magnitude from 0 to  $\infty$ , not only is  $\nu_f$  self-similar [37], the self-similarity equation has multiple scales. Self-similar functions, especially ones with multiple scales, usually turn out to be fractals. For example, Weierstrass’s nowhere-differentiable but continuous functions, which are commonly used examples of fractal graphs, satisfy  $f(x) = \lambda^{s-2} \sin(\lambda t) + \lambda f(\lambda t)$  with  $1 < s < 2$ ,  $\lambda > 1$ , and  $\lambda$  large enough [17]. Repetition of the same structure at finer scales and an irregular appearance in Figure 4 suggest that  $\nu_f$  too may be a type of fractal.

In Section 4, we use Furstenberg’s formula and the invariant measure  $\nu_f$  given in Section 3, and arrive at Theorem 4.2 ( $e^{\gamma_f} = 1.13198824\dots$ ). The proof of Theorem 4.2 depends on a computer calculation. Thus its correctness depends not only upon mathematical arguments that can be checked line by line, but upon a program that can also be checked line by line and the correct implementation of various software and hardware components of the computer system. The most famous of theorems whose proofs depend on computer calculations is the four color theorem. The first proof of the four color theorem (all planar graphs can be colored using only four colors so that no two adjacent vertices have the same color) by Appel, Haken and Koch caused controversy and aroused great interest because it relied on producing and checking 1834 graphs using 1200 hours of 1976 computer time [28], [2]. In spite of improvements (for example, the number 1834 was brought down to 1482 soon afterwards by Appel and Haken themselves), all proofs of the four color theorem still rely on the computer.

Computer assisted proofs are more common now. Our computation uses floating point arithmetic, which is inexact owing to rounding errors. Thus it becomes necessary to bound the effect of the rounding errors, which we do in the appendix. An early example of rigorous use of floating point arithmetic is due to Brent [9]. Lanford’s proof of Feigenbaum’s conjecture about the period doubling route to chaos used interval arithmetic [29]. The computer assisted proof of chaos in the Lorenz equations announced by Mischaikow and Mrozek [32] [33] is another notable example. We will discuss the use of floating point arithmetic and other issues related to our Theorem 4.2 in Section 4.

Besides random matrix products, random Fibonacci sequences are connected to many areas of mathematics. For example, the invariant measure  $\nu_f$  is also the distribution of the continued fractions

$$\pm 1 + \frac{1}{\pm 1 + \frac{1}{\pm 1 + \dots}}$$

with each  $\pm 1$  independent and either  $+1$  or  $-1$  with probability  $1/2$ . The matrices  $A$  and  $B$  in (1.1) can both be thought of as Möbius transformations of the complex plane; then the random matrix product and the exponential growth of  $|t_n|$  in (1.2) and (1.3) would correspond to the dynamics of complex numbers acted upon by a composition of the Möbius transformations  $A$  and  $B$  [7, p. 38]. Also, the random walk on slopes  $m \rightarrow \pm 1 + 1/m$  can be thought of as a random dynamical system [3]. These different interpretations amount merely to a change of vocabulary as far as

the computation of  $\gamma_f$  is concerned; but each interpretation offers a different point of view.

The study of random matrix products, initiated by Bellman [4, 1954], has led to many deep results and applications. Applications have been made to areas as diverse as Schrödinger operators, image generation, and demography [14], [15], [40]. Furstenberg and Kesten [19, 1960], Furstenberg [18, 1963], Oseledec [34, 1968], Kingman [26, 1973], and Guivarc'h and Raugi [21, 1985] are some of the profound contributions to this area. We enthusiastically recommend the lucid, elegant and well-organized account by Bougerol [7]. For a more modern treatment, see [5]. For the basics of probability, our favorite is Breiman [8].

Our interest in random recurrences was aroused by their connection to random triangular matrices [42]. The asymptotic behaviour as  $n \rightarrow \infty$  of the condition number of a triangular matrix of dimension  $n$  whose entries are independent, identically distributed random variables can be deduced from the asymptotic behaviour of a random recurrence. In particular, let  $L_n$  be a lower triangular matrix of dimension  $n$  whose diagonal entries are all 1 and whose subdiagonal entries are  $+1$  or  $-1$  with probability  $1/2$ . Consider the random recurrence

$$\begin{aligned} r_1 &= 1 \\ r_2 &= \pm r_1 \\ r_3 &= \pm r_1 \pm r_2 \\ &\vdots \\ r_n &= \pm r_1 \pm r_2 \pm \cdots \pm r_{n-1}, \end{aligned}$$

where each  $\pm$  is independent, and  $+$  or  $-$  with probability  $1/2$ . Unlike the random Fibonacci recurrence, this recurrence has infinite memory. The limits

$$\lim_{n \rightarrow \infty} \sqrt[n]{\|L_n\|_2 \|L_n^{-1}\|_2} \quad \text{and} \quad \lim_{n \rightarrow \infty} \sqrt[n]{|r_n|}$$

are equal if either of the limits is a constant almost surely. Unable to find these limits, we considered random Fibonacci sequences as a simplification. But the limit

$$\lim_{n \rightarrow \infty} \sqrt[n]{\|L_n\|_2 \|L_n^{-1}\|_2}$$

is determined when entries of  $L_n$  are drawn from various other distributions, including normal and Cauchy distributions, in [42]. For a conjecture about random recurrences along the lines of Furstenberg's theorem, see [41].

## 2. FURSTENBERG'S FORMULA

To determine  $\gamma_f$ , we use a formula from the theory of random matrix products that complements (1.2). Three things that will be defined below — the notation  $\bar{x}$  for directions in the real plane  $R^2$ ,  $\text{amp}(\bar{x})$ , which is a smooth function of  $\bar{x}$  (the diagram just after (2.2)), and  $\nu_f(\bar{x})$  which is a probability measure over directions  $\bar{x}$  (Figure 4) — combine to give a formula for  $\gamma_f$ :

$$(2.1) \quad \gamma_f = \int \text{amp}(\bar{x}) d\nu_f(\bar{x}).$$

This formula, derived by Furstenberg [7, p. 77], is the basis of our determination of  $\gamma_f$ .




FIGURE 2. By (2.2),  $\nu_f(I) = \nu_f(I_1)/2 + \nu_f(I_2)/2$ .

Directions  $\bar{x}$  can be parameterized using angles,  $\bar{x} = \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}$  with  $\theta \in (-\pi/2, \pi/2]$ , or using slopes,  $\bar{x} = \begin{pmatrix} 1 \\ m \end{pmatrix}$  with  $m \in (-\infty, \infty]$ . Slopes  $m$  and angles  $\theta$  are related by  $m = \tan \theta$  and  $\theta = \arctan m$ . We use slopes in all places except Figure 4. In our notation,  $x$  is a vector in the direction  $\bar{x}$ , and  $\bar{x}$  is the direction of the vector  $x$  for  $x \neq 0$ .

To define  $\nu_f$ , consider the  $\mu_f$ -induced random walk on directions that sends  $\bar{x}_0$  to  $\bar{x}_1 = \overline{Ax}_0$  or to  $\bar{x}_1 = \overline{Bx}_0$  with probability  $1/2$ , and then sends  $\bar{x}_1$  to  $\bar{x}_2$  similarly, and so on. In terms of slopes, the slope  $m$  is mapped by the random walk to  $1 + 1/m$  or to  $-1 + 1/m$  with probability  $1/2$ . The measure  $\nu_f$  is the unique invariant probability measure over  $\bar{x}$  for this random walk, i.e.,

$$\nu_f(S) = \frac{1}{2}\nu_f(\overline{A^{-1}S}) + \frac{1}{2}\nu_f(\overline{B^{-1}S}),$$

where  $S$  is any Borel measurable set of directions. We also say that  $\nu_f$  is  $\mu_f$ -invariant. For the existence and uniqueness of  $\nu_f$ , see [7, p. 10, p. 32]. It is also known that  $\nu_f$  must be continuous [7, p. 32], i.e.,  $\nu_f(\{\bar{x}\}) = 0$  for any fixed direction  $\bar{x}$ .

Since the bijections  $\bar{x} \rightarrow \overline{A^{-1}x}$  and  $\bar{x} \rightarrow \overline{B^{-1}x}$  (sometimes called backward maps) map the slope  $m$  to  $1/(-1+m)$  and to  $1/(1+m)$ , respectively, the condition for  $\mu_f$ -invariance in terms of slopes is

$$(2.2) \quad \nu_f([a, b]) = \frac{1}{2}\nu_f\left(\left[\frac{1}{-1+b}, \frac{1}{-1+a}\right]\right) + \frac{1}{2}\nu_f\left(\left[\frac{1}{1+b}, \frac{1}{1+a}\right]\right),$$

where  $[a, b]$  is any interval in the real line with  $\pm 1 \notin (a, b)$ . See Figure 2.


The function  $\text{amp}(\bar{x})$  defined by

$$\text{amp}(\bar{x}) = \frac{1}{2} \log \frac{\|Ax\|}{\|x\|} + \frac{1}{2} \log \frac{\|Bx\|}{\|x\|}$$

gives the average amplification in the direction  $\bar{x}$  when  $x$  is multiplied by  $A$  or  $B$  with probability  $1/2$ . Recall that  $\|\cdot\|$  was taken to be the 2-norm. In terms of slopes,

$$\begin{aligned} \text{amp}(m) &= \frac{1}{4} \log \left( \frac{m^2 + (-1+m)^2}{1+m^2} \right) + \frac{1}{4} \log \left( \frac{m^2 + (1+m)^2}{1+m^2} \right) \\ &= \frac{1}{4} \log \left( \frac{1+4m^4}{(1+m^2)^2} \right). \end{aligned}$$

The figure below plots  $\text{amp}(m)$  vs  $m$ .



Furstenberg's formula (2.1) can now be put in a concrete form using slopes to parameterize directions  $\bar{x}$ :

$$(2.3) \quad \gamma_f = \int_{-\infty}^{\infty} \text{amp}(m) d\nu_f(m) = \frac{1}{4} \int_{-\infty}^{\infty} \log \left( \frac{1+4m^4}{(1+m^2)^2} \right) d\nu_f(m).$$

If we were to use a norm other than the 2-norm for vectors in the real plane,  $\text{amp}(m)$  and  $\text{amp}(\bar{x})$  would be different functions. But Furstenberg's formula (2.1) holds for any norm, even though the measure  $\nu_f$  is independent of the norm. Our choice of the 2-norm is one of many equally suitable alternatives. For the weighted 2-norm

$$\left\| \begin{pmatrix} a \\ b \end{pmatrix} \right\| = \sqrt{a^2 + \frac{1+\sqrt{5}}{2}b^2},$$

$\text{amp}(m) > 0$  for all  $m$  except  $m = \pm\sqrt{(\sqrt{5}-1)/2}$ , and  $\text{amp}(m) = 0$  at those two points.

To illustrate how (2.3) is used, we verify quickly that  $\gamma_f > 0$ . The invariance condition (2.2) applied to the set  $[-\infty, -1] \cup [1, \infty]$  implies  $\nu_f(|m| \geq 1) \geq 1/2$ , because the image of  $[1, \infty]$  under  $m \rightarrow 1/(-1+m)$  and the image of  $[-\infty, -1]$

under  $m \rightarrow 1/(1+m)$  are  $[0, \infty]$  and  $[-\infty, 0]$ , respectively. Now,

$$\begin{aligned} \gamma_f &= \int_{-\infty}^{\infty} \text{amp}(m) \, d\nu_f(m) \\ &> \min_{|m| < 1} \text{amp}(m) \, \nu_f(|m| < 1) + \min_{|m| \geq 1} \text{amp}(m) \, \nu_f(|m| \geq 1) \\ &= -\frac{1}{4} \log\left(\frac{5}{4}\right) \nu_f(|m| < 1) + \frac{1}{4} \log\left(\frac{5}{4}\right) \nu_f(|m| \geq 1) \\ &\geq 0. \end{aligned}$$

The first inequality above is strict, because  $\nu_f$  must be continuous and  $\text{amp}(m)$  is not a constant function. Minimizing  $\text{amp}(m)$  over  $|m| < 1$  and  $|m| \geq 1$  is basic calculus: the minima occur at the points  $m = \pm 1/2$  and  $m = \pm 1$ . The final  $\geq$  is by  $\nu_f(|m| \geq 1) \geq 1/2$ . Actually, it will be shown in Section 3 that  $\nu_f(|m| \geq 1) = (\sqrt{5} - 1)/2$ .

### 3. THE STERN-BROCOT TREE AND CONSTRUCTION OF THE INVARIANT MEASURE $\nu_f$

Assuming  $\pm 1 \notin (a, b)$  as before, we write down the invariance condition once more for easy reference:

$$(3.1) \quad \nu_f([a, b]) = \frac{1}{2} \nu_f\left(\left[\frac{1}{-1+b}, \frac{1}{-1+a}\right]\right) + \frac{1}{2} \nu_f\left(\left[\frac{1}{1+b}, \frac{1}{1+a}\right]\right).$$

Our goal in this section is to find  $\nu_f$ , the unique probability measure on the real line  $R$  satisfying (3.1) for all intervals  $[a, b]$  not containing  $\pm 1$ . Since  $\nu_f$  must be continuous, it does not matter whether we take the intervals in (3.1) to be open or closed or half-closed.

The construction of  $\nu_f$  is based on the Stern-Brocot tree shown in Figure 3. The Stern-Brocot tree is an infinite binary tree that divides  $R$  recursively. Represent  $\infty$




FIGURE 3. The Stern-Brocot tree; its nodes are intervals of the real line  $R$ . The division of any interval  $[a/b, c/d]$ , except the root, into two children is done by inserting the point  $\frac{a+c}{b+d}$ .

as  $\frac{1}{0}$  and 0 as  $\frac{0}{1}$ , and write negative fractions with the numerator negative. Then the root of the Stern-Brocot tree is the real line  $[\frac{-1}{0}, \frac{1}{0}]$ . Its left and right children are  $[\frac{-1}{0}, \frac{0}{1}]$  and  $[\frac{0}{1}, \frac{1}{0}]$ , the positive and negative halves of  $R$ . The rest of the tree is defined by dividing any node  $[\frac{a}{b}, \frac{c}{d}]$  other than the root into a left child  $[\frac{a}{b}, \frac{a+c}{b+d}]$  and a right child  $[\frac{a+c}{b+d}, \frac{c}{d}]$ . For example, the root's left child  $[\frac{-1}{0}, \frac{0}{1}]$  divides into  $[\frac{-1}{0}, \frac{-1}{1}]$  and  $[\frac{-1}{1}, \frac{0}{1}]$ .

The Stern-Brocot tree was discovered and reported independently by the mathematician Moriz Stern in 1858 [36] and by the watchmaker Achille Brocot in 1860 [10]. Unaware of its existence, we found it again while trying to construct  $\nu_f$ . We summarize some basic facts about it in Lemma 3.1. The Stern-Brocot tree and its connections with continued fractions are discussed in detail by Graham, Knuth, and Patashnik [20]. Their definition of the Stern-Brocot tree is slightly different from ours. We adopt their notation  $a \perp b$  to say that integers  $a$  and  $b$  are relatively prime.

**Lemma 3.1.** (a) *The Stern-Brocot tree is symmetric about 0 with its right half positive and its left half negative.*

(b) *If  $[\frac{a}{b}, \frac{c}{d}]$  is a node in the positive half of the Stern-Brocot tree, then  $bc - ad = 1$ ,  $a \perp b$ , and  $c \perp d$ .*

(c) *Conversely, if  $a/b$  and  $c/d$  are non-negative rational numbers with zero and infinity represented as  $\frac{0}{1}$  and  $\frac{1}{0}$  respectively, and  $bc - ad = 1$  then  $[\frac{a}{b}, \frac{c}{d}]$  occurs as a node in the Stern-Brocot tree. Consequently, every rational number  $a/b$ ,  $a \perp b$ , appears as an endpoint of a Stern-Brocot interval of finite depth.*

*Proof.* (a) is obvious; see Figure 3. The proof of (b) is an easy induction on the depth of the tree. (c) is a little bit less easy. Its proof is related to Euclid's algorithm for computing the greatest common divisor of two integers. See [20].  $\square$

We adopt a labelling scheme for Stern-Brocot intervals (nodes of the Stern-Brocot tree) that differs only a bit from that in [20]. The root  $[\frac{-1}{0}, \frac{1}{0}]$  has the empty label. Its left and right children  $[\frac{-1}{0}, \frac{0}{1}]$  and  $[\frac{0}{1}, \frac{1}{0}]$  are labelled  $l$  and  $r$  respectively. The left child of  $l$ ,  $[\frac{-1}{0}, \frac{-1}{1}]$ , is labelled  $lL$ . The right child of  $lL$ ,  $[\frac{-2}{1}, \frac{-1}{1}]$ , is labelled  $lLR$ , and so on. Only the first letter of a label is in lower case, because the division of the root is special.

We use  $l\alpha$  or  $r\alpha$  to denote the labels of Stern-Brocot intervals other than the root, with  $\alpha$  being a possibly empty sequence of  $L$ s and  $R$ s. The sequence obtained by changing  $\alpha$ 's  $L$ s to  $R$ s and  $R$ s to  $L$ s is denoted  $\bar{\alpha}$ . For example, the reflection of the positive interval  $r\alpha$  about 0 is the negative interval  $l\bar{\alpha}$ . The length of  $\alpha$  is denoted by  $|\alpha|$ . We take the depth of  $l\alpha$  or  $r\alpha$  to be  $1 + |\alpha|$ .

Lemmas 3.2 and 3.3 express the maps  $m \rightarrow 1/m$  and  $m \rightarrow \pm 1 + m$  succinctly for Stern-Brocot intervals. They allow us to reduce the invariance requirement (3.1) for Stern-Brocot intervals to an infinite system of linear equations (see (3.2)). That reduction is the first step in constructing  $\nu_f$ .

**Lemma 3.2.** *The image of the interval  $[a/b, c/d]$  under the map  $m \rightarrow 1/m$  — which is  $[d/c, b/a]$  if 0 is not an interior point — is given by the following rules for Stern-Brocot intervals:*

$$l\alpha \rightarrow l\bar{\alpha}, \quad r\alpha \rightarrow r\bar{\alpha}.$$

*Proof.* We give the proof for intervals of type  $r\alpha$  using induction on the depth of  $r\alpha$  in the Stern-Brocot tree. The proof for intervals  $l\alpha$  is similar.

The base case  $r \rightarrow r$  is true because  $m \in [0, \infty]$  if and only if  $1/m \in [0, \infty]$ .

For the inductive case, note that  $[\frac{a}{b}, \frac{c}{d}]$ , its left child  $[\frac{a}{b}, \frac{a+c}{b+d}]$ , and its right child  $[\frac{a+c}{b+d}, \frac{c}{d}]$  are mapped by  $m \rightarrow 1/m$  to  $[\frac{d}{c}, \frac{b}{a}]$ , its right child  $[\frac{b+d}{a+c}, \frac{b}{a}]$ , and its left child  $[\frac{d}{c}, \frac{b+d}{a+c}]$ , respectively. Therefore, if  $r\alpha \rightarrow r\bar{\alpha}$  then  $r\alpha L \rightarrow r\bar{\alpha}R$  and  $r\alpha R \rightarrow r\bar{\alpha}L$ .  $\square$

Unlike the inversion operation  $m \rightarrow 1/m$  in the previous lemma, both the operations  $m \rightarrow \pm 1 + m$  in the following lemma change the depth of Stern-Brocot intervals.

**Lemma 3.3.** *The image of Stern-Brocot intervals under the map  $m \rightarrow -1 + m$  is given by the following rules:*

$$l\alpha \rightarrow lL\alpha, \quad rL\alpha \rightarrow lR\alpha, \quad rR\alpha \rightarrow r\alpha.$$

*Similarly, the image of Stern-Brocot intervals under the map  $m \rightarrow 1 + m$  is given by the following rules:*

$$lL\alpha \rightarrow l\alpha, \quad lR\alpha \rightarrow rL\alpha, \quad r\alpha \rightarrow rR\alpha.$$

*Proof.* Similar to the previous proof. We will outline the proof for  $m \rightarrow 1 + m$  only.

The base cases, adding 1 to the intervals  $lL$ ,  $lR$  and  $r$ , are easy to check.

For the induction, we note that  $[\frac{a}{b}, \frac{c}{d}]$  is divided in the Stern-Brocot tree at the point  $\frac{a+c}{b+d}$ , and its map under  $m \rightarrow 1 + m$ ,  $[1 + \frac{a}{b}, 1 + \frac{c}{d}]$ , is divided in the Stern-Brocot tree at the point  $1 + \frac{a+c}{b+d}$ . Thus  $[\frac{a}{b}, \frac{c}{d}]$ , its left child, and its right child map to  $[1 + \frac{a}{b}, 1 + \frac{c}{d}]$ , its left child, and its right child, respectively.  $\square$

By Lemma 3.3, subtraction and addition of 1 to intervals in the Stern-Brocot tree correspond to left and right rotation of the tree. Tree rotations are used to implement balanced trees in computer science [13].

Thanks to Lemmas 3.2 and 3.3, the backward maps  $m \rightarrow 1/(\pm 1 + m)$  can be performed on Stern-Brocot intervals easily. For example,  $1/(1 + lLRL) = 1/lRL = lLR$ . The invariance requirement (3.1) for Stern-Brocot intervals becomes an infinite set of linear equations for  $\nu_f(I)$ ,  $I$  being any Stern-Brocot interval:

$$\begin{aligned}
 \nu_f(l) &= \frac{1}{2}\nu_f(lR) + \frac{1}{2}(\nu_f(l) + \nu_f(rR)), \\
 \nu_f(r) &= \frac{1}{2}(\nu_f(r) + \nu_f(lL)) + \frac{1}{2}\nu_f(rL), \\
 \nu_f(lL\alpha) &= \frac{1}{2}\nu_f(l\overline{L\alpha}) + \frac{1}{2}\nu_f(l\bar{\alpha}), \\
 \nu_f(lR\alpha) &= \frac{1}{2}\nu_f(l\overline{R\alpha}) + \frac{1}{2}\nu_f(r\bar{\alpha}), \\
 \nu_f(rL\alpha) &= \frac{1}{2}\nu_f(l\overline{R\alpha}) + \frac{1}{2}\nu_f(r\overline{L\alpha}), \\
 \nu_f(rR\alpha) &= \frac{1}{2}\nu_f(r\bar{\alpha}) + \frac{1}{2}\nu_f(r\overline{R\alpha}).
 \end{aligned}
 \tag{3.2}$$

We guessed the solution of (3.2). Even though the linear system (3.2) has only rational coefficients, its solution involves  $\sqrt{5}$ , an irrational number! Let  $g = (1 + \sqrt{5})/2$ . Since  $\nu_f$  is a probability measure, we require that  $\nu_f([-\infty, \infty]) = 1$ .




FIGURE 4. (a), (b), (c) show the measure  $\nu_f$  over directions in  $R^2$ . In these figures, the interval  $[0, \infty]$  is divided into  $2^3$ ,  $2^5$ , and  $2^8$  Stern-Brocot intervals of the same depth, and then slopes are converted to angles in the interval  $[0, \pi/2]$ . The area above an interval gives its measure under  $\nu_f$ . Because of symmetry,  $\nu_f$  in the directions  $[-\pi/2, 0]$  can be obtained by reflecting (a), (b) and (c). Some of the spikes in (c) were cut off because they were too tall. (d) is the distribution function for  $\nu_f$  with directions parameterized using angles.

The solution is:

$$\begin{aligned}
 \nu_f(r) &= 1/2, \\
 \nu_f(r\alpha L) &= \begin{cases} \frac{1}{1+g}\nu_f(r\alpha) & \text{if } |\alpha| \text{ is even,} \\ \frac{g}{1+g}\nu_f(r\alpha) & \text{if } |\alpha| \text{ is odd,} \end{cases} \\
 \nu_f(r\alpha R) &= \begin{cases} \frac{g}{1+g}\nu_f(r\alpha) & \text{if } |\alpha| \text{ is even,} \\ \frac{1}{1+g}\nu_f(r\alpha) & \text{if } |\alpha| \text{ is odd,} \end{cases} \\
 \nu_f(l\alpha) &= \nu_f(r\bar{\alpha}).
 \end{aligned}
 \tag{3.3}$$

For example,  $\nu_f(r) = 1/2$ ,  $\nu_f(rL) = (1+g)^{-1}/2$ ,  $\nu_f(rLL) = g(1+g)^{-2}/2$ . Since  $\nu_f(l\alpha) = \nu_f(r\bar{\alpha})$  by (3.3), the measure  $\nu_f$  is symmetric about 0. The same features of  $\nu_f$  repeat at finer and finer scales. See Figure 4.

**Theorem 3.4.** *The measure  $\nu_f$  defined by (3.3) satisfies the invariance requirement (3.1) for every Stern-Brocot interval. Further, with directions parameterized by slopes,  $\nu_f$  defined by (3.3) gives the unique  $\mu_f$ -invariant probability measure over directions in the real plane  $R^2$ .*

*Proof.* To show that  $\nu_f$  is  $\mu_f$ -invariant, it is enough to show that  $\nu_f$  satisfies the invariance conditions (3.2) for Stern-Brocot intervals. The reason is that  $\nu_f$  is obviously a continuous measure, every rational appears in the Stern-Brocot tree at a finite depth by Lemma 3.1c, and the rationals are dense in  $R$ . For the uniqueness of  $\nu_f$ , see [7, p. 31].

It is enough to prove the invariance condition for positive intervals  $r\alpha$ . The validity of the invariance condition for negative Stern-Brocot intervals follows from symmetry. Assume the invariance condition for the interval  $rL\alpha$ :

$$\nu_f(rL\alpha) = \frac{1}{2}\nu_f(l\overline{R\alpha}) + \frac{1}{2}\nu_f(r\overline{RL\alpha}).$$

Then the invariance condition for  $rL\alpha L$ ,

$$\nu_f(rL\alpha L) = \frac{1}{2}\nu_f(l\overline{R\alpha R}) + \frac{1}{2}\nu_f(r\overline{RL\alpha R}),$$

is also true, because the three fractions

$$\frac{\nu_f(rL\alpha L)}{\nu_f(rL\alpha)}, \quad \frac{\nu_f(l\overline{R\alpha R})}{\nu_f(l\overline{R\alpha})}, \quad \frac{\nu_f(r\overline{RL\alpha R})}{\nu_f(r\overline{RL\alpha})}$$

are all either  $g/(1+g)$  or  $1/(1+g)$  according as  $|\alpha|$  is even or odd. By a similar argument, if the invariance condition (3.2) holds for all positive Stern-Brocot intervals at depth  $d \geq 2$ , then the invariance condition holds for all positive Stern-Brocot intervals at depth  $d + 1$ .

Therefore, it suffices to verify (3.2) for  $r, rL$ , and  $rR$ . For  $r$ , (3.2) requires

$$\frac{1}{2} = \frac{1}{2}\left(\frac{1}{2} + \frac{1}{2(1+g)}\right) + \frac{1}{2}\left(\frac{g}{2(1+g)}\right),$$

which is obviously true. For  $rL$ , (3.2) requires,

$$\frac{1}{2(1+g)} = \frac{g}{4(1+g)} + \frac{1}{4(1+g)^2},$$

which is true because  $g = (1 + \sqrt{5})/2$ . The invariance condition for  $rR$  can be verified similarly. Thus the invariance condition (3.2) holds for all Stern-Brocot intervals, and we can say that  $\nu_f$  is the unique  $\mu_f$ -invariant probability measure.  $\square$

Because of symmetry, the measure  $\nu_f$  over slopes given by (3.3) is invariant even for the distribution that picks one of  $\begin{pmatrix} 0 & 1 \\ \pm 1 & \pm 1 \end{pmatrix}$  with probability  $1/4$ . Moreover, Furstenberg’s integral for the Lyapunov exponent  $\gamma$  of this distribution is also given by (2.3).

According to historical remarks in [11], measures similar to  $\nu_f$  have been studied by Denjoy, Minkowski, and de Rham. But is  $\nu_f$  a fractal? To make this precise, we need the definition

$$\dim(\nu_f) = \inf\{\dim(S) \mid \nu_f \text{ is supported on } S\},$$

where  $\dim(S)$  is the Hausdorff dimension of  $S \subset R$ . To show that  $\nu_f$  is a fractal, it is necessary to prove that  $0 < \dim(\nu_f) < 1$ . It is known that  $0 < \dim(\nu_f)$  [7, p. 162]. David Allwright of Oxford University has shown us a short proof that  $\nu_f$  is singular with respect to the Lebesgue measure; Allwright’s proof relies on Theorems 30 and 31 of Khintchine [25]. The Hausdorff dimensions of very similar measures have been determined by Kinney and Pitcher [27]. We also note that Ledrappier has conjectured a formula for  $\dim(\nu_f)$  [30] [7, p. 162].

For some distributions supported on 2-dimensional matrices with non-negative entries, the infinite linear system analogous to (3.2) is triangular, or in other words, the invariance requirement for a Stern-Brocot interval involves only intervals at a lesser depth. For a typical example, choose  $\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}$  with probability  $p$ ,  $0 < p < 1$ , and  $\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$  with probability  $1 - p$ . In this example, the invariant measure over directions parameterized by slopes is supported on  $[0, \infty]$ , the slope  $m$  is mapped to  $1/(1 + m)$  and  $1 + 1/m$  respectively, and the ranges of those two maps ( $[0, 1]$  and  $[1, \infty]$ ) are disjoint. Chassaing, Letac and Mora [11] have found the invariant measure for several 2-dimensional random matrix products that fit into this framework. All their matrices have non-negative entries. Moreover, since the linear systems for finding the invariant measure are triangular for all the examples in [11], the solution can have irrational numbers only if the original problem does.

The techniques described in this section can be used to find the invariant measure corresponding to the random recurrence  $t_n = \alpha t_{n-1} + t_{n-2}$  if  $\alpha$  is distributed on the positive integers. But this situation was already covered in [11]. When the random variable  $\alpha$  takes both positive and negative values, random Fibonacci recurrence is the only example we know where the technique of using Stern-Brocot intervals for finding the invariant measure can be made to work. Numerical results about a generalization of the random Fibonacci sequences where  $\alpha$  takes both positive and negative values are given in [16].

4.  $e^{\gamma_f} = 1.13198824\dots$

Furstenberg’s integral for  $\gamma_f$  (2.3) can be written as

$$\gamma_f = 2 \int_0^\infty \frac{1}{4} \log\left(\frac{1 + 4m^4}{(1 + m^2)^2}\right) d\nu_f(m)$$

because both the integrand and  $\nu_f$  are symmetric about 0. In this section, we use this formula to compute  $\gamma_f$  with the help of a computer. Thus the determination of  $e^{\gamma_f}$  to be 1.13198824... is computer assisted. We will explain later why we report this result as a theorem (Theorem 4.2), even though it is computer assisted.

Let  $I_j^d$ ,  $1 \leq j \leq 2^d$ , be the  $2^d$  positive Stern-Brocot intervals at depth  $d + 1$ . Then,

$$(4.1) \quad p_d = 2 \sum_{j=1}^{2^d} \min_{m \in I_j^d} \text{amp}(m) \nu_f(I_j^d) < \gamma_f < q_d = 2 \sum_{j=1}^{2^d} \max_{m \in I_j^d} \text{amp}(m) \nu_f(I_j^d).$$

The inequalities above are strict because  $\text{amp}(m)$  is not constant, and  $\nu_f$  is continuous. Also, (4.1) defines  $p_d$  and  $q_d$ . Since  $\gamma_f$  is trapped in the intervals  $(p_d, q_d)$ , and the interval length  $|q_d - p_d|$  shrinks to 0 as  $d$  increases, we can find  $\gamma_f$  to any desired accuracy by computing  $p_d$  and  $q_d$  for large enough  $d$ .

We computed  $p_d$  and  $q_d$  with  $d = 28$  on a computer using IEEE double precision arithmetic (the C program used is described in the appendix). Computations in floating point arithmetic are not exact, but when done carefully, give an answer that is close to the exact answer. If  $\text{fl}(e)$  denotes the number obtained by evaluating the expression  $e$  in floating point arithmetic,  $\text{fl}(e)$  depends both on the type of floating point arithmetic used and the algorithm used to evaluate  $e$ . Our computations using IEEE double precision arithmetic [24] and an algorithm described in the appendix

gave

$$(4.2) \quad \text{fl}(p_{28}) = 0.1239755981508, \quad \text{fl}(q_{28}) = 0.1239755994406.$$

In hexadecimal code, the 64 bits of  $\text{fl}(p_d)$  and  $\text{fl}(q_d)$  in IEEE double precision format are *3fbfbcdd638f4d87* and *3fbfbcdd6919756d*, respectively. The appendix will explain the way to reproduce our computation to get exactly these two numbers. We will now upper bound the errors  $|\text{fl}(p_{28}) - p_{28}|$  and  $|\text{fl}(q_{28}) - q_{28}|$  to realize our aim of obtaining bounds for  $\gamma_f$  from (4.2).

IEEE double precision arithmetic (defined by the standard IEEE-754 [24]) can represent all real numbers of binary form  $(-1)^s b_0 . b_1 \dots b_{52} 2^{e-1023}$  exactly. Here,  $b_0 = 1$ , the bits  $b_1$  to  $b_{52}$  can be 1 or 0, the sign bit  $s$  can be 1 or 0, and the biased exponent  $e$  can be any integer in the range  $0 < e < 2047$ . The number 0 can also be represented exactly. In fact, the values  $e = 0$  and  $e = 2047$  are used to implement special features that we do not describe. From here on, floating point arithmetic always refers to IEEE double precision arithmetic, and floating point number refers to a number in that arithmetic. Thus if  $a$  is a real number in the range  $[2^{-1022}, (1 + 2^{-1} + \dots + 2^{-52})2^{1023}]$ ,  $a$  can be represented such that  $\text{fl}(a) = a(1 + E)$  with the relative error  $E$  satisfying  $|E| < 2^{-52}$  [22, p. 42].

The IEEE standard treats  $+$ ,  $-$ ,  $\times$ ,  $\div$ ,  $\sqrt{\phantom{x}}$  as basic operations. The basic operations cannot always be performed exactly. For example, the sum of two floating point numbers may not have an exact floating point representation. However, all these basic operations are performed as if an intermediate result correct to infinite precision is coerced into a representable number by rounding. We assume the “round to nearest” mode, which is the default type of rounding. Thus, if  $a$  and  $b$  are floating point numbers,

$$(4.3) \quad \begin{aligned} \text{fl}(a + b) &= (a + b)(1 + E), \\ \text{fl}(a - b) &= (a - b)(1 + E), \\ \text{fl}(a/b) &= (a/b)(1 + E), \\ \text{fl}(a \times b) &= (a \times b)(1 + E), \\ \text{fl}(\sqrt{a}) &= (\sqrt{a})(1 + E), \end{aligned}$$

where the relative error  $E$  may depend upon  $a$ ,  $b$ , and the operation performed, but  $|E| < 2^{-52}$ . For convenience, we denote  $2^{-52}$  by  $u$ .<sup>1</sup> For (4.3) to be valid, however, the operation should not overflow and produce a number that is too big to be represented, or underflow and produce a number that is too small to be represented.

The C program we give in the appendix uses a function `tlog(x)` to compute  $\log x$ . This becomes necessary because `log` is not a basic operation in the IEEE standard. However, `tlog()` is implemented so that

$$(4.4) \quad \text{fl}(\log a) = \log a(1 + E)$$

with  $|E| < u$  whenever  $a$  is a positive floating point number. For the clever ideas that go into `tlog()` and the error analysis, see the original paper by Tang [38].

The proof of the following lemma is given in the appendix.

---

<sup>1</sup>The bounds on  $|E|$  can be taken as  $2^{-53}$  [22, p. 42], but with the current choice the relative error of Tang’s log function (see (4.4)) has the same bound as that of the basic operations.

**Lemma 4.1.** *Assume that (4.3) and (4.4) hold with  $0 < u < 1/10$  for the floating point arithmetic used. Then for the algorithm to compute the sums  $p_d$  and  $q_d$  described in the appendix,*

$$\begin{aligned} |\text{fl}(p_d) - p_d| &< \frac{\log 4}{4}(e^{u(d+1)} - 1) + \frac{33}{4}ue^{u(d+1)}, \\ |\text{fl}(q_d) - q_d| &< \frac{\log 4}{4}(e^{u(d+1)} - 1) + \frac{33}{4}ue^{u(d+1)}. \end{aligned}$$

It is easy, though a bit tedious, to show that the discretization error  $|p_d - q_d|$  is  $O(1/2^d)$ . By Lemma 4.1, the rounding errors in computing  $p_d$  and  $q_d$  are roughly a small multiple of  $u$ . Thus to compute  $\gamma_f$  with an absolute error of  $\epsilon$ , the depth of the calculation has to be about  $-\log_2 \epsilon$  and the unit roundoff of the floating point arithmetic has to be at least as small as  $\epsilon$ .

In the theorem below, by 1.13198824... we mean a number in the interval  $[1.13198824, 1.13198825)$ .

**Theorem 4.2.** (a) *The constant  $\gamma_f$  lies in the interval*

$$(0.1239755980, 0.1239755995).$$

(b)  $e^{\gamma_f} = 1.13198824\dots$

(c) *As  $n \rightarrow \infty$ ,*

$$\sqrt[n]{|t_n|} \rightarrow 1.13198824\dots$$

*with probability 1.*

*Proof.* In the computation leading to  $\text{fl}(p_{28})$  and  $\text{fl}(q_{28})$ , there are no overflows or underflows, and hence (4.3) and (4.4) are always true. Therefore, we can use  $u = 2^{-52}$  and  $d = 28$  in Lemma 4.1 to get

$$|\text{fl}(p_{28}) - p_{28}| < 10^{-14}, \quad |\text{fl}(q_{28}) - q_{28}| < 10^{-14}.$$

Now the values of  $\text{fl}(p_{28})$  and  $\text{fl}(q_{28})$  in (4.2) imply (a). (b) is implied by (a). In fact, we can also say that the digit of  $e^{\gamma_f}$  after the last 4 in (b) must be an 8 or a 9. (c) follows from earlier remarks.  $\square$

Theorem 4.2 above is the main result of this paper. We arrived at Theorem 4.2 using Lemma 4.1 and rounding error analysis. An alternative is to use interval arithmetic to validate the computation [1]. Instead of rounding the computations to the nearest floating point number, interval arithmetic carefully rounds the various stages of the computation either upwards or downwards to compute a lower bound for  $p_d$  and an upper bound for  $q_d$ . As a result, were we to use interval arithmetic there would be no need for rounding error analysis. A disadvantage would be that the manipulation of rounding modes necessary for implementing interval arithmetic would make it significantly more expensive on most computers. Our approach exposes the ideas behind floating point arithmetic and shows that floating point arithmetic is rigorous too. Besides, the rounding error analysis as summarized by Lemma 4.1 gives a clear idea of the error due to rounding. This tells us, for example, that the rounding errors  $|\text{fl}(p_{28}) - p_{28}|$  and  $|\text{fl}(q_{28}) - q_{28}|$ , which are both less than  $10^{-14}$ , are much smaller than the discretization error  $|p_{28} - q_{28}|$ , which is about  $10^{-8}$ .




FIGURE 5. The Lyapunov exponent  $\gamma_f(p)$  vs.  $p$ . To obtain the curve above,  $\gamma_f(p)$  was determined by numerically approximating the correct invariant distribution for 199 values of  $p$ , equally spaced in  $(0, 1)$ . Each  $\gamma_f(p)$  is accurate to 5 decimal digits. For a description of the numerical method, sometimes called Ulam's method, see [16] or [23].

Since the proof of Theorem 4.2 relies on a computer calculation, the validity of the proof requires some comment. The construction of  $\nu_f$  in Section 2, the program and the rounding error analysis given in the appendix can all be checked line by line. However, Theorem 4.2 still assumes the correct implementation of various software and hardware components including the standard IEEE-754. We did the computation on two entirely different systems — SUN's Sparc server 670 MP, and Intel's i686 with the Linux operating system. In both cases, the results were exactly the same as given in (4.2); the hex codes for  $\text{fl}(p_d)$  and  $\text{fl}(q_d)$  matched the hex codes given below (4.2). As it is very unlikely that two systems with such different architectures may have the same bug, we feel that the correctness of Theorem 4.2 should, at worst, be doubted no more than that of tedious and intricate proofs that can be checked line by line. Though the use of floating point arithmetic to prove a theorem may be unusual, the proof of Theorem 4.2 is only as dependent on the correctness of the computer system as, say, the proof of the four-color theorem; in other words, assuming the implementation of IEEE arithmetic to be correct is just like assuming the implementation of a memory-to-register copy instruction to be correct.

Besides, all components of a computer system, like mathematical proofs, can be checked in careful line by line detail, and this is done many times during and after their implementation. However, experience has shown that some bugs can defy even the most careful scrutiny. A great deal of research has gone into developing systems to verify that hardware and software implementations meet their specification [12].

In recent work, Tsitsiklis and Blondel [39] claim that the upper Lyapunov exponent is not "algorithmically approximable." They prove that there can be no Turing machine which accepts a pair of matrices as input and returns an approximation to the upper Lyapunov exponent with bounded error. The distribution can

be anything which picks both the input matrices with nonzero probability and no others. This uncomputability result holds when the dimension of the matrices is 48 or greater.

To interpret this result properly, we think that it must be compared with similar results for easier analytic problems like computing eigenvalues of matrices and zeros of polynomials. For literature on similar analytic problems, we refer to [6]; but the model of computation used in that book is not the Turing model. In another sense the result of Tsitsiklis and Blondel is limited. It applies only when the class of problems includes distributions with singular matrices. But most of the theory of random matrix products has been developed for distributions supported on nonsingular matrices. When the distribution is supported on nonsingular matrices, we give an algorithm for computing the top Lyapunov exponent with an arbitrarily small absolute error in [41]. For this algorithm to be effective, a mild irreducibility assumption about the support of the distribution has to hold.

To conclude, we ask: Is there a short analytic description of  $\gamma_f$ ? The fractal quality of  $\gamma_f$  suggests no. But let  $\gamma_f(p)$  be the Lyapunov exponent of the obvious generalization  $t_1 = t_2 = 1$ , and for  $n \geq 2$ ,  $t_n = \pm t_{n-1} \pm t_{n-2}$  with each  $\pm$  sign independent and either  $+$  with probability  $p$  or  $-$  with probability  $1 - p$ . Unfortunately, the techniques described in this paper for  $\gamma_f(1/2)$  do not seem to generalize easily to  $\gamma_f(p)$ ,  $0 < p < 1$ . A beautiful result of Peres [35] implies that  $\gamma_f(p)$  is a real analytic function of  $p$ . See Figure 5. The analyticity of  $\gamma_f(p)$  vs.  $p$  seems to increase the possibility that there might be a short analytic description of  $\gamma_f$ .

#### APPENDIX. ROUNDING ERROR ANALYSIS

The main steps in the computation of  $p_d$  and  $q_d$  are the computation of  $\nu_f(I_j^d)$ , where  $I_j^d$ ,  $1 \leq j \leq 2^d$ , are the  $2^d$  positive Stern-Brocot intervals of depth  $d + 1$ ; the minimization and maximization of  $\text{amp}(m)$  over  $I_j^d$ ; and the summation over  $1 \leq j \leq 2^d$  as in the defining equation (4.1). We describe some aspects of the computation and then give a rounding error analysis to prove Lemma 4.1. A C program for computing  $p_d$  and  $q_d$  for  $d = 28$  is given at the end of this section so that our computation can be reproduced; its perusal is not necessary for reading this section.

Lemma 3.2 implies that the denominators of the  $2^d$  positive Stern-Brocot intervals of depth  $d + 1$  occur in an order that is the reverse of the order of the numerators. For example, the positive Stern-Brocot intervals of depth 4 are defined by divisions at the points  $\frac{0}{1}, \frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \frac{1}{1}, \frac{3}{2}, \frac{2}{1}, \frac{3}{1}, \frac{1}{0}$ , the numerators for that depth occur in the order 0, 1, 1, 2, 1, 3, 2, 3, 1, and the denominators occur in the reverse order 1, 3, 2, 3, 1, 2, 1, 1, 0. We use this fact to avoid storing the denominators of the Stern-Brocot divisions. The numerators are stored in the array `num[]` by the C program.

To compute  $p_d$  and  $q_d$ , we use (4.1) in the following form:

$$(A.1) \quad \begin{aligned} p_d &= \sum_{j=1}^{2^d} \min_{m \in I_j^d} \left( \log \frac{1 + 4m^4}{(1 + m^2)^2} \right) \frac{\nu_f(I_j^d)}{2}, \\ q_d &= \sum_{j=1}^{2^d} \max_{m \in I_j^d} \left( \log \frac{1 + 4m^4}{(1 + m^2)^2} \right) \frac{\nu_f(I_j^d)}{2}. \end{aligned}$$

By (3.3),  $\nu_f(I_j^d)/2$  is one of the  $d + 1$  numbers  $g^{d-i}(1 + g)^{-d}/4$ ,  $0 \leq i \leq d$ , where  $g = (1 + \sqrt{5})/2$ . The array `table[]` in the C program is initialized after precomputing these  $d + 1$  numbers to very high accuracy in the symbolic algebra system *Mathematica* so that `table[i] = (g^{d-i}(1 + g)^{-d}/4)(1 + E)` with the relative error  $E$  satisfying  $|E| < u$ . The index  $i$  into `table[]` for getting  $\nu_f(I_j^d)/2$  is obtained by taking the binary representation of  $j$ , flipping all the odd bits if  $d$  is even and all the even bits if  $d$  is odd with the least significant bit taken as an even bit, and then counting the number of 1s; correctness of this procedure can be proved easily using induction.

The minimization and the maximization of  $4 \operatorname{amp}(m)$  over  $I_j^d$  in (A.1) are easy to do. Since  $\operatorname{amp}(m)$  has its only local minimum for  $m \geq 0$  at  $m = 1/2$  (see the figure just after (2.2)), both the minimum and the maximum are at the endpoints of  $I_j^d$ .

The summations in (A.1) are performed pairwise, not left to right. The pairwise summation of  $2^d$  numbers is done by dividing the  $2^d$  numbers into  $2^{d-1}$  pairs of adjacent numbers, adding each pair to get  $2^{d-1}$  numbers, and then reducing the  $2^{d-1}$  numbers to  $2^{d-2}$  numbers similarly, and so on until a single number is obtained. Rounding error analysis leads to smaller upper bounds on  $|\operatorname{fl}(p_d) - p_d|$  and  $|\operatorname{fl}(q_d) - q_d|$  for pairwise summation than for term-by-term left to right summation [22, p. 92]. The bounds for left to right summation are not small enough to give  $e^{\gamma_f}$  correctly to the 8 decimal digits shown in Theorem 4.2.

Lemmas A.1 and A.2 help simplify the proof of Lemma 4.1.

**Lemma A.1.** *Assume  $0 < f_1(u) < 1 + e_1 < g_1(u)$  and  $0 < f_2(u) < 1 + e_2 < g_2(u)$ .*

(a) *If  $a > 0$ ,  $b > 0$ , and  $a(1 + e_1) + b(1 + e_2) = (a + b)(1 + E)$ , then  $\min(f_1(u), f_2(u)) < 1 + E < \max(g_1(u), g_2(u))$ .*

(b) *If  $1 + E = (1 + e_1)(1 + e_2)$ , then  $f_1(u)f_2(u) < 1 + E < g_1(u)g_2(u)$ .*

(c) *If  $1 + E = (1 + e_1)/(1 + e_2)$ , then  $f_1(u)/g_2(u) < 1 + E < g_1(u)/f_2(u)$ .*

*Proof.* To prove (a), note that  $1 + E$  is the weighted mean of  $1 + e_1$  and  $1 + e_2$ . (b) and (c) are trivial.  $\square$

Consider the computation  $\operatorname{fl}(m^2)$ :

$$\operatorname{fl}(m^2) = \operatorname{fl}(m) \operatorname{fl}(m)(1 + e') = m^2(1 + e')(1 + e'')^2,$$

where  $e''$  is the relative error in representing  $m$ , and  $e'$  is the relative error caused by rounding the multiplication. By (4.3) and remarks in the paragraph preceding it,  $1 - u < 1 + e'$ ,  $1 + e'' < 1 + u$ . Lemma A.1b allows us to gather the factors  $1 + e'$  and  $(1 + e'')^2$  together and write

$$(A.2) \quad \operatorname{fl}(m^2) = m^2(1 + e_0),$$

with  $(1 - u)^3 < 1 + E < (1 + u)^3$ .

Consider the computation  $\operatorname{fl}(1 + m^2)$ :

$$\operatorname{fl}(1 + m^2) = (1 + \operatorname{fl}(m^2))(1 + e''') = (1 + m^2(1 + e')(1 + e'')^2)(1 + e'''),$$

where  $e'''$  is the relative error in the addition  $1 + m^2$ , and  $e''$ ,  $e'$  are, as before, the relative errors in representing  $m$  and the multiplication  $m \times m$ , respectively. As it was with  $1 + e'$  and  $1 + e''$ ,  $1 - u < 1 + e''' < 1 + u$  by (4.3), and we can use Lemma A.1a to pull  $(1 + e')(1 + e'')^2$  out of the sum  $1 + m^2$ , and Lemma A.1b to multiply

$(1 + e')(1 + e'')^2(1 + e''')$  to get

$$(A.3) \quad \text{fl}(1 + m^2) = (1 + m^2)(1 + e'_0),$$

with  $(1 - u)^4 < 1 + e'_0 < (1 + u)^4$ .

Thus Lemma A.1 allows us to pull factors like  $(1 + e_i)$  out of sums (Lemma A.1a), or to multiply them together (Lemma A.1b), or to divide between them (Lemma A.1c). Rounding error analyses of simple computations, like the analyses of  $\text{fl}(m^2)$  and  $\text{fl}(1 + m^2)$  given above, feature three steps. First, relative errors  $e_i$  caused by rounding are assigned to all the basic operations. Second,  $1 + e_i$  are bounded using (4.3) or (4.4). Third, factors like  $(1 + e_i)$  are gathered together using Lemma A.1. In the proof of Lemma 4.1, we always spell out the first step in detail, but sometimes omit details for the second and third steps.

The inequalities in Lemma A.2 below are used in the proof of Lemma 4.1.

**Lemma A.2.** (a) *If  $0 < u < 1/4$ , then  $\log \frac{1+u}{1-u} < 3u$ .*

(b)  $(1 + \alpha)^d < e^{\alpha d}$  for  $\alpha > 0$  and  $d$  a positive integer.

*Proof.* It is easy to prove (a) by expanding  $\log((1 + u)/(1 - u))$  in a series. (b) can be proved by comparing the binomial expansion of  $(1 + \alpha)^d$  with the series expansion of  $e^{\alpha d}$ .  $\square$

The summations in the proof below are all over  $1 \leq j \leq 2^d$ .

*Proof of Lemma 4.1.* We will prove the upper bound only for  $|\text{fl}(p_d) - p_d|$ . The proof for  $|\text{fl}(q_d) - q_d|$  is similar.

First, consider the computation of  $4 \text{amp}(m) = \log \frac{1+4m^4}{(1+m^2)^2}$ :

$$\begin{aligned} & \text{fl}\left(\log \frac{1 + 4m^4}{(1 + m^2)^2}\right) \\ &= \log\left(\frac{(1 + 4m^4(1 + e_0)^2(1 + e_1)(1 + e_2))(1 + e_3)}{(1 + m^2)^2(1 + e'_0)^2(1 + e_4)}(1 + e_5)\right) (1 + e_6), \end{aligned}$$

where  $e_0$  and  $e'_0$  are the relative errors in  $\text{fl}(m^2)$  and  $\text{fl}(1 + m^2)$  as in (A.2) and (A.3) respectively,  $e_1, e_2$  are the relative errors of the two multiplications  $(4 \times m^2) \times m^2$ ,  $e_3$  of the addition  $1 + 4m^4$ ,  $e_4$  of the multiplication  $(1 + m^2) \times (1 + m^2)$ ,  $e_5$  of the division  $(1 + 4m^4)/(1 + m^2)^2$ , and  $e_6$  of taking the log. By assumptions (4.3) and (4.4),  $1 - u < 1 + e_i < 1 + u$  for  $1 \leq i \leq 6$ . Lemma A.1 gives

$$(A.4) \quad \text{fl}\left(\log \frac{1 + 4m^4}{(1 + m^2)^2}\right) = \left(\log \frac{1 + 4m^4}{(1 + m^2)^2}\right)(1 + E_1) + E_2,$$

with  $1 - u < 1 + E_1 < 1 + u$  and  $|E_2| < (1 + u) \log((1 + u)^{10}(1 - u)^{-9})$ . A weaker, but simpler, bound is  $|E_2| < 10(1 + u) \log((1 + u)/(1 - u))$ . Now, the assumption  $u < 1/10$  implies  $10(1 + u) < 11$ , which, together with Lemma A.2b, gives the simple bound  $|E_2| < 33u$ .

Second, recall that  $\nu_f(I_j^d)/2$  is obtained by precomputing  $g^{d-i}(1 + g)^{-d}/4$  to high precision. Therefore

$$(A.5) \quad \text{fl}(\nu_f(I_j^d)/2) = \frac{\nu_f(I_j^d)}{2}(1 + E_3),$$

with  $|E_3| < u$ .

Finally, consider the pairwise summation to compute  $p_d$ . Let  $m_j$  be the endpoint of  $I_j^d$  where  $\text{amp}(m)$  is minimum. Then,

$$\text{fl}(p_d) = \sum \left( \log \frac{1 + 4m_j^4}{(1 + m_j^2)^2} (1 + E_1^j) + E_2^j \right) \left( \frac{\nu_f(I_j^d)}{2} (1 + E_3^j) \right) (1 + E_4^j)$$

where  $E_1^j$  and  $E_2^j$  are the relative errors in computing  $\log((1 + 4m_j^4)(1 + m_j^2)^{-2})$ , and therefore are bounded like  $E_1$  and  $E_2$  in (A.4);  $E_3^j$  is the relative error in computing  $\nu_f(I_j^d)/2$  and is bounded like  $E_3$  in (A.5); and the factors  $1 + E_4^j$  take up the errors in the pairwise summation. By Higham [22, p. 91],  $E_4^j$  can be chosen so that  $(1 - u)^d < 1 + E_4^j < (1 + u)^d$ . Lemma A.1 gives

$$(A.6) \quad \text{fl}(p_d) = \frac{1}{2} \sum \log \frac{1 + 4m_j^4}{(1 + m_j^2)^2} \nu_f(I_j^d) (1 + E_a^j) + \frac{1}{2} \sum \nu_f(I_j^d) E_b^j$$

with  $(1 - u)^{d+2} < 1 + E_a^j < (1 + u)^{d+2}$  and  $|E_b^j| < 33u(1 + u)^{d+1}$ .

Bounding  $|\text{fl}(p_d) - p_d|$  is now a simple matter:

$$\begin{aligned} |\text{fl}(p_d) - p_d| &< \frac{1}{2} \sum \left| \log \frac{1 + 4m_j^4}{(1 + m_j^2)^2} \nu_f(I_j^d) |E_a^j - 1| + \frac{1}{2} \sum \nu_f(I_j^d) |E_b^j| \right. \\ &< \frac{\log 4}{4} ((1 + u)^{d+2} - 1) + \frac{33}{4} u(1 + u)^{d+1} \\ &< \frac{\log 4}{4} (e^{u(d+2)} - 1) + \frac{33}{4} u e^{u(d+1)}. \end{aligned}$$

The second inequality above uses  $\sum \nu_f(I_j^d) = 1/2$ ,  $|\log \frac{1+4m_j^4}{(1+m_j^2)^2}| < \log 4$ ,  $|E_a^j - 1| < (1 + u)^{d+2} - 1$ , and  $|E_b^j| < 33u(1 + u)^{d+1}$ . The bound on  $|E_a^j - 1|$  can be derived easily from  $(1 - u)^{d+2} < 1 + E_a^j < (1 + u)^{d+2}$ . The final inequality follows from Lemma A.2b.  $\square$

Upper bounding  $|\text{fl}(q_d) - q_d|$  involves a small, additional detail. For the rightmost positive Stern-Brocot interval  $I_j^d$ ,  $\text{amp}(m)$  is maximum at  $m = \infty$ . This causes no difficulty, however, because  $\log((1 + 4m^4)/(1 + m^2)^2)$  is taken as  $\log 4$  at  $m = \infty$  by the computation, and as a result, the bounds in (A.4) still hold.

A program to compute  $p_d$  and  $q_d$  is given below so that the computation leading to (4.2) can be easily reproduced. The program uses up 1.1 gigabytes of memory. It can be written using only a small amount of memory, but then it would be harder to read. For finding logs, we used the version of Tang's algorithm [38] that does not precompute and store  $1/F$  for  $F = 1 + j2^{-7}$ ,  $0 \leq j \leq 128$ . Though we do not give the code here because it is machine dependent, the guidelines given in [38] are enough to reproduce that log function (called `tlog()` in the program) exactly.

```
#include <stdlib.h>
#include <stdio.h>
#define D 28
#define N 268435456
#define NRT 16384
unsigned int filter = 0xAAAAAAA;
```

```
#define bitcount(x,b) \
```

```

{b = 0;                \
 for( ; x!=0; x&=(x-1)) \
   b++;                \
}
double tlog(double);
double sum(double *, int);

static double table[D+1] = {
  3.51792099313013395856e-7,
  2.17419474349120812252e-7,
  1.34372624963892583604e-7,
  8.30468493852282286483e-8,
  5.13257755786643549553e-8,
  3.17210738065638736930e-8,
  1.96047017721004812623e-8,
  1.21163720344633924307e-8,
  7.48832973763708883155e-9,
  4.62804229682630359918e-9,
  2.86028744081078523237e-9,
  1.76775485601551836682e-9,
  1.09253258479526686555e-9,
  6.75222271220251501272e-10,
  4.17310313575015364275e-10,
  2.57911957645236136997e-10,
  1.59398355929779227278e-10,
  9.85136017154569097184e-11,
  6.08847542143223175599e-11,
  3.76288475011345921584e-11,
  2.32559067131877254014e-11,
  1.43729407879468667570e-11,
  8.88296592524085864439e-12,
  5.48997486270600811265e-12,
  3.39299106253485053174e-12,
  2.09698380017115758091e-12,
  1.29600726236369295083e-12,
  8.00976537807464630088e-13,
  4.95030724556228320737e-13};

void main()
{
  int i,j,*num;
  double lower,upper,larray1[NRT],larray2[NRT],
          uarray1[NRT],uarray2[NRT];
  unsigned int *lptr, *uptr;

  num = (int *)malloc(sizeof(int)*(N+1));
  num[0] = 1; num[1]=1;

```

```

for(i=2;i<N;i=i+2){
    num[i] = num[i/2];
    num[i+1] = num[i/2]+num[i/2+1];}
num[N] = 1;

for(i=0; i<NRT; i++){
    unsigned int k,b,x; double m, m2, m2p1,
        left, right, measure;

    k = i*NRT; m =(double)num[k]/(double)num[N-k];
    m2 = m*m; m2p1 = m2+ 1.0;
    left = tlog((1+4*m2*m2)/(m2p1*m2p1));

    if (i < NRT/4)
        for(j=0; j<NRT; j++){
            k = i*NRT+j;
            m = (double)num[k+1]/(double)num[N-k-1];
            m2 = m*m;
            m2p1 = 1 + m2;
            right = tlog((1+4*m2*m2)/(m2p1*m2p1));
            x = k^filter;
            bitcount(x,b);
            measure = table[b];
            larray1[j] = measure*right; uarray1[j] = measure*left;
            left = right;}
    else if(i < NRT-1)
        for(j=0;j<NRT;j++){
            k = i*NRT+j;
            m = (double)num[k+1]/(double)num[N-k-1];
            m2 = m*m;
            m2p1 = 1 + m2;
            right = tlog((1+4*m2*m2)/(m2p1*m2p1));
            x = k^filter;
            bitcount(x,b);
            measure = table[b];
            larray1[j] = measure*left; uarray1[j] = measure*right;
            left = right;}
    else /* i == NRT-1 */
        for(j=0; j<NRT;j++){
            k = i*NRT+j;
            if(j==NRT-1)
                right = tlog(4.0);
            else{
                m = (double)num[k+1]/(double)num[N-k-1];
                m2 = m*m;
                m2p1 = 1 + m2;
                right = tlog((1+4*m2*m2)/(m2p1*m2p1));}
            x = k^filter;
            bitcount(x,b);

```

```

    measure = table[b];
    larray1[j] = measure*left; uarray1[j] = measure*right;
    left = right;}
larray2[i] = sum(larray1,NRT); uarray2[i] = sum(uarray1,NRT);}

lower = sum(larray2,NRT);
upper = sum(uarray2,NRT);

lptr = (unsigned int *)&lower;
uptr = (unsigned int *)&upper;
printf("(l,r)= (%.17E, %.17E)\n",lower, upper);
printf("(l,u) in hex = (%x %x, %x %x)\n",*lptr,*lptr+1,*uptr,
      *(uptr+1));
}

/* sums a list, length being a power of 2 */
double sum(double *list, int length)
{
    int i,step;

    for(step = 1; step < length; step = 2*step)
        for(i=0; i < length; i += 2*step)
            list[i]+= list[i+step];

    return list[0];
}

```

#### ACKNOWLEDGMENTS

I thank Nick Trefethen for showing great interest in this work. Discussions with Persi Diaconis, Harry Kesten, Bob Strichartz, and Nick Trefethen were immensely helpful. I thank Jason Hickey for letting me use his Linux machine.

#### REFERENCES

- [1] G. Álefeld and J. Herzberger, *Introduction to Interval Computations*, Academic Press, New York, 1983. MR **85d**:65001
- [2] L. Appel and W. Haken, The solution of the four-color map problem, *Scientific American* *237*(4) (Oct. 1977), 108-121. MR **58**:27598e
- [3] L. Arnold and H. Crauel, Random dynamical systems, in *Lyapunov Exponents*, eds. L. Arnold, H. Crauel, J.-P. Eckmann, Lecture Notes in Math. 1486, Springer-Verlag, Berlin, 1991, 1-22. MR **93m**:60116
- [4] R. Bellman, Limit theorems for non-commutative operations, *I. Duke Math. J.* *21* (1954), 491-500. MR **15**:969e
- [5] M.A. Berger, *An Introduction to Probability and Stochastic Processes*, Springer-Verlag, Berlin, 1993. MR **93k**:60001
- [6] L. Blum, F. Cucker, M. Shub and S. Smale, *Complexity and Real Computation*, Springer-Verlag, New York, 1998. MR **99a**:68070
- [7] P. Bougerol and J. Lacroix, *Random Products of Matrices with Applications to Schrödinger Operators*, Birkhauser, Basel, 1985. MR **88f**:68013
- [8] L. Breiman, *Probability*, SIAM, Philadelphia, 1992. MR **93d**:60001

- [9] R.P. Brent, On the zeros of the Riemann zeta function in the critical strip, *Math. of Comp.* 33 (1979), 1361-1372. MR 80g:10033
- [10] A. Brocot, Calcul des rouages par approximation nouvelle méthode, *Revue Chronometrique* 6 (1860), 186-194.
- [11] P. Chassaing, G. Letac and M. Mora, Brocot sequences and random walks on  $SL_2(\mathbb{R})$ , in *Probability measures on groups* 7, ed. H. Heyer, Lecture Notes in Math. 1064, Springer-Verlag, Berlin, 1984, 36-48. MR 86g:60012
- [12] E. Clarke and J. Wing, Formal methods: state of the art and future directions, *ACM Computing Surveys* 28(4) (1996), 626-643.
- [13] T.H. Cormen, C.E. Leiserson and R.L. Rivest, *Introduction to Algorithms*, MIT Press, Cambridge, Massachusetts, (1990). MR 91i:68001
- [14] A. Crisanti, G. Paladin and A. Vulpiani, *Products of Random Matrices in Statistical Physics*, Springer-Verlag, Berlin, 1993. MR 95d:82031
- [15] P. Diaconis and M. Shahshahani, Products of random matrices and computer image generation, in *Random Matrices and their Applications*, eds. J.E. Cohen, H. Kesten and C.M. Newman, Contemp. Math., vol. 50, American Mathematical Society, Providence, 1986, 173-182. MR 88d:60153
- [16] M. Embree and L.N. Trefethen, *Growth and decay of random Fibonacci sequences*, Proc. Royal Soc. London Series A, to appear.
- [17] K. Falconer, *Fractal Geometry, Mathematical Foundations and Applications*, John Wiley and Sons, New York, 1990. MR 92j:28008
- [18] H. Furstenberg, Non-commuting random products, *Trans. Amer. Math. Soc.* 108 (1963), 377-428. MR 29:648
- [19] H. Furstenberg and H. Kesten, Products of random matrices, *Ann. Math. Stat.* 31 (1960), 457-469. MR 22:12558
- [20] R. Graham, D. Knuth and O. Patashnik, *Concrete Mathematics*, Addison-Wesley, Reading, Massachusetts, 1994. MR 97d:68003
- [21] Y. Guivarc'h and A. Raugi, Frontière de Furstenberg, propriétés de contraction et théoèmes de convergence, *Zeit. für Wahrsch. und Verw. Gebiete.* 69 (1985), 187-242. MR 86h:60126
- [22] N.J. Higham, *Accuracy and Stability of Numerical Algorithms*, SIAM, Philadelphia, 1996. MR 97a:65047
- [23] F.Y. Hunt and W.M. Miller, On the approximation of invariant measures, *J. Stat. Phys.* 66(1/2) (1992), 535-548. MR 93a:58105
- [24] *IEEE Standard for Binary Floating-Point Arithmetic, ANSI/IEEE Standard 754-1985*, Institute of Electrical and Electronics Engineers, New York, 1985. Reprinted in *SIGPLAN Notices* 22(2) (1987), 9-25.
- [25] A. Ya. Khintchine, *Continued Fractions*, P. Noordhoff, Groningen, 1963. MR 28:5038
- [26] J.F.C. Kingman, Subadditive ergodic theory, *Ann. Prob.* 1 (1973), 883-909. MR 50:8663
- [27] J.R. Kinney and T.S. Pitcher, The dimension of some sets defined in terms of  $f$ -expansions, *Zeit. für Wahr. und Verw. Gebiete* 4 (1966), 293-315. MR 33:6670
- [28] V. Krishnamurthy, The four color theorem, Appendix IV in F. Harary, *Graph Theory*, Indian student edition, Narosa/Addison-Wesley, New Delhi, 1988.
- [29] O.E. Lanford, A computer assisted proof of the Feigenbaum conjectures, *Bull. Amer. Math. Soc.* 6 (1982), 427-434. MR 83g:58051
- [30] F. Ledrappier, Quelques propriétés des exposants caractéristiques, in *Ecole d'été de Saint-Flour 12-1982*, ed P.L. Hennequin, Lecture Notes in Math. 1097, Springer-Verlag, Berlin, 1984. MR 88b:58081
- [31] R. Lima and M. Rahibe, Exact Lyapunov exponent for infinite products of random matrices, *J Phys. A: Math. Gen.* 27 (1994), 3427-3437. MR 95d:82004
- [32] K. Mischaikow and M. Mrozek, Chaos in the Lorenz equations: a computer assisted proof, *Bull. Amer. Math. Soc. (N.S.)* 33 (1995), 66-72. MR 95e:58121
- [33] K. Mischaikow and M. Mrozek, Chaos in the Lorenz equations: a computer assisted proof. Part II: details, *Math. Comp.* 67 (1998), 1023-1046. MR 98m:58095
- [34] V.I. Oseledec, A multiplicative ergodic theorem, *Trans. Moscow Math. Soc.* 19 (1968), 197-231. MR 39:1629
- [35] Y. Peres, Analytic dependence of Lyapunov exponents on transition probabilities, in *Lyapunov Exponents*, ed L. Arnold, Lecture Notes in Math. 1486, Springer-Verlag, Berlin, 1986, 64-80. MR 94c:60116

- [36] M.A. Stern, Ueber eine zahlentheoretische Funktion, *J. für die reine und angewandte Mathematik* 55 (1858), 193-220.
- [37] R.S. Strichartz, A. Taylor and T. Zhang, Densities of self-similar measures on the line, *Experimental Mathematics* 4(2) (1995), 101-128. MR 97c:28014
- [38] P.T.P. Tang, Table-driven implementation of the logarithm function for IEEE floating-point arithmetic, *ACM Trans. Math. Soft.* 16(4) (1990), 378-400.
- [39] J.N. Tsitsiklis and V.D. Blondel, The Lyapunov exponent and joint spectral radius of pairs of matrices are hard — when not impossible — to compute and to approximate, *Mathematics of Control, Signals and Systems* 10 (1997), 31-40. CMP 98:05
- [40] S. Tuljapurkar, Demographic applications of random matrix products, in *Random Matrices and their Applications*, eds. J.E. Cohen, H. Kesten and C.M. Newman, Contemp. Math., vol. 50, American Mathematical Society, Providence, 1986, 319-326. MR 87i:92050
- [41] D. Viswanath, *Lyapunov Exponents from Random Fibonacci Sequences to the Lorenz Equations*, Ph.D. thesis, Dept. of Computer Science, Cornell University, 1998.
- [42] D. Viswanath and L.N. Trefethen, Condition numbers of random triangular matrices, *SIAM J. Matrix Anal. Appl.* 19 (1998), 564-581. MR 99b:65061

MATHEMATICAL SCIENCES RESEARCH INSTITUTE, 1000 CENTENNIAL DRIVE, BERKELEY, CALIFORNIA 94720

*E-mail address:* divakar@cs.cornell.edu