

EVALUATION OF ZETA FUNCTION OF THE SIMPLEST CUBIC FIELD AT NEGATIVE ODD INTEGERS

HYUN KWANG KIM AND JUNG SOO KIM

ABSTRACT. In this paper, we are interested in the evaluation of the zeta function of the simplest cubic field. We first introduce Siegel's formula for values of the zeta function of a totally real number field at negative odd integers. Next, we will develop a method of computing the sum of a divisor function for ideals, and will give a full description for a Siegel lattice of the simplest cubic field. Using these results, we will derive explicit expressions, which involve only rational integers, for values of a zeta function of the simplest cubic field. Finally, as an illustration of our method, we will give a table for zeta values for the first one hundred simplest cubic fields.

1. INTRODUCTION

Using finite dimensionality of elliptic modular forms of weight h , Siegel [7] developed an ingenious method of computing $\zeta_K(b)$, where K is a totally real algebraic number field, $\zeta_K(s)$ is the Dedekind zeta function of K , and b is a negative odd integer. However, evaluation of values of a zeta function by means of Siegel's formula requires complicated computations in algebraic number theory, since the formula involves terminology of algebraic number theory, such as norm, trace and different of K . The problem of expressing zeta values in terms of elementary functions was first studied by Zagier [10]. Siegel's formula has been exploited by Zagier to give an elementary expression for $\zeta_K(1-2s)$, where K is a real quadratic field and s is a positive integer, which involves only rational integers and not algebraic numbers or norm of ideals. In this paper, we will be interested in expressing zeta values of a certain class of totally real cyclic cubic fields, which are called the simplest cubic fields, in terms of elementary functions.

It is well known (cf. [5, Appendix A.3]) that every cyclic cubic field can be obtained by adjoining to \mathbb{Q} a root of an irreducible polynomial

$$f(x) = x^3 + mx^2 - (m+3)x + 1,$$

where m runs over the set of rational numbers. Let K_m (or simply K) denote the cyclic cubic field corresponding to $m \in \mathbb{Q}$. Since K_m and K_{-m-3} represent the same field, we may assume that $m \geq -\frac{3}{2}$. The discriminant of the polynomial $f(x)$

Received by the editor July 25, 2000 and, in revised form, September 26, 2000.

2000 *Mathematics Subject Classification.* Primary 11R42; Secondary 11R16.

Key words and phrases. The simplest cubic field, zeta function, Siegel lattice.

The present studies were supported by the Korea Research Foundation Grant (KRF-97-001-D00011-D1101) and Com²MaC-KOSEF.

is D^2 , where $D = m^2 + 3m + 9$. Let ρ be the negative root of $f(x)$. Then

$$\rho' = \frac{1}{1 - \rho}, \quad \rho'' = 1 - \frac{1}{\rho}$$

are the other roots of $f(x)$ so that $K = \mathbb{Q}(\rho)$ is a cyclic cubic field. The terminology “simplest cubic field” goes back to a work of Shanks [6]. He studied the arithmetic of a family of cyclic cubic fields which corresponds to $m \in \mathbb{Z}$ such that $D = m^2 + 3m + 9$ is a prime, and he called these fields the simplest cubic fields. The notion was extended by Washington [8] in which he studied the arithmetic of a family of cyclic cubic fields which corresponds to $m \in \mathbb{Z}, m \not\equiv 3 \pmod{9}$. The simplest cubic field in the sense of this paper means that it corresponds to $m \in \mathbb{Z}$ such that $D = m^2 + 3m + 9$ is square-free. In this case, we have

Proposition 1.1. *Let $m(\geq -1)$ be an integer such that $D = m^2 + 3m + 9$ is square-free. Then $\{1, \rho, \rho^2\}$ forms an integral basis of K and $\{-1, \rho, \rho'\}$ generates the full unit group of K .*

Proof. See [8]. □

In this paper, we shall apply Siegel’s formula to the simplest cubic field K to obtain an elementary expression of $\zeta_K(1 - 2s)$. In Section 2, we will introduce Siegel’s formula and the notion of a Siegel lattice. In Section 3, we will express the sum of an ideal divisor function $\sigma_r(\mathfrak{A})$ in terms of the usual sum of divisor function $\sigma_r(n)$. In Section 4, we shall describe a Siegel lattice for the simplest cubic field. In Section 5, we will obtain a formula for the values of the zeta function of K which involves only rational integers. Finally, as an illustration of our computation, we will compute $\zeta_K(-1)$, $\zeta_K(-3)$, and $\zeta_K(-5)$ for the first one hundred values of corresponding m ’s.

2. SIEGEL’S FORMULA AND A SIEGEL LATTICE

In this section, we first state Siegel’s formula for values of the zeta function of a totally real algebraic number field at negative odd integers. Next, we discuss what is needed to apply Siegel’s formula for the computation of values of the zeta function. Finally, we introduce the notion of a Siegel lattice which will be crucial in our computation.

Let K be an algebraic number field and \mathcal{O}_K be the ring of integers of K . For an ideal \mathfrak{A} of \mathcal{O}_K , we define the sum of divisors function $\sigma_r(\mathfrak{A})$ by setting

$$(1) \quad \sigma_r(\mathfrak{A}) = \sum_{\mathfrak{B}|\mathfrak{A}} N_{K/\mathbb{Q}}(\mathfrak{B})^r,$$

where \mathfrak{B} runs over all ideals of \mathcal{O}_K which divide \mathfrak{A} . Note that, if $K = \mathbb{Q}$ and $\mathfrak{A} = (n)$, our definition coincides with the usual sum of the divisor function

$$(2) \quad \sigma_r(n) = \sum_{\substack{d|n \\ d>0}} d^r.$$

Now let K be a totally real algebraic number field. For $l, s = 1, 2, \dots$, we define

$$(3) \quad S_l^K(2s) = \sum_{\substack{\nu \in \delta^{-1} \\ \nu \gg 0 \\ \text{tr}(\nu) = l}} \sigma_{2s-1}((\nu)\delta),$$

where δ denotes the different of K . Later we shall study the sum (3) intensively. At this moment, we remark that this is a finite sum.

We now state Siegel’s formula.

Theorem 2.1 (Siegel). *Let $s = 1, 2, \dots$, be a natural number, K a totally real algebraic number field of degree n , and $h = 2sn$. Then*

$$(4) \quad \zeta_K(1 - 2s) = 2^n \sum_{l=1}^r b_l(h) S_l^K(2s).$$

The numbers $r \geq 1$ and $b_1(h), \dots, b_r(h) \in \mathbb{Q}$ depend on h . In particular,

$$(5) \quad r = \dim_{\mathbb{C}} \mathfrak{M}_h,$$

where \mathfrak{M}_h denotes the space of modular forms of weight h . Thus by a well-known formula,

$$r = \begin{cases} \lfloor \frac{h}{12} \rfloor & \text{if } h \equiv 2 \pmod{12}, \\ \lfloor \frac{h}{12} \rfloor + 1 & \text{if } h \not\equiv 2 \pmod{12}. \end{cases}$$

Proof. See [7] or [10]. □

Remark. By applying (4) to the simplest cubic field K , we obtain

$$(6) \quad \zeta_K(-1) = 2^3 * b_1(6) * S_1^K(2),$$

$$(7) \quad \zeta_K(-3) = 2^3 * [b_1(12) * S_1^K(4) + b_2(12) * S_2^K(4)],$$

$$(8) \quad \zeta_K(-5) = 2^3 * [b_1(18) * S_1^K(6) + b_2(18) * S_2^K(6)].$$

Zagier [10] contains a table for values of Siegel coefficients $b_l(h)$ for $4 \leq h \leq 40$. We quote the values of Siegel coefficients which will be necessary in our computation:

$$(9) \quad b_1(6) = -\frac{1}{504},$$

$$(10) \quad b_1(12) = -\frac{1}{8190}, \quad b_2(12) = \frac{1}{196560},$$

$$(11) \quad b_1(18) = -\frac{22}{3591}, \quad b_2(18) = -\frac{1}{86184}.$$

The essence of Siegel’s formula is that it transforms an infinite series (i.e., the value of a zeta function) into finite sums involving $S_l^K(2s)$ which itself is a finite sum of powers of divisors of ideal $((\nu)\delta)$ over the ν ’s in K which satisfy the Siegel conditions described in (3). Therefore we need to establish the following two items to compute $S_l^K(2s)$:

- (i) the method of computing the sum of a divisor function $\sigma_r(\mathfrak{A})$ for an integral ideal \mathfrak{A} ,
- (ii) the description of ν ’s in K which satisfy Siegel conditions described in (3).

In Section 3, we shall develop a method of computing the sum of divisor function $\sigma_r(\mathfrak{A})$ when K is a cyclic extension of \mathbb{Q} of prime degree. In Section 4, we shall give a full description of ν ’s in K which satisfy Siegel conditions when K is the simplest cubic field. At this moment, we examine the sum in equation (3) more closely for an arbitrary totally real algebraic number field K , and introduce the notion of a Siegel lattice which is first studied in [3].

Let K be a totally real algebraic number field of degree n and S_K (or simply S) be the set of elements in K which satisfy Siegel conditions described in (3). Fix an integral basis $\{\alpha_1, \dots, \alpha_n\}$ of K . For $\nu \in K$, we can write

$$(12) \quad \nu = x_1\alpha_1 + \dots + x_n\alpha_n, \quad x_i \in \mathbb{Q},$$

and we have an embedding $\phi : K \rightarrow \mathbb{R}^n$ given by

$$(13) \quad \phi(\nu) = (x_1, \dots, x_n).$$

The condition $\nu \in \delta^{-1}$ implies that the denominator of $x_i, i = 1, \dots, n$, is bounded by D_K where D_K denotes the discriminant of K . The condition $\text{tr}(\nu) = l$ is equivalent to saying that $\phi(\nu)$ lies in the hyperplane

$$(14) \quad a_1x_1 + \dots + a_nx_n = l,$$

where $a_i = \text{tr}_{K/\mathbb{Q}}(\alpha_i)$. Finally the condition $\nu \gg 0$ becomes n distinct linear inequalities defined over K in the variables x_1, \dots, x_n . Therefore the elements ν in S can be put in one-to-one correspondence to the lattice points in a bounded $(n-1)$ -dimensional region under ϕ . We shall call this lattice (or any set which can be put in one-to-one correspondence with this set under a suitable linear transformation) as a Siegel lattice for K and denote it by T_K (or simply T). Notice that equation (3) expresses $S_l^K(2s)$ as a weight sum of divisor functions over a Siegel lattice. Hence the description of the Siegel lattice is of crucial importance in the computation of $S_l^K(2s)$.

3. COMPUTATION OF THE SUM OF DIVISORS

In this section, we develop a method of computing the sum of the divisor function of K when K is a cyclic extension of \mathbb{Q} of prime degree.

Let K be a cyclic field of prime degree q and W denote the group of q th roots of unity and ζ be a primitive q th root of unity. We define an arithmetic function $\chi : \mathbb{N} \rightarrow W \cup \{0\}$ in the following manner.

For a prime p , we set

$$\chi(p) = \begin{cases} 0 & \text{if } p \text{ is ramified in } K/\mathbb{Q}, \\ 1 & \text{if } p \text{ splits completely in } K/\mathbb{Q}, \\ \zeta & \text{if } p \text{ is inert in } K/\mathbb{Q}, \end{cases}$$

and extend χ multiplicatively. We put χ^j by χ_j for $j = 0, 1, 2, \dots, q - 1$.

Lemma 3.1. *Let ζ be a primitive q th root of unity. Then we have*

$$\sum_{\substack{s_1 + \dots + s_{q-1} = t \\ s_1, \dots, s_{q-1} \geq 0}} \zeta^{s_1 + 2s_2 + \dots + (q-1)s_{q-1}} = \begin{cases} 1 & \text{if } t \equiv 0 \pmod{q}, \\ -1 & \text{if } t \equiv 1 \pmod{q}, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Consider the polynomial

$$\phi_i(x) = 1 + \zeta^i x + \zeta^{2i} x^2 + \dots = \sum_{k=0}^{\infty} \zeta^{ik} x^k,$$

and put

$$\phi(x) = \prod_{i=1}^{q-1} \phi_i(x).$$

By simple computation, we have

$$\phi(x) = \sum_{t=0}^{\infty} a_t x^t,$$

where

$$a_t = \sum_{\substack{s_1 + \dots + s_{q-1} = t \\ s_1, \dots, s_{q-1} \geq 0}} \zeta^{s_1 + 2s_2 + \dots + (q-1)s_{q-1}}.$$

Note that $\phi_i(x) = \frac{1}{1 - \zeta^i x}$ and

$$\begin{aligned} \phi(x) &= \prod_{i=1}^{q-1} \frac{1}{1 - \zeta^i x} \\ &= \frac{1-x}{\prod_{i=0}^{q-1} (1 - \zeta^i x)} = \frac{1-x}{1-x^q} = (1-x)(1+x^q+x^{2q}+\dots), \quad |x| < 1. \end{aligned}$$

By comparison of coefficients, we obtain

$$a_t = \begin{cases} 1 & \text{if } t \equiv 0 \pmod{q}, \\ -1 & \text{if } t \equiv 1 \pmod{q}, \\ 0 & \text{otherwise.} \end{cases}$$

□

Theorem 3.2. *Let \mathfrak{A} be an integral ideal of K . Then, for any $r \geq 0$,*

$$(15) \quad \sigma_r(\mathfrak{A}) = \sum_{(j_1 \dots j_{q-1})^2 | \mathfrak{A}^q} \chi_1(j_1) \dots \chi_{q-1}(j_{q-1}) j_1^r \dots j_{q-1}^r \sigma_r\left(\frac{N}{j_1^2 \dots j_{q-1}^2}\right),$$

where $N = \text{Norm}_{K/\mathbb{Q}}(\mathfrak{A})$ denotes the norm of \mathfrak{A} , the function σ_r on the right-hand side is the usual sum of divisors function defined by equation (2) in Section 2, and the summation is over all positive integers j_1, \dots, j_{q-1} such that $(j_1 \dots j_{q-1})^2$ divides \mathfrak{A}^q , i.e., $((j_1 \dots j_{q-1})^2) \supset \mathfrak{A}^q$.

Proof. We put $\widetilde{\sigma}_r(\mathfrak{A})$ to be the right-hand side of (15). Since $\sigma_r(\mathfrak{A})$ and $\widetilde{\sigma}_r(\mathfrak{A})$ are both multiplicative, we may assume that \mathfrak{A} is a power \mathfrak{P}^m of a prime ideal \mathfrak{P} . Let p be the rational prime lying below \mathfrak{P} . Then

$$N(\mathfrak{P}) = p^f,$$

where f is the inertial degree of \mathfrak{P} in K/\mathbb{Q} . We have

$$(16) \quad \sigma_r(\mathfrak{A}) = \sigma_r(\mathfrak{P}^m) = \sum_{s=0}^m N(\mathfrak{P}^s)^r = \sum_{s=0}^m p^{f sr} = \sigma_{fr}(p^m).$$

To evaluate $\widetilde{\sigma}_r(\mathfrak{A})$, we must distinguish three cases, according to the value of $\chi(p)$.

Case 1. $\chi(p) = 1$. Since $[K : \mathbb{Q}] = q$ is a prime, (p) splits completely in K/\mathbb{Q} , and $f = 1$. Write

$$(p) = \mathfrak{P}_1 \dots \mathfrak{P}_q,$$

where $\mathfrak{P}_1 = \mathfrak{P}$ and $\mathfrak{P}_i, i = 2, \dots, q$, are the conjugates of \mathfrak{P} . If $j | \mathfrak{A}^q$, then $j | \mathfrak{P}_i^{mq}$ for each i . So,

$$j^q | p^{mq}.$$

This means that j is a power of p . Since p splits completely, we must have $j = 1$. Hence $j_1 = \dots = j_{q-1} = 1$ is the only term on the right-hand side of (16). Therefore we have

$$\widetilde{\sigma}_r(\mathfrak{A}) = \widetilde{\sigma}_r(\mathfrak{P}^m) = \sigma_r(N) = \sigma_r(p^m),$$

and this coincides with (16) since $f = 1$.

Case 2. $\chi(p) = 0$. Since p is ramified in K/\mathbb{Q} , $(p) = \mathfrak{P}^q$ and $f = 1$. If $j|\mathfrak{A}$, then $j|\mathfrak{P}^m$. So $j^q|p^m$, which implies that j is a power of p . Since $\chi(p) = 0$, the only term in $\widetilde{\sigma}_r(\mathfrak{A})$ that does not vanish is the term corresponding to $j_1 = \dots = j_{q-1} = 1$, namely, $\sigma_r(N)$. Therefore, we have

$$\widetilde{\sigma}_r(\mathfrak{A}) = \sigma_r(N) = \sigma_r(p^m),$$

and this coincides with (16) since $f = 1$.

Case 3. $\chi(p) = \zeta$, a primitive q th root of unity. Since p is inert in K/\mathbb{Q} , $\mathfrak{P} = (p)$, and $f = q$. By definition,

(17)

$$\widetilde{\sigma}_r(\mathfrak{A}) = \widetilde{\sigma}_r((p^m)) = \sum_{(j_1 \cdots j_{q-1})^2 | p^{mq}} \chi_1(j_1) \cdots \chi_{q-1}(j_{q-1}) j_1^r \cdots j_{q-1}^r \sigma_r\left(\frac{N}{j_1^2 \cdots j_{q-1}^2}\right).$$

Write $j_i = p^{s_i}$, $i = 1, \dots, q - 1$. Then (17) becomes

(18)

$$\widetilde{\sigma}_r(\mathfrak{A}) = \sum_{\substack{2(s_1 + \cdots + s_{q-1}) \leq mq \\ s_i \geq 0}} p^{r(s_1 + \cdots + s_{q-1})} \zeta^{s_1} \zeta^{2s_2} \cdots \zeta^{(q-1)s_{q-1}} \sigma_r(p^{mq-2(s_1 + \cdots + s_{q-1})}).$$

Furthermore, from (18) it follows that

$$(19) \quad \widetilde{\sigma}_r(\mathfrak{A}) = \sum_{t=0}^{\lfloor \frac{mq}{2} \rfloor} \sum_{\substack{s_1 + \cdots + s_{q-1} = t \\ s_i \geq 0}} p^{rt} \sigma_r(p^{mq-2t}) \zeta^{s_1 + 2s_2 + \cdots + (q-1)s_{q-1}}.$$

Finally, we get

$$(20) \quad \widetilde{\sigma}_r(\mathfrak{A}) = \sum_{t=0}^{\lfloor \frac{mq}{2} \rfloor} p^{rt} \sigma_r(p^{mq-2t}) \sum_{\substack{s_1 + \cdots + s_{q-1} = t \\ s_i \geq 0}} \zeta^{s_1 + 2s_2 + \cdots + (q-1)s_{q-1}}.$$

Now we consider two cases, say m is even or m is odd. We only give a proof for the case that m is even since the other case can be treated similarly. Write $m = 2m'$. Then $\lfloor \frac{mq}{2} \rfloor = m'q$. By Lemma 3.1, (20) becomes

$$\begin{aligned} \widetilde{\sigma}_r(\mathfrak{A}) &= \sum_{t=0}^{m'q} p^{rt} \sigma_r(p^{2m'q-2t}) a_t \\ &= \sum_{\substack{t=qt' \\ 0 \leq t' \leq m'}} \frac{(p^r)^{2m'q-qt'+1} - (p^r)^{qt'}}{p^r - 1} - \sum_{\substack{t=qt'+1 \\ 0 \leq t' \leq m'-1}} \frac{(p^r)^{2m'q-qt'} - (p^r)^{qt'+1}}{p^r - 1}. \end{aligned}$$

So, we get

$$\widetilde{\sigma}_r(\mathfrak{A}) = \frac{1}{p^r - 1} \left\{ \sum_{l=0}^{2m'} (p^r)^{ql+1} - \sum_{l=0}^{2m'} (p^r)^{ql} \right\} = \sigma_{rq}(p^m).$$

This agrees with (16) since $f = q$. □

Remark 3.1. When $q = 2$, i.e., K is a real quadratic field, the equation (15) becomes the formula obtained by Zagier [10].

4. DESCRIPTION OF A SIEGEL LATTICE FOR THE SIMPLEST CUBIC FIELDS

In this section, we shall give a full description of a Siegel lattice for the simplest cubic field. As a result, we derive a formula for the number of points in a Siegel lattice.

Let $m(\geq -1)$ be an integer such that $m^2 + 3m + 9$ is square-free and K be the simplest cubic field defined by the irreducible polynomial

$$(21) \quad f(x) = x^3 + mx^2 - (m + 3)x + 1.$$

Recall that the discriminant d_K , the ring of integers \mathcal{O}_K , and the different δ_K of K are given, respectively, by

$$(22) \quad d_K = D^2 = (m^2 + 3m + 9)^2,$$

$$(23) \quad \mathcal{O}_K = \mathbb{Z}[\rho] = \mathbb{Z} \oplus \mathbb{Z}\rho \oplus \mathbb{Z}\rho^2,$$

$$(24) \quad \delta_K = (f'(\rho)) = -(m + 3) + 2m\rho + 3\rho^2,$$

where ρ denotes the negative root of $f(x)$. Let ν be an element of K . We can write

$$(25) \quad \nu = \alpha + \beta\rho + \gamma\rho^2, \quad \alpha, \beta, \gamma \in \mathbb{Q}.$$

Now suppose that ν satisfies Siegel conditions, i.e.,

$$(26) \quad \nu \in \delta^{-1}, \quad \nu \gg 0, \quad \text{tr}(\nu) = l.$$

1. $\nu \in \delta^{-1}$

$$\nu \in \delta^{-1} \iff \nu(-(m + 3) + 2m\rho + 3\rho^2) \in \mathcal{O}_K.$$

Hence we can write

$$(27) \quad \nu(-(m + 3) + 2m\rho + 3\rho^2) = A + B\rho + C\rho^2,$$

with $A, B, C \in \mathbb{Z}$.

From (25),(27), we obtain the following system of linear equations:

$$(28) \quad -(m + 3)\alpha - 3\beta + m\gamma = A,$$

$$(29) \quad 2m\alpha + 2(m + 3)\beta + (-m^2 - 3m - 3)\gamma = B,$$

$$(30) \quad 3\alpha - m\beta + (m^2 + 2m + 6)\gamma = C.$$

Using Cramer’s rule, it follows that

$$(31) \quad \alpha = \frac{a}{D}, \quad \beta = \frac{b}{D}, \quad \gamma = \frac{c}{D}, \quad (a, b, c) \in \Lambda,$$

where Λ is a free module of rank 3 in \mathbb{Z}^3 and $D = m^2 + 3m + 9$.

By substitution of (31) into (28),(29),(30), we finally have

$$(32) \quad -(m + 3)a - 3b + mc = DA \equiv 0 \pmod{D},$$

$$(33) \quad 2ma + 2(m + 3)b - (m^2 + 3m + 3)c = DB \equiv 0 \pmod{D},$$

$$(34) \quad 3a - mb + (m^2 + 2m + 6)c = DC \equiv 0 \pmod{D}.$$

2. $\text{tr}(\nu) = l$

$$\text{tr}(\nu) = l \iff 3\alpha - m\beta + (m^2 + 2m + 6)\gamma = l.$$

From (31),(34), it follows that

$$(35) \quad C = l, b = \frac{3a + (m^2 + 2m + 6)c - lD}{m}.$$

By substitution of (35) into (32), we have

$$(36) \quad -a + 3l - 2c = mA.$$

In particular, m divides $a + 2c - 3l$. Now we introduce a new variable t by the formula

$$(37) \quad t = \frac{a + 2c - 3l}{m}.$$

By substitution of (37) into (35), we have

$$b = 3t + (m + 2)c - l(m + 3).$$

3. $\nu \gg 0$

$$\nu \gg 0 \iff D\nu = a + b\rho + c\rho^2 \gg 0.$$

This condition becomes three linear inequalities in the variables a, b, c defined over K . Using (35),(37), we have the following system of linear inequalities in the variables c, t defined over K :

$$(38) \quad (\rho^2 + (m + 2)\rho - 2)c + (m + 3\rho)t + l(3 - (m + 3)\rho) > 0,$$

$$(39) \quad (\rho'^2 + (m + 2)\rho' - 2)c + (m + 3\rho')t + l(3 - (m + 3)\rho') > 0,$$

$$(40) \quad (\rho''^2 + (m + 2)\rho'' - 2)c + (m + 3\rho'')t + l(3 - (m + 3)\rho'') > 0.$$

Let L_1 (resp., L_2, L_3) denote the line in (c, t) -plane defined by the left-hand side of (38) (resp., (39),(40)). By simple computation, we obtain

$$(41) \quad ((-\rho + \rho')l, \frac{l}{\rho''}) \text{ is the point of intersection of } L_1 = L_2 = 0,$$

$$(42) \quad ((-\rho' + \rho'')l, \frac{l}{\rho}) \text{ is the point of intersection of } L_2 = L_3 = 0,$$

$$(43) \quad ((-\rho'' + \rho)l, \frac{l}{\rho'}) \text{ is the point of intersection of } L_3 = L_1 = 0,$$

We summarize the above computation as in the following proposition.

Proposition 4.1. *Let $m(\geq -1)$ be an integer such that $D = m^2 + 3m + 9$ is square-free, and K be the simplest cubic field defined by equation (21). Let S be the set of elements in K which satisfy Siegel conditions described by equation (26) and T be the set of integral points in (c, t) -plane which lie inside of the triangle surrounded by the lines $L_1 = 0, L_2 = 0,$ and $L_3 = 0$. For $\nu \in S$, by equation (31), we can write*

$$(44) \quad \nu = \frac{a}{D} + \frac{b}{D}\rho + \frac{c}{D}\rho^2, \quad a, b, c \in \mathbb{Z}.$$

Then the mapping $\eta : S \rightarrow T$ given by $\eta(\nu) = (c, t)$, where

$$(45) \quad c = c, t = \frac{a + 2c - 3l}{m}$$

gives a one-to-one correspondence between S and T . The inverse mapping $\tau : T \rightarrow S$ is given by

$$\tau(c, t) = \nu = \frac{a}{D} + \frac{b}{D}\rho + \frac{c}{D}\rho^2,$$

where $a = mt - 2c + 3l$, $b = 3t + (m + 2)c - l(m + 3)$. □

Remark 4.1. A straightforward calculation shows that $\nu = \tau(c, t)$ satisfies equation (27) with A, B, C in \mathbb{Z} .

Example 4.1. As an illustration of our discussion, we describe the Siegel lattice T for the simplest cubic field K with $m = 8$. We first note that $g_l(x) = x^3 - l^2Dx + l^3D$ (resp., $h_l(x) = x^3 - l(m + 3)x^2 + ml^2x + l^3$) is the cubic polynomial whose roots are the conjugates of $(-\rho + \rho')l$ (resp., $\frac{l}{\rho}$). By estimating the roots of $g_l(x)$ and $h_l(x)$, we can find the rough location of vertices of the triangle. For the roots of $g_l(x)$, we have

$$\begin{aligned} (-m - 3)l < (-\rho'' + \rho)l < (-m - 2)l \\ < (-\rho' + \rho'')l < (m + 1)l < (-\rho + \rho')l < (m + 2)l. \end{aligned}$$

Similarly, we obtain

$$-l < \frac{l}{\rho} < 0 < \frac{l}{\rho''} < l \leq (m + 2)l < \frac{l}{\rho'} < (m + 3)l.$$

For $(c, t) \in T$, the corresponding ν in S is given by

$$(46) \quad \nu = \frac{a}{D} + \frac{b}{D}\rho + \frac{c}{D}\rho^2,$$

where $a = mt - 2c + 3l$, $b = 3t + (m + 2)c - (m + 3)l$, $c = c$. Since

$$\delta = (m + 3 - 2m\rho - 3\rho^2),$$

it follows from a simple computation that

$$(47) \quad (\nu)\delta = (t + (-2t - c + 2l)\rho - l\rho^2),$$

where $b = 2t + c - 2l$. Let $N(c, t)$ denote the norm function $N_{K/\mathbb{Q}}((\nu)\delta)$. By elementary computation, we obtain

$$(48) \quad \begin{aligned} N(c, t) = & [lt^2 + (c - l)lt]m^2 \\ & + [-2t^3 + (-3c + 6l)t^2 + (-c^2 + 3lc)t + (-lc^2 + 3l^2c - 2l^3)]m \\ & + [-3t^3 + (3c^2 - 9lc + 9l^2)t + (c^3 - 6lc^2 + 9l^2c - 3l^3)]. \end{aligned}$$

Note that a point (c, t) in the plane near the boundary the triangle lies inside the triangle if and only if $N(c, t) > 0$. Combining these data, we conclude that the Siegel lattice for K is given as in Figures 1 and 2.

We now describe the Galois action on a Siegel lattice. We start from the following simple observation.

Lemma 4.2. *Let K be a totally real Galois extension of \mathbb{Q} with Galois group G . If $\nu \in K$ satisfies the Siegel conditions described in equation (26), then so does $\sigma(\nu)$ for $\sigma \in G$.*

Proof. This is clear! The most important thing is to realize that this is an important fact. □




FIGURE 1. Siegel lattice for $m = 8$ with $\text{tr}(\nu) = 1$.




FIGURE 2. Siegel lattice for $m = 8$ with $\text{tr}(\nu) = 2$.

By Lemma 4.2, the Galois group $G = \text{Gal}(K/\mathbb{Q})$ acts on the set S and S can be put into one-to-one correspondence with the Siegel lattice T . Therefore, we have the induced Galois action on T . Now we return to the simplest cubic field case and describe the Galois action on T .

Proposition 4.3 (Galois action on a Siegel lattice). *Let $m(\geq -1)$ be an integer such that $D = m^2 + 3m + 9$ is square-free, and let K be the simplest cubic field*

defined by equation (21). Then the Galois group $G(= \langle \sigma \rangle)$ induces an action on T given by

$$(49) \quad \sigma(c, t) = (-2c - 3t + (m + 3)l, c + t).$$

If l is not divisible by 3, then every G -orbit contains three points. In particular, N_l is divisible by 3, where N_l denotes the number of lattice points in T which corresponds to $\text{tr}(\nu) = l$.

Proof. Let $(c, t) \in T$. By Proposition 4.1, it corresponds to $\nu \in S$ where ν is given by

$$D\nu = (mt - 2c + 3l) + \{3t + (m + 2)c - (m + 3)l\}\rho + c\rho^2.$$

By an actual computation, we obtain

$$\begin{aligned} D\nu' &= \{(m + 6)t + (m + 4)c - (2m + 3)l\} \\ &\quad + \{-3(m + 1)t - (2m + 1)c + (m^2 + 4m + 3)l\}\rho \\ &\quad + \{-3t - 2c + (m + 3)l\}\rho^2. \end{aligned}$$

From the transformation formula (45), it follows that $\eta(\nu') = (c', t')$, where $c' = -2c - 3t + (m + 3)l$ and $t' = c + t$. This proves the first assertion. Now suppose that the Galois action on T has a fixed point, say (c, t) . Then it follows from (49) that

$$(c, t) = (-2c - 3t + (m + 3)l, c + t).$$

Thus we must have $c = 0$ and $(m + 3)l = 3t$. Since m is not divisible by 3, we conclude that l is divisible by 3. □

We now prove the main result of this section.




FIGURE 3. Galois action on the Siegel lattice for $m = 8$ with $\text{tr}(\nu) = 1$.

Theorem 4.4. *Let $m(\geq -1)$ be an integer such that $D = m^2 + 3m + 9$ is square-free, and let K be the simplest cubic field defined by (21). Let N_l denote the number of Siegel lattice points for K which corresponds to $\text{tr}(\nu) = l$. Then we have*

$$N_1 = \begin{cases} 3\left(\frac{3s^2 + 5s + 4}{2}\right), & \text{if } m = 3s + 1, \\ 3\left(\frac{3s^2 + 7s + 6}{2}\right), & \text{if } m = 3s + 2 \end{cases}$$

and

$$N_2 = \begin{cases} 3(6s^2 + 10s + 9), & \text{if } m = 3s + 1, \\ 3(6s^2 + 14s + 13), & \text{if } m = 3s + 2. \end{cases}$$

Proof. We only give a detailed proof for the case $m = 3s + 2$, since the other case can be treated in the same manner. We assume that $m \geq 5$. (The case of $m = 2$ can be treated by direct computation.) The basic idea of the proof is to find a set of representatives of “good shape” for the Galois action on T .

First, we consider the case of $\text{tr}(\nu) = 1$. First note that $(1, 0)$ is the only point in the Siegel lattice with $t = 0$. Let l_0 be the point $(1, 0)$. Then $\sigma(l_0)$ (resp., $\sigma^2(l_0)$) is the point $(m + 1, 1)$ (resp., $(-m - 2, m + 1)$). For $1 \leq i \leq s + 1$, let l_i be the line joining $(3s + 5 - 3i, i)$ and $(1, i)$. By simple computation, we know that $\sigma(l_i)$ is the line joining $(-3s - 5 + 3i, 3s + 5 - 2i)$ and $(3s + 2 - 3i, 1 + i)$, and $\sigma^2(l_i)$ is the line joining $(0, i)$ and $(-3s - 2 + 3i, 3s + 3 - 2i)$ (see Figure 3). This proves that the set of lattice points on $\bigcup_{i=0}^{s+1} l_i$ becomes a set of representatives for the Galois action (see Figure 5). Therefore,

$$N_1 = 3\left\{1 + \sum_{i=1}^{s+1} (3s + 5 - 3i)\right\} = 3\left(\frac{3s^2 + 7s + 6}{2}\right).$$




FIGURE 4. Galois action on the Siegel lattice for $m = 8$ with $\text{tr}(\nu) = 2$.




FIGURE 5. A set of representatives for the Galois action on the Siegel lattice for $m = 8$ with $\text{tr}(\nu) = 1$.




FIGURE 6. A set of representatives for the Galois action on the Siegel lattice for $m = 8$ with $\text{tr}(\nu) = 2$.

Secondly, we consider the case of $\text{tr}(\nu) = 2$. Let t_0 be the line joining $(2, 0)$ and $(3, 0)$ and t_1 be the line joining $(1, 1)$ to $(3s + 6, 1)$. For $2 \leq i \leq 2s + 3$, let t_i be the line joining $(6s + 10 - 3i, i)$ and $(1, i)$. As in the case of $\text{tr}(\nu) = 1$, the lattice points on $\bigcup_{i=0}^{2s+3} t_i$ becomes a set of representatives for the Galois action on T (see Figures 4 and 6). Therefore,

$$N_2 = 3(2 + 3s + 6 + \sum_{i=2}^{2s+3} (6s + 10 - 3i)) = 3(6s^2 + 14s + 13).$$

□

5. VALUES OF THE ZETA FUNCTIONS

In this section, we apply the previously discussed result to the evaluation of the zeta function of the simplest cubic field. We shall derive explicit expressions for $\zeta_K(-1), \zeta_K(-3), \zeta_K(-5)$ which are elementary in the sense that they involve only rational integers and not algebraic numbers or ideals. As an illustration, we present Table 1 for values of $-21\zeta_K(-1), 8190\zeta_K(-3)$, and $-3591\zeta_K(-5)$ for the first one hundred simplest cubic fields.

Recall the definition of $S_i^K(2s)$:

$$(50) \quad S_i^K(2s) = \sum_{\substack{\nu \in \delta^{-1} \\ \nu \gg 0 \\ \text{tr}(\nu) = i}} \sigma_{2s-1}((\nu)\delta).$$

By virtue of Theorem 3.2, we have

$$(51) \quad \sigma_{2s-1}((\nu)\delta) = \sum_{(jj')^2 | ((\nu)\delta)^3} \chi(j)\overline{\chi(j')}(jj')^{2s-1} \sigma_{2s-1}\left(\frac{N}{(jj')^2}\right),$$

where N denotes the norm of ideal $(\nu)\delta$.

Lemma 5.1. *Let p be a prime or $p = 1$ such that $p^2 | ((\nu)\delta)^3$. Then only p 's such that $p | (\nu)\delta$ contribute in the sum (51).*

Proof. Suppose that $p^2 | ((\nu)\delta)^3$. If p is inert in K/\mathbb{Q} , then $p | (\nu)\delta$. If p splits in K/\mathbb{Q} , we can write $(p) = \mathfrak{P}\mathfrak{P}'\mathfrak{P}''$, and

$$(\nu)\delta = \mathfrak{P}^a \mathfrak{P}'^b \mathfrak{P}''^c \prod \mathfrak{B}_i,$$

where $(\mathfrak{B}_i, p) = 1$. Since $p^2 | ((\nu)\delta)^3$, it follows that $\min(3a, 3b, 3c) \geq 2$, and consequently we have $p | (\nu)\delta$. Finally, if p is ramified in K/\mathbb{Q} , p does not contribute in the sum (51) since the character value on the right-hand side of (51) vanishes. □

From the unique factorization of ideals into prime ideals, it follows that

$$(52) \quad \sigma_{2s-1}((\nu)\delta) = \sigma_{2s-1}((\nu')\delta), \quad \text{for every } \nu \in \delta^{-1}.$$

Now we shall compute $S_1^K(2s)$. Let T be the Siegel lattice of K which is computed in Section 4 and corresponds to $\text{tr}(\nu) = l = 1$. In Section 4, we have a one-to-one correspondence between points (c, t) in T and ideals $(\nu)\delta$, where ν is an element of K which satisfies the Siegel conditions, $\nu \in \delta^{-1}, \nu \gg 0$ and $\text{tr}(\nu) = 1$. For $(c, t) \in T$, by equation (47), the corresponding ideal is given by

$$(53) \quad (\nu)\delta = (t + (-2t - c + 2)\rho - \rho^2).$$

Let $f_m(c, t)$ be the norm of the ideal $(\nu)\delta$. By (48), it can be explicitly expressed by

$$(54) \quad \begin{aligned} f_m(c, t) = & [t^2 + (c - 1)t]m^2 \\ & + [-2t^3 + (-3c + 6)t^2 + (-c^2 + 3c)t + (-c^2 + 3c - 2)]m \\ & + [-3t^3 + (3c^2 - 9c + 9)t + (c^3 - 6c^2 + 9c - 3)]. \end{aligned}$$

Note that $p|(\nu)\delta$ if and only if $p = 1$. By Lemma 5.1 we have

$$\sigma_{2s-1}((\nu)\delta) = \sigma_{2s-1}(f_m(c, t)).$$

Thus we have

$$(55) \quad S_1^K(2s) = \sum_{(c,t) \in T} \sigma_{2s-1}(f_m(c, t)) = 3 \sum_{(c,t) \in T_0} \sigma_{2s-1}(f_m(c, t)),$$

where T_0 denotes a set of representatives for the Galois action described in the proof of Theorem 4.4.

Next we shall compute $S_2^K(2s)$. Let T denote the Siegel lattice which corresponds to $\text{tr}(\nu) = l = 2$. As in the case $l = 1$, for $(c, t) \in T$ the corresponding ideal is given by

$$(56) \quad (\nu)\delta = (t + (-2t - c + 4)\rho - 2\rho^2).$$

Therefore $p|(\nu)\delta$ if and only if $p = 1$ when c or t is odd, and $p|(\nu)\delta$ if and only if $p = 1$ or $p = 2$ when both c and t are even. Let $g_m(c, t)$ denote the norm of the ideal in (56), which is given explicitly by

$$(57) \quad \begin{aligned} g_m(c, t) = & [2t^2 + 2(c - 2)t]m^2 \\ & + [-2t^3 + (-3c + 12)t^2 + (-c^2 + 6c)t + (-2c^2 + 12c - 16)]m \\ & + [-3t^3 + (3c^2 - 18c + 36)t + (c^3 - 12c^2 + 36c - 24)]. \end{aligned}$$

If either c or t is odd, then

$$\sigma_{2s-1}((\nu)\delta) = \sigma_{2s-1}(g_m(c, t)).$$

If both c and t are even, then

$$\sigma_{2s-1}((\nu)\delta) = \sigma_{2s-1}(g_m(c, t)) + [\chi(2) + \bar{\chi}(2)]2^{2s-1}\sigma_{2s-1}\left(\frac{g_m(c, t)}{4}\right).$$

Since $f_m(x)$ is irreducible over $GF(2)$, 2 is inert in K/\mathbb{Q} . Hence $\chi(2) + \bar{\chi}(2) = -1$ by definition of χ . Therefore we have

$$(58) \quad S_2^K(2s) = 3\left[\sum_{(c,t) \in \tilde{T}_0} \sigma_{2s-1}(g_m(c, t)) - 2^{2s-1} \sum_{\substack{(c,t) \in \tilde{T}_0 \\ c, t: \text{even}}} \sigma_{2m-1}\left(\frac{g_m(c, t)}{4}\right) \right],$$

where \tilde{T}_0 denotes a set of representatives for the Galois action. By (6), (7), (8), (55), and (58), we finally have the following theorem.

Theorem 5.2. *Let $m(\geq -1)$ be an integer such that $D = m^2 + 3m + 9$ is square-free, and let K be the simplest cubic field defined by equation (21) in Section 4.*

Define elementary functions $f_m(c, t)$ and $g_m(c, t)$ by (54), (57), respectively. Then we have

$$\begin{aligned}
 -21\zeta_K(-1) &= \sum_{(c,t) \in T_0} \sigma_1(f_m(c, t)), \\
 8190\zeta_K(-3) &= -24 * \sum_{(c,t) \in T_0} \sigma_3(f_m(c, t)) \\
 &\quad + \sum_{(c,t) \in \tilde{T}_0} \sigma_3(g_m(c, t)) - 8 * \sum_{\substack{(c,t) \in \tilde{T}_0 \\ c, t: \text{even}}} \sigma_3\left(\frac{g_m(c, t)}{4}\right), \\
 -3591\zeta_K(-5) &= 528 * \sum_{(c,t) \in T_0} \sigma_5(f_m(c, t)) \\
 &\quad + \sum_{(c,t) \in \tilde{T}_0} \sigma_5(g_m(c, t)) - 32 * \sum_{\substack{(c,t) \in \tilde{T}_0 \\ c, t: \text{even}}} \sigma_5\left(\frac{g_m(c, t)}{4}\right),
 \end{aligned}$$

where T_0 (resp., \tilde{T}_0) denotes a set of representatives for the Galois action on the Siegel lattice for $\text{tr}(\nu) = l = 1$ (resp., $\text{tr}(\nu) = l = 2$). □

From Theorem 5.2, we can easily compute values of the zeta function of the simplest cubic field. As an illustration, we give Table 1 for values of $-21\zeta_K(-1)$, $8190\zeta_K(-3)$, and $-3591\zeta_K(-5)$ for the first one hundred simplest cubic fields.

We first give some remarks on our computation.

Remark 5.1. Halbritter and Pohst [2] developed a method of computing special values of a class zeta function of a totally real cubic field. Byeon [1] exploited this result to give

$$\zeta_K(-1, C) = -\frac{p(m)}{2^3 * 3^3 * 5 * 7},$$

where $p(m) = m^6 + 9m^5 + 55m^4 + 195m^3 + 544m^2 + 876m + 840$, K is the simplest cubic field corresponding to m , and C denotes the principal ideal class. We remark that $\zeta_K(-1, C) = \zeta_K(-1)$, if K has class number 1. For $m = -1, 1, 2, 4, 7, 8, 10$ which are all the values of m such that K has class number 1, our result coincides with Byeon’s result.

Remark 5.2. In [7], Siegel gave three examples for the zeta values of totally real number fields. In the last example, Siegel computed that

$$\zeta_K(2) = \frac{2^3}{3 * 7^4} \pi^6,$$

where K is the maximal real subfield of cyclotomic field, $\mathbb{Q}(\zeta_7)$. We note that this field is the same as the simplest cubic field with $m = -1$. By our computation, we have

$$\zeta_K(-1) = -\frac{1}{21}.$$

By the functional equation, our result and Siegel’s result coincide. More generally, we have (cf. [4]) that if $D = m^2 + 3m + 9 = p$ is a prime, our simplest cubic field corresponding to m is the cubic subfield of $\mathbb{Q}(\zeta_p)$. Therefore our computation contains a table of zeta values of cubic subfield of $\mathbb{Q}(\zeta_p)$, where p runs over primes of the form $m^2 + 3m + 9$.

TABLE 1. Values of zeta functions of the first one hundred simplest cubic fields

m	D	$-21\zeta_K(-1)$	$8190\zeta_K(-3)$	$-3591\zeta_K(-5)$
-1	7	1	$3^2 * 337$	$3 * 19 * 7393$
1	13	7	$2^2 * 11 * 43 * 113$	$2^2 * 11 * 8459933$
2	19	$3^2 * 7$	$13 * 37 * 53 * 127$	$733 * 33830759$
4	37	$3 * 7^2$	$3^4 * 29 * 43 * 3433$	$3^2 * 769 * 5478511391$
7	79	$7^2 * 199$	$3^2 * 5 * 12491 * 124799$	$11 * 138582878707283203$
8	97	$7^2 * 367$	$5 * 23 * 1783 * 1439381$	$3 * 13 * 214201811 * 9547468279$
10	139	$5^2 * 7 * 43$	$3^3 * 13 * 3659 * 2851327$	$2^2 * 31 * 3709558534651158701$
11	163	$2^2 * 7 * 491$	$2^2 * 7 * 19 * 89 * 193 * 1226857$	$149 * 76308073 * 941711501441$
13	$7^2 * 31$	$3^2 * 89 * 113$	$5 * 16562502882041$	$77158038781 * 7803617145641$
14	13 * 19	$3^2 * 19 * 109$	$23 * 263 * 33871899353$	
16	313	$7 * 11 * 1307$	$43 * 109 * 81553 * 2826371$	$5^2 * 113 * 8747 * 554117 * 145617822941$
17	349	$2^6 * 3 * 643$	$1747 * 9649 * 136679953$	$3^2 * 7 * 291060286406388642904463$
19	$7^2 * 61$	$3^2 * 5^2 * 7^2 * 61$	$3 * 67 * 47053638267793$	$670333 * 35680459 * 2151776396311$
20	$7^2 * 67$	$3^2 * 32999$	18239333904161593	$2^2 * 16087 * 13650960177131550855541$
22	$13 * 43$	$2^2 * 71 * 6977$	$5 * 11 * 491 * 2307105606107$	$5 * 2221 * 2719 * 61493 * 693223 * 275750779$
23	607	$2^2 * 5 * 7 * 11 * 23 * 119923 * 26135957$		$2^2 * 16087 * 13650960177131550855541$
25	709	$2^4 * 63313$	$199 * 283 * 9807732170543$	$7 * 23 * 37 * 59^2 * 1187 * 197033309231552533$
26	$7^2 * 109$	$2^3 * 3 * 5 * 11987$	$7 * 17 * 2378473 * 5150532091$	$29 * 277 * 1259 * 1075352008183959644609$
28	877	1954357	$199 * 283 * 9807732170543$	$1163 * 1184303536883 * 36523018721741$
29	937	$2 * 3^2 * 7 * 11 * 1667$	$2 * 11^2 * 1613 * 5933822299921$	$2^2 * 3 * 104092933933 * 334151629777440823$
31	1063	$3^2 * 61 * 6947$	$2^2 * 3 * 13 * 127 * 283880002753163$	$7 * 59 * 68045729 * 101323217 * 284306417491$
32	1129	4092029	$719 * 1187809893357673$	$2 * 3^2 * 73 * 2346733 * 3260563 * 286291222738333$
34	$7^2 * 181$	$37^2 * 43^2 * 3719$	$2 * 3 * 7 * 456028428993552851$	$2^2 * 3 * 43 * 13054889 * 504745450802855360303$
35	$13 * 103$	$27 * 3 * 41 * 443$	$2^2 * 3 * 13 * 89 * 109 * 18630840320461$	$2 * 3^2 * 73 * 2346733 * 3260563 * 286291222738333$
37	1489	$5^2 * 11 * 37369$	$2 * 3 * 61 * 3769193 * 43027272209$	$2 * 3 * 43 * 13054889 * 504745450802855360303$
38	1567	$3^2 * 3663089$	$1311847 * 3364513 * 19201327$	$5 * 13^2 * 317 * 35999 * 3091476759796261264391$
40	$7^2 * 13 * 19$	$3^2 * 5 * 241 * 4127$	$5 * 690629 * 4888245098879$	$4473971 * 19663857147270005853525581$
43	1987	$3^2 * 11 * 257 * 863$	$2 * 3 * 11^2 * 89 * 337 * 571 * 3449 * 7109671$	$2 * 11 * 37 * 67 * 73 * 9349 * 70912913 * 153913652563969$
44	$31 * 67$	$2^3 * 173 * 19373$	$2 * 8311 * 98028449 * 274128131$	$2^2 * 3 * 323058826847 * 56864372241656924759$
46	$31 * 73$	$7^2 * 29 * 25943$	$9649 * 66721 * 213263 * 8118853$	$2^2 * 3 * 43 * 67 * 809 * 321631 * 82442490325966470781$
47	$7^2 * 337$	$3^2 * 5^2 * 7 * 37 * 643$	$3 * 7 * 13 * 3121 * 1743502925737697$	$3 * 311 * 21844013 * 66661255861 * 1974861474997$
49	2557	$3^2 * 7^2 * 439 * 5059$	$47 * 55541728640468168369$	$19 * 29 * 582719 * 20278712979050892925969543$
50	$19 * 151$	$2 * 27613801$	$2 * 3 * 563 * 324720829 * 3130337051$	$2 * 3 * 43 * 67 * 809 * 321631 * 82442490325966470781$
52	$13 * 229$	$3^3 * 5^3 * 19681$	$4872547 * 1199410637635117$	$5 * 7^2 * 59 * 7282052597 * 21947107898623329581$
53	$7^2 * 457$	$3^2 * 5 * 7 * 131 * 5387$	$3 * 89 * 2089344583 * 13569011087$	$3 * 13 * 667673 * 1332146506670126336212698037$
56	3313	$2^2 * 41 * 1148177$	$3^2 * 1392032587525412966413$	$3^2 * 167329 * 2978321 * 4220323 * 4042430674898279$
58	3547	$3^2 * 73 * 171929$	$2^2 * 59 * 8849 * 7691695039188067$	$2^2 * 233 * 130631 * 923960723340243327410960881$
59	$19 * 193$	$2 * 3^3 * 7^2 * 50359$	$2 * 3^2 * 1436274899 * 999085267501$	$2 * 3 * 13^2 * 433 * 542712625267643149137794099129$
		$2 * 3^3 * 59 * 44221$	$2 * 19 * 523 * 176153 * 9304374089077$	$2 * 4447 * 15271 * 1059986933 * 2386540073717358073$

TABLE 1. Values of zeta functions of the first one hundred simplest cubic fields (continued)

m	D	$-2\zeta_K(-1)$	$8190\zeta_K(-3)$	$-3591\zeta_K(-5)$
61	$7^3 \cdot 13^4 \cdot 43$	$2^3 \cdot 7 \cdot 1361 \cdot 2539$	$2 \cdot 11 \cdot 19 \cdot 557 \cdot 130604017 \cdot 1694844391$	$2 \cdot 11 \cdot 29 \cdot 47 \cdot 968490011 \cdot 24170128718591566624981$
62	$7^5 \cdot 77$	$47 \cdot 4077299$	$3 \cdot 17 \cdot 3083 \cdot 407546982398304031$	$3 \cdot 11 \cdot 59 \cdot 510814649940548529242657677008589$
64	$4 \cdot 297$	$22 \cdot 37 \cdot 5923 \cdot 12823 \cdot 1350477 \cdot 5669501$	$22 \cdot 3 \cdot 17 \cdot 5923 \cdot 12823 \cdot 1350477 \cdot 5669501$	$22 \cdot 23 \cdot 47 \cdot 23173 \cdot 71257 \cdot 376039 \cdot 731987945025979501$
65	$43 \cdot 103$	$5^3 \cdot 3803 \cdot 13669$	$29 \cdot 37 \cdot 203921 \cdot 878387 \cdot 636038869$	$13 \cdot 31 \cdot 317 \cdot 2043773003 \cdot 325180992724547059981117$
67	$37 \cdot 127$	$22 \cdot 487 \cdot 148859$	$2^2 \cdot 11 \cdot 400485790711 \cdot 104857345427$	$2^2 \cdot 5 \cdot 13 \cdot 1709 \cdot 4219 \cdot 295319 \cdot 94941639754441146519461$
68	$7 \cdot 691$	$2^4 \cdot 20852521$	$2^2 \cdot 8831 \cdot 6409766466827819957$	$2^2 \cdot 3433 \cdot 4547 \cdot 50993 \cdot 247734713 \cdot 9162170167019153$
70	5119	395061967	$5 \cdot 13 \cdot 19 \cdot 29 \cdot 293 \cdot 32070120083484583$	$11 \cdot 19 \cdot 277 \cdot 7349 \cdot 200497 \cdot 1128697545302716610226713$
71	$19 \cdot 277$	$2^3 \cdot 3^2 \cdot 29 \cdot 1259 \cdot 2089$	$2 \cdot 29 \cdot 467 \cdot 209771 \cdot 736657 \cdot 97987523$	$2 \cdot 2729 \cdot 17099 \cdot 379910561649181 \cdot 937933283457641$
73	5557	48613047	$2 \cdot 563 \cdot 530837595019567524541$	$3^5 \cdot 3623 \cdot 1537947554229723139091402802533393$
74	$13^4 \cdot 439$	$5 \cdot 105141151$	$2^5 \cdot 33768545449 \cdot 666517986487$	$2^3 \cdot 3 \cdot 199 \cdot 7146507571928846478343$
76	$7 \cdot 859$	$113 \cdot 6228709$	$3 \cdot 199 \cdot 74099 \cdot 187417 \cdot 248309$	$3 \cdot 5 \cdot 7 \cdot 103 \cdot 1933 \cdot 95987 \cdot 2156784490871 \cdot 18298248941341$
77	$31 \cdot 199$	$5 \cdot 133569811$	$2^3 \cdot 3 \cdot 23 \cdot 257 \cdot 2539 \cdot 74099 \cdot 187417 \cdot 248309$	$2^3 \cdot 3 \cdot 11 \cdot 132 \cdot 23 \cdot 4363 \cdot 8423 \cdot 10631 \cdot 26176876349533965283$
79	$13 \cdot 499$	851224189	$5 \cdot 1381337 \cdot 255345403 \cdot 1002547561$	$821 \cdot 2215927245348689 \cdot 100278474963067212523$
80	$61 \cdot 109$	$2^3 \cdot 5 \cdot 11^3 \cdot 23 \cdot 677$	$2^2 \cdot 3^2 \cdot 131 \cdot 239 \cdot 11441887 \cdot 162717310997$	$2^2 \cdot 3 \cdot 5 \cdot 439 \cdot 563 \cdot 10711 \cdot 1506532989667378018879889443$
82	$7 \cdot 997$	$7 \cdot 1723^2 \cdot 81023$	$3 \cdot 761 \cdot 14593 \cdot 88450324216204751$	$3 \cdot 211287341 \cdot 505440799 \cdot 1272556442751141143053$
83	$7 \cdot 1021$	$2^8 \cdot 13^2 \cdot 5976947$	$22 \cdot 1279 \cdot 20323 \cdot 3348565004376611$	$2^2 \cdot 11 \cdot 12037233969541386189917635679611287083$
85	7489	$2^8 \cdot 11 \cdot 17 \cdot 29 \cdot 863$	$79 \cdot 61089110061312639682529$	$7 \cdot 31 \cdot 2287 \cdot 28097 \cdot 5670167297219 \cdot 11201950091570507$
86	$79 \cdot 97$	$2^3 \cdot 3^2 \cdot 7 \cdot 19 \cdot 1844207$	$2^8 \cdot 233 \cdot 3055527910211944058537$	$2^3 \cdot 941263 \cdot 6395143312067 \cdot 22964195433294933523$
88	8017	$31^2 \cdot 1511273$	$11^2 \cdot 140527 \cdot 457252338043234607$	$4508771281021307 \cdot 415609747066992341305171$
89	$7 \cdot 1171$	$11 \cdot 146880147$	$3 \cdot 7 \cdot 337 \cdot 2347523 \cdot 546968036265713$	$3^2 \cdot 41 \cdot 10007 \cdot 647874887252786071132152969854657$
91	8563	$3^5 \cdot 11 \cdot 11828293$	$2^2 \cdot 7 \cdot 193 \cdot 1231177 \cdot 1391189 \cdot 1337318531$	$2^2 \cdot 5 \cdot 193435090239370960677626089659653542847$
92	$13^6 \cdot 73$	$229 \cdot 8567723$	$19^2 \cdot 353 \cdot 443 \cdot 42743 \cdot 49639 \cdot 1196990407$	$29 \cdot 1307 \cdot 129265607569746936676799219000646107$
94	9127	$2 \cdot 3 \cdot 5^2 \cdot 14530823$	$2 \cdot 3^2 \cdot 3019 \cdot 11411 \cdot 19489 \cdot 210407 \cdot 7580051$	$2 \cdot 3 \cdot 89 \cdot 4038289 \cdot 3618126365534773923284310078919$
95	9319	$2^2 \cdot 3^2 \cdot 7 \cdot 9090947$	$2 \cdot 2 \cdot 3 \cdot 23 \cdot 4567 \cdot 35374410576344175397$	$2 \cdot 3^5 \cdot 20185492307257657676699655862201968689$
97	$7 \cdot 19^2 \cdot 73$	$2731 \cdot 961033$	$3 \cdot 5 \cdot 17 \cdot 61 \cdot 67^2 \cdot 109 \cdot 3904483830648433$	$3^2 \cdot 5 \cdot 17 \cdot 97919 \cdot 205592831335243722144072088351789$
98	9907	$17 \cdot 457 \cdot 354791$	$2^4 \cdot 3 \cdot 17 \cdot 23^2 \cdot 137 \cdot 57855893979838143$	$3^2 \cdot 3 \cdot 17 \cdot 19 \cdot 97 \cdot 20609717 \cdot 124085049678527961591142803$
101	10513	$2^8 \cdot 3 \cdot 19 \cdot 32407373$	$2^2 \cdot 3 \cdot 331 \cdot 677 \cdot 3863 \cdot 71119 \cdot 70459213369$	$2^2 \cdot 3 \cdot 23^2 \cdot 607 \cdot 395611 \cdot 35005189 \cdot 692544145041062423953$
104	$7 \cdot 37^2 \cdot 43$	$9871 \cdot 399727$	$3 \cdot 1083719 \cdot 23840944791443319923$	$3 \cdot 43 \cdot 44782939 \cdot 144556052208881 \cdot 8343123581468893$
106	$31^3 \cdot 373$	$5 \cdot 19 \cdot 107^2 \cdot 473353$	$5 \cdot 71 \cdot 2917 \cdot 241973 \cdot 5724853 \cdot 40509583$	$89 \cdot 70750299 \cdot 15354927047269503 \cdot 15288111433181$
107	11779	$5 \cdot 7^2 \cdot 281 \cdot 490579$	$17 \cdot 37 \cdot 293 \cdot 1018481089 \cdot 612853210009$	$5^3 \cdot 109 \cdot 7883 \cdot 42239 \cdot 28449433686441097449132066461$
109	$19^2 \cdot 643$	$2^6 \cdot 3^2 \cdot 5^3 \cdot 17 \cdot 613$	$2^3 \cdot 19 \cdot 97 \cdot 191 \cdot 52694742964789700339$	$2^8 \cdot 5^2 \cdot 7 \cdot 421 \cdot 83490123413 \cdot 14210937656088181$
110	$7 \cdot 1777$	$5 \cdot 73 \cdot 15036509$	$2^2 \cdot 17 \cdot 1303 \cdot 5669 \cdot 15661 \cdot 2140408245153$	$2^2 \cdot 2752 \cdot 11 \cdot 13337384836208063510416613292901761$
112	12889	$7^3 \cdot 83^2 \cdot 29^2 \cdot 753$	$2 \cdot 107 \cdot 100879779217989642012333$	$2 \cdot 773 \cdot 7068577 \cdot 3180414766618890249818740403039$
113	$13 \cdot 1009$	$2^8 \cdot 3 \cdot 3685259$	$2 \cdot 73 \cdot 89 \cdot 487 \cdot 36445741 \cdot 1058171951659$	$2 \cdot 17 \cdot 245277829556100359 \cdot 50842392001722969955391$
115	$37 \cdot 367$	$3^2 \cdot 793909517$	$1759 \cdot 71971 \cdot 245639826949277927$	$8093 \cdot 20173 \cdot 6848789 \cdot 627157393537 \cdot 879607085547023$
116	$19 \cdot 727$	$947 \cdot 1721^2 \cdot 5237$	$397 \cdot 10123805161 \cdot 87557778932609$	$13 \cdot 31 \cdot 41 \cdot 2837447687141181961372180943339791141217$
121	15013	$2 \cdot 3^2 \cdot 227 \cdot 653 \cdot 1931$	$2^4 \cdot 1483835047 \cdot 26577931740532803$	$3 \cdot 225402687410705729 \cdot 329093984334811525889781$
122	15259	$2^2 \cdot 3^2 \cdot 227 \cdot 653 \cdot 1931$	$3 \cdot 7^2 \cdot 1611428063939 \cdot 297046253679$	$2^4 \cdot 3 \cdot 110967014115947068405778374120360190687371$
127	16519	$2^3 \cdot 3^2 \cdot 223 \cdot 271 \cdot 2903$	$2^4 \cdot 3^3 \cdot 7 \cdot 19^2 \cdot 191 \cdot 1567 \cdot 112592326068652129$	

TABLE 1. Values of zeta functions of the first one hundred simplest cubic fields (continued)

m	D	$-21\zeta_K(-1)$	$8190\zeta_K(-3)$	$-3591\zeta_K(-5)$
130	17299	$2^*691*10685029$	$2^*179*1193*851229091*2329032723197$	$2^2*129998635880941*17015528241242907885387499001$
133	18097	$2^*1761*269*30223$	$7*317*1747*598994083883237615721$	$22422527167*164280574059307*3944708488276557173$
134	18367	17748484291	$2*31*41579*999277978409878150031$	$2*11161*76612778111278214348391073812463709713$
136	18913	$2^2*5*239*4529563$	$3174678346344873370589816419$	$67*863*34250691863*40991015742997*290804843257813$
140	20029	$3^*7*13*17*4838063$	$2^2*3^2*71*184780690774287462772183$	$2^2*3*3*97*197293*2398243*80520645458171421266738272043$
142	20599	$2^4*3*131*311*13523$	$5*7*571*1213*179909*1319533658614187$	$97*431*3259*33773*131282827414382010881591945190163$
143	20887	$2^5*2713*954847$	$2^3*11*1753*8035793*5110448131339981$	$2^4*653*22229*87317*8094756331*428542865890674920449$
155	24499	$2^2*101*104228587$	$5^2*275107*7115791*391966716864499$	$7*13*67*6989293*55307177005420027*172549789664464673$
158	25447	$223*2017*104911$	$5*2757037*5407561*9290233*36450241$	$5*17^2*427315325862435484252362369926770131512869$
163	27067	$2^2*14626374283$	$7*157*42178439*138575623*865517501$	$7*4569406109*38063279570318234990797412471959643$
164	27397	$3^*69247*279557$	$5*7*197*2099*35743339*81808157288513$	$239*307*51202087*37027328252363604705350933056517$
169	29077	$2^3*43*53*5160767$	$3*11*17*73*1567641825047747473525643$	$3^2*13147*6669983*726057550873961*469422685830334319$
172	30109	$2*3^4*5*5*277169$	$2^2*3*101747*67109061379176186974579$	$2^2*3*293*3163*8089*43674699384570741974762236223409271$
175	31159	$5^*7*2424763087$	$3^3*19*47*79*54682554396150587382661$	$3^3*17*151*373*86509*6900841*37116756006254642622403129$
176	31513	$2^2*3^3*11*17*443*21611$	$2*3*1091*38867*444778684474815104501$	$2*59*52027961640131*267437762901247*5711626833065897$
179	32587	$2^2*11*43*353*146801$	$2*7^2*428735646347*3392424912471912502820921$	$71*1672381*3079124513*36885904429362021984809827111$
182	33679	$239*1627*293453$	$468509*38324912471912502820921$	$1787*21757*52218007*653240873*57219691957603126607837$
197	39409	$2^5*7*13*149*394633$	$5*13*1879*2017*22511*80363*1209969999751$	$2^2*79*118096859687*989118903346943*2859469032020339041$
200	40609	$2^2*7*4327*1619249$	$2^2*29*43*4297*207061*6895901*217742339627$	$2^3*2680991952303795365657*2109475765541940340229423$
205	42649	$3^*7*10864049089$	$2^4*2851*1445699*142309602670323219$	$2^3*11*823*33641*41989797991*787031379684399222464801657$
206	43063	252696953417	$2*373*458099774023*2946831153113677$	$2^3*11*159855*12405168581731*17122246624369*126100416816631$
212	45589	$2*19*7447442731$	$2^5*23*1171*79537*38644787*564943119089$	

REFERENCES

1. D. Byeon, Special values of zeta functions of the simplest cubic fields and their applications, Proc. Japan Acad. Ser. A. Math. Sci. **74** (1988), 13-15. MR **99c**:11141
2. U. Halbritter and M. Pohst, On the computation of the values of zeta functions of totally real cubic fields, J. Number Theory **36** (1990), 266-288. MR **92b**:11080
3. H. K. Kim and H. J. Hwang, Values of zeta functions and class number 1 criterion for the simplest cubic fields, Nagoya Math. J. **160** (2000), 161-180. CMP 2001:06
4. J. S. Kim, Determination of class numbers of the simplest cubic fields, to appear in Comm. of the Korean Math. Soc.
5. A. J. Lazarus, The class number and cyclotomy of simplest quartic fields, PhD thesis, University of California, Berkeley, 1989.
6. D. Shanks, The simplest cubic fields, Math. Comp. **28** (1974), 1137-1152. MR **50**:4537
7. C. L. Siegel, Berechnung von Zetafunktionen an ganzzahligen Stellen, Nachr. Akad. Wiss. Göttingen Math.-Phys. Kl. II (1969), 87-102. MR **40**:5570
8. L. C. Washington, Class numbers of the simplest cubic fields, Math. Comp. **48** (1987), 371-384. MR **88a**:11107
9. L. C. Washington, Introduction to cyclotomic fields, Graduate Texts in Mathematics, **83**, Springer-Verlag, New York 1982. MR **85g**:11001
10. D. B. Zagier, "On the values at negative integers of the zeta function of a real quadratic field", Enseignement Math. **22** (1976), 55-95. MR **53**:10742

DEPARTMENT OF MATHEMATICS, POHANG UNIVERSITY OF SCIENCE AND TECHNOLOGY, POHANG 790-784, KOREA; AND SCHOOL OF MATHEMATICS, KOREA INSTITUTE FOR ADVANCED STUDY, SEOUL 130-012, KOREA

E-mail address: `hkkim@postech.ac.kr`

DEPARTMENT OF MATHEMATICS, POHANG UNIVERSITY OF SCIENCE AND TECHNOLOGY, POHANG 790-784, KOREA

E-mail address: `integer@euclid.postech.ac.kr`