

# Synthesis and *In Vitro* Anti-Lung Cancer Activity of Novel 1, 3, 4, 8-Tetrahydropyrrolo [4, 3, 2-de]quinolin-8(1H)-one Alkaloid Analogs

Dwayaja H. Nadkarni<sup>1,‡</sup>, Feng Wang<sup>2,‡</sup>, Wei Wang<sup>2,‡</sup>, Elizabeth R. Rayburn<sup>2,3</sup>, Scharri J. Ezell<sup>2</sup>, Srinivasan Murugesan<sup>1</sup>, Sadanandan E. Velu<sup>1,4,\*</sup> and Ruiwen Zhang<sup>2,3,4,‡,\*</sup>

<sup>1</sup>Department of Chemistry, <sup>2</sup>Department of Pharmacology and Toxicology, Division of Clinical Pharmacology, <sup>3</sup>Gene Therapy Center, <sup>4</sup>Comprehensive Cancer Center, and University of Alabama at Birmingham, Birmingham, AL 35294, USA

**Abstract:** The high mortality rate and lack of effective therapies make lung cancer an ideal target for novel therapeutic agents. The present study was designed to implement a novel chemical synthesis pathway and to determine the biological activities of synthetic makaluvamine analogs in human lung cancer. Seventeen compounds were synthesized and purified, and their chemical structures were elucidated on the basis of physicochemical constants and NMR spectra. Their *in vitro* activity was determined in human lung cancer cell lines. Based on initial screens, compound 1c was found to be the most potent, and was therefore used as a model for further studies in lung cancer cells. 1c induced both apoptosis and S-phase cell cycle arrest. Furthermore, it activated p53 and induced cleavage of PARP and caspases 8 and 9. Our preclinical data indicate that the makaluvamine analogs are potential therapeutic agents against lung cancer, providing a basis for further development of 1c (and perhaps other analogs) as a novel anti-cancer agent.



**Key Words:** Makaluvamine, marine alkaloid, lung cancer, apoptosis.

**# Author Profile:** Ruiwen Zhang is Professor of Pharmacology and Toxicology, Director of the Cancer Pharmacology Laboratory, Senior Scientist in the Comprehensive Cancer Center, Center for AIDS Research, and Gene Therapy Center at the University of Alabama at Birmingham. He has extensive experience in molecular therapeutics, clinical pharmacology, and toxicology.

## INTRODUCTION

Cancer, particularly lung cancer, poses a major health problem worldwide. Although significant progress has been seen with early detection, treatment, and prevention of recurrence, new therapies are urgently needed in order to improve the clinical outcome for cancer patients. Natural products from marine animals and their derivatives have proven to be a rich source of novel, medicinally valuable compounds [1-4]. The toxic substances produced by marine invertebrates such as sponges, bryozoans, tunicates and ascidians, assist the organisms in the deterrence of predators and the paralysis of prey [5]. As such, marine natural products exhibit potent biological activities.

Marine sponges of the genera *Latrunculia*, *Batzella*, *Prianos* and *Zyzya* are a rich source of alkaloids bearing a pyrrolo[4,3,2-de]quinoline skeleton [6, 7]. This series of alkaloids, comprised of approximately 60 metabolites, including makaluvamines [8-13], exhibit numerous biological activities [8, 11] and exert cytotoxicity against numerous tumor cell lines [11, 14, 15]. Many also have anti-fungal [8] or antimicrobial activity [16]. A number of excellent reviews have

highlighted the chemistry and bioactivity of this class of compounds [6, 17-19], prompting us to investigate the potential role of chemically synthesized derivatives of marine alkaloids as novel anti-cancer agents.

We recently synthesized a number of makaluvamine analogs and evaluated their anti-cancer activity against several human cancer cell lines [20, 21]. This manuscript describes the synthesis of two additional groups of analogs, the first group (Fig. (1)) contains a p-toluenesulfonyl (tosyl) group on the pyrrole ring N, while the second group (Fig. (2)) contains a free pyrrole N atom. Both groups contain substituents at the 6-position of the pyrroloiminoquinone ring of makaluvamines. Herein, we provide a comparison of the biological activity of these new analogs in human lung cancer cell lines.

## MATERIALS AND METHODS

### Synthesis and Characterization of the Compounds

#### Reagents and Instrumentation


Anhydrous solvents used for reactions were purchased from Aldrich Chemical Company, and solvent evaporations were carried out *in vacuo* using a rotary evaporator. Thin layer chromatography (TLC) was performed on silica gel plates with a fluorescent indicator (Whatmann, silica gel, UV254, 25  $\mu$ m plates). Spots were visualized by UV light (254 and 365nm) and purification by column and flash chromatography was carried out using BAKER silica gel (40 $\mu$ m) in the solvent systems indicated. Proton nuclear magnetic resonance (<sup>1</sup>H-NMR) and carbon nuclear magnetic

\*Address correspondence to these authors at the University of Alabama at Birmingham, Department of Pharmacology and Toxicology, 1670 University Blvd, 113 Volker Hall, Birmingham, AL 35294, USA; Tel: (205)-934-8558; Fax: (205)-975-9330; E-mail: ruiwen.zhang@ccc.uab.edu

University of Alabama at Birmingham, Department of Chemistry, 280 Chemistry Building, 901 14<sup>th</sup> Street South, Birmingham, AL 35294, USA; Tel: (205)-975-2478; Fax: (205)-934-2543; E-mail: svelu@uab.edu

<sup>‡</sup>These authors contributed equally to this project.

resonance ( $^{13}\text{C}$ -NMR) spectra were recorded on a Bruker DPX-300 spectrometer using tetramethylsilane (TMS) as an internal standard. Mass spectra results were recorded on the Micromass Platform LCC instrument. Chemical shift values ( $\delta$ ) are reported as parts per million (ppm) and coupling constants (J) as hertz (Hz).




**Fig. (1).** Makaluvamine analogs containing a tosyl group on the pyrrole Nitrogen.

#### Amination of the Pyrroloiminoquinone Derivative, Ia

A schematic of the procedure for synthesizing the compounds appears in Scheme 1. To generate the compounds in Group I, a solution of amine (7) in anhydrous MeOH was added drop-wise to a solution of the pyrroloiminoquinone derivative, Ia, over a 15 min period. The resulting solution was stirred at room temperature for 18 hr. Upon completion of the reaction, the mixture was cooled to 0°C and quenched by addition of trifluoroacetic acid (TFA). The crude product was obtained by removal of the solvent solution under reduced pressure and purified by column chromatography over silica gel using MeOH/CHCl<sub>3</sub>.

#### Detosylation

In order to generate detosylated compounds (for Group II), sodium methoxide was added to a MeOH solution of *N*-tosyl makaluvamine derivatives (Group I) and stirred at room temperature for 4-6 hr. The resulting solution was




**Fig. (2).** Makaluvamine analogs containing an un-substituted pyrrole Nitrogen.

quenched at 0°C with TFA. The solvent was completely removed under reduced pressure and the resulting residue was co-evaporated with CHCl<sub>3</sub> to remove traces of TFA. The remaining crude compound was purified by chromatography over a column of silica gel (20×2cm) using MeOH/CHCl<sub>3</sub>.

#### Biological Experiments

##### Reagents

All chemicals and solvents used were of the highest analytical grade available. Cell culture media, phosphate-buffered saline (PBS), fetal bovine serum (FBS), sodium pyruvate, non-essential amino acids, penicillin-streptomycin, and other



**Scheme 1.** Synthesis of makaluvamine analogs **Ib-g** and **IIa-j**.

cell culture supplies were obtained from the Comprehensive Cancer Center Media Preparation Shared Facility, University of Alabama at Birmingham (Birmingham, AL). The anti-human p53 (Ab-6) antibody was acquired from EMD Chemicals, Inc (Gibbstown, NJ). The anti-human MDM2 (SMP14) and PARP (H-250) antibodies were purchased from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA), and the caspase-8 (9746) and caspase-9 (9502) antibodies were purchased from Cell Signaling Technology, Inc (Danvers, MA).

#### Cell Culture


Human cancer cells and normal lung epithelial cells (BEAS-2B) were obtained from the American Type Culture Collection (Rockville, MD) and cultured according to their instructions. The human primary fibroblast cell line IMR90-EEA (IMR90), which was cultured in DMEM, was a gift from Dr. S. Lee (Harvard University, Boston, MA). All culture media contained 10% fetal bovine serum and 1% penicillin-streptomycin. A549 cells were grown in Ham's F12K medium supplemented with 2 mM L-glutamine and 1.5g/L sodium bicarbonate. H838 cells were grown in RPMI 1640 supplemented with 1.5 g/L sodium bicarbonate, 4.5 g/L glucose, 10 mM HEPES buffer, 1 mM sodium pyruvate and 2 mM L-glutamine. H1299 cells were grown in DMEM media.

#### Cell Survival Assay

The effects of test compounds on human cancer cell growth, expressed as the percentage of cell survival, were determined using the MTT assay [22, 23]. In brief, the cells were grown in 96-well plates at  $4-5 \times 10^3$  cells per well and exposed to the test compounds (0, 0.1, 0.25, 0.5, 1, 2.5  $\mu\text{M}$ ) for 72 hr. After incubation with the compounds, 10  $\mu\text{L}$  of the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) solution (5mg/mL; Sigma; St. Louis, MO) were added to each well. The plates were incubated for 2-4 hr at 37°C. The supernatant was then removed, and the formazan crystals were dissolved in 100  $\mu\text{L}$  of DMSO. The absorbance at 570nm was recorded using an OPTImax microplate reader (Molecular Devices; Sunnyvale, CA). The cell survival percentages were calculated by dividing the mean OD of compound-containing wells by that of DMSO-treated control wells. Three separate experiments were accomplished to determine the  $\text{IC}_{50}$ .

#### Cell Cycle Measurements

To determine the effects of the compound on the cell cycle, cells ( $2-3 \times 10^5$ /well) were exposed to the test compound (0, 0.1, 0.5 or 0.75  $\mu\text{M}$ ) and incubated for 24 hr prior to analysis. Cells were then trypsinized, washed with PBS,



**Fig. (3).** Growth inhibitory activity of makaluvamine analogs Ia and Ic in lung cancer and normal cells. Cells were exposed to various concentrations of the compounds for 72 hr followed by MTT assay. All assays were performed in triplicate.

Table 1. <sup>1</sup>H-NMR, <sup>13</sup>C-NMR, and MS Spectral Data of 1,3,4,8-Tetrahydropyrrolo[4,3,2-de]quinolin-8(1H)-one Alkaloid Analogs

Compd.	<sup>1</sup> H-NMR δ Values (ppm) <sup>a</sup>	<sup>13</sup> C-NMR δ Values (ppm) <sup>a</sup>	MS (ES+) <sup>b</sup> , m/z (M+)
<b>Ib</b>	(CDCl <sub>3</sub> ) δ 2.45 (s, 3H), 2.98 (t, 2H, J = 7.2 Hz), 3.96 (t, 2H, J = 7.0 Hz), 4.45 (d, 2H, J = 5.7 Hz), 6.01 (s, 1H), 7.33–7.43 (m, 7H), 7.69 (s, 1H) and 8.01 (d, 2H, J = 10.5 Hz).	(CDCl <sub>3</sub> ) δ 18.0, 21.8, 42.6, 48.3, 87.8, 118.0, 123.2, 127.7, 128.2, 128.3, 128.8, 129.0, 129.3, 130.2, 133.0, 134.2, 147.3, 150.3, 156.9 and 166.6.	432
<b>Ic</b>	(CDCl <sub>3</sub> ) δ 2.46 (s, 3H), 2.93–3.04 (m, 2H), 3.92–4.04 (m, 2H), 4.45 (bs, 2H), 6.14 (bs, 1H), 7.07 (bs, 1H), 7.20–7.27 (m, 2H), 7.35–7.40 (m, 4H), 7.67 (s, 1H), 8.02 (d, 2H, J = 8.1 Hz).	(CDCl <sub>3</sub> ) δ 18.0, 21.8, 42.7, 47.5, 88.0, 117.9, 123.1, 127.6, 128.2, 129.0, 129.5, 129.7, 130.2, 132.7, 132.9, 134.9, 147.4, 150.3, 157.1 and 166.5.	466
<b>Id</b>	(CDCl <sub>3</sub> ) δ 2.45 (s, 3H), 2.97 (bt, 2H, J = 6.6 Hz), 3.88 (s, 6H), 3.98 (bt, 2H, J = 4.5 Hz), 4.39 (d, 2H, J = 5.1 Hz), 6.80–6.89 (m, 3H), 7.06 (bs, 1H), 7.38 (d, 2H, J = 8.4 Hz), 7.67 (s, 1H) and 8.02 (d, 2H, J = 8.4 Hz).	(CDCl <sub>3</sub> ) δ 18.0, 21.8, 29.7, 42.5, 48.2, 56.0, 56.1, 87.8, 111.6, 111.7, 118.0, 121.0, 123.2, 126.6, 127.8, 128.1, 129.0, 130.2, 133.0, 147.3, 149.5, 149.5, 150.1, 156.9 and 166.7.	492
<b>Ie</b>	(CDCl <sub>3</sub> ) δ 2.46 (s, 3H), 2.93–2.99 (m, 4H), 3.54–3.62 (m, 2H), 3.98 (bt, 2H, J = 6.9 Hz), 6.20 (bs, 1H), 6.84 (bs, 1H), 7.02 (t, 2H, J = 8.5 Hz), 7.19 (dd, 2H, J <sub>1</sub> = 8.7, J <sub>2</sub> = 5.7 Hz), 7.39 (d, 2H, J = 8.4 Hz), 7.67 (s, 1H) and 8.02 (d, 2H, J = 8.4 Hz).	(CDCl <sub>3</sub> ) δ 18.0, 21.8, 29.7, 33.3, 42.5, 45.0, 87.5, 115.8, 116.0, 118.0, 123.1, 127.8, 128.2, 129.0, 130.1, 130.2 (2C), 132.4, 132.5, 133.0, 147.3, 150.6, 156.8, 160.4 and 166.4.	464
<b>If</b>	(CDCl <sub>3</sub> ) δ 2.46 (s, 3H), 2.89–2.96 (m, 4H), 3.52–3.58 (m, 2H), 3.81 (s, 3H), 3.97 (bt, 2H, J = 6.9 Hz), 6.19 (bs, 1H), 6.87 (d, 2H, J = 8.4 Hz), 7.13 (d, 2H, J = 8.4 Hz), 7.38 (d, 2H, J = 8.4 Hz), 7.65 (s, 1H) and 8.02 (d, 2H, J = 8.4 Hz).	(CDCl <sub>3</sub> ) δ 18.0, 21.8, 33.2, 42.4, 45.2, 55.3, 87.5, 114.5, 118.0, 123.2, 127.9, 128.1, 128.7, 128.9, 129.7, 130.2, 133.1, 147.2, 150.6, 156.7, 158.8 and 166.5.	476
<b>Ig</b>	(CDCl <sub>3</sub> ) δ 2.45 (s, 3H), 2.88 (t, 2H, J = 6.6 Hz), 2.97 (t, 2H, J = 6.6 Hz), 3.57–3.49 (m, 2H), 3.96 (t, 2H, J = 7.2 Hz), 5.95 (s, 2H), 6.06 (bs, 1H), 6.64–6.67 (m, 2H), 6.75 (d, 1H, J = 7.8 Hz), 6.91 (bs, 1H), 7.38 (d, 2H, J = 8.4 Hz), 7.66 (s, 1H) and 8.01 (d, 2H, J = 8.4 Hz).	(CDCl <sub>3</sub> ) δ 18.0, 21.8, 29.7, 33.7, 42.5, 45.1, 87.3, 101.1, 108.7, 108.8, 118.0, 121.7, 123.1, 127.3, 127.8, 128.2, 128.9, 130.3, 133.0, 146.8, 147.3, 148.2, 150.8, 156.8 and 166.3.	490
<b>Ilg</b>	(CD <sub>3</sub> OD) δ 2.87 (t, 2H, J = 7.5 Hz), 2.94 (t, 2H, J = 7.8 Hz), 3.54 (t, 2H, J = 6.9 Hz), 3.83 (t, 2H, J = 7.5 Hz), 5.38 (s, 1H), 6.71 (d, 2H, J = 8.4 Hz), 7.06 (d, 2H, J = 8.4 Hz) and 7.14 (s, 1H).	(CD <sub>3</sub> OD) δ 19.5, 34.4, 44.2, 46.5, 85.2, 116.5 (2C), 120.2, 124.0, 127.0, 127.2, 130.0, 130.9 (2C), 155.0, 157.4, 159.8 and 168.5.	307
<b>Ili</b>	(CD <sub>3</sub> OD) δ 2.95 (t, 2H, J = 7.8 Hz), 3.12 (t, 2H, J = 7.2 Hz), 3.68 (t, 2H, J = 6.9 Hz), 3.84 (t, 2H, J = 7.8 Hz), 5.42 (s, 1H), 7.15 (s, 1H), 7.51 (d, 2H, J = 9.0 Hz) and 8.17 (d, 2H, J = 9.0 Hz).	(CD <sub>3</sub> OD) δ 19.5, 34.8, 44.2, 45.3, 85.4, 120.2, 123.8, 124.8 (2C), 127.2, 129.8, 131.1 (2C), 147.5, 148.4, 155.1, 159.9 and 168.5.	337
<b>Iij</b>	(CD <sub>3</sub> OD) δ 2.86 (t, 2H, J = 6.8 Hz), 2.95 (t, 2H, J = 7.6 Hz), 3.55 (t, 2H, J = 7.6 Hz), 3.85 (t, 2H, J = 7.6 Hz), 5.38 (s, 1H), 7.15 (s, 1H) and 7.39 (s, 2H).	(CD <sub>3</sub> OD) δ 19.5, 33.6, 44.2, 45.9, 85.4, 112.3, 120.2, 123.9, 125.5, 127.2 (2C), 133.5, 133.8 (2C), 151.3, 155.1, 159.7 and 168.5.	466

<sup>a</sup> <sup>1</sup>H-NMR and <sup>13</sup>C-NMR spectra were recorded on a Bruker DPX-300 spectrometer. The chemical shift values are reported as parts per million (ppm) relative to tetramethylsilane as internal standard.

<sup>b</sup> Mass spectra were recorded on a Micromass Platform LCC Electrospray instrument.

and fixed in 1.5mL of 95% ethanol at 4°C overnight, followed by incubation with RNase and staining with propidium iodide (Sigma). The DNA content was determined by flow cytometry.

#### Detection of Apoptosis

Cells in early and late stages of apoptosis were detected using an Annexin V-FITC apoptosis detection kit from Bio-Vision (Mountain View, CA) [22, 23]. In apoptosis experiments, 2–3×10<sup>5</sup> cells were exposed to the test compound (0, 0.1, 0.5 or 0.75μM) and incubated for 24 hr prior to analysis. Cells were collected and washed with serum-free media, then re-suspended in 500μL of Annexin V binding buffer followed by addition of 5μL of Annexin V-FITC and 5μL of propidium iodide (PI). The samples were incubated in the dark for 5 min at room temperature and analyzed with a Becton Dickinson FACSCalibur instrument (Ex=488 nm;

Em=530 nm). Cells that were positive for Annexin V-FITC alone (early apoptosis) and Annexin V-FITC and PI (late apoptosis) were counted.

#### Western Blot Analysis

The protein expression levels were assessed following exposure of cells to various concentrations of the compounds for 24 hr [22, 23]. Whole cell lysates were fractionated with identical amounts of protein by SDS-PAGE and transferred to Bio-Rad trans-Blot nitrocellulose membranes (Bio-Rad Laboratories, Hercules, CA). The nitrocellulose membrane was incubated in blocking buffer (Tris-buffered saline containing 0.1% Tween 20 and 5% nonfat milk) for 1 hr at room temperature. Then the membrane was incubated with the appropriate primary antibody overnight at 4°C or 2 hr at room temperature with gentle shaking. The membrane was

washed three times with rinsing buffer (Tris-buffered saline containing 0.1% Tween 20) for 15 min, then incubated with goat anti-mouse/rabbit IgG-horseradish peroxidase-conjugated antibody (Bio-Rad) for 1 hr at room temperature. After repeating the washes in triplicate, the protein of interest was detected by enhanced chemiluminescence reagents from PerkinElmer LAS, Inc (Boston, MA).

## RESULTS

### Synthesis and Characterization of the Compounds

All of the test compounds were synthesized in two steps starting from the tricyclic pyrroloiminoquinone compound, Ia (see Scheme 1). This compound was prepared following the 4,6,7-trimethoxyindole approach described previously [24]. Treatment of Ia with a solution of amine (7) in anhydrous methanol at room temperature provided the tosylated members of the first group of analogs, Ib-g (Fig. (1)) in 30-51% yields. Compounds IIa-j (Fig. (2)) were prepared by treatment of appropriately substituted group I compounds with NaOMe in MeOH, which resulted in 48-62% yields of the detosylated compounds. All final products were completely characterized, and purity was confirmed by <sup>1</sup>H-NMR, <sup>13</sup>C-NMR, and MS spectroscopy (Table 1).

### Initial Screening for Growth Inhibition in Cancer Cells


Seventeen compounds were tested for their *in vitro* anti-cancer activity using the MTT assay. Three human lung cancer cell lines were cultured with test compounds Ia-g and IIa-j at concentrations ranging from 0.01 μM to 10 μM for 72 hr, then cell viability was determined. The inhibitory effects of these compounds on cell growth are illustrated in Table 2. Most of the analogs in both series showed impressive activity against all three of the tested cell lines. However, Ia, which was used as a control for the core structure of these compounds, did not show any major activity. This finding indicates that the amino N atom at the 6-position of the pyrroloiminoquinone ring enhances the activity of the makaluvamine analogs.

The analog that showed the greatest activity was the N-tosyl chlorobenzyl analog, Ic (Mean IC<sub>50</sub> 0.79 μM). Among the compounds that contained a N atom at the 6-position of the ring, the least active analogs were the hydroxyphenethyl analog, IIg, and the structurally similar 2,6-dibromohydroxyphenethyl analog, IIj (Mean IC<sub>50</sub> > 10 μM). Similar effects of the compounds were also observed in human MCF-7 breast cancer cells (data not shown). The presence of the N-tosyl group did not seem to affect the activity of the compounds


**Table 2. Growth Inhibitory Activity of Makaluvamine Analogs in Human Lung Cancer Cells. Cells were Exposed to Various Concentrations of the Compounds for 72 hr Followed by MTT Assay. All Assays were Performed in Triplicate**

	A549			H838			H1299		
	IC <sub>20</sub> <sup>a</sup>	IC <sub>50</sub>	IC <sub>80</sub>	IC <sub>20</sub>	IC <sub>50</sub>	IC <sub>80</sub>	IC <sub>20</sub>	IC <sub>50</sub>	IC <sub>80</sub>
Ia	>10	>10	>10	>10	>10	>10	>10	>10	>10
Ib	1.11	1.39	1.76	1.60	2.05	2.69	0.39	0.88	1.72
Ic	0.30	0.39	0.53	0.50	1.41	3.07	0.29	0.58	1.16
Id	1.08	1.55	2.14	1.72	2.55	3.75	0.22	0.77	2.03
Ie	2.30	2.71	3.18	2.53	2.96	3.54	1.64	2.15	3.00
If	2.50	3.10	3.72	2.56	2.93	3.44	2.14	2.81	3.87
Ig	0.29	3.01	5.53	1.74	2.61	3.92	0.84	1.69	3.26
IIa	1.32	2.21	3.37	0.39	1.05	2.27	0.35	0.67	1.19
IIb	1.08	2.10	3.48	0.64	0.93	1.35	0.37	0.91	1.87
IIc	0.85	1.95	3.63	0.56	1.51	3.25	0.40	0.81	1.48
IId	1.09	3.33	5.56	0.79	1.29	2.04	0.88	1.84	3.06
IIe	0.73	3.01	5.81	0.84	1.80	3.35	0.35	1.22	2.93
IIf	1.18	2.84	5.20	0.69	1.71	3.46	0.51	1.16	2.26
IIg	3.16	>10.00	>10.00	4.00	7.94	>10.00	4.07	>10.00	>10.00
IIh	0.77	1.99	3.57	0.43	1.23	2.76	0.36	0.69	1.20
Iii	0.18	0.56	1.29	0.42	0.59	0.83	0.20	0.42	0.78
IIj	3.22	8.82	>10.00	7.47	9.71	>10	7.70	>10.00	>10.00

<sup>a</sup> IC, inhibitory concentration (μM). IC<sub>20</sub>, IC<sub>50</sub>, and IC<sub>80</sub> are the concentrations of drug that inhibit growth by 20%, 50%, and 80%, respectively, relative to the control.




**Fig. (4).** Effects of the makaluvamine analogs on cell cycle progression. Effects of Ic (A) and Ia (B) on human lung cancer cells. (C) Effects of Ic on human non-malignant lung epithelial cells (BEAS-2B) and fibroblasts (IMR90). Cells were exposed to various concentrations of Ic or Ia for 24 hr, followed by determination of cell cycle distribution. All assays were performed in triplicate.



**Fig. (5).** Induction of apoptosis by makaluvamine analogs. Effects of Ic (A) and Ia (B) on human lung cancer cells. (C) Effects of Ic on human non-malignant lung epithelial cells (BEAS-2B) and fibroblasts (IMR90). The cells were exposed to various concentrations of Ic or Ia for 24 hr, followed by assessment of apoptosis. All assays were performed in triplicate.

because analogs from both Groups I (Ib-g) and II (IIa-j) were active against the tested cell lines. Since Ic consistently exhibited potent activity against the lung cancer cells, we com-

pared the effects of Ic and Ia on the survival of lung cancer (A549 and H1299) cell lines and normal (non-malignant) (IMR90 and BEAS-2B) cells, and found that there was a



**Fig. (6).** Effects of the makaluvamine analog Ic on the expression of various apoptosis-related proteins in human lung cancer cells. Cells were exposed to various concentrations of Ic for 24 hr, and the target proteins were detected by immunoblotting with specific antibodies.

dose-dependent decrease in cell growth following Ic exposure in all of the cells, while Ia had minimal effects (Fig. (3)).

#### Examination of the Effects of the Compounds on the Cell Cycle

To determine whether the decrease in growth was due to changes in the cell cycle, we investigated whether Ic could induce cell cycle arrest. As shown in Fig. (4A), Ic induced considerable S-phase arrest in A549 cells, but exerted a lesser effect on H1299 cells. As expected, Ia did not exert any appreciable effect on cell cycle progression (Fig. (4B)). To determine whether the effect of Ic was specific for cancer cells, we also examined whether the compound altered cell cycle progression in the “normal” cells. As shown in Fig. (4C), Ic induced modest cell cycle arrest in both cells lines, with IMR90 exhibiting a slight increase in S phase cells, and the BEAS-2B cells showing an increase in cells in G2/M. Because A549, IMR90 and BEAS-2B cells express wild-type p53, while H1299 cells do not express the protein [25], this indicates that p53 may play a role in Ic-induced cell cycle arrest.

#### Induction of Apoptosis in Human Cancer Cells

Because the growth inhibition of H1299 cells appeared to be independent of the cell cycle, we examined whether Ic also could induce apoptosis. As illustrated in Fig. (5A), Ic induced apoptosis in a dose-dependent manner in both A549 (p53 wild-type) and H1299 (p53 null) cell lines, indicating that Ic can induce apoptosis, regardless of the p53 status of

the cells. On the other hand, Ia did not increase apoptosis in either cell line (Fig. (5B)). The normal cells also exhibited an increase in apoptosis following exposure to Ic, although to a lesser extent than the lung cancer cells (Fig. (5C)).

#### Alterations in the Expression of Proteins Related to Apoptosis

We next investigated the possible mechanism(s) responsible for the pro-apoptotic activity of Ic by evaluating their effects on the expression level of various proteins involved in apoptosis. Ic activated p53 (in A549 cells), down-regulated MDM2, and increased cleavage of PARP and caspases 8 and 9 (Fig. (6)).

#### DISCUSSION

Marine natural products have attracted the attention of scientists from many disciplines, including organic chemistry, medicinal chemistry, pharmacology, and biology. As a result of improvements in deep-sea sample collection and large-scale drug production through aquaculture, the number of biologically active alkaloids isolated from marine sources has increased significantly [26]. There are approximately a dozen marine alkaloids currently undergoing various phases of human clinical trials for the treatment of multiple cancers [27, 28].

The majority of bioactive marine alkaloids have been isolated from marine sponges [29] and exhibit numerous therapeutic properties including anti-inflammatory, anti-tumor, immunosuppressive, antiviral, anti-malarial and anti-biotic characteristics [8, 11, 16]. Unfortunately, a number of



these alkaloids are only available in minute quantities, thus precluding their use in biological investigations. Despite improvements in aquaculture, laboratory synthesis of these compounds and their analogs is the only practical way to obtain these compounds in larger quantities.

We recently developed novel methods that allowed us to synthesize several analogs of makaluvamines, a subclass of marine alkaloids. In addition to developing the compounds, we also accomplished initial evaluations of their activity against several human cancer cell lines [20, 21]. Further studies with the most active of these compounds are currently underway. However, during our studies, we observed that the pyrroloiminoquinone moiety of makaluvamines is crucial for their activity; therefore, no major changes can be made to the tricyclic ring system. Since further modifications may be needed to enhance the efficacy or "drug-like" properties of these compounds, we thought it necessary to generate compounds that could undergo more extensive modifications. Two areas of the makaluvamine molecule are amenable to modifications: 1) the 6-position of the pyrroloiminoquinone ring, and 2) the pyrrole N atom. Therefore, as a continuing effort to generate new potential therapeutics, we synthesized two classes of compounds by modifying each of these areas. As many pyrroloiminoquinone alkaloids with proven anti-cancer activities have substitutions at the 6-position of the ring, we decided to explore simple substitutions with increased steric bulk, hydrophobicity, and hydrophilicity at this position on the pyrroloiminoquinone ring. To our knowledge, the anti-cancer activities of N-tosylpyrroloiminoquinones have never been explored, thus making this report the first of its kind. Moreover, this study represents the first attempt to evaluate the anti-cancer activities of these synthetic makaluvamine analogs, and the first demonstration that their biological activities are related to specific modifications of the core structure.

The results of our study indicate that the presence of a N atom at the 6-position of pyrroloiminoquinone is crucial for the anti-cancer activity of the makaluvamine analogs. Although the most active analog identified from these studies was a N-tosyl analog (Ic), in most cases, N-tosyl substitution did not significantly alter compound activity. Analogs with OH substituents on the side chain phenyl ring were the least active against lung cancer.

The present investigations demonstrated at least four major points: 1) The anti-cancer activity of the makaluvamine derivatives is dose-, structure-, and cell type-dependent; 2) Induction of apoptosis is the major mechanism by which Ic exerts its cytotoxicity, although cell cycle arrest and effects on proliferation may also be involved; 3) Ic regulates the expression of cell cycle- and apoptosis-related proteins; and 4) The anti-cancer activity of compound Ic appears to occur *via* both p53-dependent and -independent pathways, as the viability of both A549 (p53 wt) and H1299 (p53 null) cells was affected by the compound.

It is generally believed that p53 activation is one of the major mechanisms of action for many DNA damaging agents, including irradiation and numerous chemotherapeutic agents. However, approximately 50% of all human malignancies lack functional p53 [30], rendering them un-

responsive to these DNA-damaging therapies. As all of the human cell lines tested in the present study were responsive to the synthetic marine alkaloids, and were particularly sensitive to Ic, we hypothesize that these compounds may prove to be efficacious against a broad spectrum of human cancers, including those lacking functional p53. Moreover, we observed that there was a decrease in MDM2 in both cell lines. Since decreasing MDM2 has been shown to result in anti-cancer effects (decreased proliferation, increased apoptosis and decreased cell viability) independent of the p53 status of the cells or tumors [31], it is possible that Ic exerts its effects at least partially through MDM2. Additional molecular and pharmacological investigations are warranted to further elucidate the underlying mechanisms of action and to further establish Ic, and potentially other analogs, as novel anti-cancer agents.

## ACKNOWLEDGEMENTS

This work was supported by NIH grants R01 CA112029 and R01 CA121211. E. Rayburn was supported by a DoD grant W81XWH-06-1-0063 and a T32 fellowship from the NIH/UAB Gene Therapy Center (CA075930). S. E. Velu was supported by a UAB Comprehensive Cancer Center, Breast Spore pilot grant and a translational research grant from the UAB Center for Clinical and Translational Science and Council of University-Wide Interdisciplinary Research Centers.

## REFERENCES

- [1] Haefner, B. Drugs from the deep: marine natural products as drug candidates. *Drug Discov. Today*, **2003**, *8*, 536-544.
- [2] Blunt, J. W.; Copp, B. R.; Munro, M. H.; Northcote, P. T.; Prinsep, M. R. Marine natural products. *Nat. Prod. Rep.*, **2006**, *23*, 26-78.
- [3] Marris, E. Marine natural products: drugs from the deep. *Nature*, **2006**, *443*, 904-905.
- [4] Schwartzmann, G.; Brondani da Rocha, A.; Berlinck, R. G.; Jimeno, J. Marine organisms as a source of new anticancer agents. *Lancet Oncol.*, **2001**, *2*, 221-225.
- [5] Williams, D. H.; Stone, M. J.; Hauck, P. R.; Rahman, S. K. Why are secondary metabolites (natural products) biosynthesized? *J. Nat. Prod.*, **1989**, *52*, 1189-1208.
- [6] Antunes, E. M.; Copp, B. R.; Davies-Coleman, M. T.; Samaai, T. Pyrroloiminoquinone and related metabolites from marine sponges. *Nat. Prod. Rep.*, **2005**, *22*, 62-72.
- [7] Faulkner, D. J. Marine natural products. *Nat. Prod. Rep.*, **2002**, *19*, 1-48.
- [8] Carney, J. R.; Scheuer, P. J.; Kelly-Borges, M. A new bastadin from the sponge *Psammaphysilla purpurea*. *Tetrahedron*, **1993**, *49*, 8483-8486.
- [9] Casapullo, A.; Cutignano, A.; Bruno, I.; Bifulco, G.; Debitus, C.; Gomez-Paloma, L.; Riccio, R. Makaluvamine P, a new cytotoxic pyrroloiminoquinone from *Zyzzya cf. fuliginosa*. *J. Nat. Prod.*, **2001**, *64*, 1354-1356.
- [10] Hu, J. F.; Schetz, J. A.; Kelly, M.; Peng, J. N.; Ang, K. K.; Flotow, H.; Leong, C. Y.; Ng, S. B.; Buss, A. D.; Wilkins, S. P.; Hamann, M. T. New anti-infective and human 5-HT<sub>2</sub> receptor binding natural and semisynthetic compounds from the Jamaican sponge *Smenospongia aurea*. *J. Nat. Prod.*, **2002**, *65*, 476-480.
- [11] Radisky, D.C.; Radisky, E.S.; Barrows, L.R.; Copp, B.R.; Kramer, R.A.; Ireland, C.M. Novel cytotoxic topoisomerase II inhibiting pyrroloiminoquinones from Fijian sponges of the genus *Zyzzya*. *J. Am. Chem. Soc.*, **1993**, *115*, 1632-1638.
- [12] Schmidt, E. W.; Harper, M. K.; Faulkner, D. J. Makaluvamines H-M and damirone C from the pohpeian sponge *Zyzzya fuliginosa*. *J. Nat. Prod.*, **1995**, *58*, 1861-1867.
- [13] Venables, D. A.; Concepcion, G. P.; Matsumoto, S. S.; Barrows, L. R.; Ireland, C. M. Makaluvamine N: a new pyrroloiminoquinone from *Zyzzya fuliginosa*. *J. Nat. Prod.*, **1997**, *60*, 408-410.

- [14] Gunasekera, S. P.; Zuleta, I. A.; Longley, R. E.; Wright, A. E.; Pomponi, S. A. Discorhabdins S, T, and U, new cytotoxic pyrroloiminoquinones from a deep-water Caribbean sponge of the genus *Batzella*. *J. Nat. Prod.*, **2003**, *66*, 1615-1617.
- [15] Dijoux, M. G.; Schnabel, P. C.; Hallock, Y. F.; Boswell, J. L.; Johnson, T. R.; Wilson, J. A.; Ireland, C. M.; van Soest, R.; Boyd, M. R.; Barrows, L. R.; Cardellina, J. H.; 2<sup>nd</sup>. Antitumor activity and distribution of pyrroloiminoquinones in the sponge genus *Zyzya*. *Bioorg. Med. Chem.*, **2005**, *13*, 6035-6044.
- [16] Perry, N. B.; Blunt, J. W.; Munro, M. H. G. Cytotoxic pigments from new zealand sponges of the genus *Iatrunculia*: discorhabdins a, b and c. *Tetrahedron*, **1988**, *44*, 1727-1734.
- [17] Ding, Q.; Chichak, K.; Lown, J. W. Pyrroloquinoline and pyridoacridine alkaloids from marine sources. *Curr. Med. Chem.*, **1999**, *6*, 1-27.
- [18] Urban, S.; Hickford, S. J. H.; Blunt, J. W.; Munro, M. H. G. Bioactive marine alkaloids. *Curr. Org. Chem.*, **2000**, *4*, 778-821.
- [19] Harayama, Y.; Kita, Y. Pyrroloiminoquinone alkaloids: discorhabdins and makaluvamines. *Curr. Org. Chem.*, **2005**, *9*, 1567-1588.
- [20] Shinkre, B. A.; Raisch, K. P.; Fan, L.; Velu, S. E. Analogs of the marine alkaloid makaluvamines: synthesis, topoisomerase II inhibition, and anticancer activity. *Bioorg. Med. Chem. Lett.*, **2007**, *17*, 2890-2893.
- [21] Shinkre, B. A.; Raisch, K. P.; Fan, L.; Velu, S. E. Synthesis and antiproliferative activity of benzyl and phenethyl analogs of makaluvamines. *Bioorg. Med. Chem.*, **2008**, *16*, 2541-2549.
- [22] Li, M.; Zhang, Z.; Hill, D.; Chen, X.; Wang, H.; Zhang, R. Genistein, a dietary isoflavone, down-regulates the MDM2 oncogene at both transcriptional and posttranslational levels. *Cancer Res.*, **2005**, *65*, 8200-8208.
- [23] Wang, W.; Rayburn, E.R.; Hao, M.; Zhao, Y.; Hill, D.L.; Zhang, R.; Wang, H. Experimental therapy of prostate cancer with novel natural product anti-cancer ginsenosides. *Prostate*, **2008**, *68*, 809-819.
- [24] Velu, S.E.; Pillai, S. K.; Lakshminantham, M. V.; Billimoria, A. D.; Culpepper, J. S.; Cava, M. P. Efficient syntheses of the marine alkaloids makaluvamine D and discorhabdin C: The 4,6,7-trimethoxyindole approach. *J. Org. Chem.*, **1995**, *60*, 1800-1805.
- [25] Mitsudomi, T.; Steinberg, S.M.; Nau, M.M.; Carbone, D.; D'Amico, D.; Bodner, S.; Oie, H.K.; Linnoila, R.I.; Mulshine, J.L.; Minna, J.D.; et al. p53 gene mutations in non-small-cell lung cancer cell lines and their correlation with the presence of ras mutations and clinical features. *Oncogene*, **1992**, *7*, 171-180.
- [26] Pomponi, S. A. The bioprocess-technological potential of the sea. *J. Biotechnol.*, **1999**, *70*, 5-13.
- [27] Newman, D. J.; Cragg, G. M. Marine natural products and related compounds in clinical and advanced preclinical trials. *J. Nat. Prod.*, **2004**, *67*, 1216-1238.
- [28] Simmons, T. L.; Andrianasolo, E.; McPhail, K.; Flatt, P.; Gerwick, W. H. Marine natural products as anticancer drugs. *Mol. Cancer Ther.*, **2005**, *4*, 333-342.
- [29] Sipkema, D.; Franssen, M. C.; Osinga, R.; Tramper, J.; Wijffels, R. H. Marine sponges as pharmacy. *Mar. Biotechnol. (NY)*, **2005**, *7*, 142-62.
- [30] Lain, S.; Lane, D. Improving cancer therapy by non-genotoxic activation of p53. *Eur. J. Cancer*, **2003**, *39*, 1053-1060.
- [31] Zhang, Z.; Li, M.; Wang, H.; Agrawal, S.; Zhang, R. Antisense therapy targeting MDM2 oncogene in prostate cancer: Effects on proliferation, apoptosis, multiple gene expression, and chemotherapy. *Proc. Natl. Acad. Sci. USA*, **2003**, *100*, 11636-11641.