

Synthesis and dynamic behaviour of 1-(dipropylboryl)cyclohepta-2,4-diene

Ilya D. Gridnev, Mikhail E. Gurskii, Alexey V. Geiderikh and Yurii N. Bubnov*

N.D. Zelinsky Institute of Organic Chemistry, Russian Academy of Sciences, 117913 Moscow, Russian Federation.
Fax: +7 095 135 5328

Two sequential [1,3] boron shifts in 1-(dipropylboryl)cyclohepta-2,4-diene **1** result in a degenerate rearrangement ($E_A = 98.9 \pm 3.3$ kJ mol⁻¹, $\ln A = 39.3 \pm 1.4$, $\Delta G^\ddagger_{298} = 118.3 \pm 0.3$ kJ mol⁻¹); reaction of **1** with acetone affords non-conjugated alcohol **2**.

Reactions of [1,3] sigmatropic migration are prohibited by orbital symmetry and are usually characterised by high activation barriers.¹ Thus, the activation energy of [1,3] sigmatropic shifts of silicon and germanium is over 150 kJ mol⁻¹ for the corresponding allylic compounds.² For the allylic derivatives of zinc and cadmium,^{3,4} the second order of [1,3] migration was experimentally confirmed, while [1,3] sigmatropic shifts of tin and mercury are observed only in the presence of Lewis acids,^{5,6} which indicates the intermolecular character of these processes. In a series of this type of reactions, one can clearly distinguish the intramolecular [1,3] boron shift in allylic boron derivatives, the so-called ‘permanent allylic rearrangement’, which is characterised by relatively low activation barriers (the activation energy is between 20 and 100 kJ mol⁻¹).⁷

Continuing our investigations into the influence of structural factors on the regularities of [1,3] boron migration in allylic type triorganoboranes,⁷ we synthesized a new cyclic compound of this type, 1-(dipropylboryl)cyclohepta-2,4-diene **1**, and using NMR spectroscopy investigated its dynamic properties and reaction with acetone.


Compound **1** was prepared in 40% yield by the reaction of lithium cycloheptadienide⁸ with chloro(dipropyl)borane (Scheme 1).[†]

Dienic borane **1** is a liquid which spontaneously ignites in the air. Its chemical shift ($\delta_{\text{H}}^{\text{11B}} = 83.5$) is characteristic of triorganoboranes. Similarly to other allylboranes,⁹ compound **1** reacts with acetone with the allylic rearrangement to give a non-conjugated dienic alcohol **2** (Scheme 2).


It should be pointed out that this reaction has a remarkable feature, *i.e.*, the selective formation of 1,4-dienic compound **2** involves the destruction of the system of conjugated double bonds in the starting borane **1**. A similar phenomenon was observed in the reactions of linear penta-2,4-dienylboranes.¹⁰

[†] Cycloheptadienyl(dipropyl)borane **1** was obtained in 40% yield by borylation of lithium cycloheptadienide⁸ at -30 to -20 °C. bp 58–59 °C (1 mmHg), $n_D^{16} = 1.4895$.

2-(Cyclohepta-1,4-dien-3-yl)propan-2-ol **2** was prepared by the reaction of **1** with acetone followed by oxidation with H₂O₂ and column chromatography on SiO₂. Yield 48%. ¹H NMR (400 MHz, CDCl₃, 298 K): 1.24 (s, 6H, 2CH₃); 1.7–1.8 (br.s, 1H, OH); 2.05–2.45 (m, 4H, H-4); 3.25 (m, 1H, H-1); 5.68 (dd, 2H, ³J = 11.5, ³J = 4.2, H-2); 5.80 (dm, 2H, ³J = 11.5, H-3). ¹³C NMR (400 MHz, CDCl₃, 298 K): 26.24 (C-4); 27.12 (CH₃); 49.84 (C-1); 73.71 (C-OH); 130.04, 131.05 (C-2, C-3).



Scheme 1



Scheme 2 Reagents: i, Me_2CO ; ii, H_2O_2 , OH^- .

The assignment of signals in the NMR spectra of compound **1** was performed using two-dimensional correlation NMR spectroscopy (see Table 1). On increasing the temperature to 100°C , ^1H and ^{13}C NMR spectra exhibit a reversible broadening of the signals, which indicates the dynamic behaviour of borane **1**. Figure 1 shows the ^{13}C 2D EXSY spectrum of compound **1**. This spectrum contains three cross-peaks corresponding to the exchange between C^1 and C^5 , C^2 and C^4 , and between C^6 and C^7 . The presence of these three cross-peaks attests to the degenerate rearrangement in **1** occurring as a migration of the boryl group from position 1 to position 5.

One can propose two mechanisms for the migration of the dipropylboryl group in **1**, and both are in agreement with all of the obtained experimental data, *viz.*, [1,5] sigmatropic boron shift (see Scheme 3) and two sequential [1,3] boron shifts involving the formation of a thermodynamically unstable isomer **1a** as an intermediate. We believe the

Table 1 Parameters for the ^1H and ^{13}C NMR spectra of **1**.

Atom	δ_{H}	J/Hz	δ_{C}
1	2.55	^a	44.11
2	5.91	5.9 (H^1, H^2), 11.8 (H^2, H^3)	136.45
3	5.82	^a	123.67
4	5.82	^a	125.63
5	5.82	^a	132.71
6	1.77, 1.98	13.4 (2J), 9.0 ($\text{H}^{6\text{a}}, \text{H}^{7\text{a}}$), 4.0 ($\text{H}^{6\text{a}}, \text{H}^{7\text{b}}$), 3.0 ($\text{H}^1, \text{H}^{6\text{a}}$), 7.5 ($\text{H}^1, \text{H}^{6\text{b}}$), 7.5 ($\text{H}^{6\text{b}}, \text{H}^{7\text{b}}$), 3.1 ($\text{H}^{6\text{b}}, \text{H}^{7\text{a}}$)	27.38
7	2.17, 2.28	17.4 (2J)	32.24
Pr	0.9, 1.2, 1.4	—	17.51, 17.71, 29.63

^aUnresolved multiplet.






Figure 1 ^{13}C EXSY spectrum of **1** (100 MHz, 298 K, CDCl_3), matrix size 1024×1024 , mixing time τ_m 1 s.



Scheme 3

mechanism of two sequential [1,3] boron migrations to be more valid, because we have previously shown that it is the [1,3] boron shift that is the most rapid process in linear penta-2,4-dienyl(dipropyl)borane **3**^{7b} (Scheme 4).



Scheme 4 $\Delta G^\ddagger(3\mathbf{a} \rightarrow 3\mathbf{a}) = 91.5 \text{ kJ mol}^{-1}$; $\Delta G^\ddagger(3\mathbf{a} \rightarrow 3\mathbf{b}) = 85.5 \text{ kJ mol}^{-1}$; $\Delta G^\ddagger(3\mathbf{b} \rightarrow 3\mathbf{b}) = 82.3 \text{ kJ mol}^{-1}$ (values at 366 K).

Kinetic analysis of the degenerate rearrangement in **1** (Figure 2) was carried out using a series of ^1H and ^{13}C EXSY spectra recorded at different temperatures. The rate constants were calculated by equation (1)¹¹ using the intensities of diagonal and cross-peaks corresponding to the exchange between H^1 and H^5 , H^6 and H^7 , C^6 and C^7 . The analysis gave the following values of activation parameters:




Figure 2 Arrhenius plot for the degenerate rearrangement in **1**. Squares correspond to rate constants obtained from ^{13}C EXSY spectra (exchange between C^6 and C^7), circles and triangles to the rate constants obtained from ^1H EXSY spectra (exchange between H^1 and H^5 and averaged data for the exchanges in two pairs of H^6 and H^7). Goodness of fit is 0.997.

$$E_A = 98.9 \pm 3.3 \text{ kJ mol}^{-1}, \quad \ln A = 39.3 \pm 1.4, \quad \Delta G_{298}^{\neq} = 118.3 \pm 0.3 \text{ kJ mol}^{-1}$$

$$k = \frac{1}{t_m} \ln \frac{r+1}{r-1}, \quad r = \frac{I_{AA} + I_{BB}}{I_{AB} + I_{BA}} \quad (1)$$

where I_{AA} , I_{BB} are the intensities of the diagonal peaks; I_{AB} , I_{BA} are intensities of the cross-peaks.

The activation barrier of the [1,3] boron shift obtained for compound **1** ($\Delta G_{298}^{\neq} = 118.3 \text{ kJ mol}^{-1}$) is the highest of those measured for a series of allylic type organoboranes.⁷ The most close values were obtained for *cis,cis*-penta-2,4-dienyl(dipropyl)borane **3a** (Scheme 4),^{7b} which indicates that the system of conjugated double bonds in the molecule of allylic type triorganoborane retards the [1,3] boron shift.

This work was partly supported by the Russian Foundation for Basic Research (grant no. 94-03-08857).

References

- (a) R. B. Woodward and R. Hoffmann, *Angew. Chem., Int. Ed. Engl.*, 1969, **8**, 781; (b) C. W. Spangler, *Chem. Rev.*, 1976, **76**, 187.
- (a) M. Kobayashi and N. Kobayashi, *Chem. Lett.*, 1986, 385; (b) H. Kwart and T. Slutsky, *J. Am. Chem. Soc.*, 1972, **94**, 2515.
- K. H. Thiele and J. Kohler, *J. Organomet. Chem.*, 1967, **7**, 365.
- E. G. Hoffman, H. Nehl, H. Lemmkuhl, K. Seevogel and W. Stempfle, *Chem. Ber.*, 1984, **117**, 1364.
- (a) Yo. Naruta, Yu. Nishigaiishi and K. Maruyama, *Tetrahedron*, 1985, **45**, 1067; (b) M. J. Hails, B. E. Mann and C. M. Spencer, *J. Chem. Soc., Dalton Trans.*, 1983, 729.
- P. D. Sleezer, S. Winstein and W. G. Young, *J. Am. Chem. Soc.*, 1963, **85**, 1890.
- (a) Yu. N. Bubnov, M. E. Gurskii, I. D. Gridnev, A. V. Ignatenko, Yu. A. Ustynyuk and V. I. Mstislavsky, *J. Organomet. Chem.*, 1992, **424**, 127; (b) M. E. Gurskii, I. D. Gridnev, A. V. Geiderich, A. V. Ignatenko, V. I. Mstislavsky and Yu. A. Ustynyuk, *Organometallics*, 1992, **11**, 4056; (c) I. D. Gridnev, M. E. Gurskii and Yu. N. Bubnov, in *Current Topics in the Chemistry of Boron*, Royal Society of Chemistry, Cambridge, 1994, pp. 56–59.
- R. B. Bates, W. H. Deines, D. A. McCombs and D. E. Potter, *J. Am. Chem. Soc.*, 1969, **91**, 4608.
- B. M. Mikhailov and Yu. N. Bubnov, *Organoboron Compounds in Organic Synthesis*, Harwood, Chur, London, New York, 1984.
- (a) M. G. Hutchings, W. E. Paget and K. Smith, *J. Chem. Res. (M)*, 1983, 342; (b) M. Schlosser, *Abstracts of the 7th IUPAC Conference on Organic Synthesis*, Nancy, France, 1988, p. 9; (c) Yu. N. Bubnov, A. V. Geiderikh, S. B. Golovin, M. E. Gurskii, M. Yu. Etinger, L. I. Lavrinovich, P. Yu. Meleshkin, S. V. Ponomarev and V. I. Zheludeva, *Mendeleev Commun.*, 1994, 55; (d) Yu. N. Bubnov, A. V. Geiderikh, A. V. Ignatenko, P. Yu. Meleshkin and M. E. Gurskii, *Mendeleev Commun.*, 1994, 214.
- C. L. Perrin and T. J. Dwyer, *Chem. Rev.*, 1990, **90**, 935.

*Received: Moscow, 22nd February 1996
Cambridge, 20th May 1996; Com. 6/0138C*